

Original Research Article

Exploring the Relationship Between Endophytic Bacteria and Biochemical Compounds in Pearl Millet

ABSTRACT

Pearl millet (*Pennisetum glaucum*) is an important cereal crop in Asia and Africa, being one of the oldest and most important cereal crops. This study focuses on quantifying endophytic bacteria in several parts of the pearl millet plant—roots, stems, and leaves. Total phenolic content, orthodihydroxy phenols, total soluble sugars, protein levels, and enzymatic activity (peroxidase, polyphenol oxidase, and catalase) were all measured and linked with the amount of endophytic bacteria. The leaves have the highest total phenolic content, followed by the stem. Orthodihydroxy phenols were also significantly greater in leaves (1.82 mg catechol $\mu\text{equi./g}$) than in stems and roots. Total soluble sugars showed a similar trend, with higher levels in stems (9.21 mg glucose $\mu\text{equi./g}$) and leaves compared to roots. Protein content exhibited significant variation among the plant parts, with leaves (10.63%) containing the highest protein content followed by stems and roots. Enzymatic activities, crucial for plant defence and growth, varied across the plant sections. Peroxidase activity was highest in roots (359.98 OD/g), followed by stems and leaves. Polyphenol oxidase activity was highest in the roots (1.63 OD/g) and lowest in the leaves, while catalase activity followed a similar pattern, with the highest activity in the roots. The number of Pearl millet endophytic bacteria in the roots is 16, whereas the amount in the stem is 12. This in-depth analysis sheds light on future research targeted at better understanding plant-microbe interactions and developing ways to improve plant health and productivity.

Keywords: Orthodihydroxy phenols, Total soluble sugars, Peroxidase, Polyphenol oxidase, and Catalase.

INTRODUCTION

Plants can be thought of complex ecosystems in which a wide variety of microorganisms inhabit various niches. Such niches are made up of internal tissues where endophytic bacteria can thrive without harming the host (Kumar et al., 2013). Plant-associated bacteria serve a crucial role in nutrient cycling and preventing phytopathogens from causing harm. Plant microbe interactions occur mostly in the rhizosphere and within plant tissues. Microbial endophytes are microorganisms that complete their one phase of life cycle within plant tissues without generating visible symptoms, according to Rangjaroen et al., (2014). Endophytes can be defined, in a generalist manner, as a group of microorganisms that infect internal tissues of plant without causing any immediate symptom of infection and/ or visible manifestation of

Comment [A1]: 1. Journal format not followed
2. Check every level of headings
3. Need latest references (within 5 years)
4. Materials and methods need well presentation
5. Result section need improvement for explaining findings
6. Discussion section not well presented, lack of justification of your findings
7. Add scientific name of Pearl Millet in title
8. 'Isolation' word should be included in title

Comment [A2]: This sentence should be in Introduction section with reference. Never give any statement in abstract.

Comment [A3]: Check carefully to write unit,

Formatted: Highlight

Comment [A4]: Follow previous comment

Formatted: Not Highlight

Comment [A5]: Mention numeric result also

Formatted: Not Highlight

Formatted: Not Highlight

Comment [A6]: Same here like previous comment

Formatted: Not Highlight

Formatted: Not Highlight

Comment [A7]: Follow previous comment

Formatted: Not Highlight

Formatted: Not Highlight

Comment [A8]: Improve this line

Comment [A9]: 1. Check level of headings
2. Please replace old citations with recent publications
3. Please justify problem statement in the introduction and some hypothesis

Formatted: Not Highlight

Comment [A10]: Write the citation correct way

disease and live in mutualistic association with plants for at least a part of their life cycle (Bacon and White 2000). Pearl millet was developed as a fodder and cereal crop in Africa around 3000 years ago and subsequently introduced into India. Appa and De Wet (1999) identified India as the world's second largest hub of pearl millet diversity. Pearl millet has a better level of heat tolerance and is more adept at using soil moisture than maize and sorghum. It is commonly used as a pasture crop in the United States, Australia, Southern Africa, and South America. It is mostly grown in India during the Kharif season, with modest production in Andhra Pradesh, Karnataka, and Tamil Nadu during the Rabi season. Only

about 8% of India's pearl millet farming land is irrigated (Chandrashekar *et al.*, 2007).

Endophytic bacteria are microorganism that live within plant tissues without inflicting noticeable harm. They've been identified as important contributors to plant health and growth. This study investigates the complex link between endophytic bacteria and biochemical substances found in plant parts such as roots, stems, and leaves. Endophytic

bacteria specifically alter the quantities and activities of phenols, flavanols, total soluble sugars, proteins, peroxidase enzyme, orthodihydroxy phenols, and catalase enzyme. These chemicals, which include phenols and flavanols, have been linked to a variety of plant physiological activities, including growth, development, and defence against biotic and abiotic challenges (Carvalho *et al.*, 2014). Understanding these interactions can provide valuable insights into the potential applications of endophytic bacteria in agriculture, biocontrol, and sustainable plant production systems. One study by Hardoim *et al.*, (2008)

investigated endophytic bacterial communities in different plant compartments and found that the roots and stems of plants tend to host a greater diversity and abundance of endophytic bacteria compared to leaves and flowers. Another study by Bulgarelli *et al.*, (2013) focused on the *Arabidopsis thaliana* plant and found that the roots harboured a higher abundance of endophytic bacteria compared to the aerial parts of the plant. The researchers used culture-independent techniques to analyze the bacterial communities and identified a diverse range of endophytic bacteria in the roots.

Endophytic bacteria are important for the synthesis and accumulation of phenolic chemicals in plant tissues. Endophytic bacteria can create particular enzymes that increase phenolic compound synthesis in plants (Li *et al.*, 2014). Sugar availability in plants can also influence endophytic bacteria colonisation and activity. Endophytic bacteria use sugars as a carbon source from plants, which can alter their growth, metabolism, and interactions with the plant host (Daubechet *et al.*, 2021). Peroxidases are a class of enzymes involved in various physiological processes in plants, including the metabolism of reactive oxygen species

Formatted: Not Highlight

Comment [A11]: Too old reference! Include latest one

Formatted: Not Highlight

Comment [A12]: This sentence should be in the last line of introduction

Formatted: Not Highlight

Formatted: Not Highlight

Comment [A13]: Unnecessary words like one study, another study should avoid

Comment [A14]: Italic

Formatted: Not Highlight

(ROS), lignin synthesis, and defense against pathogens. The peroxidase enzyme activity can vary among different plant parts depending on various factors such as the plant species, developmental stage, and environmental conditions. Rizwan *et al.*, 2012 investigated the peroxidase activity in different plant parts of *Brassica juncea* (Indian mustard) under heavy metal stress. They found that the roots exhibited significantly higher peroxidase activity compared to stems and leaves, suggesting a greater role of peroxidases in root tissues under stress conditions. Keeping in view, the importance of endophytic bacteria, the present study was undertaken entitled “Exploring the Relationship Between Endophytic Bacteria and Biochemical Compounds in Pearl Millet”.

Comment [A15]: Revise sentence

Formatted: Not Highlight

MATERIALS AND METHODS

Comment [A16]: 1. Materials and Methods have subsections and numbers accordingly.
2. Contact any expert to know how to write Methodology sections.

Comment [A17]: Need GPS

Formatted: Not Highlight

The present investigation was carried out in the laboratory and field area of Department of Plant Pathology, CCS Haryana Agricultural University, Hisar. Plant samples were collected from pearl millet grown at Plant Pathology experimental area of CCS Haryana Agricultural University, Hisar of the *kharif* 2021. The Root, stem and leaves of pearl millet grown in Plant Pathology experimental area of CCS Haryana Agricultural University Hisar were collected with latitude 29°14'N and longitude of 75°70'E, having sandy loam soil texture. Roots, leaves and stems were washed with running tap water and then surface sterilized sequentially in 75 % (v/v) ethanol for 2 min, 2.6 % (w/v) sodium hypochlorite solution for 5 min and 75 % (v/v) ethanol for 1 min. For isolation of bacterial endophytes one g of plant tissue was crushed in pestle and mortar with 10 ml sterile distilled water to get homogenous paste and allowed to settle down for 20 min. The supernatant was diluted and approximately 10 µl was placed on NA plates and incubated at 28±2°C for 3 days (Janse, 2005). Bacterial colonies appearing on the plates were considered to be endophytes. Colonies were characterized according to different visual observations and finally purified using streak plate technique. The bacterial colonies were maintained at 4 ±1°C for further studies.

Comment [A18]: How do you know its sandy loam soil texture? mention reference?

Comment [A19]: ml or mL??

Comment [A20]: Same as previous comment

Comment [A21]: Spell out

Formatted: Not Highlight

Comment [A22]: There must be a section (eg. Isolation Procedure of Endophytic Bacteria)

Comment [A23]: 1. Check level of heading
2. Subsections have number or not? Follow format

Prevalence of endophytic bacteria in relation to biochemical compounds of pearl millet

Endophytic bacteria were isolated from different plant parts and their colony number were counted. One gram sample of roots, leaves and stems were washed with running tap water and then surface sterilized sequentially in 75 % (v/v) ethanol for 2 min, 2.6 % (w/v) sodium hypochlorite solution for 5 min and 75 % (v/v) ethanol for 1 min. Finally, the plant parts were thoroughly washed six times with sterile distilled water. For isolation of bacterial endophytes one g of plant tissue was crushed in pestle and mortar with 10 ml sterile distilled water to get homogenous paste and allowed to settle down for 20 min. The supernatant was diluted and approximately 10 µl was placed on NA plates and incubated at 28±2°C for 3 days.

Comment [A24]: Spell full

Fifteen petri plates from 5 different samples of each roots, leaves and stem respectively were taken. Bacterial colonies appearing on the plates were considered to be endophytes. Similarly, biochemical compounds were recorded from different plant parts of pearl millet and their correlation were measured with the number of endophytic bacteria.

Extraction and estimation of total phenols

Extraction: Total phenols and different phenols were extracted from leaves using 80% hot alcohol. Dry powdered sample (0.1 g) of susceptible and resistant genotype was homogenized with 80% ethanol (total 10 ml) and centrifuged at 5000 rpm for 15 minutes, supernatant was taken and final volume was made upto 10 ml with 80% ethanol.

Estimation of phenols: Total phenolic content was estimated by the method of Swain and Hillis (1959) using Folin- Ciocalteu reagent.

Reagents

- i. Folin- Ciocalteu reagent: Folin- Ciocalteu reagent was diluted 1:1(v/v) before use
- ii. Saturated sodium carbonate solution: Dissolved anhydrous sodium carbonate (35 g) in 500 ml of distilled water by heating on a water bath at 70-80°C and then cooled the contents overnight and used the supernatant.

Method :The alcohol extract (0.5 ml) was taken in test tube and diluted with 3.5 ml of distilled water. To this added 0.5 ml Folin- Ciocalteu reagent and 2 ml saturated sodium carbonate solution and then test tubes were heated in boiling water for 2-3 minutes. Cooled the test tubes and read the absorbance at 650 nm using spectrophotometer. The blank was prepared by taking ethanol in place of extract and rest of procedure was same.

Calculations: The amount of phenol in the sample was determined from the standard curve prepared simultaneously by taking Catechol as the standard phenol and the data was expressed as mg/g dry weight.

Estimation of Orthodihydroxy phenols

The Orthodihydroxy phenols were determined by the method of Johnson and Schaal (1952).

Reagent

- i. Arrow's reagent was prepared by dissolving sodium nitrite 10 g and sodium molybdate 10 g in distilled water and made volume to 100 ml with distilled water
- ii. 0.5 N HCl
- iii. 0.5 N NaOH

Method :Two ml of extract was taken in test tube and added 2 ml of 0.5 N HCl, 1 ml of Arrow's reagent and 4 ml of distilled water in succession. After this 2 ml of 0.5 N NaOH was

added and the solution was shaken till, pink colour appeared. Blank was prepared without the addition of Arrow's reagent

Calculation: The amount of Orthodihydroxy phenol in the sample was determined from the standard curve prepared simultaneously by taking Catechol as the standard phenol and the data was expressed as mg/g dry weight.

Formatted: Not Highlight

Estimation of Flavanol

The Flavanols were determined by the method of Balbaaet *al.*, (1974) with slight modification.

Reagent: Aluminium chloride (0.1M): 24.143 g of aluminium chloride was dissolved in one litre of distilled water.

Procedure: One ml of the extract was taken in a test tube and then 5ml of 0.1 M Aluminium chloride solution was added to it and thereafter, the absorbance of the solution was read at 420 nm using spectrophotometer. A blank was prepared simultaneously and calculate the amount of flavanols by taking Catechol as standard and the data was expressed as mg/g dry weight.

Estimation of total soluble sugars

Reagent

1. Concentrated H₂SO₄.
2. Anthrone reagent: 0.2 g in 100 ml conc. H₂SO₄.

Procedure: 0.5 ml of the extract and 1.5 ml of distilled water was taken in a test tube. Then 4 ml of anthrone reagent was added. The tubes were shaken and allowed to cool for 30 minutes and read the absorbance at 625 nm on spectrophotometer. The concentration of total sugars was calculated from the standard curve of glucose prepared simultaneously and the data was expressed as mg glucose equi.g .

Estimation of protein content

Total nitrogen in the samples was estimated by conventional Micro-Kjeldahl's method. Multiplied nitrogen content with suitable factor (3.25) to calculate crude protein content.

Comment [A25]: Reference???

Comment [A26]: Not clear

Reagent

- i. Concentrated sulphuric acid
- ii. Mixed indicator :-Bromocresol green (0.1% in 95% alcohol) and Methyl red (0.1% in 95% alcohol)
- iii. Boric acid solution (2%)
- iv. Sodium hydroxide solution(40%)

- v. Digestion mixture: Mix CuSO_4 and K_2SO_4 in 1:10 ratio
- vi. Standard hydrochloric acid solution (0.01 N)

Procedure

Digestion: 250 mg dried leaves sample was weighed and taken into clean and dry digestion flask. In this 0.5g of digestion mixture and 10.0 ml of concentrated sulphuric acid was added. The flask was heated on a hot plate until the solution became colorless/very light green. Cooled it and after cooling made the final volume up to 100 ml with distilled water.

Comment [A27]: Spell out

Distillation and titration

1. Ten milliliter of 4% boric acid was taken in a 100 ml conical flask. Added 2-3 drops of mixed indicator and then the flask was placed under the condenser and end of the condenser was dipped in boric acid solution.
2. Pipetted out 10 ml of digested material into the distilling portion (vacuum jacket) of the set followed by 10 ml 40% NaOH.
3. Steam distilled the mixture by opening the steam inlet and keeping the other outlets closed except the receiving end of the condensor. The liberated ammonia was collected in the boric acid which resulted in the colour change of the solution from pink to blue. Distilled for 5-6 min till the volume in the conical flask increased by 2 to 2.5 folds (30-40 ml).
4. Closed the steam inlet, washed down the tip with a few ml of distilled water, removed the conical flask containing the distillate and titrated against standard 0.01N HCl until the colour changed to light pink. Noted the titre value.

Run a blank preparation which was identically prepared except that it does not contain the sample.

Calculation: Crude protein content was calculated by multiplying the nitrogen content with a suitable factor (3.25).

Formatted: Not Highlight

Enzymatic study

Preparation of extract

Plant material (roots and leaves separately) were taken and washed with cold distilled water and dabbed dry with several folds of filter paper. Extraction conditions were standardized with respect to molarity and pH of buffer to achieve maximum extraction of enzymes in leaves. All the steps of extraction were carried out at 0-4°C. 200 mg tissue was macerated in chilled pestle and mortar in the presence of 10 ml 0.1 M phosphate buffer (pH 7.5). The homogenate was centrifuged at 10,000 g for 20 minutes at 4°C. The supernatant was carefully decanted and used as the crude enzyme preparation.

Peroxidase

The enzyme activity was estimated by the method of Shannon *et al.*, (1966). The reaction mixture contained 2.7 ml of 50 mM phosphate buffer (pH 6.5), 0.1 ml of 0.5% hydrogen peroxide, and 0.1 ml of 0.2% O- dianisidine and 0.01 ml of enzyme extract. The reaction was initiated by the addition of 0.1 ml of H₂O₂. The assay mixture without H₂O₂ served as blank. Change in absorbance was followed at 430 nm for 3 min. One unit of peroxidase was defined as amount of enzyme required to cause change in O.D. of 1 unit per minute.

Catalase

Catalase activity was determined by the procedure of Sinha (1972). The reaction mixture (1 ml) consisted of 0.5 ml of 0.2 M phosphate buffer (pH 7.0), 0.4 ml of 0.2 M hydrogen peroxide and 0.1 ml of properly diluted enzyme extract. After incubating at 37°C for 2 min, the reaction was terminated by adding 3 ml mixture of 5% (w/v) potassium dichromate and glacial acetic acid (1:3 v/v) to the reaction mixture. The tubes were heated in boiling water bath for 10 min. A control was run under similar conditions where enzyme extract was added after stopping the reaction. After cooling the tubes, absorbance of test and control was measured at 570 nm. The amount of residual H₂O₂ in the reaction mixture was determined by subtracting the absorbance of test samples from that of control. One unit of enzyme activity is defined as the amount of enzyme which catalyzed the oxidation of 100 µmole H₂O₂ per minute under assay conditions.

Polyphenol oxidase

The enzyme activity was estimated by the method of Taneja and Sachar (1974).

Reagents:

1. Assay buffer : 0.1 M phosphate buffer (pH 6.2)
2. Catechol solution : 0.11g Catechol dissolved in 100 ml phosphate buffer.

Procedure: The reaction mixture consisted of 0.1 ml enzyme extract and 2.9 ml catechol solution. The reaction mixture was incubated at 37°C for 1 h. Run a blank. Simultaneously consisting of catechol solution. The absorbance was measured at 410 nm on a spectrophotometer. The enzyme activity was expressed as change in 0.01 absorbance min g protein.

RESULTS

Bacterial endophytes provide various benefits to the host plant, including growth stimulation, protection from pathogens, and the ability to live in a variety of environmental conditions. Bacterial endophytes are better capable of communicating and interacting with the plant than rhizospheric bacteria. Samples from root, stem and leaves of pearl millet were collected from

Comment [A28]: Results have subsections and numbers accordingly

Plant Pathology experimental area of CCS Haryana Agricultural University Hisar with latitude 29°14'N and longitude of 75°70'E, having sandy loam soil texture of the *kharif* 2021. A total of 38 pearl millet bacterial endophytes (PMEB) were obtained. Out of which, 16 bacterial endophytes were obtained from Root (PMREB 1, PMREB 2, PMREB 3, PMREB 4, PMREB 5, PMREB 6, PMREB 7, PMREB 8, PMREB 9, PMREB 10, PMREB 11, PMREB 12, PMREB 13, PMREB 14, PMREB 15 and PMREB 16), 12 obtained from stem (PMSEB 1, PMSEB 2, PMSEB 3, PMSEB 4, PMSEB 5, PMSEB 6, PMSEB 7, PMSEB 8, PMSEB 9, PMSEB 10, PMSEB 11 and PMSEB 12) and 10 from the leaves (PMLEB 1, PMLEB 2, PMLEB 3, PMLEB 4, PMLEB 5, PMLEB 6, PMLEB 7, PMLEB 8, PMLEB 9 and PMLEB 10).

Table 1:List of endophytic bacterial isolates retrieved from root, stem and leaf of pearl millet

Sr.No.	Source	Endophytic bacteria
1	Pearl millet root endophytic bacteria	PMREB 1, PMREB 2, PMREB 3, PMREB 4, PMREB 5, PMREB 6, PMREB 7, PMREB 8, PMREB 9, PMREB 10, PMREB 11, PMREB 12, PMREB 13, PMREB 14, PMREB 15 and PMREB 16.
2	Pearl millet stem endophytic bacteria	PMSEB 1, PMSEB 2, PMSEB 3, PMSEB 4, PMSEB 5, PMSEB 6, PMSEB 7, PMSEB 8, PMSEB 9, PMSEB 10, PMSEB 11 and PMSEB 12.
3.	Pearl millet leaves endophytic bacteria	PMLEB 1, PMLEB 2, PMLEB 3, PMLEB 4, PMLEB 5, PMLEB 6, PMLEB 7, PMLEB 8, PMLEB 9 and PMLEB 10.

These endophytic bacteria of pearl millet were streaked on nutrient agar plates retrieved during *kharif* crop 2021. The bacterial colonies were maintained at 4 ±1°C for further studies.

Formatted: Not Highlight




Plate1: Pearl millet bacterial endophytes (*kharif 2021*)

Comment [A29]: Plate 1 or Figure 1??

Prevalence of endophytic bacteria in relation to biochemical compounds of pearl millet

Endophytic bacteria were isolated and enumerated from different plant sections. Samples of roots, stems, and leaves were collected from pearl millet. The plant material was processed and analysed for total phenolic content, Orthodihydroxy phenols, flavanols, total soluble sugars, protein levels Peroxidase, Polyphenol oxidase and Catalase. The analyses were performed using established methods, including spectrophotometry and biochemical assays. Similarly, biochemical substances were extracted from several plant portions of pearl millet and their correlation with the quantity of endophytic bacteria was determined.

Statistical analysis was conducted to determine significant differences among the plant parts, using CD (critical difference) values at a significance level of 0.05.

Formatted: Not Highlight

Comment [A30]: This should be in Materials and Methods section

Total Phenols

Comment [A31]: Table?

The leaves exhibited the highest total phenolic content (8.3 mg catechol **equi./g**), followed by the stem (6.2 mg catechol **equi./g**) and the root (3.1 mg catechol **equi./g**). Minimum number of Pearl millet endophytic bacteria in the leaves is 10 followed by stem and maximum found in roots.

Ortho-dihydroxy Phenol and Flavanol

Orthodihydroxy phenols were also found in varying amounts, with the leaves (1.82 mg catechol **equi./g**) containing higher levels compared to the stem (1.59 mg catechol **equi./g**) and root (1.02 mg catechol **equi./g**) but minimum number of Pearl millet endophytic bacteria in the leaves is 10 followed by stem and maximum found in roots. Regarding flavanols, the stem had the highest concentration (11.9 mg catechol **equi./g**), followed by the leaves (10.10 mg catechol **equi./g**) and the root (8.2 mg catechol **equi./g**) (Table 2).

4.2.3 Total soluble sugars

The analysis of total soluble sugars indicated a similar trend, with the stem (9.21 mg glucose **equi./g**) and leaves (9.97 mg glucose **equi./g**) having higher levels compared to the root (6.83 mg glucose **equi./g**). Maximum number of Pearl millet endophytic bacteria in the roots is 16 followed by stem that is 12.

Table 2: Estimation of Total Phenol, Orthodihydroxy Phenol, Flavanol, Total Soluble Sugar and protein (%) from different parts in Pearl millet

Sr. No.	Source	No. of Pearl millet endophytic bacteria	Total phenolic content (mg catechol equi./g)	Orthodihydroxy phenol (mg catechol equi./g)	Flavanol (mg catechol equi./g)	Total soluble sugars (mg glucose equi./g)	Protein (%)
1.	Root	16	3.1	1.02	8.2	6.83	6.65
2.	Stem	12	6.2	1.59	11.9	9.21	8.47
3.	Leaves	10	8.3	1.82	10.10	9.97	10.63
CD (P=0.05)			0.578	0.139	0.955	0.819	1.521
SE(m)			0.164	0.039	0.271	0.232	0.431

Protein content

Protein levels were also found to vary significantly among the plant parts. The leaves exhibited the highest protein content (10.63%), followed by the stem (8.47%) and the root (6.65%). but minimum number of Pearl millet endophytic bacteria in the leaves is 10 followed by stem and maximum found in roots.

Enzymatic activities

Enzymes play a crucial role in various physiological processes in plants, including **defense** against pathogens, oxidative stress management, and regulation of plant growth and

Comment [A32]: Reference?? Which table?

Comment [A33]: Reference?

Formatted: Not Highlight

Formatted: Not Highlight

development. In this study, we investigated the activities of three important enzymes, peroxidase, polyphenol oxidase, and catalase, in different plant parts. The root, stem, and leaves of a specific plant species were analyzed, and the changes in enzyme activity were measured. The results provide valuable insights into the distribution and potential functions of these enzymes in different plant organs. The results of the enzyme activity assays are presented in Table 3. Maximum number of Pearl millet endophytic bacteria in the roots is 16 followed by stem that is 12. Whereas, peroxidase activity was highest in the root (359.98 OD/g), followed by the stem (349.22 OD/g) and leaves (327.05 OD/g). Polyphenol oxidase activity showed highest activity in the leaves (2.19 OD/g) and the lowest in the roots (1.63 OD/g). Catalase activity was highest in the root (17.79 OD/g), followed by the stem (14.63 OD/g) and leaves (14.33 OD/g). The CD values indicate significant differences among the plant parts for all three enzymes (Table 3).

Table 3: Estimation of Peroxidase, Polyphenol oxidase and Catalase from different parts in Pearl millet

Sr. No.	Source	No. of Pearl millet endophytic bacteria	Peroxidase (Change in OD/g)	Polyphenol oxidase (Change in OD/g)	Catalase (Change in OD/g)
1.	Root	16	359.98	1.63	17.79
2.	Stem	12	349.22	1.87	14.63
3.	Leaves	10	327.05	2.19	14.33
CD (P=0.05)			9.077	0.178	1.463
SE(m)			8.161	0.05	0.415

DISCUSSION

Endophytic bacteria were isolated from different plant tissues *viz.* roots, stems and leaves. The plant material was processed and analysed for total phenolic content, Orthodihydroxy phenols, flavanols, total soluble sugars, protein levels, Peroxidase, Polyphenol oxidase and Catalase. Phenolic compounds play various roles in plants, including defense against pathogens, UV radiation, and herbivores. Similarly, the leaves exhibited the highest total phenolic content (8.3 mg catechol equi./g), followed by the stem (6.2 mg catechol equi./g) and the root (3.1 mg catechol equi./g). According to Strobelet *al.*, (2002), the leaves have the highest phenol concentration. Endophytic bacteria use sugars as a carbon

Comment [A34]: 1. Please provide some more comparisons with similar study done by others
2. The authors should finish the discussion with what are the main points (take home message).
3. The deeper scientific interpretations of your findings are strongly suggested
4. No discussion presented about protein content, Ortho-dihydroxy phenols, and Flavanols
5. No discussion presented about Endophytic bacteria isolation
6. Please replace old citations with recent publications. Add more latest citations

Formatted: Not Highlight

Formatted: Not Highlight

Comment [A35]: Table?

source from plants, which can alter their growth, metabolism, and interactions with the plant host (Daubechet *et al.*, 2021). The analysis of total soluble sugars indicated that stem (9.21 mg glucose equi./g) and leaves (9.97 mg glucose equi./g) having higher levels compared to the root (6.83 mg glucose equi./g). In comparison to roots, stems, and leaves have higher TSS content. Rather than storing sugar, these plant parts provide different purposes such as nutrient intake, structural support, and photosynthesis. Some plants, however, may store starch or other carbs in their storage roots or stems, resulting in somewhat higher TSS content than ordinary roots or stems (Powell and Matthews, 2018).

Enzymes play a crucial role in various physiological processes in plants, including defense against pathogens, oxidative stress management, and regulation of plant growth and development. In this study, we investigated the activities of three important enzymes, peroxidase, polyphenol oxidase, and catalase, in different plant parts. The root, stem, and leaves of a specific plant species were analyzed, and the changes in enzyme activity were measured. The results provide valuable insights into the distribution and potential functions of these enzymes in different plant organs. Peroxidase activity was highest in the root (359.98 OD/g), followed by the stem (349.22 OD/g) and leaves (327.05 OD/g). Rizwan *et al.*, 2012 investigated the peroxidase activity in different plant parts of *Brassica juncea* (Indian mustard) under heavy metal stress. They found that the roots exhibited significantly higher peroxidase activity compared to stems and leaves, suggesting a greater role of peroxidases in root tissues under stress conditions. Rennenberget *et al.*, 2006, found that the roots showed the highest peroxidase activity, followed by stems and leaves. The study suggested that peroxidases in roots might play a crucial role in the defense mechanisms of trees against environmental stresses in different parts of *Quercus ilex* (holm oak) trees.

Catalase activity was highest in the root (17.79 OD/g), followed by the stem (14.63 OD/g) and leaves (14.33 OD/g). A study on *Glycine max* (soybean) found that catalase activity was higher in the leaves compared to the roots and stems (Nakano and Asada, 1981). Polyphenol oxidase activity showed a similar trend, with the highest activity in the root (1.63 OD/g) and the lowest in the leaves (2.19 OD/g). However, the specific plant part with the maximum PPO enzyme activity can vary depending on the plant species and the physiological state of the plant. Murata and Galis (2014), emphasizes that leaves often exhibit higher PPO activity compared to other plant parts. This is because PPO plays a significant role in plant defense mechanisms against herbivory and pathogen attack, and leaves are the primary site of interaction with these threats.

CONCLUSION

Formatted: Highlight

Formatted: Not Highlight

Comment [A36]: Too old citation

Formatted: Highlight

Comment [A37]: 1. The conclusion must be self-explanatory and highlight novelty and implication of your study
2. In conclusion, please explain how the findings can be used for the government or relevant sectors.

The study of endophytic bacteria in pearl millet sheds light on their presence within the plant's tissues, emphasizing their role in influencing biochemical compounds and enzymes.

Pearl millet, vital in Asian and African agriculture, showcases diverse endophytic bacteria across its roots, stems, and leaves. These microorganisms, without causing visible symptoms, impact the plant's total phenolic content, soluble sugars, and enzymatic activities. The investigation reveals significant variations in biochemical compounds and enzyme activities among plant parts, indicating the crucial influence of endophytic bacteria. Understanding these complex interactions between endophytes and biochemical substances within pearl millet presents potential insights for agricultural practices, offering prospects to enhance plant health and productivity in sustainable crop production systems.

REFERENCES

1. Appa, R. S. and de Wet, J. M. J. (1999). Taxonomy and evolution. In: *Pearl Millet Breeding*, I. S. Khairwal, K. N. Rai, D. J. Andrews and G. Harinarayana, Eds. Science Publishers, Enfield, NH, pp. 29–47.
2. Bacon, C. W. and White, J. F. (2000). Microbial endophytes. *Marcel Dekker, CRC Press, New York*. (<https://doi.org/10.1201/9781482277302>)
3. Balbaa, S. I., Zaki, A. Y., and El Shamy, A. M. (1974). Total Flavonoid and Rutin Content of the Different Organs of *Sophora japonica* L. *Journal of the Association of official Analytical Chemists*, **57**(3): 752-755.
4. Bulgarelli, D., Rott, M., Schlaeppi, K., VerLoren V. T. E., Ahmadinejad, N., Assenza, F. and Schulze, L. P. (2013). Revealing structure and assembly cues for Arabidopsis root-inhabiting bacterial microbiota. *Nature*, **488**(7409): 91-95.
5. Carvalho, T. L. G., Balsemao-Pires, E., Saraiva, R. M., Ferreira, P. C. G. and Hemerly, A. S. (2014). Nitrogen signalling in plant interactions with associative and endophytic diazotrophic bacteria. *Journal of experimental botany*, **65**(19): 5631-5642.
6. Chandrashekhara, N. S., Deepak, S. A., Amruthesh, K. N., Shetty, N. P. and Shetty, H. S. (2007). Endophytic bacteria from different plant origin enhance growth and induce downy mildew resistance in pearl millet. *Asian Journal of Plant Pathology*, **1**: 1-11.
7. Daubech, Hirkala, D. L. M. and Nelson, L. M. (2021). Influence of plant genotype and sugar availability on root endophytic bacterial communities. *Plant and Soil*, 468, 75–88. [DOI: 10.1007/s11104-020-04762-5].

Formatted: Not Highlight

Comment [A38]: This sentence should be in introduction or Discussion sections with reference

Formatted: Not Highlight

Comment [A39]: Findings should be presented here with results

Comment [A40]: References should be in Format.

8. Hardoim, P. R., Van Overbeek, L. S. and Elsas, J. D. V. (2008). Properties of bacterial endophytes and their proposed role in plant growth. *Trends in Microbiology*, **16**(10): 463-471.
9. Janse, J. D. (2005). Phytobacteriology: Principles and Practices. *Centre for Agriculture and Bioscience International Publishing*. Pp. 366.
10. Kumar, V., Pathak, D. V., Dudeja, S. S., Saini, R., Giri, R., Narula, S. and Anand, R. C. (2013). Legume nodule endophytes more diverse than endophytes from roots of legumes or nonlegumes in soils of Haryana, India. *Journal of Microbiology and Biotechnology Research*, **3**(3): 83-92.
11. Li, Y., Uddin, W. and Kaminski, J. E. (2014). Effects of relative humidity on infection, colonization and conidiation of *Magnaportheorhizae* on perennial ryegrass. Department of Plant Pathology and Environmental Microbiology, The Pennsylvania State University, University Park 16802. **63**: 590- 597.
12. Murata, M. and Galis, I. (2014). Polyphenol oxidase in leaves: is there any significance to the chloroplastic localization. *Journal of Experimental Botany*, **65**(4): 1093-1100.
13. Nakano, Y. and Asada, K. (1981). Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant and Cell Physiology*, **22**(5): 867-880.
14. Powell, A. A. and Mathews, S. (2018). Seed Treatments: Developments and Prospects. *Sage Journal*, **10**(3): 61-86.
15. Rangjaroen, C., Rerkasem, B., Teamroong, N., Sungthong, R. and Lumyong, S. (2014). Comparative study of endophytic and endophytic diazotrophic bacterial communities across rice landraces grown in the highlands of Northern Thailand. *Archives of Microbiology*. **196**(1): 35-49.
16. Rennenberg, H., Mathews, S. and Nonoh, J. (2006). Respiration of the root apoplast and vitality of root cells. *Environmental and Experimental Botany*, **56**: 54-62.
17. Rizwan, Muhammad., Abaid, U. M. and Hassan, M. (2012). A critical review on effects, tolerance mechanisms and management of cadmium in vegetables. *Plant Physiology and Biochemistry*, **51**: 109-115.
18. Shannon, L. M., Key, E. and Law, J. Y. (1966). Peroxidase isoenzymes from horse reddish roots: isolation and physical properties. *Journal of Biological Chemistry*, **241**: 2166-2172.

19. Sinha, A. K. (1972). Colorimetric assay of catalase. *Analytical biochemistry*, **47**(2): 389-394.
20. Strobel, G.A. (2002). Microbial gifts from rain forests. *Canadian Journal of Plant Pathology*, **24**: 14-20.

UNDER PEER REVIEW