

Original Research Article

Impact of biofertilizers and organic manures on Herbage Production of KasuriMethi (*Trigonellacorniculata* L.) CV. – Pusa Kasuri

ABSTRACT

A study was conducted at the Research field, Department of Horticulture, College of Agriculture, Gwalior M.P. during rabi 2021-22. The purpose of the study is to evaluate the plant in terms of various parameters such as AGR, CGR, leaf area, fresh leaf yield and dry leaf yield. The results of the study indicated that the application of organic manures and biofertilizers improved the plant growth and yield. The highest median AGR (0.311 and 0.395 g/day) at 30-60 and 60-90 DAS, CGR (7.99 and 12.21 g/m²/day) at 30-60 and 60-90 DAS, leaf area (93.90 cm²), fresh leaf yield (5.78 Kg/plot) and (96.26 q/ha), dry leaf yield (0.49 Kg/plot) and (8.18 q/ha) were observed in the plants treated with FYM (16 t/ha) + Vermicompost (4 t/ha) + *Rhizobium* (10 ml/kg seed) + PSB (10 ml/kg seed) + KSB (10 ml/kg seed) as compared with control.

1. INTRODUCTION

Kasurimethi (*Trigonellacorniculata* L.) member of the family *Fabaceae* is an herbaceous, bushy, slow growing annual spice crop mainly grown for herbage that to dry herb. It is mainly grown as leafy vegetable and seeds in the plains of north India. It contains important nutrients required for growth and maintenance of human body [1]. Kasurimethi is a semi-arid crop, grows up to 30 cm height and its leaves are pinnate with the size of leaflets being 1.25-2.0 cm and consists bright orange-yellow colored flowers. Pods are 1.2-2.2 cm long, sickle shaped with 4-8 seeds per pod. Kasurimethi is mainly grown for herbage as well as for seed and used as a spice to add aroma and flavour to the food products.

Crop nutrients play a pivotal role in crop production, influencing plant growth, development, and overall productivity. Essential nutrients, including nitrogen, phosphorus, potassium, sulfur, calcium, magnesium, and various micronutrients, are crucial for the synthesis of proteins, enzymes, and other essential compounds within plants. Adequate nutrient availability ensures proper root development, robust vegetative growth, efficient flowering, and successful reproduction, ultimately leading to higher yields and improved crop quality [2]. Balanced nutrient management is essential for preventing deficiencies or excesses, both of which can compromise plant health and reduce agricultural productivity. By understanding and optimizing nutrient levels in the soil, farmers can enhance nutrient uptake by crops, promote sustainable agriculture, and contribute to global food security. Efficient nutrient management practices not only benefit crop production but also help conserve soil fertility, minimize environmental impact, and support long-term agricultural sustainability [3].

Bio-fertilizers are one of the important nutrient sources in the integrated nutrient management system. Application of *bio-fertilizers* helps in fixing the nitrogen, mobilizes the availability of nutrients and also helps in buildup of the micro flora. Combination effect of organic manures and nitrogen fixing biofertilizers and phosphate solubilizing bacterium helps to increase the availability of nutrients [4]. Biofertilizers play a crucial role in sustainable agriculture by promoting soil fertility and enhancing plant growth without causing harm to the environment [5]. Unlike chemical fertilizers that can have negative impacts on soil health and water quality, biofertilizers consist of living microorganisms such as bacteria, fungi, and algae that form symbiotic relationships with plants [6]. These microorganisms contribute to nutrient cycling, fix atmospheric nitrogen, and solubilize minerals, making essential nutrients more accessible to plants [7]. By fostering a healthy soil ecosystem, biofertilizers not only improve crop yields but also enhance soil structure, water retention, and overall resilience. Additionally, the use of biofertilizers can reduce the reliance on synthetic inputs, mitigating environmental pollution and contributing to a more sustainable and ecologically balanced agricultural system [8].

In crop production, especially for crops like methi (fenugreek), various organic amendments and beneficial microorganisms play vital roles in enhancing soil fertility and promoting plant growth. Vermicompost, produced through the decomposition of organic matter by earthworms, is a rich source of nutrients and improves soil structure, water retention, and microbial activity [9]. It enhances methi's nutrient uptake efficiency, leading to healthier plants and improved yields.

Farm yard manure (FYM) serves as a valuable organic amendment, providing a balanced mix of nutrients essential for plant growth [10]. Its incorporation into the soil enhances soil fertility, stimulates microbial activity, and improves methi's nutrient absorption. Rhizobium, a nitrogen-fixing bacterium, forms a symbiotic relationship with leguminous crops like methi. By converting atmospheric nitrogen into a plant-available form, rhizobium enhances methi's nitrogen supply, promoting vigorous growth and reducing the need for synthetic nitrogen fertilizers. Phosphorus solubilizing bacteria (PSB) play a crucial role in making phosphorus more accessible to plants [11]. Methi requires phosphorus for root development, flowering, and seed formation. PSB help solubilize insoluble phosphorus in the soil, making it available for methi plants, thereby improving overall crop productivity. Similarly, potassium solubilizing bacteria (KSB) contribute to the availability of potassium, another essential nutrient for methi. Potassium is vital for flowering, fruiting, and overall stress tolerance in plants. KSB enhance potassium uptake by solubilizing insoluble forms of potassium in the soil [6, 8].

In summary, the use of vermicompost, farm yard manure, rhizobium, phosphorus solubilizing bacteria, and potassium solubilizing bacteria in methi cultivation promotes

sustainable agriculture by improving soil fertility, nutrient availability, and overall crop health, contributing to increased methi yields and quality.

Keeping the above beneficial facts of biofertilizers and organic manures in mind the present study was carried out with the objective to find the effect of biofertilizers and organic manures on absolute growth rate (AGR), crop growth rate (CGR), leaf area and various leaf yields of kasurimethi.

2. MATERIAL AND METHODS

The experiment was conducted Experimental Field, Department of Horticulture, College of Agriculture, Gwalior (M.P.). The experiment was laid out in the Randomized Block Design with three replications. Each replication was comprised of sixteen treatments consisting organic manures i.e. FYM and Vermicompost and bio-fertilizers i.e. *Rhizobium*, PSB and KSB were applied for enhancing the crop physiological parameters and herbage yield of kasurimethi (*Trigonellacorniculata* L.) cv. Pusa Kasuri. The details of treatment combination used are T₀ - Control, T₁ - FYM (16 t/ha), T₂ - Vermicompost (4 t/ha), T₃ - *Rhizobium* (30 ml/kg seed), T₄ - PSB (30ml/kg seed), T₅ - KSB (30 ml/kg seed), T₆ - FYM (16 t/ha) + *Rhizobium* (30 ml/kg seed), T₇ - FYM (16 t/ha) + PSB (30 ml/kg seed), T₈ - FYM (16 t/ha) + KSB (30 ml/kg seed), T₉ - Vermicompost (4 t/ha) + *Rhizobium* (30ml/kg seed), T₁₀ - Vermicompost (4 t/ha) + PSB (30 ml/kg seed), T₁₁ - Vermicompost (4 t/ha) + KSB (30 ml/seed), T₁₂ - FYM (16 t/ha) + *Rhizobium* (10 ml/kg seed) + PSB (10 ml/kg seed) + KSB (10 ml/kg seed), T₁₃ - Vermicompost (4 t/ha) + *Rhizobium* (10 ml/kg seed) + PSB (10 ml/kg seed) + KSB (10 ml/kg seed), T₁₄ - FYM (16 t/ha) + Vermicompost (4 t/ha) + *Rhizobium* (10 ml/kg seed) + PSB (10 ml/kg seed) + KSB (10 ml/kg seed), T₁₅ - *Rhizobium* (10 ml/kg seed) + PSB (10 ml/kg seed) + KSB (10 ml/kg seed).

The experimental plot was ploughed thrice by tractor drawn cultivator and leveled. The clods were crushed weeds were removed and brought to fine tilt. The land was divided into plots of required size (1.90 m x 2.70 m). Provision was made for bunds and irrigation channels. The seeds of the variety Pusa Kasuri were used with the seed rate of 18-20 Kg/ha.

Observation were done at different stages of growth period such as AGR (g/day) at 30-60 and 60-90 DAS, CGR (g/m²/day) at 30-60 and 60-90 DAS, leaf area (cm²) at 60 DAS and yield parameter like fresh leaf yield (kg/plot), fresh leaf yield (q/ha), dry leaf yield (kg/plot), dry leaf yield (q/ha).

3. RESULT & DISCUSSION

There was significantly effect of various treatments on AGR and CGR at different interval from day after sowing. Among Different level of treatment T₁₄ was excellent treatment for enhancing AGR (g/day) and CGR (g/m²/day) in kasurimethi plant at different development stages and it also approved the maximum AGR (0.311 g/day) and CGR (7.99 g/m²/day) at 30-60 DAS and AGR (0.395 g/day) and CGR (12.21

g/m²/day) at 60-90 DAS, whereas minimum AGR and CGR was recorded in T₀ – Control. It may be due to higher nutrient uptake and bioactive substances which have a similar effect of GA₃ and cytokinin. This helps in breaking of apical dominance and accelerated higher number of branches, which increased the AGR and CGR of the plant. The results are in confirmation with the results achieved by Kusuma *et al.* [12], Babalshaware *et al.* [4] and Raghuvanshi *et al.* [13].

The result revealed that the different organic manures and bio-fertilizers significantly enlarge the leaf area of kasurimethi and the treatment T₁₄ was outstanding treatment for extension of leaf area in kasurimethi, it also established the maximum leaf area (3.90 cm²) at 60 DAS, treatment T₀ – Control was documented the minimum leaf area (2.00 cm²) at 60 DAS. This may be due to production of more number of branches and plant height and which enhanced availability of nutrients at the appropriate time, which has increased the leaf area. Findings are in agreement with those of Raiyani *et al.* [14], Suman *et al.* [15], Aakash *et al.* [5] and Sahu *et al.* [16].

Among the different levels of treatment T₁₄ was recorded maximum fresh leaf yield (5.78 kg/plot). It was at par to treatment T₉, T₁₀, T₁₁, T₁₂ and T₁₃. Otherwise, treatment T₀ – Control was recorded the minimum fresh leaf yield. Among the different levels of treatment T₁₄ was recorded maximum fresh leaf yield (96.26 q/ha) as the minimum fresh leaf yield (80.57 q/ha) was recorded in T₀ – Control, respectively. Similar findings were also reported by Meena *et al.* [17], Solanki *et al.* [8], Babalshaware *et al.* [4] and Chandan *et al.* [18].

Among the different levels of treatment T₁₄ was recorded significantly maximum dry leaf yield (0.49 kg/plot), as the minimum dry leaf yield (0.41 kg/ha) was recorded in treatment T₀ – Control, respectively. Among the different levels of treatment T₁₄ was recorded significantly maximum dry leaf yield (8.1 q/ha), as the minimum dry leaf yield (6.85 q/ha) was recorded in treatment T₀ – Control, respectively. It might be due to combined application of *Rhizobium*, PSB and KSB with different organic manures, which resulted in better utilization of inorganic nitrogen, greater biological N fixation and more synthesis of plant growth hormones. It enhances the dry matter accumulation in leaves and dry leaf yield in kasurimethi. Similar results for most of the characters were also reported by Murali *et al.* [19], Raiyani *et al.* [14], Babalshaware *et al.* [20] and Singh *et al.* [11].

Table 1: Effect of organic manures and bio-fertilizers on AGR (g/day) and CGR (g/m²/day) at 30-60 and 60-90 DAS of kasurimethi

Treatments detail	AGR (g/day) at 30-60 DAS	AGR (g/day) at 60-90 DAS	CGR (g/m ² /day) at 30-60 DAS	CGR (g/m ² /day) at 60-90 DAS
T ₀	0.225	0.295	6.95	9.20
T ₁	0.248	0.326	7.37	10.42
T ₂	0.256	0.326	7.46	10.55
T ₃	0.241	0.311	7.13	9.69
T ₄	0.243	0.317	7.28	9.87
T ₅	0.227	0.297	7.01	9.46
T ₆	0.260	0.330	7.54	10.74
T ₇	0.267	0.337	7.60	10.99

T ₈	0.256	0.326	7.46	10.55
T ₉	0.274	0.350	7.80	11.52
T ₁₀	0.280	0.369	7.85	11.56
T ₁₁	0.272	0.344	7.75	11.43
T ₁₂	0.299	0.380	7.89	11.72
T ₁₃	0.310	0.389	7.90	12.01
T ₁₄	0.311	0.395	7.99	12.21
T ₁₅	0.247	0.318	7.32	10.22
SEm ±	0.005	0.005	0.047	0.077
CD 5%	0.014	0.014	0.136	0.222

Table 2;Effect of organic manures and bio-fertilizers on leaf area (cm²) at 60 DAS,fresh leaf yield (kg/plot), (q/ha),dry leaf yield (kg/plot) and (q/ha) of kasurimethi

Treatments detail	Leaf area (cm ²) at 60 DAS	Fresh leaf yield (kg/plot)	Fresh leaf yield (q/ha)	Dry leaf yield (kg/plot)	Dry leaf yield (q/ha)
T ₀	2.00	4.84	80.57	0.41	6.85
T ₁	2.63	5.15	85.72	0.44	7.29
T ₂	2.74	5.19	86.54	0.44	7.36
T ₃	2.22	5.00	83.36	0.43	7.09
T ₄	2.34	5.08	84.67	0.43	7.20
T ₅	2.15	4.95	82.40	0.42	7.00
T ₆	2.69	5.29	88.19	0.45	7.50
T ₇	2.95	5.26	87.70	0.45	7.45
T ₈	2.82	5.26	87.55	0.45	7.44
T ₉	3.34	5.43	90.46	0.46	7.69
T ₁₀	3.53	5.51	91.79	0.47	7.80
T ₁₁	3.16	5.37	89.38	0.46	7.60
T ₁₂	3.71	5.58	92.92	0.47	7.90
T ₁₃	3.82	5.71	95.17	0.48	8.09
T ₁₄	3.90	5.78	96.26	0.49	8.18
T ₁₅	2.41	5.13	85.39	0.44	7.26
SEm ±	0.155	0.157	2.612	0.013	0.222
CD 5%	0.447	0.453	7.544	0.038	0.641

4. CONCLUSION

According to the current research, the use of organic manures and biofertilizers had a significantly positive impact on the growth and herbage yield of kasurimethi. Among the various treatments that were evaluated, T₁₄ – FYM (16 t/ha) + Vermicompost (4 t/ha) + *Rhizobium* (10 ml/kg seed) + PSB (10 ml/kg seed) + KSB (10 ml/kg seed) yielded the most favorable results in terms of growth viz., AGR (0.311 and 0.395 g/day) at 30-60 and 60-90 DAS, CGR (7.99 and 12.21 g/m²/day) at 30-60 and 60-90 DAS, leaf area (93.90 cm²) and herbage yield viz., fresh leaf yield (5.78 Kg/plot) and (96.26 q/ha), dry leaf yield (0.49 Kg/plot) and (8.18 q/ha).

REFERENCES

1. Singh, S., Choudhary ,M., R., Garhwal , O., P., Jakhar, M., L and Yadav, B., L. (2012) 'Effect of *biofertilizers* and inorganic sources of Nitrogen and Phosphorus on quality production of kasurimethi (*Trigonellacorniculata*)' *International J. Seed Spices* 2(2),38-40
2. Aakash, Thakur N. S., Singh M. K., Bhayal L., Meena K., Choudhary S. K., Kumawat N., Singh R. K., Singh U. P., Singh S. K., Sustainability in Rainfed Maize (*Zea mays* L.) Production Using Choice of Corn Variety and Nitrogen Scheduling. *Sustainability*. 2022;14(5):3116.
3. Kumar, V., Singh, M. K., Lakshmi, D. U., Aakash, Saikia, N. and Kumari, A. (2022). Response of N P S doses and urea foliar spray on lentil under guava (*Psidiumguajava*) + lentil (*Lens culinaris*) based agri-horti system. *The Indian Journal of Agricultural Sciences*, 92(11): 1410–1412.
4. Babaleshwar, S., Shetty, G.R., Pooja, D.A. and Bhat, R. (2020). Performance of KasuriMethi (*Trigonellacorniculata*L) under Hill Zone of Karnataka. *International Journal of Current Microbiology and Applied Sciences*, 9(6):1758-1762.
5. Aakash, Singh M. K., Saikia N., Bhayal L., Bhayal D., Effect of integrated nutrient management on growth, yield attributes and yield of green pea in humid subtropical climate of Indo-Gangetic Plains. *Annals of Agricultural Research*. 2023;44,2(Sep. 2023):190–196.
6. Solanki, K., Choudhary , S. K., Aakash, Singh , V., Singh , A. and Birla, D. (2023). Response of *Bacillus megaterium* and *Bacillus mucilaginosus* Strains on Yield and Quality of Soybean. *International Journal of Environment and Climate Change*, 13(11), 776–783.
7. Bhayal L., Aakash Jain M., P., Bhayal D., Meena K., Impact of foliar spray of NPK and Zn on soybean for growth, yield, quality, energetics and carbon foot print under dryland condition at Indore. *Legume Research*; 2022.
8. Solanki, K., Choudhary , S. K., Aakash, Singh , V., Nath , H., &Anshuman , K. (2023). Response of *Bacillus megaterium* and *Bacillus mucilaginosus* Strains on Growth and Nutrient Uptake of Soybean. *International Journal of Plant & Soil Science*, 35(21), 267–276.
9. Singh, B., Meena, R.N., Meena, A., Meena, K., Aakash, Meena, R. and Kumar, S. (2022). Effect of sowing date and inter-row annual legume green-manuring on growth, yield and quality of desi cotton (*Gossypiumarboreum*). *Indian Journal of Agronomy*. 67(3): 289-291.
10. Bhayal, L., Kewat, M.L., Bhayal, D., Aakash, Jha, A.K., and Badkul, A.J. (2022). Influence of Different Sowing Dates and Nutrient Management on Yield Attributes and Yield of Wheat (*Triticumaestivum* L.). *International Journal of Plant & Soil Science*, 34(22), 362-367.
11. Singh S. R., Singh M., K., Aakash, Meena K., Vishwakarma S., P. Effect of Different NPK Levels on Fodder Production of Sudan Grass (*Sorghum bicolor* var. *Sudanese*). *International Journal of Bio-resource and Stress Management*. 2021;12(3):199-204.

12. Kusuma, M.V., Venkatesha, J., Ganghadarappa, P.M., Hiremath, J.S., Mastiholi, A.B. and Manjunatha, G. (2019). Effect of integrated nutrient management on growth and yield of fennel (*Foeniculumvulgare Mill.*).
13. Raghuwanshi, D., K., Tomar, K., S. (2021) 'Study of organic production techniques of fenugreek (*Trigonellafoenum-graecum L.*)' M. Sc (Ag/Horti) Thesis, RVSKVV Gwalior, College of Agriculture, Gwalior (M.P.).
14. Raiyani, V., N., Kathiriya, R., K., Thummer, V., M. and Rupareliya V., V. (2018) 'Effect of FYM and biofertilizers on growth, yield attributes and yield of fenugreek (*Trigonellafoenum-graecum L.*)' *International Journal of Chemical Studies*, 6(4): 746-748.
15. Suman, P., Lakshminarayana, D., Prasanth, P. and SaidaNaik, D. (2019). Effect of integrated nutrient management on yield parameters of coriander (*CoriandrumsativumL.*) cultivars under Telangana conditions. *International Journal of Chemical Studies*,7(1): 392-394.
16. Sahu, P.K., Naruka, I.S., Haldar, A., Chundawat, R.S. and Kumar, L. (2020). Studies on the effects of integrated nutrient management on fenugreek (*Trigonellafoenum-graecumL.*). *International Journal of Chemical Studies*, 8(2): 1082-1089.
17. Meena, R.D., Lal, G., Mehta, P., Sharma, Y.K., Meena, S.S., Meena, R.D., Meena, N.K., Mishra, B.K. R., Meena, R.L., Meena, N.L. and Tripathi, G.K. (2018). Efficacy of soil solarization on growth and yield of cumin (*CuminumcyminumL.*) under arid conditions. *International Journal of Seed Spices*,8(1): 74-79.
18. Chandan, T.K., Lakshminarayana, D., Seenivasan, N., Joshi, V. and Kumar, S.P. (2021). Growth and yield of KasuriMethi (*TrigonellacorniculataL.*) var. PusaKasuri as influenced by different organic manures and biofertilizers under Telangana conditions. *International Journal of Horticulture and Food Science*, 3(2): 26-30.
19. Murali, M., Rajiv, U. and Hemant, K. (2018). Effect of different levels of organic manure on the growth and yield of mustard (*Brassica juncea L.*) under jatropha (*Jatropha circus L.*) based agroforestry system. *Journal of Pharmacognosy and Phytochemistry*, 7(4): 955-958.
20. Babaleshwar, S.B., Shetty, G. R., Shivakumar, S. J. and Nadukeri, S. (2017). Influence of integrated nutrient management on growth and physiological attributes of kasurimethi (*TrigonellacorniculataL.*) under hill zone of Karnataka. *Environment & Ecology*, 35(2): 661-665.