

## Serological Evidence of West Nile Virus among individuals with febrile illness in a Tertiary Hospital in Port Harcourt, Nigeria

### ABSTRACT

**Aims:** West Nile Virus (WNV) infection can cause severe illness. Very little is known about the seroepidemiology of WNV infection in patients with febrile illness in Nigeria and many other developing countries. This study was carried out to determine the seroprevalence of WNV in patients with febrile illness attending a tertiary hospital in Port Harcourt, Nigeria and to determine if there was an association between WNV infection in relation to age and sex.

**Study Design:** Cross Sectional Study

**Place and Duration of Study:** The University of Port Harcourt Teaching Hospital (UPTH), in Rivers State, Nigeria from September, 2019 to December 2019.

**Methods:** Human sera were obtained and WNV IgG was determined using Enzyme Linked Immunosorbent Assay (ELISA) technique.

**Results:** Of the 90 study subjects tested, WNV IgG antibodies were present in 27 (30.0%) study participants while 63 (70.0%) study participants were seronegative for WNV IgG antibody. In relation to age, higher prevalence of WNV occurred among 61-70 years (31.3%, n= 5) compared to 41-60 (30.8%, n= 12) and 20-40 (28.6%, n= 10). Higher prevalence of WNV IgG antibodies occurred in males (34.3%, n=12) than their female counterparts (30.9%, n= 17). This study indicated that there is no association between WNV infection in relation to age and sex.

**Conclusion and Recommendations:** These results show that WNV is circulating in Rivers State and has accounted for malaria-like infection in the region. It is recommended that WNV serological testing for malaria patients should be included as a routine test since they are most likely to present similar symptoms of WNV fever. Also, proper hygiene which includes eliminating mosquito breeding sites is recommended to mitigate the spread of West Nile Virus infection.

**Keywords:** West Nile Virus (WNV), Antibodies, IgG, seroprevalence, ELISA.

### 1. INTRODUCTION

“West Nile virus (WNV) is an emerging and re-emerging zoonotic flavivirus first identified in and endemic to Africa” (Olufemi et al., 2021). “West Nile Virus (WNV) is a mosquito-borne viral pathogen that is the causative agent of West Nile fever and encephalitis” (Kolawole et al., 2018). It is one of the most prevalent arboviruses in the world and has been documented as a pathogen that has an impact on both human and animal health (Shomaker et al., 2013).

“The virus is transmitted between birds by biting mosquitoes, with equids and humans being incidental hosts” (Olufemi et al., 2021). “The majority of infected incidental hosts display no or only mild clinical signs, but a fraction develop encephalitis” (Olufemi et al., 2021).

“First discovered in 1937, the West Nile virus (WNV) was found in a patient who had a febrile illness and lived in the West Nile district of northern Uganda” (Hughes et al., 1940; Castro-Jorge et al., 2019). The WNV has been isolated across several continents of the world including Africa, Asia, Europe, the Mediterranean region, the Middle East, Australia and the Americas (Barakat et al., 2004), and in 1951, the first known WNV human epidemic was reported in Israel, with young children accounting for the majority of cases (Qian et al., 2014). “It was the first time that the main clinical signs were properly described, and they primarily included fever, headache, anorexia, exanthema, myalgia, abdominal discomfort, and vomiting and the condition was minor, with no known fatalities and infections appears to be sporadic with symptoms of lymphadenopathy, sore throat, and diarrhoea” (Castro-Jorge et al., 2019).

The West Nile virus (WNV) is a neurovirulent, zoonotic, mosquito-borne virus that belongs to the family *Flaviviridae* in the genus *Flavivirus* and is well-known to cause encephalitis and meningitis outbreaks (Opaleye et al., 2014; Blázquez et al., 2015; Sule & Oluwayelu, 2016). The virus can infect and cause disease in horses and humans while also maintaining an enzootic cycle between ornithophilic mosquitoes and birds (Lim et al., 2011). Infection in humans is initiated following a bite from a female mosquito (e.g., *Culex* mosquito species) infected with the WNV virus (Qian et al., 2014; Sule & Oluwayelu, 2016).

Studies carried out in the Nile Delta region revealed that WNV was endemic along the Nile, with a 60% seroprevalence rate in people, and that it was contagious in a wide variety of animals, including birds and non-human mammals (Hughes et al., 1956; Castro-Jorges et al., 2019). WNV's arthropod-borne nature was first hypothesized in 1943, and it has subsequently been identified as one of the most pervasive arboviruses since it can infect more than 65 different species of mosquitoes (Colpitts et al., 2012).

In Northern Nigeria, WNV has been linked to mosquitoes and specific antibodies have been isolated in patients with febrile illness (Baba et al., 2015). Seroprevalence studies can be used to monitor WNV activity in any population. As a result, it has been reported that camels, goats, cattle, sheep, and horses in Nigeria and Romania have antibodies against WNV that prevent hemagglutination (Olaleye et al., 1990; Hubalek & Halouzka, 1999).

“Blood serves as a vehicle for transmission of blood-borne pathogens including hemoparasities” (Okonko et al., 2012). “Malaria is one of the most prevalent and deadly widespread of all parasitic diseases in the world. It is mosquito-borne and one of the killer diseases of the world” (Ibekwe et al., 2009). Malaria is a life-threatening disease caused by parasites that are transmitted to people through the bites of infected female *Anopheles* mosquitoes (Ejike et al., 2020a; Okonko et al., 2023a, b, c; Innocent-Adiele et al., 2023).

Malaria remains a significant health threat worldwide (Okonko et al., 2023c, d). Almost half of the world populations are at the risk of malaria (Ali et al., 2010; Okonko et al., 2019), although malaria transmission is unstable throughout the year in these reports (Okonko et al., 2012b). Over 50% of the people on earth are vulnerable to the malaria infection and they have estimated

over 300 million malaria cases annually in the tropics (Okonko et al., 2019; Ejike et al., 2020b; John et al., 2020). An estimated 300 million malaria cases occur annually in the tropics, with 90% of these in the sub-Saharan, a region that already suffers the tremendous burden of HIV-1 infection (Okonko et al., 2023b; Innocent-Adiele et al., 2023). Co-infection with malaria and viruses such as HIV, West Nile virus, Dengue virus, among others, have led to several deaths especially in sub-Saharan Africa (Okonko et al., 2021).

“Diagnostic tools for WNV infection in Nigeria are not well established hence the current prevalence rate of WNV infection in Nigeria is unknown” (Kolawole et al., 2018). In a recent study reported by Sule and Oluwayelu (2016), the study showed that, there is no correlation between West Nile Virus (WNV) infection and febrile illness in people of Southwestern Nigeria. Although very few studies have shown the prevalence of WNV in Nigeria (Baba et al., 2013), none have shown its presence in the study area and there is no evidence of WNV exposure among humans in this area. In this study we report the prevalence of West Nile Virus (WNV) among patients with febrile illness attending the University of Port Harcourt Teaching Hospital and also show that there is no correlation between age and WNV infection among study subjects.

## **MATERIALS AND METHODS**

### **2.1. Study Area**

The study area is University of Port Harcourt Teaching Hospital, located in municipal area of Port Harcourt. Port Harcourt is the capital city of Rivers State, also known as the Garden City, it is located in the forest zone of South Southern Nigeria. Port Harcourt city lies on the longitude  $7^{\circ}$  East of Greenwich meridians and latitude  $4^{\circ}75'$  North of the Equator. Port Harcourt heaviest precipitation occurs during September with an average of 370mm of rain. December on average is the driest month of the year; with an average rainfall of 20mm. Temperatures throughout the

year in the city is relatively constant, showing little variation throughout the course of the year. Average temperature is typically between 25°C - 28°C in the city. The city is an important trade majorly petroleum and educational centre and houses one of the largest and foremost teaching hospitals in Africa.

## **2.2 Study Population**

The study population constituted of 90 malaria patients with apparent symptoms of fever in the University of Port Harcourt Teaching Hospital (UPTH).

## **2.3 Blood Sample Collection, Plasma Preparation and Storage**

About 5ml of blood sample was aseptically collected by venipuncture from patients. Each blood sample was dispensed into appropriately labeled EDTA-treated blood sample tube, screw-capped and left at room temperature for about 40 min, after which it was spun at 3,000 rpm for 10 min to separate plasma from the blood. The plasma was dispensed into labeled Eppendorf tubes and stored at -20°C until analyzed for WNV antibodies. Samples were clearly identified with codes. Haemolysed and visibly hyperlipemic samples, as well as those containing residues of fibrin or heavy particles were discarded as they could generate false serologic results. Plasma samples were stored at +2°C - 8°C up to five days after collection. For longer storage periods, samples were stored frozen at -20°C.

## **2.4. Serological Analysis of West Nile virus Antibody**

Microplate were ~~are~~ coated with a highly purified immune dominant West Nile virus (WNV) antigen. In the first incubation, the solid phase is treated with diluted samples and anti-WNV antibodies are captured, it presents by the antigens. After washing out all the components of the sample, in the second incubation bound, anti-WNV were detected by the addition of a mix of both anti-hlgG antibody labeled with peroxidase (HRP). The enzyme captured on the solid

phase, acting on the substrate/chromogen mixture generates an optical signal that is proportional to the amount of WNV antibodies present in the sample. Test results were interpreted as ratio sample OD450nm (S) and the cut-off value (Co), mathematically  $S/Co \leq 0.9$  as Negative, 0.9-1.1 as Equivocal and  $>1.2$  as Positive. The results were then used to evaluate the sociodemographic characters obtained from the questionnaires administered to participants that were enrolled in the study.

## 2.5. Data Analysis

Data analysis was carried out using Microsoft Excel 2016 version to calculate the international unit (IU) from optical density (OD). Values less than or equal to 0.9 were considered negative while values greater than or equal to 1.2 were considered positive. Results are expressed as numbers and percentages.

## 3. RESULTS

### 3.1. Patients Characteristics

The total number of patients with febrile illness included in this study was 90 with age range of 20-70yrs. The age group (41-60yrs) constituted the largest populations making up 43.3%, followed by age group (20-40yrs) (38.9%) while age group (61-70yrs) were the least (17.8%). Characteristics taken into consideration were age and sex as shown in Table 1.


**Table 1: Participants Characteristics**

Characteristics	No. Tested (%)
<b>Age groups (years)</b>	
20-40	35 (38.9)
41-60	39 (43.3)

61 and above	16 (17.8)
<b>Sex</b>	
Males	35 (38.9)
Females	55 (61.1)
<b>Total</b>	<b>90 (100.0)</b>

### 3.2. Overall Seroprevalence of West Nile Virus (WNV)


A total number of 90 samples were tested, of which 27(30.0%) were seropositive and 63(70.0%) were seronegative for WNV as shown in Figure 1.



**Figure 1: Overall Seroprevalence of West Nile Virus (WNV) IgG antibodies**

### 3.3. Seroprevalence of West Nile Virus (WNV) IgG antibodies in Relation to Age


In relation to age, higher prevalence of West Nile Virus (WNV) IgG antibodies was observed in the elderly (61-70yrs) (31.3%, n= 5) compared to middle aged subjects (41-60yrs) (30.8%, n= 12) and the young adults (20-40yrs) (28.6%, n= 10) as presented in Figure 2. This difference was not statistically associated ( $X^2 = 0.971947$ ,  $df= 2$ ,  $p>0.05$ ).



**Figure 2: Seroprevalence of West Nile Virus (WNV) IgG antibodies in relation to age**

### **3.4 Seroprevalence of West Nile Virus (WNV) IgG antibodies in Relation to Sex**

Higher prevalence of WNV IgG antibodies was detected in male subjects (34.3%, n=12) than in female subjects (30.9%, n= 17) as shown in Figure 3. This difference was not statistically associated ( $X^2 = 0.738258$ , df= 1,  $p>0.05$ ).



**Figure 3: Seroprevalence of West Nile Virus (WNV) IgG antibodies in relation to sex**

#### 4. DISCUSSION

This study provides scientific data on the seroprevalence of **West Nile Virus (WNV) IgG antibodies** among patients who were recruited in the study with febrile illness in Port Harcourt, Rivers State, **Nigeria**. The seroprevalence of West Nile Virus in certain parts of Nigeria is well documented (Baba et al., 2006; Sule & Oluwayelu, 2016). In this study, we examined a total of 90 study participants who were accessing healthcare services in a tertiary teaching hospital in Port Harcourt who were recruited in this study with clinical presentation of febrile illness. Of this number, more females participated in the study. This observation agrees favorably with Ahmed and Hamedelnil (2015), and Kolawole et al. (2018) findings. This higher proportions in females shifted the population density in favor of the female subjects (Kolawole et al., 2018). Also, higher proportions of the participants were in age group 41-60 years age group. This observation differs from that of Kolawole et al. (2018) who observed a higher proportion in age group 21-30 years. This higher proportions in 41-60 years age groups shifted the population density in favor of this age groups.

Of these 90 participants, 30.0% were positive to anti-WNV IgG antibodies. In recent years, more seroprevalence studies have been reported, similar to the findings of this study (Baba et al., 2006), however, this figure (30.0%) is higher than the 19.4% and 7.5% reported by Kolawole et al. (2015, 2018), respectively, in Ilorin, Nigeria and 25.0% reported by Baba et al. (2013) in Maiduguri, Nigeria. It is higher than the 9.5% and 13.2% obtained in Kenya and Sudan, respectively (Mease et al., 2011; Ahmed & Hamedelnil, 2015); the 24.0% reported in another region of Kenya (Sutherland et al., 2011)

The 30.0% reported here is lower than the 66.0% reported in Democratic Republic of Congo (Nur et al., 1999), the 80.2% reported by Baba et al. (2006) among patients with febrile illness suspected of malaria/typhoid fever in Maiduguri, Nigeria, the 35.0% and 85.0% reported by Oderinde et al. (2020) in Maiduguri, Nigeria, and the 76.5% reported in Nigeria (Abdullahi et al., 2020). Baba et al. (2006) concluded that their study area was endemic for flavivirus and that the peak of WNV activity in the semi-arid zone in Nigeria appeared to be in November.

In another study, it was reported that 45.0% of febrile participants had WNV antibodies but were negative for malaria parasite and Widal tests, thereby accounting for undifferentiated febrile illness (Sule et al., 2016). Increasingly, surveys suggest a high seroprevalence of WNV antibodies in Nigerian populations. The differences in the results obtained from this study and other studies in Nigeria and overseas could be largely influenced by environmental conditions (Baba et al., 2006; Kolawole et al., 2018). This in turn indicates that WNV is highly endemic

within Nigeria and further investigations for the presence and distribution of virus in reservoir avian species and vector mosquito species are needed (Sule et al., 2018).

In this present study, the elderly (61 years and above) were more susceptible to WNV infection (suggesting underlining diseases or immunosuppressed state) than young adults (20-40 years) and middle-aged (41-60 years) but the difference was not statistically significant. This is different from what was reported by Ahmed and Hamedelnil (2015) and Kolawole et al. (2018). They (Baba et al., 2006; Ahmed & Hamedelnil, 2015; Kolawole et al., 2018) reported that age has not been shown to be a potential risk factor in the acquisition of WNV infection. The distribution of WNV with respect to age in their studies were widespread and not limited to a particular age group.

Also, this study indicated that higher prevalence of WNV IgG antibodies occurred in male subjects than in female subjects but the difference was not statistically significant. This showed that gender is not a risk factor important for infection with WNV. Hence, WNV infection and gender are independent or have no association. This is in agreement with previous report on gender and acquisition of WNV infection (Baba et al., 2013; Kolawole et al., 2018). **However, our results differ from what was reported by Ahmed and Hamedelnil (2015) and Kolawole et al. (2018) who reported higher prevalences in females than males.**

Factors which could influence the acquisition of WNV infection and its prevalence may include housing and roofing type, abundance of surrounding bushes and trees which can become a favorable habitat for mosquitoes and the presence of stagnant water which is an established

factor that contributes significantly as a breeding site for mosquitoes, thereby increasing the numbers of mosquitoes and consequently the potential risk for WNV infection.

## **5. CONCLUSION**

In this study, low prevalence of West Nile Virus (WNV) IgG antibodies was shown among febrile patients in Rivers State, Nigeria. This study indicated that there is no association between WNV IgG antibodies in relation to age and sex. Although, estimating the global burden of this infection is important for appreciating the scale of an epidemic, distribution of resources to those most affected which includes identifying testing and counselling those at risk is strongly recommended. Also, local healthcare providers should encourage personal hygiene practice and the correct use of mosquito treated nets as this will significantly reduce exposure to mosquito bites and the risk for WNV infection in people living in mosquito endemic regions.

### ***Acknowledgements***

The authors would like to acknowledge the support obtained from the management and staff of the University of Port Harcourt Teaching Hospital (UPTH), Port Harcourt, Rivers State, Nigeria, Nigeria during the enrollment and collection of samples used in this study. The authors are grateful to the participants for their willingness to be part of the study.

### ***Disclosure of conflict of interest***

The authors have declared that no competing interests exist.

### **Ethical Approval and Consent**

Clearance from the health research ethical committee of University of Port Teaching Hospital (UPTH) was obtained in accordance with the code of ethics for biomedical research involving

human subjects. Signed informed consent was also obtained from each of the subjects after carefully explaining the concept of the study to them and questionnaires were distributed.

## REFERENCES

- Abdullahi, I. N., Emeribe, A. U., Ghamba, P. E., Omosigho, P. O., Bello, Z. M., Oderinde, B. S., Fasogbon, S. A., Olayemi, L., Daneji, I. M., Musa, M. H., et al. (2020). Distribution pattern and prevalence of West Nile virus infection in Nigeria from 1950 to 2020: A systematic review. *Epidemiological Health*, 42, e2020071.
- Ahmed, S. B. Y. & Hamedelnil, Y. F. (2015). Serodetection of West Nile Virus among febrile non-malarial patients at Wad Madeni Teaching Hospital. *Intern J Adv Sci Tech Res*. 5(3): 307–341.
- Alli, J. A., Okonko, I. O., Abraham, O. A., Kolade, A. F., Ogunjobi, P. I., Tonade, O. O., Ojezele, M. O. & Nwanze, J. C. (2010). A serosurvey of blood parasites (Plasmodium, Microfilaria, HIV, HBsAg, HCV antibodies) in prospective Nigerian blood donors. *Research Journal in Medical Sciences* 4(4):255-275.
- Baba, M. M., Marie-Francois, S. & Adeniji, J.A. (2006). The seasonal Effect of West Nile Virus in a Semi-arid Zone, Nigeria. *Trends in Medical Research*, 1, 31-38.
- Baba, M., Logue, C. H., Oderinde, B., Abdulmaleek, H., Williams, J., Lewis, J., Laws, T. R., Hewson, R., Marcello, A. & D'Agaro, P. (2013). Evidence of arbovirus co-infection in suspected febrile malaria and typhoid patients in Nigeria. *Journal of Infection in Developing countries*, 7(1), 051-059.

- Barakat, A. J., Reese, D. J., Barbey-Morel, C. & Evans, C. (2004). West Nile virus disease also affects children. *Case Report Clinical Practice Revision*, 5, 269-272.
- Blázquez, A-B., Martín-Acebes, M. A. & Saiz, J-C. (2015). Amino acid substitutions in the non-structural proteins 4A or 4B modulate the induction of autophagy in West Nile virus infected cells independently of the activation of the unfolded protein response. *Frontier Microbiology*, 5, 797.
- Castro-Jorge, L. A., Siconelli, M. J., Ribeiro, B., Moraes, J. M., Moraes, J. B., Agostinho, M. R., Klein, T. M., Floriano, V. G. & Fonseca, B. A. (2019). West Nile virus infections are here! Are we prepared to face another flavivirus epidemic? *Journal of the Brazilian Society of Tropical Medicine*, 52, e20190089
- Colpitts, T. M., Conway, M. J., Montgomery, R. R., & Fikrig, E. (2012). West Nile Virus: biology, transmission, and human infection. *Clinical Microbiology Reviews*, 25(4), 635–648. <https://doi.org/10.1128/CMR.00045-12>
- Ejike, I. U., Kparobo, M. O., Innocent-Adiele, H. C. & Okonko, I. O. (2020a). HIV/*Plasmodium falciparum* Coinfection among HIV-1 Infected Individuals in Uyo, Nigeria. *New York Science Journal*, 13(6):1-8
- Ejike, U. I., Cooney, T. I., Innocent-Adiele, H. C. & Okonko, I. O. (2020b). Analysis of HIV/Malaria Coinfections among HIV-1 Infected Individuals in Two Tertiary Hospitals in Old Cross River State, Nigeria. *International Journal of TROPICAL DISEASE & Health*, 41(10): 31-41
- Hubalek, Z. & Halouzka, J. (1999). West Nile fever - a re-emerging mosquito-borne viral disease in Europe. *Emerging Infectious Diseases*, 5(5), 643-50.

- Hughes, T. P., Paul, J. H., Smithburn, K. C. & Burke, A. W. (1940). A Neurotropic Virus Isolated from the Blood of a Native of Uganda 1. *American Journal of Tropical Medicine and Hygiene*, 1-20 (4), 471-92.
- Ibekwe, A. C., Okonko, I. O., Onunkwo, A. I., Ogun, A. A. & Udeze, A. O. (2009). Comparative Prevalence Level of Plasmodium in Freshmen (First Year Students) of Nnamdi Azikwe University in Awka, South-Eastern, Nigeria. *Malaysian Journal of Microbiology*, 5 (1): 51-54
- Innocent-Adiele, H. C., Onyedibia, G. C., Emeruem, K. E., Ogbuji, C. C. & Okonko, I. O. (2023). Serological evidence of HIV and Malaria co-infection among febrile HIV-Infected patients attending a tertiary Hospital in Port Harcourt, Nigeria. *International Journal of Multidisciplinary Research and Technology (IJMRT)*, 4(3): 18-24
- John, P.C., Cooney, T.I., Innocent-Adiele, H.C., Stanley, C.N. & Okonko, I.O. (2020). HIV/Malaria Coinfection among HIV-Infected Individuals in Calabar, Nigeria. *International Journal of Virology & Molecular Biology* 9(1):6-10
- Kolawole, O. M., Adelaiye, G. & Ogah, J. I. (2018). Emergence and Associated Risk Factors of Vector Borne West Nile Virus Infection in Ilorin, Nigeria. *Journal of Arthropod-Borne Diseases*, 12(4), 341-350.
- Lim, P.Y., Behr, M. J., Chadwick, C. M., Shi, P. Y. & Bernard, K. A. (2011). Keratinocytes are cell targets of West Nile virus in vivo. *Journal of Virology*, 85(10), 5197–5201.
- Mease, L. E., Coldren, R. L., Musila, L. A., Prosser, T., Ogolla, F., Ofula, V. O., Schoepp, R. J., Rossi, C. A., & Adungo, N. (2011). Seroprevalence and distribution of arboviral infections among rural Kenyan adults: a cross-sectional study. *Virology journal*, 8, 371. <https://doi.org/10.1186/1743-422X-8-371>

- Nur, Y. A., Groen, J., Heuvelmans, H., Tuynman, W., Copra, C., & Osterhaus, A. D. (1999). An outbreak of West Nile fever among migrants in Kisangani, Democratic Republic of Congo. *The American Journal of Tropical Medicine and Hygiene*, 61(6), 885–888. <https://doi.org/10.4269/ajtmh.1999.61.885>
- Oderinde, B. S., Mora-Cárdenas, E., Carletti, T., Baba, M. M., & Marcello, A. (2020). Prevalence of locally undetected acute infections of Flaviviruses in North-Eastern Nigeria. *Virus Research*, 286, 198060. <https://doi.org/10.1016/j.virusres.2020.198060>
- Okonko, I. O., Adejuwon, O. A., Okerentugba, P. O. & Innocent-Adiele, H. C. (2012a). Circulating *Plasmodium falciparum* and HIV 1/2 as Co-infections among Blood Donors in Ibadan, Southwestern Nigeria. *Nature and Science*, 10(9):42-47
- Okonko, I. O., Adejuwon, A. O., Okerentugba, P. O. & Frank-Peterside, N. (2012b). *Plasmodium falciparum* and HIV- 1/2 Coinfection among children presenting at the Out-patient clinic of Oni Memorial Children Hospital in Ibadan, Southwestern Nigeria. *Nature and Science*, 10(8):94-100.
- Okonko, I. O., Onwubuche, K., Cooney, T. I., Agbagwa, O. E., & Chijioke-Nwauche, I. N. (2021). Dual Infection of HIV and Malaria among HIV-Infected Individuals in Port Harcourt, Nigeria. *South Asian Journal of Parasitology*, 5(1), 26-31
- Okonko, I. O., Jerry, G. A., Okonko, B. J. & Baeka, G. B. (2023a). Serological Prevalence of Malaria and Hepatitis coinfection among HIV-Infected Patients presenting at a Tertiary Hospital in Port Harcourt, Rivers State, Nigeria. *Biomedicine and Nursing*, 9(1), 1-12
- Okonko, I. O., Shaibu, N., Okonko, B. J., Mba, E. I., Ogbuji, C. C. & Frank-Peterside, N. (2023b). HIV/Malaria Co-Infection and Its Determinants Among People Living With

- HIV/AIDS In Yenagoa, Bayelsa State, Nigeria. *Magna Scientia Advanced Biology and Pharmacy (MSABP)*, 09 (01):020-038
- Okonko, I. O., Shaibu, N., Mbah, E. I., Okonko, B. J. & Cookey, T. I. (2023c). Triple infections of HBV, HCV and Malaria *Plasmodium falciparum* among HIV-infected Individuals in Yenagoa, Bayelsa State, Nigeria. *Research Square PREPRINT (Version 1):1-18*. <https://doi.org/10.21203/rs.3.rs-2711727/v1>. 21 March 2023
- Okonko, I. O., Shaibu, N., Mbah, E. I., Okonko, B. J. & Cookey, T. I. (2023). Triple Infections of HBV, HCV and Malaria *Plasmodium falciparum* Among HIV-Infected Individuals in Yenagoa, Bayelsa State, Nigeria. *Annals of Experimental Biology*, 11(2), 52-63
- Okonko, I., Osadebe, A., Okoli, E., & Eke, E. (2019). Malaria and HIV seropositivity: study on selected individuals at a tertiary healthcare centre in Port Harcourt, Nigeria. *Microbiologia Medica*, 34(8160):32-36
- Olaleye, O. D., Omilabu, S. A., Ilomechina, E. N. & Fagbami, A. H. (1990). A survey for hemagglutination inhibiting antibody to West Nile virus in human and animal Sera in Nigeria. *Elsevier*, 13, 35-38.
- Olufemi, O. T., Barba, M. & Daly, J. M. (2021). A Scoping Review of West Nile Virus Seroprevalence Studies among African Equids. *Pathogens*, 10, 899. <https://doi.org/10.3390/pathogens10070899>
- Opaleye, O. O., Olawumi, B. G., Odewale, G. B., & Ojo, J. A. (2014). Absence of Serological Evidence of West Nile Virus in Blood Donors in South Western Nigeria. *Sudan Journal of Medical Science*, 9(4), 235-237.

- Qian, F., Thakar, J., Yuan, X., Nolan, M., Murray, K. O., Lee, W. T., et al. (2014). Immune Markers Associated with Host Susceptibility to Infection with West Nile Virus. *Viral Immunology*, 27(2), 39-47.
- Shomaker, T. S., Green, E. M. & Yandow, S. M. (2013). Perspective: One Health: a compelling convergence. *Academic Medicine*, 88 (1), 49–55.
- Sule, W. F., Oluwayelu, D. O. & Johnson, N. (2018). Epidemiology and ecology of West Nile virus in sub-Saharan Africa. *Parasites and Vectors*, 11, 414.
- Sule, W.F. & Oluwayelu, D.O. (2016). Serologic Evidence of exposure to West Nile virus among humans in Ibadan, southwestern Nigeria. *Sudan Journal of Medical Sciences*, 11(3), 111–124.
- Sutherland, L. J., Cash, A. A., Huang, Y. J., Sang, R. C., Malhotra, I., Moormann, A. M., King, C. L., Weaver, S. C., King, C. H., & LaBeaud, A. D. (2011). Serologic evidence of arboviral infections among humans in Kenya. *The American journal of tropical medicine and hygiene*, 85(1), 158–161. <https://doi.org/10.4269/ajtmh.2011.10-0203>