

Genetic diversity and phylogenetic relationships among *Jatropha curcas* L. genotypes from eastern India

Abstract:

Genetic diversity and relationships among 31 genotypes collected from wide geographical range in eastern India was studied employing Random Amplified Polymorphic DNA (RAPD) markers. Identified 19 markers produced 139 bands in total. Of these 131 bands were polymorphic exhibiting a high polymorphism of 94.24%. Similarity indices estimated on the basis of RAPD primers ranged widely from 0.44 to 0.83 which suggests that these accessions represent genetically diverse population possibly due to predominance of cross pollination and seed source variability. Genotyping data obtained for RAPD primer across collected accessions were used to generate the UPGMA - based phylogenetic tree which shows two major clusters. The cluster I was consisted 16 genotypes from two states Jharkhand and Odisha (Chotanagpur and Eastern plateau region) while 15 accessions from West Bengal and Bihar (Indo Gangetic plains region) were grouped together in the Cluster II. This is clear alignment of accessions as per geographical regions was possibly due to dissimilar growing conditions in different regions.

Key words: Affinity, molecular characterization, polymorphism, seed oil, variability

1. Introduction

Jatropha curcas Linn. (Family-Euphorbiaceae) plant is remarkable for its drought hardiness, rapid growth, easy propagation, small gestation period, wide adaptability and optimum plant size for convenient seed collection (Jones and Miller, 1991; Francis et al., 2005). Worldwide it is deemed a promising species for biodiesel production having a high non-edible seed oil content. Seeds contain 46-58% of oil on kernel weight and 30-40% on seed weight (Subramanian et al. 2005). In fact the potentially high yield of oil per unit land area in *J. curcas* is second only to oil palm (Fairless, 2007). Furthermore, the seed oil quality is suitable for production of biodiesel as it contains more than 75% unsaturated fatty acids (Biello, 2009). Apart from being a

potential biofuel crop *J. curcas* has multiple utilities. Preparations of all plant parts, including seeds, leaves and bark are used in traditional medicine and for veterinary purposes. The latex of *J. curcas* latex has also been reported to be an abortifacient and is efficacious in dropsy, sciatica and paralysis (Frienvis 2008). It contains an alkaloid similar to quinine in properties called “Jatrophine” which is believed to have anti-cancerous properties and used for external application on skin diseases and rheumatism (Thomas *et al.* 2008). Nath and Dutta (1992) demonstrated the wound-healing properties of ‘cur-cain’, a proteolytic enzyme isolated from the latex. Kone-Bamba *et al.* (1987) reported coagulating effects of the latex on blood plasma. The oil has a strong purgative action and is widely used for skin diseases and to soothe pain. A decoction of leaves is used against cough and as an antiseptic after birth. Extract from all plant parts have insecticidal properties (Grainage and Ahmed 1988).

Limited success has been achieved in worldwide introduction of *J. curcas* for varied purposes due to unreliable and low seed set and oil yields resulting in poor economic returns (Singh *et al.*, 2010). The Species is characterized by variable and unpredictable yield due to unidentified reasons (Ginwal *et al.*, 2004). Cultivation of *J. curcas* involves risk because it is a wild species and no varieties with desirable traits for specific growing conditions are available (Jongschaap *et al.*, 2007). The major constraint in achieving higher quality oil yield of *J. curcas* is lack of information about its genetic variability, oil composition and absence of suitable ideotypes for different cropping systems. Thus, the positive attributes of this plant are also not fully understood in terms of breeding and utilization (Fairless, 2007) which limits its large-scale cultivation and warrants the need for genetic improvement and breeding in the species.

J. curcas is a highly cross-pollinated species which is anticipated to contain wide genetic variability offering significant scope for selecting superior genotypes for enhanced productivity. The promise of this valuable crop can only be realized completely through development and release of high yielding variety/clones which needs sufficient information regarding the extent and pattern of genetic variation in *J. curcas* populations. Modern techniques have accelerated characterization of *J. curcas* germplasm at molecular level (Singh *et al.*, 2010; Xu *et al.*, 2012) and even whole genome sequencing is available (Sato *et al.*, 2010). However, information regarding the extent and pattern of genetic variation in *J. curcas* population is limited (Kaushik *et al.*, 2007; Rao *et al.*, 2008; Gairola *et al.*, 2011; Tripathi *et al.*, 2013; Brasileiro *et al.*, 2013). Overall, scanty information on genetic base, genetic makeup and the patterns of genetic diversity

of *J. curcasis* a bottleneck in development of varieties/ clones with superior traits of commerce. Thus, genetic diversity and relationships among 31 genotypes collected from wide geographical range in eastern India was studied employing Random Amplified Polymorphic DNA (RAPD) markers.

2. Material and Methods

2.1 Plant material

Germplasm of 31 genotypes of *J. curcasc* collected from different locations of Eastern India in the states of Bihar, Jharkhand, Odisha and West Bengal (Table 1) maintained as a germplasm garden at Institute of Forest Productivity, Ranchi (India) was utilized.

2.2 DNA isolation

Total genomic DNA was extracted from leaves using the CTAB method (Doyle and Doyle, 1987) with minor modifications. 0.5 g of leaves was weighed (mid-rib removed) for each sample and crushed in pre-cooled mortar with a pestle using liquid nitrogen. The crushed leaves were put in a tube and 1 ml of extraction buffer was added to each sample. 0.3ul of mercaptoethanol was then added and the tubes were incubated in hot water bath (“Sonar-Associated Scientific Technologies”) at 60⁰ C for 45 minutes with intermittent shaking every 5 minutes. Tubes were then removed from water bath, cooled and equal volume of chloroform isoamyl alcohol (24:1) was added. The contents were gently mixed and centrifuged in centrifuge unit (Hermle Labortechnik-Z216MK) at 12000 rpm at 4⁰C for 10 minutes. This formed three layers upper aqueous layer, middle protein layer and bottom chloroform layer of plant debris. The uppermost layer was carefully transferred to a fresh clean eppendorf tube. To the supernatant 3 µl of RNase was added and the tubes were incubated at 37⁰C for 30 minutes. Tubes were then cooled. Chloroform: isoamyl alcohol (24:1) in equal volume was again added and tubes were inverted gently about 30 times. Centrifugation was carried out at 12000 rpm for 10 minutes at 4⁰C. Upper layer was carefully transferred to a fresh tube. 2/3rd volume of cold isopropanol was then added to the above and was incubated for 10 minutes at room temperature to precipitate DNA. The tubes were then centrifuged at 12000 rpm at 4⁰C for 15 minutes. The supernatant was then discarded and the DNA pellet was washed with 70% ethanol for which ice cold ethanol was added and the tubes were spun at 6000rpm for 3 minutes. The process was repeated twice. Tubes were air dried and DNA was resuspended in 50 µl of TE buffer and stored at -20⁰C.

The yield and purity of DNA were measured using UV spectrophotometer (Spectroquant Pharo 300 Merck). The purity of DNA was determined by calculating the ratio of absorbance at 260 nm to that of 280 nm.

2.3 PCR amplifications

A set of 25 random decamer oligonucleotides (Bangalore Genei, India) were used as single primers for the amplification of RAPD fragments (Table 2). The G=C content varied from 60-70% while the melting temperature (T_m) was 32-34 °C.

The RAPD was carried out in thermocycler (Gene Amp-PCR System, Applied Biosystems). The cocktail of PCR reaction was prepared in a final volume of 18µl with 1 x PCR buffer, 1.5 mM MgCl₂, taq DNA polymerase 0.3 U, 0.2 mM dNTPs, 2 µM of primer, 40 ng template DNA. PCR reaction mixture was loaded into 0.2 ml thin walled clear PCR tubes. The RAPD-PCR was programmed as I-94⁰C for 5 minutes- Initial denaturation; II-94⁰C for 1 minute- Denaturation; III-36⁰C for 1 minute- Annealing; IV-72⁰C for 2 minutes- Extension; V-72⁰C for 5 minutes- final extension followed by 4⁰C for infinity to hold the samples. The steps II, III and IV were programmed to run for 45 cycles.

PCR fragments were separated electrophoretically on 1.5% agarose gel containing ethidium bromide using 1 x TAE buffer. The PCR products were visualized under UV light in a Gel Doc Unit (ULTRACAM). The banding pattern were photographed and stored as digital pictures (Fig. 1). The reproducibility of the amplification was confirmed by repeating experiment as required.

To analyze RAPD data, the total numbers of unique bands were counted for each primer used. The presence or absence of each individual band was recorded for each lane on the gel representing a different plant sample. For this a matrix was created with each sample representing one column and each band in one row. The presence of a band was recorded as a one (1) and the absence as a zero (0). Only clear, unambiguous and reproducible bands were considered for scoring. The RAPD bands obtained from different primers in the clones of *J. curcas* were scored in the form of binary data of 1 and 0 which were put in the form of data matrix. Recorded data was put in a specified format into software for further assessment.

The population genetic diversity parameters like percentage of polymorphic loci (Kimura and Crow, 1964), genetic diversity (Nei, 1973), Shannon's Information Index (Lewontin, 1972) and genetic distance (Nei, 1972) were calculated using Popgene version 1.31 (Yeh et al., 1999). The total genetic diversity (HT), mean genetic diversity within population (HS) and the relative degree of genetic diversity between populations (GST) were measured using Nei's genetic diversity (Nei, 1987).

3. Results and discussion

Information of existing genetic diversity constitutes the basis of genetic improvement programmes and sustainable management of genetic resources. It depends upon multiple factors such as breeding systems, biological characteristics, evolutionary history, and natural selection impart genetic diversity and (Hamrick and Godt 1996; Oyundelger et al., 2021). Since variability is a prerequisite for selection for improvement, it is indispensable to detect and document the amount of variation existing within and between the populations. Populations with low genetic variability have a reduced potential to adapt to environmental changes (Ellstrand and Elam, 1993). Therefore, genetic variation is important for the long term survival of a species (Falk and Holsinger, 1991). *J. curcas* being a naturalized species, its ability to grow in varied eco-climatic zones and its wide range of distribution embraces a considerable scope of genetic variation. Characterization of the available germplasm is needed for breeding programs to be efficient (King et al., 2015; Mastan et al., 2012). However, the degree of genetic diversity in and within natural populations in and outside the Centre of origin is poorly studied for the species, hampering domestication of *J. curcas* for large-scale cultivation.

High efficiency and low expense RAPD markers having the capability to scan and detect multiple loci across all the regions of genome have been found highly suited for genetic variation and phylogeny studies in tree species. In the present study RAPD provided large number of reliable and reproducible fingerprint profiles for a collection of 31 accessions of *J. curcas*. However, the availability of unique or rare fragments present in different accessions were limited. The details of RAPD primers, number of amplified products and percentage polymorphism obtained by analyzing primers have been provided in Table 3. 139 bands in total were observed. Of these 131 bands were polymorphic exhibiting a high polymorphism of 94.24%.

The genetic variation statistics for the loci have been summarized in the Table 4. The observed numbers of alleles (N_a) ranged from 1.80 to 2.00 with a mean value of 1.94 while the range of the effective numbers of alleles (N_e) was found to be 1.44 to 1.85 with a mean value of 1.64. The Nei's gene diversity (H) was 0.36 ± 0.15 and Shannon's information index (I) was recorded to be 0.53 ± 0.20 . Both values are comparatively higher than those reported for RAPD markers in *J. curcas* in earlier studies (Gupta et al. 2008; Raffi et al. 2012). This reflects the heterozygous nature of the population under study. The high value of Shannon's information index in the present study (>0.5) is an indication of the presence of high genetic diversity. Overall the markers were very informative in recording highly divergent genotypes of *J. curcas* from eastern India.

Similarity indices estimated on the basis of 19 primers ranged widely from 0.44 to 0.83 which suggests that these accessions represent genetically diverse population possibly due to predominance of cross pollination and seed source variability (Table 4). The highest genetic distance (0.79) was found between AK6 and JCRCKG that have come from West Bengal and Jharkhand, respectively while the lowest genetic distance (0.19) was there between JCGLBB and JCGLPD, both from Jharkhand. The high diversity revealed by RAPD markers is in agreement with the conclusion that out-breeding plant species retain considerable variability (Hamrick et al 1990). Still in absence of the pedigree data, the higher probability of origin of all such accessions from the same source and eventual distribution to other locations in a region/state cannot be ruled out.

To understand genetic diversity in the germplasm analyzed, genotyping data obtained for all the nineteen RAPD primer across 31 collected accessions were used to generate the UPGMA (Unweighted Pair Group Method with Arithmetic Mean) -based phylogenetic tree which shows two major clusters (Fig. 2). The accessions JBT-45/5 and JBT-45/13 were seen to be at two ends of the phylogenetic tree exhibiting high diversity. The cluster I consisted 16 genotypes from two states Jharkhand and Odisha (Chotanagpur and Eastern plateau region) while 15 accessions from West Bengal and Bihar (Indo Gangetic plains region) were grouped together in the Cluster II. This is clear alignment of accessions as per geographical regions possibly due to dissimilar growing conditions in different regions. Grouping according to geographical differences has been observed in other studies also (Gohil and Pandya 2008; Gupta et al., 2008; Senthilkumaret al., 2009). The Cluster I formed two sub-clusters with one of them

with accessions only from Jharkhand while the other sub-cluster had accessions both from Jharkhand and Orissa. Cluster II formed three sub-clusters among which one formed of only two accessions JBT-45/12 and JBT-45/13. The clusters seen in the dendrogram were also analyzed for their genetic distinctiveness. Coefficient of genetic differentiation (Nei 1973) which is the measure of diversity within groups in population genetics was calculated which was 1 indicating that the population in study is highly diverse.

The prior assessments of genetic variations among *Jatropha* germplasms using molecular markers show the presence of high genetic diversity for the Central and South American regions and insignificant genetic variation from Asia and Africa. The RAPD technique has been used for assessment of genetic diversity for *J. curcas* to revealed low to moderate level of genetic diversity of Indian germplasms (Basha and Sujatha, 2007; Reddy et al., 2007; Pamidiamarri et al., 2009; Basha et al., 2009). Several other molecular studies later employing variety of DNA markers have also underlined widespread genetic monomorphism and low genetic variability in *J. curcas* (Vásquez-Mayorga et al. 2017; Anggraeni et al. 2018; Gangapur et al. 2018; Carneiro et al. 2022). The possible reason may be absence of the anthropogenic and environmental influences in generating genetic variability mainly in the introduced conditions. It seems that adaptive genomic characters imparting tolerance of growth in wide disturbed scenarios have most probably been acquired by the species before its global distribution. Richards *et al.*, (2006) observed that a pronounced phenotypic plasticity is in itself a genotypic trait that allows the plant to respond to different environments through morphological and physiological changes for its survival. Further a limited stock has been vegetatively and apomictically propagated, since *J. curcas* is known to exhibit apomixis (Bhattacharya et al., 2005). Another reason behind low variability seems to be the introduced nature of *J. curcas* in countries (like India) where these molecular studies have been conducted with limited numbers of accessions in most of the cases.

Variation in genetic diversity within the species is usually related with geographical range, mode of reproduction, mating system, seed dispersal and fecundity. The genetic diversity in the present investigation might be due to geographical difference and seed source variability as well as high level of cross pollination in the species. Gupta et al (2008) have also recorded wide genetic base in RAPD evaluation of 13 Indian accessions from different agroclimatic zones. Likewise Rafii et al. (2012) found accessions from same and near states or regions to be grouped together according to their geographical origin. According to Pamidiamarri Reddy (2014),

Portuguese seafarers introduced accessions from Mexico and Central America to India through two dispersal routes: one brought *J. curcas* through Africa, Madagascar and finally to India, while the other passed through Spain on its way to India. This may also be a reason for geographical separation of the accessions in these studies.

Conclusion

Estimating genetic diversity is essential to ensure success in the management of genetic resources, planning and adoption of strategies for genetic breeding in promising species like *J. curcas*. RAPD markers with their high level of polymorphism and ability to decipher genetic variability and relationships appeared to be a viable system for undertaking such studies in *J. curcas*. In contrast to most studies we found ample genetic diversity among the analyzed accessions from four states of eastern India indicating that these accessions represent genetically diverse populations due to predominance of cross-pollination and seed source variability. A clear-cut clustering pattern of the accessions as per geographic regions was recorded which may possibly be due to different genetic makeup of the accessions as a result of evolution in markedly different growing conditions.

References

- Anggraeni T. D., Satyawan D., Kang Y. J., Ha J., Kim M.Y., Chitikineni A., Varshney R. K. and Lee S. (2018). Genetic diversity of *Jatropha curcas* collections from different islands in Indonesia. *Plant Genetic Resources*, 16(4), 334-342.
- Basha S. D. and Sujatha M. (2007). Inter and intra-population variability of *Jatropha curcas*(L.) characterized by RAPD and ISSR markers and development of population specific SCAR markers, *Euphytica*, 156: 375-386.
- Basha S. D., Francis G., Makkar H. P. S., Becker K. and Sujatha M. (2009). Comparative study of biochemical traits and molecular markers for assessment of genetic relationships

between *Jatropha curcas* L. germplasm from different countries, *Plant Science*, 176: 812-823.

Bhattacharya A., Datta K. and Datta S. K. (2005). Floral biology, floral resource constraints and pollination limitation in *Jatropha curcas* L. *Pakistan Journal of Biological Sciences* 8: 456-460.

Biello D. (2009). Green fuels for jets. *Sci. Am.* 19:68-69.

Brasileiro B.P., Silva S.A., Souza D.R., Santos P.A., Oliveira R.S. and Lyra D.H. (2013). Genetic diversity and selection gain in the physic nut (*Jatropha curcas*). *Genet. Mol. Res.* 12: 2341-2350.

Carneiro A., Santos A., Laviola B. G., Teodoro L.P.R., Teodoro P.E. and Rodrigues E.V. (2022). Genetic diversity and population structure in *Jatropha* (*Jatropha curcas* L.) based on molecular markers. *Genetic Resources and Crop Evolution*, 69, 245–254.

Doyle J.J. and Doyle J.L. (1990). Isolation of plant DNA from fresh tissue. *Focus*, 12: 13–15.

Ellstrand N. C. and Elam D. R. (1993). Population genetic consequences of small population size: implications for plant conservation. *Annual Review of Ecology and Systematics* 24: 217-242.

Fairless D. (2007). Biofuel: the little shrub that could maybe. *Nat.* 449: 652-655.

Falk D. A. and Holsinger K. E. (1991). *Genetics and Conservation of Rare Plants*. Oxford University Press, New York, 283 pp.

Francis G., Edinger R. and Becker K. (2005). A concept for simultaneous wasteland reclamation, fuel production and socio-economic development in degraded areas in India: need, potential and perspective of *Jatropha* plantations. *Nat. Resour. Forum* 29:12-24.

Frienvis (2008). ENVIS Centre of Forestry. Available online: <http://www.frienvis.nic.in>

Gairola K.C., Nautiyal A.R., Sharma G., Dwivedi A.K. (2011). Variability in Seed Characteristics of *Jatropha curcas* Linn. From Hill Region of Uttarakhand. *Bull. Env. Pharmacol. Life Sci.* 1:64-69.

- Gangapur D. R., Agarwal P. and Agarwal P.K. (2018). Molecular markers for genetic diversity studies in *Jatropha* (*Jatropha curcas* L.). *AIMS Environmental Science*, 5(5): 340-352.
- Ginwal HS, Rawat PS, Srivastava RL (2004). Seed source variation in growth performance and oil yield in *Jatropha curcas* Linn. in central India. *Silvae Genet.* 53:186-192.
- Gohil R. H. and Pandya J. B. (2008). Genetic diversity assessment in physic nut (*Jatropha curcas* L.). *International Journal of Plant Production* 2: 321-326.
- Grainge M. and Ahmad S. (1988). *Handbook of Plant and Pest-control Properties*. John Willey and Sons, New York, 470 pp.
- Gupta S., Srivastava M., Mishra G.P., Naik P.K. and Chauhan R. S. (2008). Analogy of ISSR and RAPD markers for comparative analysis of genetic diversity among different *Jatropha curcas* genotypes. *African Journal of Biotechnology* 7: 4230-4243.
- Hamrick J. L. (1990). Isozymes and analysis of genetic structure in plant populations. In: Soltis E. D. and Soltis P.S. (eds.), *Isozymes in Plant Biology*. Chapman and Hall, London, pp. 87-105.
- Hamrick J. L. and Godt M. J.W. (1996). Effects of life history traitson genetic diversity in plant species. *Philosophical Transactionsof the Royal Society B-Biological Sciences*, 351,1291–1298.
- Jones N. and Miller J.H. (1991). *Jatropha curcas*- a multipurpose species for problematic sites. *Land Resour. Series* 1:1-12.
- Jongschaap R.E.E., Corre W.J., Bindraban P.S. and Bradenburg W.S. (2007). Claim and Fact on *Jatropha curcas*L. Plant Research International B. V., Wageningen, The Netherlands.
- Kaushik N., Kumar K., Kumar S., Kaushik N. and Roy S. (2007). Genetic variability and divergence studies in seed traits and oil content of *Jatropha* (*Jatropha curcas*L.) accessions. *Biomass Bioener.* 31:497-502.
- Kimura M. and Crow J. F. (1964). The Number of alleles that can be maintained in a finite population. *Genetics* 49(4): 725-38.

- King A.J., Montes L.R., Clarke J.G., Itzep J., Perez C.A., Jongschaap R.E., Visser R.G., Van Loo E.N. and Graham I. (2015). Identification of QTL markers contributing to plant growth, oil yield and fatty acid composition in the oilseed crop *Jatropha curcas* L. *Biotechnology for Biofuels* 8:1-17.
- Kone- Bamba D., Pelissier Y., Ozoukou Z. F. and Ouap D. (1987). Etude de l'activitehemostatique de quinze plantesmedicinales de la "PharmacopeeTraditionnelleIvoirienne". *Plant Medicus Phytotherapie* 21: 12-130.
- Nath L. K. and Datta S. K. (1992). Wound healing response of the proteolytic enzyme curcain. *Indian Journal of Pharmacology* 24: 114-115.
- Lewontin R.C. (1972). The Apportionment of Human Diversity. In: Dobzhansky, T., Hecht, M.K., Steere, W.C. (eds.) *Evolutionary Biology*. Springer, New York, NY. https://doi.org/10.1007/978-1-4684-9063-3_14 pp 381–398
- Mastan S., Sudheer P., Rahman H, Ghosh A., Rathore M., Ravi Prakash C. and Chikara J. (2012). Molecular characterization of intra-population variability of *Jatropha curcas*L. using DNA based molecular markers. *Molecular Biology Reports* 39(4):4383-4390.
- Nei M. (1972). Genetic distance between populations. *American Naturalist*, 106, 283-292.
- Nei M. (1973). Analysis of gene diversity in subdivided populations. *Proc. Natl. Acad. Sci. USA*. 70(12):3321-3.
- Nei M. (1987). *Molecular Evolutionary Genetics*. Columbia University Press, New York.
- Oyundelger K., Herklotz V., Harpke D., Oyuntsetseg B. and Ritz C. M. (2021). Contrasting effects of local environment and grazing pressure on the genetic diversity and structure of *Artemisia frigida*. *Conservation Genetics*, 39(2), 364–375.
- Pamidimarri D. V. N. S., Pandya N., Reddy M. P. and Radhakrishnan T. (2009). Comparative study of interspecific genetic divergence and phylogenetic analysis of genus *Jatropha* by RAPD and AFLP. *Molecular Biology Reports*, 36: 901-907.

- Pamidimarri DVNS and Reddy MP (2014) Phylogeography and molecular diversity analysis of *Jatropha curcas* L. and the dispersal route revealed by RAPD, AFLP and nrDNA-ITS analysis. *Molecular Biology Reports* 41: 3225–3234.
- Rafii M.Y., Shabanmofrad M., Puteri-Edaroyati M.W., et al. (2012) Analysis of the genetic diversity of physic nut, *Jatropha curcas* L. accessions using RAPD markers. *Mol Biol Rep* 39: 6505–6511.
- Rao G. R., Korwar G. R., Shanker A. K. and Ramakrishna Y. S. (2008). Genetic associations, variability and diversity in seed characters, growth, reproductive phenology and yield in *Jatropha curcas* (L.) accessions. *Trees (Berl.)*, 22: 697-709.
- Reddy M. P., Chikara J., Patolia J. S. and Ghosh A. (2007). Genetic improvement of *J. curcas* adaptability and oil yield. In: Fact Seminar on *J. curcas* L., Agronomy and Genetics, 26-28 March 2007, Wageningen the Netherlands, Published by FACT Foundation.
- Richards C. L., Bossdorf O., Muth N.Z., Gurevitch J. and Pigliucci M. (2006). Jack of all trades, master of some? On the role of phenotypic plasticity in plant invasions. *Ecology Letters* 9: 81-993.
- Senthil Kumar R., Parthiban K.T., Hemlatha P., Kalaiselvi T. and Govinda Rao M. (2009). Investigation cross-compatibility barriers in the biofuel crop *Jatropha curcas* L. with wild *Jatropha* species. *Crop Sciences* 49: 1667-1774.
- Singh P., Sanjay Singh, Mishra S.P., Bhatia S.K. (2010). Molecular characterization of genetic diversity in *Jatropha curcas* L. *Gene, Genome Genomics* 4:1-8.
- Subramanian K. A., Singhal S. K., Saxena M. and Singhal S. (2005). Utilization of liquid biofuels in automotive diesel engines: An Indian perspective. *Biomass and Bioenergy* 29: 65-72.
- Thomas R., Sah N. K. and Sharma P. B. (2008). Therapeutic biology of *Jatropha curcas*: a mini review. *Current Pharmaceutical Biotechnology* 9: 315-324.

Tripathi A., Mishra D.K. and Shukla J.K. (2013). Genetic variability, heritability and genetic advance of growth and yield components of *Jatropha* (*Jatropha curcas*Linn.) genotypes. *Trees* 27:1049-1060.

Vásquez-Mayorga M., Fuchs E.J., Hernández E.J., Herrera F., Hernández J., Moreira I., Arnáez E. and Barboza N. M. (2017). Molecular characterization and genetic diversity of *Jatropha curcas* L. in Costa Rica. *PeerJ*. 9;5:e2931. doi: 10.7717/peerj.2931.

Xu W., Sujatha M. and Liu A. (2012). Genetic diversity in the *Jatropha* genus and its potential application. *CAB Reviews* No. 59. Available online <http://www.cabi.org/cabreviews>.

Yeh F., Boyle C.T., Ronagcai Y. and Xian J.M. (1999). Popgene version 3.1 (<http://www.vulberta.ca/~fyeh/fyah>).

Table 1: Details of *J. curcas* genotypes from different areas of eastern India.

S. No.	Accessions	Location	District	State
1	JBT-44/5	Chandwa	Latehar	Jharkhand
2	JBT-44/6	Satwarwa	Palamu	Jharkhand
3	JBT-44/15	Sonpurwa	Garhwa	Jharkhand
4	JBT-44/19	Tildag	Garhwa	Jharkhand
5	JBT-44/20	Chaenpur	Palamu	Jharkhand
6	JCRCKG	Burmu	Ranchi	Jharkhand
7	JCGLBB	Basia	Gumla	Jharkhand
8	JCGLPD	Dhauntatoli	Gumla	Jharkhand
9	JCGLSH	Hundratoli	Gumla	Jharkhand
10	JCBKDT	Dantu	Bokaro	Jharkhand

11	JCBKBD	Bardih	Bokaro	Jharkhand
12	JCSDFP	Kolebira	Simdega	Jharkhand
13	JCSDBK	Karanjtoli	Simdega	Jharkhand
14	JBT-45/2	Nandangarg	W. Champaran	Bihar
15	JBT-45/4	Santpur	E.Champaran	Bihar
16	JBT-45/6	Siwan	E.Champaran	Bihar
17	JBT-45/9	Begusarai	Begusarai	Bihar
18	JBT-45/10	Balia	Begusarai	Bihar
19	JBT-45/11	Mokama	Patna	Bihar
20	JBT-45/12	Nalanda	Biharsharif	Bihar
21	JBT-45/13	Rajauli	Nawada	Bihar
22	AK-2	Midnapur	Midnapur	W.Bengal
23	AK-3	Kusto Colony	Durgapur	W.Bengal
24	AK-5	Saltora	Bamkura	W.Bengal
25	AK-6	Jhalgram	Midnapur	W.Bengal
26	BNDAK-7	Matha	Purulia	W.Bengal
27	BNDAK-8	Kulgoda	Purulia	W.Bengal
28	AK-9	Pathalmada	Purulia	W.Bengal
29	OR-1	Rourkela	Sundergarh	Odisha
30	OR-2	Talacher	Angul	Odisha
31	OR-3	Gania	Nayagarh	Odisha

Table 2: Details of the RAPD primers used in the study

Primer	Nucleotide sequence (5'-3')	G+C content (%)	T _m °C
RAPD-1	AAAGCTGCGG	60	32
RAPD-2	AACGCGTCGG	70	34
RAPD-3	AAGCGACCTG	60	32
RAPD-4	AATCGCGCTG	60	32
RAPD-5	AATCGGGCTG	60	32
RAPD-6	ACACACGCTG	60	32
RAPD-7	ACATCGCCCA	60	32
RAPD-8	ACCACCCACC	70	34
RAPD-9	ACCGCCTATG	60	32
RAPD-10	ACGATGAGCG	60	32
RAPD-11	ACGGAAGTGG	60	32
RAPD-12	ACGGCAACCT	60	32
RAPD-13	ACGGCAAGGA	60	32
RAPD-14	ACTTCGCCAC	60	32
RAPD-15	ACCCTGAGCC	70	34
RAPD-16	AGGCGGCAAG	70	34
RAPD-17	AGGCGGGAAC	70	34
RAPD-18	AGGCTGTGTC	60	34
RAPD-19	AGGTGACCGT	60	32
RAPD-20	AGTCCGCCTC	70	34
RAPD-21	CACGAACCTC	60	32
RAPD-22	CATAGAGCGG	60	32

RAPD-23	CCAGCAGCTA	60	32
RAPD-24	CCAGCCGAAC	70	34
RAPD-25	GAGCGCCTTC	70	34

UNDER PEER REVIEW

Table 3:Primer identity with total number of bands obtained and polymorphism percentage.

Primer	Polymorphic bands	Monomorphic bands	Total bands	% Polymorphism
RAPD-3	7	0	7	100
RAPD-4	8	1	9	88.88
RAPD-6	6	0	6	100
RAPD-7	7	1	8	87.85
RAPD-8	6	0	6	100
RAPD-9	4	0	4	100
RAPD-10	12	1	13	92.3
RAPD-12	9	2	11	81.8
RAPD-13	7	0	7	100
RAPD-15	5	0	5	100
RAPD-16	6	0	6	100
RAPD-17	5	0	5	100
RAPD-18	4	1	5	80
RAPD-19	6	0	6	100
RAPD-21	11	2	13	84.6
RAPD-22	6	0	6	100
RAPD-23	8	0	8	100
RAPD-24	5	0	5	100
RAPD-25	9	0	9	100
Total	131	8	139	94.24

Table 4: Summary of genetic variation statistics for all loci

Locus	Sample Size	Na	Ne	H	I
RAPD-3	31	2.00	1.78	0.43	0.62
RAPD-4	31	2.00	1.56	0.34	0.52
RAPD-6	31	2.00	1.83	0.45	0.65
RAPD-7	31	2.00	1.73	0.40	0.57
RAPD-8	31	2.00	1.83	0.45	0.64
RAPD-9	31	2.00	1.85	0.46	0.65
RAPD-10	31	2.00	1.58	0.33	0.50
RAPD-12	31	2.00	1.63	0.35	0.51
RAPD-13	31	2.00	1.44	0.29	0.46
RAPD-15	31	2.00	1.63	0.38	0.57
RAPD-16	31	2.00	1.48	0.30	0.47
RAPD-17	31	1.80	1.60	0.31	0.44
RAPD-18	31	1.80	1.47	0.26	0.39
RAPD-19	31	1.83	1.54	0.33	0.48
RAPD-21	31	1.92	1.67	0.36	0.52
RAPD-22	31	2.00	1.63	0.36	0.53
RAPD-23	31	1.88	1.69	0.38	0.55
RAPD-24	31	1.80	1.62	0.34	0.49
RAPD-25	31	1.89	1.55	0.33	0.50
Mean	31	1.94	1.64	0.36	0.53
Standard Deviation		0.22	0.32	0.15	0.20

Table 5: Nei's analysis of genetic similarity (above diagonal) and genetic distance (below diagonal) in 31 accessions of *J. curcas*.

pop ID	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31
1	****	0.76	0.71	0.65	0.63	0.68	0.73	0.70	0.70	0.61	0.65	0.69	0.71	0.64	0.64	0.62	0.61	0.64	0.64	0.59	0.54	0.65	0.62	0.58	0.63	0.59	0.61	0.65	0.68	0.60	0.58
2	0.27	****	0.80	0.67	0.58	0.58	0.63	0.56	0.62	0.60	0.60	0.65	0.63	0.62	0.58	0.54	0.55	0.58	0.60	0.61	0.56	0.60	0.60	0.56	0.63	0.58	0.66	0.57	0.63	0.57	0.63
3	0.35	0.22	****	0.67	0.60	0.60	0.66	0.66	0.56	0.65	0.61	0.62	0.58	0.58	0.55	0.58	0.60	0.60	0.55	0.59	0.60	0.58	0.63	0.61	0.62	0.58	0.60	0.63	0.60	0.63	0.60
4	0.43	0.40	0.40	****	0.71	0.55	0.60	0.58	0.59	0.59	0.56	0.65	0.65	0.59	0.55	0.54	0.59	0.56	0.50	0.61	0.55	0.59	0.58	0.50	0.59	0.55	0.56	0.54	0.55	0.55	0.58
5	0.47	0.55	0.50	0.35	****	0.61	0.61	0.64	0.68	0.60	0.58	0.62	0.72	0.60	0.65	0.59	0.60	0.55	0.53	0.59	0.60	0.58	0.63	0.53	0.63	0.59	0.55	0.56	0.60	0.63	0.66
6	0.38	0.55	0.50	0.60	0.49	****	0.67	0.70	0.68	0.60	0.65	0.63	0.71	0.68	0.55	0.56	0.54	0.57	0.67	0.53	0.50	0.51	0.55	0.58	0.45	0.56	0.53	0.55	0.63	0.63	0.59
7	0.32	0.46	0.41	0.50	0.49	0.40	****	0.83	0.73	0.61	0.67	0.65	0.63	0.65	0.65	0.60	0.58	0.63	0.64	0.55	0.61	0.67	0.68	0.64	0.64	0.62	0.60	0.60	0.68	0.59	0.62
8	0.36	0.58	0.41	0.55	0.45	0.36	0.19	****	0.71	0.65	0.73	0.68	0.62	0.67	0.61	0.59	0.63	0.58	0.64	0.56	0.57	0.68	0.63	0.64	0.55	0.66	0.60	0.58	0.65	0.65	0.59
9	0.36	0.48	0.58	0.53	0.38	0.38	0.32	0.34	****	0.71	0.65	0.68	0.69	0.64	0.65	0.59	0.64	0.63	0.61	0.62	0.61	0.57	0.56	0.50	0.54	0.53	0.60	0.56	0.68	0.60	0.56
10	0.49	0.50	0.43	0.53	0.52	0.52	0.49	0.42	0.34	****	0.76	0.72	0.65	0.54	0.63	0.53	0.60	0.61	0.57	0.66	0.55	0.65	0.62	0.55	0.60	0.52	0.63	0.69	0.68	0.66	0.66
11	0.42	0.50	0.37	0.58	0.54	0.42	0.40	0.32	0.42	0.28	****	0.75	0.60	0.61	0.63	0.62	0.60	0.58	0.67	0.59	0.50	0.67	0.56	0.61	0.55	0.56	0.61	0.66	0.65	0.68	0.59
12	0.37	0.42	0.42	0.42	0.48	0.46	0.43	0.39	0.39	0.33	0.29	****	0.64	0.60	0.68	0.61	0.65	0.62	0.62	0.64	0.55	0.63	0.70	0.58	0.59	0.58	0.68	0.68	0.71	0.68	0.70
13	0.35	0.47	0.49	0.42	0.33	0.35	0.46	0.48	0.37	0.43	0.50	0.45	****	0.60	0.65	0.57	0.62	0.58	0.60	0.68	0.58	0.56	0.61	0.53	0.59	0.55	0.58	0.57	0.59	0.61	0.68
14	0.45	0.48	0.48	0.53	0.52	0.38	0.42	0.40	0.45	0.62	0.49	0.50	0.50	****	0.73	0.78	0.64	0.65	0.73	0.59	0.64	0.65	0.65	0.70	0.63	0.65	0.63	0.60	0.60	0.58	0.56
15	0.45	0.55	0.55	0.60	0.42	0.59	0.42	0.49	0.42	0.47	0.47	0.39	0.43	0.32	****	0.81	0.71	0.67	0.71	0.65	0.67	0.70	0.73	0.71	0.76	0.65	0.68	0.69	0.67	0.65	0.66
16	0.48	0.62	0.54	0.62	0.53	0.58	0.50	0.53	0.53	0.63	0.48	0.49	0.57	0.25	0.22	****	0.71	0.63	0.75	0.60	0.65	0.71	0.67	0.69	0.66	0.65	0.60	0.65	0.56	0.63	0.57
17	0.49	0.60	0.60	0.53	0.52	0.62	0.54	0.47	0.45	0.52	0.52	0.43	0.48	0.45	0.34	0.35	****	0.74	0.73	0.63	0.64	0.67	0.72	0.64	0.65	0.60	0.68	0.65	0.58	0.63	0.58
18	0.45	0.55	0.55	0.58	0.59	0.57	0.47	0.54	0.47	0.49	0.54	0.48	0.55	0.42	0.40	0.46	0.30	****	0.80	0.68	0.71	0.64	0.71	0.67	0.65	0.58	0.63	0.69	0.63	0.60	0.56
19	0.45	0.50	0.50	0.69	0.64	0.40	0.45	0.45	0.49	0.57	0.40	0.48	0.50	0.32	0.34	0.29	0.32	0.22	****	0.66	0.67	0.67	0.72	0.70	0.67	0.66	0.67	0.71	0.61	0.60	0.63
20	0.53	0.49	0.52	0.49	0.53	0.63	0.60	0.58	0.48	0.41	0.53	0.45	0.38	0.53	0.43	0.52	0.46	0.39	0.41	****	0.78	0.60	0.61	0.58	0.68	0.60	0.62	0.64	0.58	0.54	0.63
21	0.62	0.58	0.60	0.60	0.52	0.70	0.49	0.57	0.49	0.59	0.70	0.60	0.55	0.45	0.40	0.43	0.45	0.34	0.40	0.25	****	0.67	0.65	0.60	0.70	0.62	0.61	0.63	0.55	0.56	0.59
22	0.42	0.50	0.53	0.53	0.54	0.67	0.40	0.38	0.57	0.42	0.40	0.46	0.58	0.42	0.36	0.35	0.40	0.45	0.40	0.50	0.40	****	0.73	0.67	0.71	0.60	0.67	0.72	0.63	0.65	0.63
23	0.48	0.52	0.52	0.54	0.46	0.60	0.39	0.46	0.58	0.48	0.58	0.36	0.49	0.43	0.31	0.40	0.33	0.35	0.33	0.49	0.43	0.31	****	0.72	0.76	0.67	0.73	0.74	0.62	0.63	0.68
24	0.54	0.58	0.55	0.69	0.64	0.54	0.45	0.45	0.70	0.59	0.49	0.55	0.63	0.36	0.34	0.37	0.45	0.40	0.36	0.55	0.52	0.40	0.33	****	0.73	0.72	0.64	0.68	0.58	0.60	0.59
25	0.47	0.46	0.46	0.53	0.47	0.79	0.45	0.59	0.62	0.52	0.59	0.53	0.53	0.47	0.28	0.41	0.42	0.42	0.40	0.39	0.36	0.34	0.27	0.32	****	0.71	0.71	0.75	0.58	0.62	0.68
26	0.53	0.54	0.49	0.59	0.53	0.58	0.48	0.41	0.63	0.66	0.58	0.54	0.59	0.43	0.43	0.42	0.50	0.55	0.41	0.52	0.48	0.50	0.40	0.33	0.35	****	0.69	0.65	0.60	0.64	0.70
27	0.49	0.41	0.48	0.58	0.59	0.64	0.52	0.52	0.52	0.47	0.49	0.39	0.55	0.47	0.38	0.50	0.38	0.47	0.40	0.48	0.49	0.40	0.31	0.45	0.34	0.37	****	0.78	0.73	0.66	0.73
28	0.43	0.57	0.54	0.62	0.58	0.60	0.50	0.55	0.58	0.37	0.41	0.38	0.57	0.50	0.37	0.42	0.43	0.37	0.35	0.45	0.46	0.33	0.30	0.39	0.29	0.42	0.25	****	0.69	0.65	0.71
29	0.38	0.46	0.50	0.60	0.52	0.47	0.38	0.42	0.38	0.38	0.42	0.35	0.53	0.52	0.40	0.58	0.54	0.47	0.49	0.55	0.59	0.47	0.48	0.54	0.54	0.50	0.32	0.37	****	0.75	0.69

30	0.50	0.57	0.47	0.59	0.46	0.46	0.53	0.43	0.50	0.41	0.39	0.38	0.49	0.55	0.43	0.47	0.46	0.50	0.50	0.62	0.58	0.43	0.47	0.50	0.48	0.45	0.41	0.42	0.29	****	0.73
31	0.55	0.47	0.52	0.54	0.41	0.53	0.48	0.53	0.58	0.41	0.53	0.36	0.38	0.58	0.41	0.57	0.55	0.58	0.46	0.47	0.53	0.46	0.38	0.53	0.39	0.36	0.31	0.34	0.37	0.32	****

UNDER PEER REVIEW




Figure 1. RAPD patterns of 31 accession of *Moringa oleifera* generated by primer RPI 12 (above) and RPI 19 (below): M – 100 bp molecular weight ladder.

UNDER PEER REVIEW




Figure 2. Cluster generated from the Jaccard similarity coefficient and UPGMA clustering using RAPD markers between 31 *J. curcas* accessions.