

***Original Research Article***

**VIRAL LOAD PATTERN OF HIV POSITIVE CHILDREN AND ADOLESCENTS ATTENDING A PAEDIATRIC INFECTIOUS DISEASES CLINIC IN OWERRI, NIGERIA.**

**Abstract.**

**Background.**

Viral load (VL) tests measure the number of copies of HIV RNA in one milliliter of blood and is the preferred antiretroviral therapy (ART) monitoring strategy. Viral suppression is defined as patients having  $\leq 1000$  copies of viral RNA/ml of blood.

**Method/material.**

This study involved the retrospective review of viral load test of patients aged 2-19 years living with HIV/AIDS and enrolled in the treatment programme of the Paediatric infectious diseases clinic of the Federal University Teaching hospital Owerri .

**Results.**

117 patients met the inclusion criteria and 103 were virally suppressed giving a suppression rate of 88%. Majority of virally suppressed were in the 15-19 age group. A logistic regression analysis to ascertain the impact of age, gender, drug regimen, and duration on ARTs on the likelihood of viral suppression showed that age, gender, and drug regimen were not significant predictors of viral suppression ( $p > 0.05$ ). However, duration on ARTs was a significant predictor ( $p < 0.01$ ), with patients who had been on ARTs for greater than 24 months having higher odds of viral suppression (OR = 5.42, 95% CI = 1.84-15.93) compared to those who had been on ARTs for 0-6 months.

**Conclusion.**

Our study provides important insights into the viral load pattern of children and adolescents living with HIV/AIDS in Owerri. It highlights the importance of early initiation of ART, adherence to ART regimen, and routine viral load monitoring to achieve and maintain viral suppression.

**Key words.** Viral load, Viral suppression, Children, Adolescents.

## **Introduction.**

HIV infection remains a disease of public health importance and a significant cause of mortality and morbidity in Low - and Middle-Income countries (LMIC) of the world.<sup>1</sup> Scaling up of HIV care and treatment programme in Nigeria improved access to Antiretroviral Therapy (ART) for more children and adolescents.<sup>2</sup> ART helps suppress viral replication within the patient as well as strengthening the immune system but it does not as yet offer a cure. This viral suppression also strengthens the capacity of the immune system to protect against opportunistic infections.<sup>1</sup>

Evidence suggests that early initiation of ART reduces HIV related mortality, morbidity and overall HIV transmission.<sup>3</sup> The UNAIDS “test and start” strategy recommends that all people living with HIV be provided with lifelong ART including children and regardless of clinical status or CD4 cell count.<sup>1</sup> Additionally the World Health Organization recommends viral load testing as the preferred antiretroviral therapy (ART) monitoring strategy.<sup>4</sup>

Viral load (VL) tests measure the number of copies of HIV RNA in one milliliter of blood . It provides a direct measure of the efficacy of antiretroviral therapy (ART), predicting immunological and clinical outcomes and the risk of transmission.<sup>4</sup> Ideally , VL testing should be carried out 6 and 12 months after ART initiation and routinely every year thereafter.<sup>1</sup> A VL count of  $\leq 1000$  copies of viral RNA/ml of blood is considered viral suppression. VL needs to be reduced below 1000–1500 copies /mL for the likelihood of transmission to approach zero.<sup>5</sup>

Nigeria accounts for 26.7% of the Paediatric HIV/AIDS globally and it is estimated that only 35% of children living with HIV in Nigeria are receiving ART with viral load monitoring as a means of assessing outcome.<sup>6,7</sup> The national viral load implementation strategy and plan was launched in 2016<sup>8</sup>, and the reported national viral load suppression rate in children is 34%.<sup>9</sup>

It is of utmost importance to determine viral load status or virological suppression status of children enrolled on ART as it allows for timely detection of treatment failures, identification of patients in need of more intensive adherence counseling and support and minimizes development of drug resistance.<sup>10</sup>

This study set out to document viral load patterns in children and adolescents living with HIV/AIDS and attending the Paediatric infectious disease clinic of Federal University Teaching hospital Owerri, Imo State.

### **Materials/methods.**

The study involved the review of medical records of patients aged 2-19 years living with HIV and receiving treatment at the Paediatric infectious diseases clinic of the Federal University Teaching hospital Owerri and who have had a viral load test within the 6 months period of January to June 2022. These medical records are electronically stored in the clinic using the Nigeria Medical Record System (NMRS) Version 1.6.2.2.

Deidentified data was extracted from NMRS and information on current age, gender, duration on ART, ART Regimen, and the current Viral load were obtained. Virological suppression is defined as clients having  $\leq 1000$  copies of viral RNA/ml of blood, in line with the National Guideline for HIV Prevention, Treatment and care 2020.

Data was analyzed using R Statistical Package, descriptive findings were presented in tables, Chi Square was used to test for association. . A p-value of less than 0.05 was considered statistically significant.

### **Ethical considerations.**

Permission to use the data was sought and received from management of the Federal University Teaching hospital Owerri through the supervising authority over the clinic. The data was not shared with any third party.

## Result.

117 patients met the inclusion criteria, 55 were females and 62 males; giving a ratio of 1:1.12, The median age of patients was 15 years.

**Table 1** shows the general characteristics of the patients. 53 (45%) of patients were in the 15-19 age group and this group also had the highest number of female 29 (55%). The 10-14 age group had the highest number of male patients 41 (63%).

It also shows that all the patients were on Dolutegravir based regimens The most widely used drug combinations was TDF/3TC/DTG drug regimen with 84 (71.7%) of patients on it.

**Table 1. General Characteristics of Patients.**

Variable	Frequency	Proportion
<b>Gender</b>		
Male	62	53.0
Female	55	47.0
<b>Age Band (years)</b>		
1 – 4	7	6.0
5 – 9	16	13.7
10 – 14	41	35.0
15 – 19	53	45.3
<b>ART Regimen</b>		
ABC/3TC/DTG	33	28.2
TDF/3TC/DTG	84	71.8

## Abbreviations.

TLD. Tenofovir. ABC. Abacavir. DTG. Dolutegravir. 3TC. Lamivudine

**Table2. Pattern of Viral Load amongst patients.**

Variable	Frequency (%)		OR	p-value
	Suppressed	Unsuppressed		
<b>Gender</b>				
Male	56 (54.4)	6 (42.9)	1.13 (0.51 -2.49)	0.914
Female	47 (45.6)	8 (57.1)		
<b>Age Band (years)</b>				
1 – 4	7 (6.8)	0		
5 – 9	15 (14.6)	1 (7.1)	0.96 (0.92-1.00)	
10 – 14	33 (32.0)	8 (57.1)		
15 – 19	48 (46.6)	5 (35.7)		
<b>ART Regimen</b>				
ABC/3TC/DTG	28 (27.2)	5 (35.7)	2.45 (1.02-5.88)	0.004
TDF/3TC/DTG	75 (72.8)	9 (64.3)		
<b>Duration on ART</b>				
0 – 6	8 (7.8)			
7 – 24	19 (18.4)			0.001
>24	76 (73.8)			

**Table 2** shows that amongst the total 117 patients, 103 were virally suppressed giving a rate of 88%. Majority of the virally suppressed patients were in the 15-19 age group, with an almost equal distribution between males and females. All the patients between in the 1-4 years age group were virally suppressed while only 33(80%) of those in the 10-14 years group were virally suppressed.

The TDF/3TC/DTG drug regimen had a higher proportion of virally suppressed patients. 76 % of patients who had been on ARTs for greater than 24 months were also virally suppressed.

More male subjects were virally suppressed but there was no statistically significant difference in viral suppression rates between males and females. (P value =0.9142). There was a statistically significant relationship between duration on ART and viral suppression as well as between drug regimen and viral suppression, with P-values of 0.001 and 0.004 respectively.

Logistic regression analysis showed that patients on TDF/3TC/DTG based regimen were more likely to be virally suppressed when compared to those on ABC/3TC/DTG (AOR: 2.45; 95%CI: 1.02–5.88).

## **Discussion.**

In this study, we retrospectively reviewed the viral load pattern of children and adolescents attending the Paediatric infectious diseases clinic of the Federal University Teaching hospital Owerri. Our study found that the majority of the patients, 103 (88%) had viral load levels of 1000 copies Viral RNA/ml of blood and below, indicating viral suppression. This is consistent with the goal of ART which is to suppress viral replication and reduce morbidity and mortality associated with HIV infection.

This viral suppression rate is higher than 48.7% reported by Elon *et al*<sup>11</sup> following a review of viral suppression rate in children and adolescents enrolled in the national ART programme in four states in Northern Nigeria and 34% in a Nigerian country wide review.<sup>12</sup> It is also higher than national viral suppression rates in children in Uganda and Cameroun.<sup>13,14</sup>

It involves ramping up health care services to identify 95% of all people living with HIV (PLWH) through testing, providing antiretroviral therapy (ART) for 95% of those who are diagnosed, and achieving viral suppression of 95% of those in treatment by 2030.<sup>1</sup> We can only intensify our efforts to reach this milestone.

Majority of the patients were between 10-19 years old (71.8%). However, there were more females (55, 47.0%) than males (62, 53.0%) in the study population, suggesting that HIV/AIDS is affecting girls and young women disproportionately. This is corroborated by an early study in Northern Nigeria.<sup>11</sup>

The study also found that the most commonly prescribed ART regimen was TDF/3TC/DTG (72.0%), followed by ABC/3TC/DTG (28.0%). Although both regimens are recommended by the National guidelines for the treatment and control of HIV/AIDS in Nigeria, the

TDF/3TC/DTG regimen has been shown to have better efficacy and tolerability compared to other regimens, and it is thus encouraging to see that it is the most commonly used regimen in our study population. <sup>15</sup>

While 95% of patients between 1-9 years were virally suppressed only 86% of those 10-19 years were. However, it is concerning to note that 92.9% of those who were not virally suppressed were between 10-19 years old, suggesting that adherence to treatment may be more challenging in this age group. This agrees with the report by Elon *et al* that adolescents had lower rates of viral suppression than children 0 - 9 years. Similar findings were also documented in Cameroon by Fokam *et al* <sup>16</sup>. It is crucial to identify and address the factors contributing to treatment failure, such as poor adherence or drug resistance, to improve the clinical outcomes of these patients.

The relationship between gender and viral load suppression was not statistically significant. This agrees with early findings in Northern Nigeria. <sup>11</sup> We found a significant association between viral suppression and age, Viral suppression and duration on ART as well as viral suppression and drug regimen. Other studies have also documented the impact of age on viral load suppression. Elo *et al* <sup>11</sup> found that Adolescents had lower rates of viral suppression than children 0 - 9 years. Similar findings were documented by Fokam *et al* <sup>16</sup>

A logistic regression analysis to investigate the impact of age, gender, drug regimen, and duration on ARTs on the likelihood of viral suppression showed that age, gender, and drug regimen were not significant predictors of viral suppression ( $p > 0.05$ ). However, duration on ARTs was a significant predictor ( $p < 0.01$ ), with patients who had been on ARTs for greater than 24 months having higher odds of viral suppression (OR = 5.42, 95% CI = 1.84-15.93) compared to those who had been on ARTs for 0-6 months.

In conclusion, our study provides important insights into the viral load patterns of HIV-infected children and adolescents attending the Paediatric infectious diseases clinic of the Federal University Teaching hospital Owerri. It highlights the importance of early initiation of ART, adherence to ART regimen, and routine viral load monitoring to achieve and maintain viral suppression.

These findings suggest that targeted interventions may be needed to improve adherence to ART, especially among older patients. Targeted interventions such as peer support groups, youth-friendly services, and involvement of caregivers in adolescent care could be implemented to improve adherence to ART.

### **Limitation.**

We only assessed viral load status within a six-month period, limiting its generalizability. The study population were patients attending a single clinic in a specific region in Nigeria, which may not fully represent the general population.

### **References.**

1. <https://www.who.int/news-room/fact-sheets/detail/hiv-aids>. Accessed May 2023
2. <https://www.childrenandaids.org/sites/default/files/2017-05/Nigeria-Integrated-National-Guidelines-For-HIV-Prevention-treatment-and-care-2014.pdf>. Accessed May 2023
3. CLINICAL GUIDELINES: ANTIRETROVIRAL THERAPY <https://www.who.int/hiv/pub/arv/chapter4.pdf>. Accessed May 2023
4. World Health Organization (2013) Consolidated Guidelines on the Use of Antiretroviral Drugs for Treating and Preventing HIV Infection: Recommendations for a Public Health Approach. Geneva.
5. Rawizza, H.E., Chaplin, B., Meloni, S.T., et al. (2011) Immunologic Criteria Are Poor Predictors of Virologic Outcome: Implications for HIV Treatment Monitoring in Resource-Limited Settings. *Clinical Infectious Diseases*, 53, 1283-1290. <https://doi.org/10.1093/cid/cir729>
6. UNAIDS Nigeria (2019) Start Free, Stay Free, AIDS Free: 2018 Progress Report.
7. UNAIDS. Nigeria. HIV and AIDS Estimates 2019 Country Fact Sheet.

8. Federal Ministry of Health (2016) National Implementation Plan for Scale-Up of Viral Load Testing in Nigeria.
9. Nigeria HIV/AIDS Indicator and Impact Survey (2019) National Summary Sheet Preliminary Findings.
10. World Health Organization (2015) Antiretroviral Therapy (ART) Coverage among All Age Groups. [http://www.who.int/gho/hiv/epidemic\\_response/ART\\_text/en/](http://www.who.int/gho/hiv/epidemic_response/ART_text/en/). Accessed May 2023
11. Isaac, E.W., Ajani, A., Iliya, J., Christianah, O. and Hassan, D.M. HIV Viral Suppression in Children in a Subnational Antiretroviral Treatment Programme in Nigeria. *World Journal of AIDS* 2020;10; 170-185.
12. UNAIDS Nigeria (2019) Start Free, Stay Free, AIDS Free: 2018 Progress Report
13. Republic of Uganda (2016) Central Public Health Laboratories Viral Load Dashboard.
14. Tchouwa, G.T., Duverny, S.E., Cournil, A., et al. Nationwide Estimates of Viral Load Suppression and Acquired HIV Drug Resistance in Cameroon. *E-Clinical Medicine*.2018;1: 21-27.
15. Federal Ministry of Health . 2020. National Guidelines For HIV Prevention, Treatment and Care.
16. Fokam, J., Sosso, S.M., Yagai, B., Billong, S.C., DjubgangMbadie, R.E., Kamgaing Simo, R., et al. (2019) Viral Suppression in Adults, Adolescents and Children Receiving Antiretroviral Therapy in Cameroon: Adolescents at High Risk of Virological Failure in the Era of “Test and Treat”. *AIDS Research and Therapy*, 16, 36. <https://doi.org/10.1186/s12981-019-0252-0>