

Effect of organic and inorganic manures on growth and yield of okra

(*Abelmoschus esculentus* L. moench) cv. Arka Anamika

Abstract

The field experiment was conducted at the Horticulture Research CRC Farm – 1 of the Department of Horticulture, School of Agriculture, ITM University Gwalior (M.P.) during the year 2022 to study the effect of organic and inorganic manure on growth, yield and quality of okra. The experiment was laid out in the randomized block design with three replications and twelve treatments viz. FYM, Vermicompost, FYM + Vermicompost, 50% RDF + FYM, 50% RDF + Vermicompost, 50% RDF + FYM + Vermicompost, 75% RDF + FYM, 75% RDF + Vermicompost, 75% RDF + FYM + Vermicompost, 100% RDF + FYM, 100% RDF + Vermicompost, 100% RDF + FYM + Vermicompost. A single standard variety (Arka Anamika) is sown with the different treatment combination with the plant to plant and row to row spacing maintained at 45 cm x 60 cm respectively. On the basis of the study, the results indicated that the application of 100% RDF + FYM + Vermicompost in Arka Anamika recorded significantly higher plant height (80.20 cm), number of branches per plant (8.58), number of leaves per plant at 90 DAS (68.83). However, higher yield (5.89 q/ha) was recorded with the application of the 100% RDF + FYM + Vermicompost. Thus, integrated application of organic and inorganic manures improves the growth and yield of okra crop.

Keywords: Okra, Growth, Yield, Quality, Organic, Inorganic, Manures.

Introduction:

“Okra (*Abelmoschus esculentus* L. moench) is a vegetable crop growing in tropical and subtropical regions of the world that is significant economically. It is generally an annual plant which belongs to genus *Abelmoschus* and species *esculentus* of family *Malvaceae*. It is also known as lady's finger or bhindi, originated in tropical Africa”. [15] Okra is one of the most well-liked vegetables among all demographic groups due to its abundance in nutrients, flavour, therapeutic benefits, and industrial use. It is cultivated for its fibres, fruits or pods. The immature fruits are collected and consumed as vegetables. Okra is propagated by seeds and has a duration of 90-100 days. Okra needs a protected, warm, and humid growing season. Okra is a good source of vitamins, minerals, calories and amino acid found in seeds and compares favourably with those in poultry, eggs and soybean, (Chattopadhyay *et al.*, 2011) all parts of okra like fresh leaves, flowers, pods, stem and seeds can be used for different purposes and hence it is a multi-purpose crop in terms of its use. (Akbasova *et al.*, 2015). Okra's

mucilage can be used as a blood thinner or to substitute plasma (Vennila and Jayanthi, 2008)

“The demand of the crop is significantly increasing with the increase in population, which emphasizes the use of chemical fertilizers; as a result growers indiscriminately use the inorganic sources of plant nutrients”. [15] “Moreover, the use of expensive commercial fertilizers as per the requirement of the crop is not much affordable to the average farmers. The application of high input technologies such as chemical fertilizers, pesticides, herbicides improved the production but there is growing concern over the adverse effects of the use of chemicals on soil productivity and environment quality. So there is need of shifting towards INM approach. Integrated Nutrient Management (INM) is a sustainable approach, which aims at maintaining the soil fertility and plant nutrient supply, by incorporating all the possible sources of nutrients like organic manures, inorganic manures in an integrated and judicious manner to get higher crop yield without hampering the soil health and the environment”. (Dick and Gregorich, 2009). Considering all these aspects, a research study was carried out to study the effect of organic and inorganic manures on growth, yield and quality attributes of okra.

Materials and methods

An experiment was conducted at horticulture research CRC Farm-1, Department of horticulture, School of agriculture, ITM University Gwalior (M.P.) during March to June of 2022. The experimental site is located situated at 26°23' N latitude and 74°11' E longitude at an elevation of 211.52 m above mean sea level falling in the sub-tropical region of India. The climate of this place is bestowed with hot and dry early summers followed by hot and humid monsoon season and cold and dry winters. The soil of the experimental field was sandy clay loam in texture, slightly alkaline (pH 7.73) in reaction, low in organic carbon (4.3 g/kg) and available nitrogen (196.6 kg/ha) but medium in available phosphorus (15.85 kg/ha) and potassium (229.6 kg/ha) with electrical conductivity in the safe range. The experiment comprises of twelve treatments were replicated three times in randomised block design. The treatment details involve FYM, Vermicompost, FYM+Vermicompost, 50% RDF+FYM, 50

%RDF+Vermicompost,50%RDF+FYM+Vermicompost,75%RDF+FYM),75%RDF + Vermicompost, 75% RDF +FYM +Vermicompost, 100% RDF +FYM), 100% RDF +Vermicompost, 100 % RDF +FYM + Vermicompost. The row to row and plant to plant distance of the experiment was maintained respectively 60cm and 45cm. The land was brought to a fine tilth through ploughing and tillage. Irrigation channels and bunds were maintained properly. Direct sowing of the seeds was done in the field. Light irrigation was given after sowing. The organic manures were applied as basal dose before sowing, for proper decomposition, full dose of phosphorous and potassium and half dose of nitrogen as pre-treatment were applied just before the sowing. 30 days after sowing, the remaining half of the nitrogen dose was applied. All cultural practices were followed regularly during crop growth and observations were recorded on vegetative characters *i.e.*, plant height, number of branches per plant, number of nodes per plant, internodal length. Flowering, fruiting and yield characters *i.e.*, days to first flowering, fruit weight, fruit length, fruit yield per plant, fruit yield per hectare and Analysis of variance was performed to determine the effect of organic and inorganic manures on growth and yield of Okra using Opstat. The interpretation of treatment effects was made on the basis of critical difference at 5% probability level.

Result and Discussion

Growth attributes

The results revealed that growth parameters of Okra were significantly influenced by integrated application of organic and inorganic manures (Table 1). Okra plants fertilized with 100% RDF+FYM+Vermicompost gave maximum plant height (80.20cm) followed by 75% RDF + FYM + Vermicompost (77.58 cm), FYM + Vermicompost (60.25 cm) and Vermicompost (57.70). “This may be due to the reason that application of nutrients in integration of organic and inorganic manures have created favourable conditions for proper growth of crop. The better efficiency of organic manures is due to the fact that they might have provided both macro and micro nutrients at their optimal levels which ultimately enhanced the early growth phases and encouraged reproductive growth as well”. (Rathava *et al.*, 2018). Further it is revealed that a full dose of organic and inorganic fertilisers outperforms a single application of fertiliser. Similar results were reported by Khetran *et al.*, 2017)

Table.1 Effect of organic and inorganic manures growth attributes on okra

Treatment	Plant height (cm)	Number of branches	Number of leaves	Yield per plot (q/ha)	Yield (q/ha)
T ₁	56.75	4.41	63.00	0.25	3.41
T ₂	57.70	4.83	64.16	0.27	3.66
T ₃	60.25	5.41	65.16	0.29	3.93
T ₄	64.20	5.91	62.08	0.29	3.88
T ₅	69.37	6.08	64.00	0.33	4.41
T ₆	65.75	6.16	63.75	0.36	4.79
T ₇	70.87	6.58	64.25	0.44	5.89
T ₈	73.87	7.00	64.91	0.40	0.25
T ₉	77.58	7.91	64.58	0.47	0.27
T ₁₀	75.66	7.50	66.08	0.62	0.29
T ₁₁	78.16	8.25	66.58	0.60	0.29
T ₁₂	80.20	8.58	68.83	0.66	0.33
SE(m)±	2.12	0.07	0.78	0.002	0.344
CD at 5%	2.46	0.44	1.50	0.074	0.993

Maximum number of branches was recorded with the application of 100% RDF + FYM + Vermicompost and 75% RDF + FYM + Vermicompost. Further, maximum number of leaves at 90 days was observed with the application of 100% RDF + FYM + Vermicompost (68.83) followed by 75% RDF + FYM + Vermicompost (64.58) respectively. This may be due to the reason

that increased dose of NPK and FYM resulted in better improvement, establishment and availability of nutrients that resulted in better growth and yield attributes of okra. These results were similar to the findings of Yadav *et al.*, 2017 and Bamboriya *et al.*, 2008)

Highest yield per plot, yield per hectare was recorded with the application of 100% RDF + FYM + Vermicompost. This may be due to availability of organic and inorganic manures which resulted in increased yield attributes of okra. Further, integrated application of FYM, Poultry manure and vermicompost application improves the yield as compared to single manure application. (Bairwa *et al.*, 2009, Sharma *et al.*, 2021, Bodamwal and Rajput, 2006).

Conclusion:

It is concluded that the application of the integrated application of organic and inorganic manures significantly influenced the different growth and yield in okra.

The application of 100 % RDF + FYM + Vermi compost recorded highest plant height, number of branches, number leaves, fruit yield per plot and fruit yield per hectare in okra. Thus, integrated application of organic and inorganic manures improves the yield of okra crop.

References:

1. Akbasova, A.D., Sainova, Aimbetova, G., Makesova, I.O. and Sunakbaeva, M. 2015 Vermicompost's effect on agricultural crop productivity. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, **6**(4):2084.
2. Bairwa, H.L., Mahawer, L.N., Shukla, A.K., Kaushik, R.A. and Mathur, S.R. 2009. Response of integrated nutrient management on growth, yield and quality of okra. *Indian Journal of Agricultural Sciences*, **79**(5):381-384.
3. Bamboriya, J.S., Naga, S.R., Sharma, S.R., Choudhary, M.R. and Bamboriya, S.D. 2018. Okra (*Abelmoschus esculentus*) production, quality, and profitability as affected by organic manures and biofertilizers. *International Journal of Bio-resource and Stress Management*, **9**(4):506-509.
4. Bodamwal, S.G. and Rajput, S.G. 2006. Influence of organic and inorganic fertilizer on seed quality and seed yield of okra Parbhani Kranti. *Journal of Maharashtra Agricultural University*, **31**(1):130-131.
5. Chattopadhyay, A., Dutta, S. and Chatterjee, S. 2011. Seed yield and quality of okra as influenced by sowing dates. *African Journal of Biotechnology*, **10**(28):5461-5467.
6. Chauhan, D.V.S. 1972. *Vegetables production in India*, 3rd Ed., Ram Prasad and Sons, Agra.
7. Dick, W.A. and Greorich, E.G. 2009. Developing and Maintaining soil organic matter levels in managing soil quality; challenges in mrdenscjoing. P Elmbolt. S and Christensen BT (EDS). CAB International, Wallingford, UK. Pp 103-120.
8. Khetran, R., Muhammad, A.K., Ali, J.Z.S., Basharat, M., Fahmid, S., Ali, N., Raza, A., Ali, A. and Raza, Q. 2017. Effect of different doses of NPK fertilizer on seed yield of okra (*Abelmoschus esculentus* L. Moench). **4**(1).
9. Naik, H. M., Rao, G.B. and Srihari G. 2012. Influence of integrated nitrogen management practices on growth and yield of okra. *Journal Research ANGRAU*, **40**(4):11-13.
10. Rathava, S.N., Verma, P., Pawar, Y. and Limbachiya, T.B. 2018. Okra yield and economics: Effect of fertilizer amounts and biofertilizers (*Abelmoschus esculentus* L. Moench) cv. GAO-5 P-ISSN:2349-8528.

11. Singh, B.K., Verma, R.B., Singh, V.K., Singh, M., Maurya, D. 2018. Effect of integrated nutrient management on growth, yield and quality of okra. *International journal of current microbiology and applied sciences*, **7**(10):1033-1041.
12. Sharma, J., Sharma, B.C., Puniya, R., Sharma, R. and Menia, M 2021. Effect of Seed Priming and Plant Geometry on Growth and Yield of Wheat in Modified System of Wheat Intensification Under Irrigated Sub Tropics of Jammu. *AMA, Agricultural Mechanization in Asia, Africa and Latin America*, **51**(03): 1663-1669.
13. Vennila, C. and Jayanthi, C. 2008. Effect of integrated nutrient management on yield and quality of Okra. *Research on Crops*, **9**(1):73-75.
14. Yadav, G. L., Singh, S. P., Jitarwal, O. P., Yadav, V. K. and Choudhary, R. 2017. Effect of nitrogen and bio-organics on growth and yield of okra [*Abelmoschus esculentus* L. Moench]. **6** (23):1515-1519.
15. Singh Smriti and R.B. Ram. Effect of Organic, Inorganic and Bio-Fertilizers on Growth, Yield and Quality Traits of Okra [*Abelmoschus esculentus* (L.) Moench]. *Int.J.Curr.Microbiol.App.Sci* (2018) **7**(4): 2388-2392

UNDER PEER REVIEW