

***In vitro* evaluation of botanicals, bio-agents and fungicides against *Rhizoctonia solani*
causing Web blight disease of mungbean**

Abstract

Mungbean [*Vigna radiata* (L.) Wilczek] is the important source of proteins, minerals, and vitamins of the predominantly vegetarian Indian diet. It belongs to the family Leguminaceae. Web blight caused by *Rhizoctonia solani* (Kuhn) is one of the most important fungal diseases which come every year with different intensity and causes huge losses in mungbean yield. The present investigations were carried out in the laboratory, Department of Plant Pathology Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya to test the efficacy of different treatments viz., Neem, Garlic, Tulsi, onion, Ginger, *T. asperellum*, *T. harzianum*, Propiconazole and Hexaconazole against *Rhizoctonia solani* Kühn under *in vitro* condition. Botanicals and Fungicides were tested through Poisoned food technique and Bio-agents were tested through dual culture technique. Pathogen was isolated from diseased mungbean plant and further tested against different treatments. Radial growth and percent inhibition were recorded. Minimum radial growth and maximum percent inhibition were recorded in Propiconazole 1.45 mm, followed by Hexaconazole (3.65 mm), Garlic (10.18 mm), Ginger (11.43 mm), Neem (12.90 mm), Onion (15.42 mm), Tulsi (17.63 mm), *T. asperellum* (18.58 mm), *T. harzianum* (23.14 mm) as compared to Control (45.17 mm) at 24 hours of incubation. Similar trends were found at 36 and 48 hours intervals.

Keywords: Mungbean, Neem, Garlic, Tulsi, onion, Ginger, *T. asperellum*, *T. harzianum* Propiconazole, Hexaconazole, *Rhizoctonia solani*.

Introduction

“Mungbean [*Vigna radiata* (L.) Wilczek] is the important source of protein, vitamins and minerals. It belongs to the family Leguminaceae. Among the pulses mungbean also called as green gram or golden gram. It is mainly grown in Rajasthan, Maharashtra, Karnataka, Andhra Pradesh, Orissa, Bihar, Tamil Nadu, Madhya Pradesh, and Uttar Pradesh. In Uttar Pradesh, it is cultivated on 93000 ha, with a production of 9480 tonnes. The productivity of mung bean in India and the U.P. is 567 kg/ha and 536 kg/ha, respectively, which is very low compared to the genetic potential of 1500–2000 kg/ha” (Anonymous, 2019). “In 1924, web blight was reported for the first time on mung bean from the Philippines” (Nacien, 1924). While in India, Dwivedi and Saksena (1974) first reported it on mung beans from Kanpur, Uttar Pradesh. (Anonymous, 2004). Web blight caused by *Rhizoctonia solani* (Kuhn) is one of the most important fungal disease which appear every year in varying intensity and causes heavy reduction in yield. Gupta

et al. (2010) reported “the losses in grain yield is more when the plants get infected earlier i.e. after 25 days after sowing (DAS) than 35 and 40 DAS . The losses in yield and lost weight were 33.40 to 37.80 per cent and 23.12 to 28.60 per cent respectively”. Though, the web blight could be managed by the use of fungicide but due to the emergence of several problems like environmental pollution, residual effect, its use should be discouraged. Many botanicals and bio-agent have been found effective against *Rhizoctonia solani* in different crops, therefore keeping in view the importance of the crop and seriousness of diseases present research work carried.

Materials and Methods

The present investigations were carried out in the laboratory, Department of Plant Pathology A.N.D.U.A. & T., Kumarganj, Ayodhya (Year 2022-23) to test the efficacy of different treatments against *Rhizoctonia solani* Kühn under *in vitro*. Pathogen was isolated from diseased mungbean plant parts (Leaf and Stem) and further tested against different treatments. In order of find out the efficacy of various plants extracts against *R. solani*, five plants extract viz., Neem (leaf), Garlic (bulb), Tulsi (leaf), onion (bulb) and Ginger (rhizome) were used. Fresh leaves, bulbs and rhizomes were collected and washed thoroughly in clean water. Equal amount of washed plant material was grinded in mortar and pestle by adding same amount of sterilized water (1:1 w/v) and boiled at 80 °C for 10 minutes in hot water bath . The extract was filtered by double layer muslin cloth followed by sterilized Whatman No.1 Filter paper. By adding the 20 ml of sterilized PDA medium and 2 ml Botanical extract, the concentrations of 10.0 percent were created. Two fungicides were tested *in vitro* against *R. solani* i.e., Propiconazole (20 ppm) and Hexaconazole (20 ppm). The flasks were thoroughly shaken to ensure an even mix of the extract and fungicide under aseptic conditions. Twenty ml of sterilized melted PDA was aseptically poured in sterilized Petri dishes and allowed to solidify. The efficacy of *T. asperellum* and *T. harzianum* against *Rhizoctonia solani* were assessed by using dual culture technique by measuring the radial growth of *R. solani* as well as that of *Trichoderma spp.* Culture of Bio-agent were collected from laboratory Department of plant pathology A.N.D.U.A. & T., Kumarganj, Ayodhya. Three-day-old *R. solani* culture discs were cut into five mm pieces with a sterilized cork borer and positioned in the centre of Petri dishes with plant extracts and fungicides added. Five mm disc of each antagonist and *R. solani* cut with the help of sterilized cork borer from the age of three days old culture and were placed in Petri dishes having solidified PDA in such a

manner that they lie opposite to each other 60 mm apart. “Control (Check) Petri dishes were inoculated only with *R. solani* bits. Each treatment and control was repeated four times to make four replications. These Petri dishes were kept in BOD incubator at 28⁰ C. The observations on radial growth were made at 24, 36, and 48 hours of incubation in Petri dishes amended with different treatments as well as in control. Per cent growth inhibition was calculated by using formula” (Vincent, 1947). The conclusion was arrived at after statistical analysis of the data. The Completely Randomized Design (CRD) method recommended by Goon *et al.* was used to conduct the statistical analysis of laboratory experiments (1931). The variance ratio test at the 5% level of probability was used to determine the significance of treatment differences. The observation of per cent inhibition of mycelial growth, were transformed in to “Arc sin Transformation” = $\sin^{-1}\sqrt{p}/100$ used for statistical analysis.

$$I = \frac{C - T}{C} \times 100$$

Where,

I = Per cent inhibition of fungal growth.

C = Radial growth of control.

T = Radial growth in treated Petri dish.

Table-1: List of botanicals and their scientific names.

S.No.	Common Name	Scientific Name	Plant part used
1.	Neem	<i>Azadirachta indica</i>	Leaves
2.	Garlic	<i>Allium sativum</i>	Bulb
3.	Tulsi	<i>Ocimum snactum</i>	Leaves
4.	Onion	<i>Allium cepa</i>	Bulb
5.	Ginger	<i>Zingiber officinale</i>	Rhizome

Table-2: Effect of botanicals, bio-agents and fungicides against *R. solani* on mycelial

Treatment	Mycelial Growth (mm)		
	Duration in Hours		
	24 Hours	36 Hours	48 Hours
Neem	12.90	18.63	19.95
Garlic	10.18	14.04	15.49
Tulsi	17.63	22.00	24.66
Onion	15.42	20.00	22.33
Ginger	11.43	16.07	18.00
<i>T. asperellum</i>	18.58	23.24	25.60
<i>T. harzianum</i>	23.14	31.58	39.55
Propiconazole	1.45	1.85	2.16
Hexaconazole	3.65	4.85	5.93
Control	45.17	71.86	85.42
CD at5%	1.18	2.27	1.30
SE(m)	0.39	0.76	0.44

growth at 24, 36 and 48 hours.

Results and Discussion

➤ At 24 hours of incubation

minimum radial growth was obtained in Propiconazole (1.45 mm) followed by Hexaconazole (3.65 mm), Garlic (10.18 mm), Ginger (11.43 mm), Neem (12.90 mm), Onion (15.42 mm), Tulsi (17.63 mm), *T. asperellum* (18.58 mm), *T. harzianum* (23.14 mm) as compared to Control (45.17 mm). Propiconazole, Hexaconazole, Garlic, Ginger, Neem, Onion and *T. harzianum* were significantly different to each other while Tulsi and *T. asperellum* were found at par to each other. (Table 2, Fig 1)

➤ At 36 hours of incubation

minimum radial growth was obtained in Propiconazole (1.85 mm) followed by Hexaconazole (4.85 mm), Garlic (14.04 mm), Ginger (16.07 mm), Neem (18.63 mm), Onion (20.00 mm), Tulsi (22.00 mm), *T. asperellum* (23.24 mm), *T. harzianum* (31.58 mm) as compared to Control (71.56 mm). Propiconazole, Hexaconazole, *T. harzianum* were significantly different to each other while Neem, Onion Tulsi and *T. asperellum*, Garlic and Ginger, were found at par to each other. (Table 2, Fig 1)

➤ **At 48 hours of incubation**

minimum radial growth was obtained in Propiconazole (2.16 mm) followed by Hexaconazole (5.93 mm), Garlic (15.49 mm), Ginger (18.00 mm), Neem (19.95 mm), Onion (22.33 mm), Tulsi (24.66 mm), *T. asperellum* (25.60 mm), *T. harzianum* (39.55 mm) as compared to Control (85.42 mm). Propiconazole, Hexaconazole, Garlic, Ginger, Neem, Onion and *T. harzianum* were significantly different to each other while Tulsi and *T. asperellum* were found at par to each other. (Table 2, Fig 1)




Fig.1: Effect of botanicals, bio-agents and fungicides against *R. solani* on mycelial growth at 24, 36 and 48 hours.

Table-3: Effect of botanicals, bio-agents and fungicides against *R. solani* on mycelial growth at 24, 36 and 48 hours.

Treatment	Percent inhibition		
	Duration in Hours		
	24 Hours	36 Hours	48 Hours
Neem	71.43 (57.67)	74.09 (59.40)	76.52 (61.07)
Garlic	77.45 (61.62)	80.42 (63.76)	81.89 (64.79)
Tulsi	60.95 (51.31)	69.39 (56.40)	71.08 (57.48)
Onion	65.79 (54.20)	72.20 (58.20)	73.81 (59.23)
Ginger	74.67 (59.77)	77.63 (61.75)	78.98 (62.64)
<i>T. asperellum</i>	58.87 (50.08)	67.65 (55.31)	70.82 (57.28)
<i>T. harzianum</i>	48.74 (44.26)	56.04 (48.51)	54.30 (47.45)
Propiconazole	96.80 (79.67)	97.42 (80.74)	97.47 (80.81)
Hexaconazole	91.92 (73.46)	93.25 (74.91)	93.05 (74.70)
Control	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
CD at 5%	2.34	2.88	1.47
SE(m)	0.78	0.96	0.49

➤ **At 24 hours of incubation**

Results clearly indicated that maximum percent inhibition was found in Propiconazole (96.80%) followed by Hexaconazole (91.92%), Garlic (77.45 %), Ginger (74.67 %), Neem (71.43 %), Onion (65.79 %), Tulsi (60.95 %), *T. asperellum* (58.87 %) and *T. harzianum* (48.74%) as compared to control (0.00%). Propiconazole, Hexaconazole, Garlic, Ginger, Neem, Onion and *T. harzianum* were significantly different to each other while Tulsi and *T. asperellum* were found at par to each other. (Table 3, Fig 2)

➤ **At 36 hours of incubation**

Maximum percent inhibition was recorded in Propiconazole (97.41%) followed by Hexaconazole (93.25%), Garlic (80.42 %), Ginger (77.63 %), Neem (74.09 %), Onion (72.20 %), Tulsi (69.39 %), *T. asperellum* (67.65 %) and *T. harzianum* (56.04 %) as compared to control (0.00%). Propiconazole, Hexaconazole and *T. harzianum* were found statistically differed to each other while Tulsi, *T. asperellum*, Onion, Neem and Ginger and Garlic were found at par to each other. (Table 3, Fig 2)

➤ **At 48 hours of incubation**

Maximum percent inhibition was recorded in Propiconazole (97.47%) followed by Hexaconazole (93.25%), Garlic (81.89 %), Ginger (78.98 %), Neem (76.52 %), Onion (73.81 %), Tulsi (71.08 %), *T. asperellum* (70.82 %) and *T. harzianum* (54.3 %) as compared to control (0.00%). Propiconazole, Hexaconazole, Garlic, Ginger, Neem Onion, and *T. harzianum* were found statistically differed to each other while Tulsi and , *T. asperellum* were found at par to each other. (Table 3, Fig 2)




Fig.2: Effect of botanicals, bio-agents and fungicides against *R. solani* on percent inhibition at 24, 36 and 48 hours.

		
Neem	Garlic	Tulsi
		
Onion	Ginger	Control

Plate: 1. Response of botanicals against *R. solani* on mycelial growth.

		
<i>Trichoderma asperellum</i>	<i>Trichoderma harzianum</i>	Control

Plate: 2. Response of bio-agents against *R. solani* on mycelial growth.

		
Propiconazole	Hexaconazole	Control

Plate: 3. Response of fungicides against *R. solani* on mycelial growth.

Conclusion

Our study very well demonstrated that, Propiconazole demonstrated the highest efficacy against *Rhizoctonia solani*, followed by Hexaconazole, while garlic and ginger extracts exhibited notable inhibitory effects. *T. asperellum* and *T. harzianum* bio-agents also displayed significant antifungal activity. These findings support the potential integration of botanicals and bio-agents for web blight disease in mungbean. Many environmental concerns linked to chemical

References

- Anonymous (2019). *All India Coordinate Research Project on MullaRP, IIPR. Kanpur*, 108 pp.
- Nacien, C.C. (1924). Studies on *Rhizoctonia blight* of beans. *Philippine Agriculturist*, 8:315-321.
- Dwivedi, R.P. and Saksena, H.K. (1974). Occurance of web blight disease caused by *Thanatephorus cucumeris* on mungbean. *Indian J. Farm Sci.* 2:100.
- Saikia, U.N. 1976. Blight of mung caused by *Corticium sasakii* a new disease recorded from Assam. *Indian Phytopath.* 29: 61-62.
- Bains, S.S., Dhaliwal, H.S. and Basandrai, A.K. 1988. A new blight of Mung and Mash in Punjab. *Ann. Biol. Ludhiana.* 4: 113–114.
- Tiwari, A. and Khare, M.N. (1998). Variability among isolates of *Rhizoctonia solani* infecting mungbean. *Indian phytopath.*, 51:334-337.
- Anonymous (2004). *Annual Report (kharif)*. All India Co-ordinated Research Project on MullaRP (ICAR), IIPR, Kanpur, 112 pp.
- Gupta, R.P. Singh, S.K. and Singh, R.V. (2010). Assessment of losses due to web blight and weather effects on disease development in mungbean. *Indian Phytopath.* 63 (1) : 108-109.

- Vincent, J.M. (1947). Distortion of fungal hyphae in the presence of certain inhibitors. *Nature* 159:850-853.
- Dubey, S.C. and Patel, B. (2001). Evaluation of fungal antagonists against *Thanatephorus Cucumeris* causing web blight of urd and mungbean. *Indian Phytopath.* 54(2) : 206- 209.
- Dubey, S.C. (2002). Evaluation of *Glioclodium virens* and *Trichoderma viride* as foliar spray against web blight of urd and mungbean. *J. Mycol. Pl. Pathol.* 32 (2):236-237
- Meena, R.L., Rathore and Mathur, K. (2002). Evaluation of fungicides and plant extracts against banded leaf and sheath blight of maize (*Zea mays*). *J. Mycol. PL. Pathol.* 32 (3):397.
- Khan, A.A. and Sinha, A.P. (2007). Screening of *Trichoderma* spp. against *Rhizoctonia solani* the causal agent of rice sheath blight. *Indian Phytopath.* 60 (4):450-456
- Kumar, Vipin., Chaudhary, V.P., Kumar, Dharmendra., Kumar, Ajay., Sagar, Sushma and Chaudhary, Sorabh. (2017) Efficacy of botanicals and fungicides against *Rhizoctonia solani* inciting sheath blight disease on Rice (*Oryza sativa* L.) *Journal of Applied and Natural Science* 9 (4): 1916 -1920(2017)
- Shinde, G.R. and Patel, R.L. (2004). Evaluation of plant extract against *R. solani* incitant of black scurf Disease in Potato. *J. Mycol. Pl. Pathol.* 34 (2) 28:4- 285