

Review Article

"Revolutionizing plant virus detection: A review on molecular breakthroughs and their implications"

Abstract:

Viruses are very minute microorganisms which are incapable of independent existence and can infect wide range of wild and cultivated plants. The plant's reaction to viral infections is diverse and depends on factors such as the specific virus- host interaction under different environmental conditions with symptoms ranging from acting as asymptomatic carrier to showing severe diseases symptoms which further results in plant mortality. Since various viral infections can result in similar symptoms, relying solely on the disease's external appearance offers only partial insights for disease diagnosis. To achieve more precise and dependable virus identification, it is essential to consider various characteristics and properties of the virus. The molecular detection of plant viruses has revolutionized the field of plant virology. This review highlights the overview of the different approaches utilized in the diagnosis of plant viruses, with a particular focus on significant advancements in molecular techniques, ranging from traditional PCR-based methods, ELISA to high-throughput next-generation sequencing (NGS) technologies. Molecular tools have enabled precise and rapid identification of plant viruses, contributing to better disease management in agriculture. The review also delves in to emerging technologies such as CRISPR-based diagnostics and nanopore sequencing, which hold promise for enhancing the precision and speed of plant virus detection.

Key words: PCR, ELISA, NGS (Next Generation Sequencing), CRISPR, Nanopore sequencing.

1. Introduction:

Plant viruses are very small size pathogenic entities and hence cannot be observed through light microscopes. They necessitate the use of transmission electron microscopes. Viruses are structured with a protective coat protein called capsid enveloping a core that carries genetic information in the form of either DNA or RNA. Their genome size is very small which encodes

only few proteins hence its very difficult to trace virus and control them (Rojas *et al.*,2008). Viral infections in crops have a detrimental impact on crop development, leading to diminished yields, affecting the survival of grafts or scions, and compromising the quality of fruits causing significant economic losses(Maksimov *et al.*,2019).Viruses induce a range of symptoms in infected plants, including mosaic patterns, yellowing, chlorosis (loss of green color), virescence (abnormal growth of chlorophyll), fasciation (abnormal flattening of plant parts), fasciculation (formation of abnormal clusters), stunting, witches' broom-like growth, and necrosis (cell death). These symptoms result in impaired growth and reproductive functions, ultimately causing a decline in the overall health and productivity of the plants (Seo *et al.*, 2018).

Identifying viral diseases in plants based on symptoms can be quite challenging as some plant viruses can infect plants without manifesting visible signs, making symptom-based diagnosis more complicated. Accurate detection using precise and sensitive techniques plays a vital role in the effective management of plant diseases. Additionally, the presence of specialized tools for virus diagnosis and identification is a fundamental requirement for the development and assessment of disease management strategies(Rubio *et al.*,2020).

Due to wide diversity in genetic material of viruses infecting plants and their high mutation rates, it's very difficult to fully rely on any single method of detection. Hence , all methods starting from preliminary visual observation to microscopic identification ,serological methods, molecular methods, and next generation sequencing must be employed to allow early identification and management of virus .These tests allow for the exact identification of viral pathogens in plants, guaranteeing a more precise diagnosis(Massart *et al.* ,2014).With advancement in scientific field and introduction of next-generation sequencing (NGS) technologies,millions or even billions of nucleotides can be sequenced simultaneously without need of prior sequence orientation leading to significant transformation in the field of plant virus diagnosis(Stobbe *et al.*, 2014).Comprehensive discussions of key diagnostic methodologies, ranging from conventional visual assessments of indicator plants to cutting-edge third-generation sequencing technologies, are outlined in the following sections.

2. Conventional morphological assessment of diseased plants

Historically, the predominant method for identifying plant pathogens involved visually inspecting infected plants and seeds based on specific symptoms, using indicator plants, assessing their host range, examining physical characteristics of virus such as their shape and size, and studying virus-vector relationships. When the symptoms are distinct and easily recognizable, visual inspection is a straightforward task. Nonetheless, several variables like the virus strain, the specific plant variety, the timing of infection, and environmental conditions can all impact the appearance of these symptoms (Matthews 1980). When dealing with a plant viral disease of established origin, the key factor in its recognition typically hinges on the symptoms evident in the host plants as it bears a distinct relation to the disease. The external indications of viral plant infections are intricately linked to irregularities in plant physiology, and these symptoms are further classified into mosaics and chlorosis (Scholthof *et al.*, 2011). Visual inspections can be carried out using various techniques, including visible-light imaging, chlorophyll fluorescence imaging, hyperspectral imaging, and thermal imaging (Badoliet *et al.*, 2017). In most cases, it is important to combine visual symptom inspection in the field with additional confirmatory tests to ensure a precise diagnosis of viral infection as in plants, to get clarity in mixed infections, in different environmental or growing conditions, and depending on species or cultivar (Bock, 1982).

2.1 A close-up view of the invisible world of virus through microscopy

Understanding the biology of viruses and the causes of viral diseases is essential for preventing these diseases, ensuring accurate and effective virus diagnosis, and implementing virus control measures. A conventional method for studying viruses in plant tissues entails using advanced light microscopy and high-resolution electron microscopy, enabling the visual inspection of viruses within plants. The exceptional ability of electron microscopy (EM) to resolve at the nanometer scale allows for the direct visualization of viruses, offering valuable images for diagnostic and research purposes. Transmission electron microscopy (TEM) has advanced the characterization of viral particle morphology in both unprocessed and purified samples. These TEM studies have led to one of the earliest proposals for classifying viruses based on their distinctive morphological and serological associations, along with certain biological features. TEM enables direct detection, involving the initial homogenization of infected tissue followed by the application of negative staining. TEM's defining characteristic is its ability to offer an

expansive and immediate view, enabling the rapid assessment of the current situation, including the quantity and configuration of viruses present, and even revealing unexpected findings (Gentile and Gelderblom, 2014). As an initial stage in identifying pathogens, TEM necessitates only small sample quantities that contain a significant viral load. TEM is impartial towards RNA or DNA genomes because it focuses on the input of proteins, the viral capsid, or ribonucleoprotein (RNP) complexes (King *et al.*, 2012). Tobacco mosaic virus is known for its characteristic production of needle-shaped and hexagonal crystals, whereas Potato virus X and wheat mosaic virus are notable for their capacity to generate spherical structures that do not crystallize. Various viruses that affect plants, including Tomato yellow leaf curl disease (TYLCD), Potato virus S (PVS), Rice stripe mosaic virus (RSMV), Tomato brown rugose fruit virus (ToBRFV), Pepino mosaic virus (PepMV), and Potato virus M were investigated and studied using electron microscopy techniques (Song *et al.*, 2017; Guo *et al.*, 2020; Klap *et al.*, 2020; Zhang *et al.*, 2020). Therefore, TEM acts as a crucial instrument for establishing which of the available techniques such as bioassays, serological methods, or molecular biology approaches should be employed next to identify the genus and species of the virus more precisely.

3. Probing the physical features of viruses

In the past, characteristics of a virus like its thermal inactivation point, dilution end point, and *in vitro* longevity were utilized as indicators of the virus's infectivity in sap extracts and for identifying plant viruses. Nevertheless, these properties have been found to be unreliable and are no longer recommended for virus diagnosis though still studied in literature (Francki, 1980).

4. Serological Assays

Serological tests are based on the interaction of specific antibody with antigen against which it is produced. One such method based on serological assay is ELISA. Traditional ELISA procedures involve the use of polystyrene plates that can bind antibodies or proteins and incorporate an enzyme-substrate reaction. The application of enzyme-linked immunosorbent assay (ELISA) is highly valuable tool for virus detection in plants. The techniques, such as ELISA (enzyme-linked immunosorbent assay), which rely on the dependable detection of protein molecules using either polyclonal or monoclonal antibodies, are extensively utilized in the realm of plant viral diagnostics. Clark and Adams were the first to employ the ELISA method for the diagnosis of plant viral diseases. The ELISA technique's practical efficiency remained consistent, irrespective

of any ratio between antibodies and antigens. Once the appropriate concentrations were established, they could be universally applied for subsequent virus detection tests, regardless of the virus concentration. The enzyme-labeled antibody's reaction was directly linked to the virus concentration, underscoring the technique's robust quantitative potential. Bar-Joseph *et al.*, 1979 conducted experiments to investigate the serological reactions of four different plant viruses, namely citrus tristeza virus (CTV), carnation mottle virus (CarMV), carnation yellow fleck virus (CYFV), and tobacco mosaic virus (TMV). They analyzed how these viruses interacted with their corresponding gamma-globulin alkaline phosphatase conjugates when applied in a sandwich format on antibody-coated microplates. Among ELISA-based approaches, direct tissue blot immunoassay (DTBIA), double antibody sandwich (DAS) ELISA, and tissue-print (TP) ELISA stand out as the most employed methods for viral identification (Xie *et al.*, 2013 ; Rani *et al.*, 2019 ; Abd El-Aziz *et al.*, 2019). Numerous variations of ELISA have been developed and applied to enable the swift detection of various plant viral diseases in a diverse range of host plants.

4.1 The Dot blot immunobinding assay (DBIA)

It is a modified blotting technique widely employed to specifically identify nucleic acids and proteins. In this method, antigen is applied onto a nitrocellulose membrane in the form of drop of a test sample. Then the test antibody is added to nitrocellulose membrane, followed by exposure to a peroxidase-conjugated secondary antibody targeting the first antibody. The final step involves the development of the assay using 4-chloro-1-naphthol. This DBIA procedure is utilized for screening the supernatants of hybridomas to detect monoclonal antibodies and to screen pathological sera for multiple antibodies (Hawkes *et al.*, 1982). This technique can detect viruses even if very small quantity of antigen or crude sap is available.

4.2 Double antibody sandwich (DAS) ELISA

It is a form of indirect enzyme-linked immunosorbent assay (ELISA) that has been created for the purpose of detecting viruses in plants. This technique involves using protein in two stages to form a sandwich-like structure with layers of antibody-antigen-antibody. The initial application of protein primes the microtiterplate for the antibody coating layer. The second layer of protein is conjugated to the enzyme and is responsible for identifying the second antibody layer by giving a particular colour on reaction with substrate.

4.3 Tissue blotting immunoassay (TBIA)

Tissue blots were created gently but firmly by pressing freshly cut plant tissue onto nitrocellulose membranes. Infected freshly cut plant tissue is directly electro-blotted on to the nitrocellulose membrane (transfer of viral antigen to membrane). The existing antigens were first reacted with primary antibodies specific to virus antigen and detected with enzyme labelled secondary antibodies also called probes that react with primary antibodies (Fegla *et al.*, 2001). Distinct reactions specific to the Faba bean necrotic yellow virus (FBNYV) antigen were noticed on tissue blots from infected plants, whereas no such reactions were observed on tissue blots from non-inoculated faba bean plants. The red staining was localized to the midrib and secondary vein regions of leaf, petiole, and stem sections, indicating that the FBNYV was confined to the vascular tissue (Lina *et al.*, 1993).

ELISA relies on the interaction between antibodies and antigens, highlighting the critical need for highly specific antibodies that can accurately detect the target substance. However, ELISA may occasionally produce incorrect positive results, mainly because of non-specific responses or the cross-reaction with specific components found in the samples. (Kfir and Genthe, 1993). In summary, although ELISA has been widely employed in diagnostics, its shortcomings have become more apparent. Hence, we need to discuss nucleic acid-based methods, CRISPR based technologies and high-throughput next-generation sequencing (NGS) technologies.

5. Nucleic acid-based methods

Nucleic acid-based methods have brought a significant transformation to the realm of viral detection. They provide excellent sensitivity and precision while enabling the identification and detailed analysis of various viruses, even including newly emerging ones. The selection of a particular technique relies on factors such as the specific virus under investigation, the type of sample, available resources, and the desired speed in obtaining results. The rapid advancements in molecular and genomics technologies have expanded the applications of these methods, particularly in the detection of infectious pathogens such as viruses. Due to their extreme higher sensitivity, nucleic acid-based techniques whether focused on DNA or RNA, have gained widespread acceptance in viral diagnostics. In general, nucleic acid-based analysis involves three crucial stages: nucleic acid extraction (either DNA or RNA), amplification, and the pivotal step of product analysis, which directly provides the test result (Klap *et al.*, 2020). Amplification can

be achieved by various PCR techniques which are discussed below and further sequencing can be carried out for virus detection.

5.1 PCR(Polymerase Chain Reaction)

DNA amplification techniques for viral detection can be analysed with polymerase chain reaction (PCR). PCR technology has made substantial strides in simplifying and enhancing the detection and diagnosis of specific genes. This technique involves creating millions of copies of a specific region of the viral genome, which can be visualized through methods like electrophoresis or by using fluorescent probes. PCR can utilize genomic DNA or complementary DNA generated through reverse transcription (RT) of viral RNA as templates. The amplification process consists of three major steps which are denaturation, annealing followed by extension. These three steps are repeated through multiple cycles (usually 20 to 40 cycles), with the newly synthesized DNA segments serving as templates for subsequent cycles (Mullis and Faloona, 1987).

5.1.1 Real-time quantitative PCR (qPCR)

It is a modified version of PCR method that tracks the advancement of the reaction by identifying a fluorescent marker that attaches to the double-stranded DNA or gets released from specific 15 to 30 nucleotide probes. This form of PCR is employed for the purpose of measuring nucleic acids accurately. Further, isothermal DNA amplification can be achieved by various methods such as i) Helicase dependent amplification (HAD) ii) Recombinase polymerase amplification (RPA) iii) Nucleic acid sequence-based amplification method (NASBA) iv) Loop-mediated isothermal amplification (LAMP)(Olmos *et al.*, 2007). More recently, novel molecular diagnostic tools have been created, utilizing the prokaryotic clustered regularly interspaced short palindromic repeats (CRISPR) immune system, which is widely recognized for its applications in genome editing (Chertow, 2018). The PCR technique is employed to oversee the effectiveness of antiviral treatment for HIV-1, HBV, HSV-1, and HSV-2. Performing uniplex RT-PCR for each virus or viroid separately can be costly and demanding in terms of resources, as it necessitates both time and materials.

5.1.2 Nested-PCR

It is a method within the realm of polymerase chain reaction (PCR) that enhances both the sensitivity and specificity of DNA amplification. This approach employs two distinct sets of amplification primers and involves a two-step amplification process. In the initial round, a single pair of primers is employed for a specified number of cycles, typically ranging from 15 to 30. The resulting amplified product from the first round serves as the template for a subsequent round of amplification, utilizing a different pair of primers. However, it is noteworthy that a notable drawback associated with nested-PCR is the increased susceptibility to contamination issues.

5.1.3 Multiplex PCR

Multiplex PCR for DNA targets and multiplex RT-PCR (mRT-PCR) for RNA targets are efficient, dependable, and cost-effective techniques that have proven effective in simultaneously identifying various pathogens in a single test. With mRT-PCR, distinct sets of specific primers are used for two or more targets within a single reaction tube, allowing for the concurrent amplification of multiple target nucleic acids in a single assay. Multiplex detection of more than three targets by fluorescent probes is carried out by the dyes with compatible range of spectra. Simultaneous detection of four pathogens has been reported for four retroviruses (Vet *et al.*, 1999).

6. DNA microarrays

DNA microarrays or biochips consist of a surface to which multiple capture probes are attached, with each probe designed to be specific for a DNA or RNA sequence of the target. Their primary function is to detect numerous sequences in a single test. Various materials are currently used to create microarrays, including glass, nylon, and different types of polymers. A single chip can accommodate up to 30,000 DNA probes, representing gene sequences. These probes can be PCR products that have been amplified to high concentrations or relatively short oligonucleotide probes, typically ranging from 30 to 50 base pairs. Once the probes are arrayed on the chip, it can be exposed to DNA/RNA from the sample to be tested, which is labeled with fluorescent markers (Lopez *et al.*, 2003).

7. Loop Mediated isothermal amplification

The LAMP method is a simple and easily executable technique, provided that the suitable primers have been prepared in advance. This process necessitates the utilization of a DNA polymerase, four primers, and a conventional laboratory water bath or heat block for the reaction. LAMP amplifies DNA with high specificity, efficiency, and rapidity under isothermal conditions. The year 2000 marked the inception of Loop-mediated isothermal amplification (LAMP) technology, with the primary goal of improving the efficiency of nucleic acid amplification in relation to sensitivity and specificity(Notomiet *al.*,2000).

A combination of Loop-mediated isothermal amplification (LAMP) and subsequent detection using a real-time fluorescence assay with CRISPR-Cas12a was employed to identify the presence of tomato yellow leaf curl virus and tomato leaf curl New Delhi virus (Mahas *et al.*, 2021).

8. CRISPR/Cas

CRISPR/Cas (clustered regularly interspaced short palindromic repeats/CRISPR- associated) is an innate immune system naturally employed by prokaryotic organisms like bacteria and archaea to protect themselves from invasive viruses and plasmids(Doudna and Charpentier, 2014; Jinek *et al.*, 2012). CRISPR/Cas has become a popular choice for genome editing because of its precise targeting and the ability to modify any gene by changing the guide RNA sequence. Its adaptability for specific targets has led to its use in interfering with viruses in a range of organisms, including plants (Mahas *et al.*, 2018,2019).

9. In situ hybridization (ISH)

In situ hybridization is a technique used to identify and pinpoint a particular genetic sequence within a single cell. This genetic sequence is specifically attached to a tissue sample through matching base pairs, or hybridization, with a visible genetic fragment known as a probe.In various scientific domains, including virus exploration, in situ hybridization (ISH) is commonly employed. The advancement of high-throughput techniques such as next-generation sequencing has led to a higher identification of novel viruses. ISH proves to be a valuable tool in verifying a potential link between a recently identified pathogen and changes in tissue (Bodewes *et al.*, 2014 ;Pfankuchet *et al.*,2016).

10. High-Throughput Sequencing

The introduction of high-throughput sequencing (HTS) methods, commonly referred to as next-generation sequencing, has brought about a significant transformation in the diagnosis of plant viruses (Maree *et al.*, 2018; Villamor *et al.*, 2019). High-throughput sequencing (HTS) does not require prior knowledge of viral genetic codes and can simultaneously sequence millions or even billions of DNA strands, allowing us to identify all the viruses in a plant (Roossinck, 2015).

High-throughput sequencing (HTS) can be categorized into two main types i.e., Second-generation sequencing and third generation sequencing. The second-generation sequencing relies on the generation of random libraries of DNA fragments using the initial DNA material or cDNA obtained through retro transcription of RNA with random primers or oligo-dT. These libraries are subsequently clonally amplified, attached to synthetic DNA adapters, and then sequenced in a parallel manner. This process generates a substantial number of short sequences reads, typically ranging from 100 to 500 nucleotides in length. These short reads are subsequently assembled by aligning overlapping sequences based on nucleotide identity, and this assembly is achieved through computational analysis, often using software tools like the geneious package. (www.geneious.com). Furthermore, high-throughput sequencing (HTS) is employed for investigating a broad spectrum of scientific inquiries in the realm of plant virology. These applications encompass the identification of novel plant viruses and viroids, epidemiological investigations, exploration of synergistic interactions among viruses, and the analysis of genetic diversity and the evolutionary mechanisms that drive virus populations (Kehoe *et al.*, 2014; Hadidi *et al.*, 2016; Pecman *et al.*, 2017; Tan *et al.*, 2019; Xu *et al.*, 2019).

Third-generation sequencing involves the direct, real-time sequencing of individual DNA molecules, eliminating the necessity for clonal amplification. This innovation not only streamlines the DNA preparation process but also yields extended reads spanning several kilobases in length. (Goodwin *et al.*, 2016; van Dijk *et al.*, 2018).

Diverse biotechnological enterprises are actively engaged in the advancement of genomic sequencing methodologies, exemplified by innovations like Single-Molecule Real-Time (SMRT) sequencing and Nanopore sequencing. In the context of SMRT sequencing, it entails the utilization of a specialized flow cell replete with an array of myriad picolitre wells, each

featuring a transparent base referred to as zero-mode waveguides. Within these wells, a DNA polymerase is immobilized. The sequencing process involves the continuous visualization of nucleotide incorporation into individual DNA molecules ensconced in the wells. A laser and camera system are employed to meticulously capture the emitted light's color and duration as nucleotides are temporarily held in abeyance at the well's bottom during the incorporation event.

Conversely, Nanopore sequencing hinges upon the controlled translocation of DNA or RNA strands through a nanopore, which can manifest within membrane proteins or artificial materials such as silicon nitride and aluminum oxide. This passage induces an ionic current that is governed by an applied voltage. The unique aspect of Nanopore sequencing lies in its capability to discern variations in the ionic current, contingent upon the structural attributes, dimensions, and length of the traversing DNA sequences. This process endows Nanopore sequencing with several merits, including cost-efficiency relative to other High-Throughput Sequencing (HTS) technologies, portability due to the compact size of the sequencing apparatus, and expeditious sample processing. Furthermore, it obviates the necessity for reverse transcription when working with RNA viruses (Kiselev *et al.*, 2020). This cutting-edge technology has been employed in recent times for the identification of various plant viruses, such as plum pox virus (PPV), the tomato yellow leaf curl virus (TYLCV), as well as for the exploration and detection of previously unknown plant viruses (BronzatoBadiale *et al.*, 2018; Chalupowicz *et al.*, 2019; Naito *et al.*, 2019).

11. Current challenges and future prospects

The accurate and rapid identification of the organisms causing plant diseases is essential for their timely management to avoid crop losses. It enables informed decision-making when it comes to selecting crops and using chemicals to control disease outbreaks. Recent advancements in molecular detection technology have led to the development of more convenient, efficient, and precise tests, which can detect various plant pathogens, including viruses. These tests offer valuable support to farmers, agricultural experts, and plant health specialists by reducing the reliance on symptom-based identification and facilitating early disease detection. These modern techniques become powerful management tools when combined with a deep understanding of the crop, knowledge of the pathogen's biology, and an awareness of the ecological factors influencing the disease. Consequently, they assist in determining the optimal timing for

implementing disease control measures. Moreover, diagnostic assays play a crucial role in programs aimed at producing plant propagation materials that are free from viruses. While molecular diagnostic tools are highly specific and sensitive, it's essential to complement them with other methods, such as traditional culture-based techniques or emerging proteomic approaches, which hold promise in revealing insights into pathogenicity and virulence factors. This, in turn, broadens the possibilities for diagnosing and safeguarding crops against diseases.

References:

- Abd El-Aziz, M.H. Three modern serological methods to detect plant viruses. *J. Plant. Sci. Phytopathol.* 2019, 3, 101–106.
- Balodi, R.; Bisht, S.; Ghatak, A.; Rao, K.H. Plant disease diagnosis: Technological advancements and challenges. *Indian Phytopathol.* 2017, 70, 275–281.
- Bar-Joseph M, Moscovitz M, Sharafi Y. Reuse of coated enzyme-linked immunosorbent assay plates. *Phytopathology.* 1979; 69: 424-426.
- Bock, K.R. (1982). The identification and partial characterization of plant viruses in the tropics. *Tropical Pest Management* 28: 399–411.
- Bodewes, R.; Lapp, S.; Hahn, K.; Habierski, A.; Förster, C.; König, M.; Wohlsein, P.; Osterhaus, A.D.; Baumgärtner, W. Novel canine bocavirus strain associated with severe enteritis in a dog litter. *Vet. Microbiol.* 2014, 174, 1–8.
- BronzatoBadial, A., Sherman, D., Stone, A., Gopakumar, A., Wilson, V., Schneider, W., et al. (2018). Nanopore sequencing as a surveillance tool for plant pathogens in plant and insect tissues. *Plant Dis.* 102, 1648–1652. doi: 10.1094/PDIS-04-17-0488-RE.

Chalupowicz, L., Dombrovsky, A., Gaba, V., Luria, N., Reuven, M., Beerman, A., et al. (2019). Diagnosis of plant diseases using the Nanopore sequencing platform. *Plant Pathol.* 68, 229–238. doi: 10.1111/ppa.12957.

Chertow, D. S. (2018). Next-generation diagnostics with CRISPR. *science*, 360(6387), 381-382.

Doudna, J.A. and Charpentier, E. (2014) The new frontier of genome engineering with CRISPR- Cas9. *Science*, 346, 1258096.

Fegla GI, El-Samra IA, Younes HA, Abd El-Aziz MH. Comparative studies for detection of Tomato Mosaic Tobamovirus (ToMV), Cucumber Mosaic Cucumovirus (CMV) and Potato Y Potyvirus (PVY). *Adv Agric Res.* 2001; 6: 239-254.

Francki, R.I.B. (1980). Limited value of the thermal inactivation point, longevity in vitro and dilution end point as criteria for the characterization, identification, and classification of plant viruses. *Intervirology* 13: 91–98.

Gentile, M., and Gelderblom, H. R. (2014). Electron microscopy in rapid viral diagnosis: an update. *New Microbiol.* 37, 403–422.

Goodwin, S., McPherson, J. D., McCombie, W. R. (2016). Coming of age: ten years of next-generation sequencing technologies. *Nat. Rev. Genet.* 17, 333–351. doi: 10.1038/nrg.2016.49.

Guo, L.; Wu, J.; Chen, R.; Hong, J.; Zhou, X.; Wu, J. Monoclonal antibody-based serological detection of Rice stripe mosaic virus infection in rice plants or Leafhoppers. *Viol. Sin.* 2020, 35, 227–234.

Hadidi, A., Flores, R., Candresse, T., Barba, M. (2016). Next-generation sequencing and genome editing in plant virology. *Front. Microbiol.* 7, 1325. doi: 10.3389/fmicb.2016.01325.

Hawkes R, Niday E, Gordon J. A dot-immunobinding assay for monoclonal and other antibodies. *Anal Biochemical.* 1982; 119: 142-147. Hawkes R, Niday E, Gordon J. A dot-immunobinding assay for monoclonal and other antibodies. *Anal Biochemical.* 1982; 119: 142-147.

Jinek, M. , Chylinski, K. , Fonfara, I. , Hauer, M. , Doudna, J.A. and Charpentier, E. (2012) A programmable dual- RNA- guided DNA endonuclease in adaptive bacterial immunity. *Science*, 337, 816–821.

Kehoe, M. A., Coutts, B. A., Buirchell, B. J., Jones, R. A. C. (2014). Plant virology and next generation sequencing: experiences with a potyvirus. *PLoS One* 9, e104580. doi: 10.1371/journal.pone.0104580.

Kfir, R. and Genthe, B. 1993. Advantages and disadvantages of the use of immunodetection techniques for the enumeration of microorganisms and toxins in water. *Water Sci. Technol.* 27: 243-252.

King, A. M. Q., Lefkowitz, E., Adams, M. J., and Carstens, E. B. (2012). *Ninth Report of the International Committee on Taxonomy of Viruses*. Elsevier.

Kiselev, D., Matsvay, A., Abramov, I., Dedkov, V., Shipulin, G., Khafizov, K. (2020). Current trends in diagnostics of viral infections of unknown etiology. *Viruses* 12, 212. doi: 10.3390/v12020211.

Klap, C.; Luria, N.; Smith, E.; Bakelman, E.; Belausov, E.; Laskar, O.; Lachman, O.; Gal-On, A.; Dombrovsky, A. The potential risk of plant-virus disease initiation by infected tomatoes. *Plants* 2020, 9, 623.

Klap, C.; Luria, N.; Smith, E.; Bakelman, E.; Belausov, E.; Laskar, O.; Lachman, O.; Gal-On, A.; Dombrovsky, A. The potential risk of plant-virus disease initiation by infected tomatoes. *Plants* 2020, 9, 623.

Lina Katul V, Naiss HJ, Makkouk KM, Leemann DE, Casper R. Characterization and serology of virus-like particles associated with faba bean necrotic yellows. *Ann Appl Biol.* 1993; 123: 629-647.

Lo´Pez, M. M., Bertolini, E., Olmos, A., Caruso, P., Gorris, M. T., Llop, P., Penyalver, R. and Cambra, M. (2003). Innovative tools for detection of plant pathogenic viruses and bacteria. *Int Microbiol* 6: 233–243.

Mahas, A. , Aman, R. and Mahfouz, M. (2019) CRISPR- Cas13d mediates robust RNA virus interference in plants. *Genome Biol.* 20, 263.

Mahas, A. , Hassan, N. , Aman, R. , Marsic, T. , Wang, Q. , Ali, Z. and Mahfouz, M.M. (2021) LAMP- coupled CRISPR- Cas12a module for rapid and sensitive detection of plant DNA viruses. *Viruses*, 13, 466.

Mahas, A. and Mahfouz, M. (2018) Engineering virus resistance via CRISPR- Cas systems. *Curr. Opin. Virol.* 32, 1–8.

Maksimov, I.V.; Sorokan, A.V.; Burkhanova, G.F.; Veselova, S.V.; Alekseev, V.Y.; Shein, M.Y.; Avalbaev, A.M.; Dhaware, P.D.; Mehetre, G.T.; Singh, B.P.; et al. Mechanisms of plant tolerance to RNA viruses induced by plant-growth-promoting microorganisms. *Plants* 2019, 8, 575.

Maree, H. J., Fox, A., Al Rwahnih, M., Boonham, N., Candresse, T. (2018). Application of HTS for routine plant virus diagnostics: state of the art and challenges. *Front. Plant Sci.* 9, 1082. doi: 10.3389/fpls.2018.01082.

Massart, S.; Olmos, A.; Jijakli, H.; Candresse, T. Current impact and future directions of high throughput sequencing in plant virus diagnostics. *Virus Res.* 2014, 188, 90–96.

Matthews, R.E.F. (1980). Host plant responses to virus infection. Pp 297–359. In: H. Fraenkel-Conrat and Wagner, R.R. (eds.). *Comprehensive virology*, vol. 16, virus-host interaction, viral invasion, persistence, and diagnosis, Plenum Press, New York, USA.

Mullis, K. B., & Faloona, F. A. (1987). [21] Specific synthesis of DNA in vitro via a polymerase-catalyzed chain reaction. In *Methods in enzymology* (Vol. 155, pp. 335-350). Academic Press.

Naito, F. Y. B., Melo, F. L., Fonseca, M. E. N., Santos, C. A. F., Chanes, C. R., Ribeiro, B. M., et al. (2019). Nanopore sequencing of a novel bipartite New World begomovirus infecting cowpea. *Arch. Virol.* 164, 1907–1910. doi: 10.1007/s00705-019-04254-5.

Notomi T, Okayama H, Masubuchi H, Yonekawa T, Watanabe K, Amino N, Hase T. Loop-mediated isothermal amplification of DNA. *Nucleic Acids Res.* 2000;28(12):E63.

Olmos, A., Capote, N., Bertolini, E., & Cambra, M. (2007). Molecular diagnostic methods for plant viruses. In *Biotechnology and plant disease management* (pp. 227-249). Wallingford UK: CAB International.

Pecman, A., Kutnjak, D., Gutiérrez-Aguirre, I., Adams, I., Fox, A., Boonham, N., et al. (2017). Next generation sequencing for detection and discovery of plant viruses and viroids: comparison of two approaches. *Front. Microbiol.* 8 1998. doi: 10.3389/fmicb.2017.01998.

Pfankuche, V.M.; Bodewes, R.; Hahn, K.; Puff, C.; Beineke, A.; Habierski, A.; Osterhaus, A.D.; Baumgärtner, W. Porcine bocavirus infection associated with encephalomyelitis in a pig, Germany. *Emerg. Infect. Dis.* 2016, 22, 1310–1312.

Rani, A.; Donovan, N.; Mantri, N. The future of plant pathogen diagnostics in a nursery production system. *Biosens. Bioelectron.* 2019, 145, 111631.

Rojas, M.R.; Gilbertson, R.L. Emerging plant viruses: A diversity of mechanisms and opportunities. In *Plant Virus Evolution*; Roossinck, M.J., Ed.; Springer: Berlin/Heidelberg, Germany, 2008; pp. 27–51.

Roossinck, M. J. (2015). Metagenomics of plant and fungal viruses reveals an abundance of persistent lifestyles. *Front. Microbiol.* 5, 767. doi: 10.3389/fmicb.2014.00767.

Rubio, L.; Galipienso, L.; Ferriol, I. Detection of plant viruses and disease management: Relevance of genetic diversity and evolution. *Front. Plant. Sci.* 2020, 11, 1092.

Scholthof, K.B.; Adkins, S.; Czosnek, H.; Palukaitis, P.; Jacquot, E.; Hohn, T.; Hohn, B.; Saunders, K.; Candresse, T.; Ahlquist, P.; et al. Top 10 plant viruses in molecular plant pathology. *Mol. Plant Pathol.* 2011, 12, 938–954. [CrossRef] [PubMed].

Seo, J.-K.; Kim, M.-K.; Kwak, H.-R.; Choi, H.-S.; Nam, M.; Choe, J.; Choi, B.; Han, S.-J.; Kang, J.-H.; Jung, C. Molecular dissection of distinct symptoms induced by tomato chlorosis virus and tomato yellow leaf curl virus based on comparative transcriptome analysis. *Virology* 2018, 516, 1–20.

Song, G.; Wu, J.Y.; Xie, Y.; Liu, Y.; Qian, Y.J.; Zhou, X.P.; Wu, J.X. Monoclonal antibody-based serological assays for detection of Potato virus S in potato plants. *J. Zhejiang Univ. Sci. B* 2017, 18, 1075–1082.

Stobbe, A.H.; Roossinck, M.J. Plant virus metagenomics: What we know and why we need to know more. *Front. Plant Sci.* 2014, 5, 150.

Tan, S. H., Osman, F., Bodaghi, S., Dang, T., Greer, G., Huang, A., et al. (2019). Full genome characterization of 12 citrus tatter leaf virus isolates for the development of a detection assay. *PLoS One* 14, e0223958. doi: 10.1371/journal.pone.0223958.

Van Brunschot, S. L., Bergervoet, J. H. W., Pagendam, D. E., de Weerd, M., Geering, A. D. W., Drenth, A., et al. (2014). A bead-based suspension array for the multiplexed detection of begomoviruses and their whitefly vectors. *J. Virol. Methods* 198, 86–94. doi: 10.1016/j.jviromet.2013.12.014.

Vet, J. A. M., Majithia, A. R., Marras, S. A. E., Tyagi, S., Dube, S., Poiesz, B. J., et al. (1999). Multiplex detection of four pathogenic retroviruses using molecular beacons. *Proc. Natl. Acad. Sci. U.S.A.* 96, 6394–6399. doi: 10.1073/pnas.96.11.6394.

Villamor, D. E. V., Ho, T., Al Rwahnih, M., Martin, R. R., Tzanetakis, I. E. (2019). High throughput sequencing for plant virus detection and discovery. *Phytopathology* 109, 716–725. doi: 10.1094/PHYTO-07-18-0257-RVW.

Xie, Y.; Jiao, X.; Zhou, X.; Liu, H.; Ni, Y.; Wu, J. Highly sensitive serological methods for detecting tomato yellow leaf curl virus in tomato plants and whiteflies. *Virol. J.* 2013, 10, 142.

Xu, Y., Li, S., Na, C., Yang, L., Lu, M. (2019). Analyses of virus/viroid communities in nectarine trees by next-generation sequencing and insight into viral synergisms implication in host disease symptoms. *Sci. Rep.* 9, 12261. doi: 10.1038/s41598-019-48714-z.

Zhang, M.; Ye, J.; He, J.S.; Zhang, F.; Ping, J.; Qian, C.; Wu, J. Visual detection for nucleic acid-based techniques as potential on-site detection methods. A review. *Anal. Chim. Acta* 2020, 1099, 1–15.