

# Gaussian Generalized Woodall Numbers

In this work, we define and investigate Gaussian generalized Woodall numbers in detail, and focus on four specific cases: Gaussian modified Woodall numbers, Gaussian modified Cullen numbers, Gaussian Woodall numbers, and Gaussian Cullen numbers. We present some identities and matrices related to these sequences, as well as recurrence relations, Binet's formulas, generating functions, Simpson's formulas, and summation formulas.

**Keywords.** Woodall numbers, Cullen numbers, Gaussian Woodall numbers, Gaussian Cullen numbers, Gaussian generalized Woodall numbers, Gaussian modified Woodall numbers, Gaussian modified Cullen numbers.

## 1. Introduction

First, we recall some properties of generalized Woodall numbers. The generalized Woodall sequence  $\{W_n\}_{n \geq 0} = \{W_n(W_0, W_1, W_2, 5, -8, 4)\}_{n \geq 0}$  is defined by the third-order recurrence relation as

$$W_n = 5W_{n-1} - 8W_{n-2} + 4W_{n-3} \quad (1.1)$$

with the initial values  $W_0, W_1, W_2$  not all being zero.

A generalized Woodall sequence  $\{W_n\}_{n \geq 0} = \{W_n(W_0, W_1, W_2, 5, -8, 4)\}_{n \geq 0}$  can be extended to negative subscripts by defining

$$W_{-n} = 2W_{-(n-1)} - \frac{5}{4}W_{-(n-2)} + \frac{1}{4}W_{-(n-3)}$$

for  $n = 1, 2, 3, \dots$ . Therefore, recurrence (1.1) holds for all integer  $n$ . For more details, see [48].

Next, we give Binet formula of generalized Woodall numbers.

**THEOREM 1.1.** [48, Theorem 1.1] *Binet formula of generalized Woodall numbers can be given as*

$$W_n = (A_1 + A_2n) \times 2^n + A_3$$

where  $A_1$ ,  $A_2$  and  $A_3$  are defined by

$$\begin{aligned} A_1 &= -W_2 + 4W_1 - 3W_0, \\ A_2 &= \frac{W_2 - 3W_1 + 2W_0}{2}, \\ A_3 &= W_2 - 4W_1 + 4W_0, \end{aligned}$$

that is,

$$W_n = ((-W_2 + 4W_1 - 3W_0) + \frac{W_2 - 3W_1 + 2W_0}{2}n) \times 2^n + (W_2 - 4W_1 + 4W_0). \quad (1.2)$$

Here,  $\alpha, \beta$  and  $\gamma$  are the roots of the cubic equation

$$x^3 - 5x^2 + 8x - 4 = (x - 2)^2 (x - 1) = 0,$$

where

$$\begin{aligned} \alpha &= \beta = 2, \\ \gamma &= 1. \end{aligned}$$

Now, the first few generalized Woodall numbers with positive subscript and negative subscript are given in the following table.

Table 1. The first few generalized Woodall numbers with positive subscript and negative subscript.

$n$	$W_n$	$W_{-n}$
0	$W_0$	$W_0$
1	$W_1$	$\frac{1}{4}(8W_0 - 5W_1 + W_2)$
2	$W_2$	$\frac{1}{4}(11W_0 - 9W_1 + 2W_2)$
3	$4W_0 - 8W_1 + 5W_2$	$\frac{1}{16}(52W_0 - 47W_1 + 11W_2)$
4	$20W_0 - 36W_1 + 17W_2$	$\frac{1}{16}(57W_0 - 54W_1 + 13W_2)$
5	$68W_0 - 116W_1 + 49W_2$	$\frac{1}{64}(240W_0 - 233W_1 + 57W_2)$
6	$196W_0 - 324W_1 + 129W_2$	$\frac{1}{64}(247W_0 - 243W_1 + 60W_2)$

Now, we define four specific cases of the sequence  $\{W_n\}$ .

The Woodall numbers  $\{R_n\}$ , sometimes called Riesel numbers, and also called Cullen numbers of the second kind, are numbers of the form

$$R_n = n \times 2^n - 1.$$

The first few Woodall numbers are:

$$1, 7, 23, 63, 159, 383, 895, 2047, 4607, 10239, 22527, 49151, 106495, 229375, 491519, 1048575, \dots$$

(sequence A003261 in the OEIS [43]). Woodall numbers were first studied by Allan J. C. Cunningham and H. J. Woodall in [12] in 1917, inspired by James Cullen's earlier study of the similarly-defined Cullen numbers.

The Cullen numbers  $\{C_n\}$  are numbers of the form

$$C_n = n \times 2^n + 1.$$

The first few Cullen numbers are:

$$1, 3, 9, 25, 65, 161, 385, 897, 2049, 4609, 10241, 22529, 49153, 106497, 229377, 491521, \dots$$

(sequence A002064 in the OEIS).

Woodall and Cullen sequences have been studied by many authors and more detail can be found in the extensive literature dedicated to these sequences, see for example, [7,8,12,21,23,25,28,33,34,35,39] and references therein.

Note that  $\{R_n\}$  and  $\{C_n\}$  hold the following relations:

$$R_n = 4R_{n-1} - 4R_{n-2} - 1,$$

$$C_n = 4C_{n-1} - 4C_{n-2} + 1.$$

Note also that the sequences  $\{R_n\}$  and  $\{C_n\}$  satisfy the following third order linear recurrences:

$$R_n = 5R_{n-1} - 8R_{n-2} + 4R_{n-3}, \quad R_0 = -1, R_1 = 1, R_2 = 7, \quad (1.3)$$

$$C_n = 5C_{n-1} - 8C_{n-2} + 4C_{n-3}, \quad C_0 = 1, C_1 = 3, C_2 = 9. \quad (1.4)$$

Modified Woodall sequence  $\{G_n\}_{n \geq 0}$  and modified Cullen sequence  $\{H_n\}_{n \geq 0}$  are defined, respectively, by the third-order recurrence relations,

$$G_n = 5G_{n-1} - 8G_{n-2} + 4G_{n-3}, \quad G_0 = 0, G_1 = 1, G_2 = 5, \quad (1.5)$$

$$H_n = 5H_{n-1} - 8H_{n-2} + 4H_{n-3}, \quad H_0 = 3, H_1 = 5, H_2 = 9. \quad (1.6)$$

The sequences  $\{G_n\}_{n \geq 0}$ ,  $\{H_n\}_{n \geq 0}$ ,  $\{R_n\}_{n \geq 0}$  and  $\{C_n\}_{n \geq 0}$  can be extended to negative subscripts by defining,

$$G_{-n} = 2G_{-(n-1)} - \frac{5}{4}G_{-(n-2)} + \frac{1}{4}G_{-(n-3)},$$

$$H_{-n} = 2H_{-(n-1)} - \frac{5}{4}H_{-(n-2)} + \frac{1}{4}H_{-(n-3)},$$

$$R_{-n} = 2R_{-(n-1)} - \frac{5}{4}R_{-(n-2)} + \frac{1}{4}R_{-(n-3)},$$

$$C_{-n} = 2C_{-(n-1)} - \frac{5}{4}C_{-(n-2)} + \frac{1}{4}C_{-(n-3)},$$

for  $n = 1, 2, 3, \dots$  respectively. Therefore, recurrences (1.3), (1.4), (1.5) and (1.6) hold for all integer  $n$ .

For all integers  $n$ , modified Woodall, modified Cullen, Woodall and Cullen numbers (using initial conditions in (1.2)) can be expressed using Binet's formulas as

$$\begin{aligned} G_n &= (n-1)2^n + 1, \\ H_n &= 2^{n+1} + 1, \\ R_n &= n \times 2^n - 1, \\ C_n &= n \times 2^n + 1, \end{aligned}$$

respectively.

Now we give some information about Gaussian sequence from literature.

- First we give Gaussian numbers with second order recurrence
  - Horadam [26] introduced Gaussian Fibonacci numbers as

$$GF_n = F_n + iF_{n-1}$$

where  $F_n = F_{n-1} + F_{n-2}$ ,  $F_0 = 0$ ,  $F_1 = 1$  (in fact, he defined these numbers as  $GF_n = F_n + iF_{n-1}$  and he called these numbers as complex Fibonacci numbers.)

- Pethe and Horadam [36] introduced generalized Gaussian Fibonacci numbers

$$GF_n = F_n + iF_{n-1}$$

where  $F_n = F_{n-1} + F_{n-2}$ ,  $F_0 = 0$ ,  $F_1 = 1$ .

- Halıcı and Öz [24] studied Gaussian Pell and Pell Lucas numbers by written, respectively,

$$\begin{aligned} GP_n &= P_n + iP_{n-1}, \\ GQ_n &= Q_n + iQ_{n-1}, \end{aligned}$$

where  $P_n = 2P_{n-1} + P_{n-2}$ ,  $P_0 = 0$ ,  $P_1 = 1$  and  $Q_n = 2Q_{n-1} + Q_{n-2}$ ,  $Q_0 = 2$ ,  $Q_1 = 2$ .

- Aşçı and Gürel [1] presented Gaussian Jacobsthal and Gaussian Jacobsthal Lucas numbers given by, respectively,

$$\begin{aligned} GJ_n &= J_n + iJ_{n-1}, \\ Gj_n &= j_n + j_{n-1}, \end{aligned}$$

where  $J_n = J_{n-1} + 2J_{n-2}$ ,  $J_0 = 0$ ,  $J_1 = 1$  and  $j_n = j_{n-1} + 2j_{n-2}$ ,  $j_0 = 2$ ,  $j_1 = 1$ .

- Taşcı [68] introduced and studied Gaussian Mersenne numbers and define by

$$GM_n = M_n + iM_{n-1}$$

where  $M_n = 3M_{n-1} - 2M_{n-2}$ ,  $M_0 = 0$ ,  $M_1 = 1$ .

- Taşçı [66] introduced and studied Gaussian balancing and Lucas Balancing numbers and given by, respectively,

$$\begin{aligned} GB_n &= B_n + iB_{n-1}, \\ GC_n &= C_n + C_{n-1}, \end{aligned}$$

where  $B_n = 6B_{n-1} - B_{n-2}$ ,  $B_0 = 0$ ,  $B_1 = 1$  and  $C_n = 6C_{n-1} - C_{n-2}$ ,  $C_0 = 1$ ,  $C_1 = 3$ .

- Ertaş and Yılmaz [2] studied Gaussian Oresme numbers and given by

$$GT_n = T_n + iT_{n-1}$$

where  $T_n = T_{n-1} - \frac{1}{4}T_{n-2}$ ,  $T_0 = 0$ ,  $T_1 = \frac{1}{2}$ .

- Now, we present Gaussian numbers with third order recurrence relations.

- Soykan, Taşdemir, Okumuş and Göcen [45] presented Gaussian generalized Tribonacci numbers by given

$$GW_n = W_n + iW_{n-1}$$

where  $W_n = W_{n-1} + W_{n-2} + W_{n-3}$ , with the initial condition  $W_0, W_1, W_2$ .

- Taşçı [67] studied Gaussian Padovan and Gaussian Pell- Padovan numbers by written, respectively,

$$\begin{aligned} GP_n &= P_n + iP_{n-1}, \\ GR_n &= R_n + iR_{n-1}, \end{aligned}$$

where  $P_n = P_{n-2} + P_{n-3}$ ,  $P_0 = 1$ ,  $P_1 = 1$ ,  $P_2 = 1$ , and  $R_n = 2R_{n-2} + R_{n-3}$ ,  $R_0 = 1$ ,  $R_1 = 1$ ,  $R_2 = 1$ .

- Cerda-Morales [9] defined Gaussian third-order Jacobsthal numbers by given

$$GJ_n = J_n + iJ_{n-1}$$

where  $J_n = J_{n-1} + J_{n-2} + 2J_{n-3}$ ,  $J_1 = 0$ ,  $J_2 = 1$ ,  $J_3 = 1$ .

## 2. Gaussian Generalized Woodall Numbers

In this chapter, we define Gaussian generalized Woodall numbers and we give some properties.

Gaussian generalized Woodall numbers  $\{GW_n\}_{n \geq 0} = \{GW_n(GW_0, GW_1, GW_2)\}_{n \geq 0}$  are defined by

$$GW_n = 5GW_{n-1} - 8GW_{n-2} + 4GW_{n-3}, \quad (2.1)$$

with the initial conditions

$$GW_0 = W_0 + i\left(\frac{1}{4}(8W_0 - 5W_1 + W_2)\right), \quad GW_1 = W_1 + iW_0, \quad GW_2 = W_2 + iW_1,$$

not all being zero. The sequences  $\{GW_n\}_{n \geq 0}$  can be extended to negative subscripts by defining

$$GW_{-n} = 2GW_{-(n-1)} - \frac{5}{4}GW_{-(n-2)} + \frac{1}{4}GW_{-(n-3)}$$

for  $n = 1, 2, 3, \dots$ . Therefore, recurrence (2.1) hold for all integer  $n$ . Note that for  $n \geq 0$ , we get

$$GW_n = W_n + iW_{n-1} \quad (2.2)$$

and

$$GW_{-n} = W_{-n} + iW_{-n-1}. \quad (2.3)$$

The first few generalized Gaussian Woodall numbers with positive subscript and negative subscript are given in the following table.

Table 2. The first few generalized Gaussian Woodall numbers.

$n$	$GW_n$	$GW_{-n}$
0	$W_0 + i\frac{1}{4}(8W_0 - 5W_1 + W_2)$	$W_0 + i\frac{1}{4}(8W_0 - 5W_1 + W_2)$
1	$W_1 + iW_0$	$\frac{1}{4}(8W_0 - 5W_1 + W_2) + i\frac{1}{4}(11W_0 - 9W_1 + 2W_2)$
2	$W_2 + iW_1$	$\frac{1}{4}(11W_0 - 9W_1 + 2W_2) + i\frac{1}{16}(52W_0 - 47W_1 + 11W_2)$
3	$4W_0 - 8W_1 + 5W_2 + iW_2$	$\frac{1}{16}(52W_0 - 47W_1 + 11W_2) + i\frac{1}{16}(57W_0 - 54W_1 + 13W_2)$

We consider four special cases of  $GW_n$  :

$GW_n(0, 1, 5 + i) = GG_n$  is the sequence of Gaussian Modified Woodall numbers,

$GW_n(3 + 2i, 5 + 3i, 9 + 5i) = GH_n$  is the sequence of Gaussian Modified Cullen numbers,

$GW_n(-1 - \frac{3}{2}i, 1 - i, 7 + i) = GR_n$  is the sequence of Gaussian Woodall numbers and

$GW_n(1 + \frac{1}{2}i, 3 + i, 9 + 3i) = GC_n$  is the sequence of Gaussian Cullen numbers.

We formally define them as follows. Four special cases of  $GW_n$  with the initial conditions are defined by

$$GG_n = 5GG_{n-1} - 8GG_{n-2} + 4GG_{n-3}, \quad GG_0 = 0, GG_1 = 1, GG_2 = 5 + i,$$

$$GH_n = 5GH_{n-1} - 8GH_{n-2} + 4GH_{n-3}, \quad GH_0 = 3 + 2i, GH_1 = 5 + 3i, GH_2 = 9 + 5i,$$

$$GR_n = 5GR_{n-1} - 8GR_{n-2} + 4GR_{n-3}, \quad GR_0 = -1 - \frac{3}{2}i, GR_1 = 1 - i, GR_2 = 7 + i,$$

$$GC_n = 5GC_{n-1} - 8GC_{n-2} + 4GC_{n-3}, \quad GC_0 = 1 + \frac{1}{2}i, GC_1 = 3 + i, GC_2 = 9 + 3i.$$

Note that for all integers  $n$ , we obtain

$$GG_n = G_n + iG_{n-1},$$

$$GH_n = H_n + iH_{n-1},$$

$$GR_n = R_n + iR_{n-1},$$

$$GC_n = C_n + iC_{n-1}.$$

The first few values of Gaussian Modified Woodall numbers, Gaussian Modified Cullen numbers, Gaussian Woodall numbers and Gaussian Cullen numbers with positive and negative subscript are given in the following table.

Table 3. The first few values of special cases of generalized Gaussian Woodall numbers.

$n$	0	1	2	3	4	5	6	7
$GG_n$	0	1	$5 + i$	$17 + 5i$	$49 + 17i$	$129 + 49i$	$321 + 129i$	$769 + 321i$
$GG_{-n}$	0	$\frac{1}{4}i$	$\frac{1}{4} + \frac{1}{2}i$	$\frac{1}{2} + \frac{11}{16}i$	$\frac{11}{16} + \frac{13}{16}i$	$\frac{13}{16} + \frac{57}{64}i$	$\frac{57}{64} + \frac{15}{16}i$	$\frac{15}{16} + \frac{247}{256}i$
$GH_n$	$3 + 2i$	$5 + 3i$	$9 + 5i$	$17 + 9i$	$33 + 17i$	$65 + 33i$	$129 + 65i$	$257 + 129i$
$GH_{-n}$	$3 + 2i$	$2 + \frac{3}{2}i$	$\frac{3}{2} + \frac{5}{4}i$	$\frac{5}{4} + \frac{9}{8}i$	$\frac{9}{8} + \frac{17}{16}i$	$\frac{17}{16} + \frac{33}{32}i$	$\frac{33}{32} + \frac{65}{64}i$	$\frac{65}{64} + \frac{129}{128}i$
$GR_n$	$-1 - \frac{3}{2}i$	$1 - i$	$7 + i$	$23 + 7i$	$63 + 23i$	$159 + 63i$	$383 + 159i$	$895 + 383i$
$GR_{-n}$	$-1 - \frac{3}{2}i$	$-\frac{3}{2} - \frac{3}{2}i$	$-\frac{3}{2} - \frac{11}{8}i$	$\frac{11}{8} - \frac{5}{4}i$	$-\frac{5}{4} - \frac{37}{32}i$	$-\frac{37}{32} - \frac{35}{32}i$	$-\frac{35}{32} - \frac{135}{128}i$	$\frac{135}{128} - \frac{33}{32}i$
$GC_n$	$1 + \frac{1}{2}i$	$3 + i$	$9 + 3i$	$25 + 9i$	$65 + 25i$	$161 + 65i$	$385 + 161i$	$897 + 385i$
$GC_{-n}$	$1 + \frac{1}{2}i$	$\frac{1}{2} + \frac{1}{2}i$	$\frac{1}{2} + \frac{5}{8}i$	$\frac{5}{8} + \frac{3}{4}i$	$\frac{3}{4} + \frac{27}{32}i$	$\frac{27}{32} + \frac{29}{32}i$	$\frac{29}{32} + \frac{121}{128}i$	$\frac{121}{128} + \frac{31}{32}i$

We now present the Binet formula for the Gaussian generalized Woodall numbers.

**THEOREM 2.1.** *The Binet's formula for the Gaussian generalized Woodall numbers is*

$$GW_n = (((-W_2 + 4W_1 - 3W_0) + \frac{W_2 - 3W_1 + 2W_0}{2}n)2^n + (W_2 - 4W_1 + 4W_0)) + i((( -W_2 + 4W_1 - 3W_0) + \frac{W_2 - 3W_1 + 2W_0}{2}(n - 1))2^{n-1} + (W_2 - 4W_1 + 4W_0)).$$

*Proof.* The proof follows from (1.2) and (2.2).  $\square$

The previous Theorem gives the following results, as special cases.

**COROLLARY 2.2.** *For all  $n$  we have the following Binet's Formulas*

- (a):  $GG_n = i2^{n-1}(n - 2) + 2^n(n - 1) + 1 + i.$
- (b):  $GH_n = 2i2^{n-1} + 2 \times 2^n + 1 + i.$
- (c):  $GR_n = i2^{n-1}(n - 1) + 2^n n - 1 - i.$
- (d):  $GC_n = i2^{n-1}(n - 1) + 2^n n + 1 + i.$

The following Theorem presents the generating function of Gaussian generalized Woodall numbers.

**THEOREM 2.3.** *The generating function of Gaussian generalized Woodall numbers is given as*

$$f_{GW_n}(x) = \sum_{n=0}^{\infty} GW_n x^n = \frac{GW_0 + (GW_1 - 5GW_0)x + (GW_2 - 5GW_1 + 8GW_0)x^2}{1 - 5x + 8x^2 - 4x^3}. \tag{2.4}$$

*Proof.* Let

$$f_{GW_n}(x) = \sum_{n=0}^{\infty} GW_n x^n$$

be generating function of Gaussian generalized Woodall numbers. Then using the definition of Gaussian Woodall numbers, and subtracting  $xf(x)$ ,  $x^2f(x)$  and  $x^3f(x)$  from  $f(x)$  we obtain (note the shift in the

index  $n$  in the third line)

$$\begin{aligned}
(1 - 5x + 8x^2 - 4x^3)f_{GW_n}(x) &= \sum_{n=0}^{\infty} GW_n x^n - 5x \sum_{n=0}^{\infty} GW_n x^n + 8x^2 \sum_{n=0}^{\infty} GW_n x^n - 4x^3 \sum_{n=0}^{\infty} GW_n x^n, \\
&= \sum_{n=0}^{\infty} GW_n x^n - 5 \sum_{n=0}^{\infty} GW_n x^{n+1} + 8 \sum_{n=0}^{\infty} GW_n x^{n+2} - 4 \sum_{n=0}^{\infty} GW_n x^{n+3}, \\
&= \sum_{n=0}^{\infty} GW_n x^n - 5 \sum_{n=1}^{\infty} GW_{n-1} x^n + 8 \sum_{n=2}^{\infty} GW_{n-2} x^n - 4 \sum_{n=3}^{\infty} GW_{n-3} x^n, \\
&= (GW_0 + GW_1 x + GW_2 x^2) - 5(GW_0 x + GW_1 x^2) + 8GW_0 x^2 \\
&\quad + \sum_{n=3}^{\infty} (GW_n - 5GW_{n-1} + 8GW_{n-2} - 4GW_{n-3}) x^n, \\
&= GW_0 + GW_1 x + GW_2 x^2 - 5GW_0 x - 5GW_1 x^2 + 8GW_0 x^2, \\
&= GW_0 + (GW_1 - 5GW_0)x + (GW_2 - 5GW_1 + 8GW_0)x^2.
\end{aligned}$$

Now, it follows that

$$f_{GW_n}(x) = \frac{GW_0 + (GW_1 - 5GW_0)x + (GW_2 - 5GW_1 + 8GW_0)x^2}{1 - 5x + 8x^2 - 4x^3}.$$

This completes the proof.  $\square$

The previous Theorem gives the following results as particular examples:

$$f_{GG_n}(x) = \frac{x + ix^2}{1 - 5x + 8x^2 - 4x^3}, \quad (2.5)$$

$$f_{GH_n}(x) = \frac{(8 + 6i)x^2 - (10 + 7i)x + 3 + 2i}{1 - 5x + 8x^2 - 4x^3}, \quad (2.6)$$

$$f_{GR_n}(x) = \frac{-(6 + 6i)x^2 + (6 + \frac{13}{2}i)x - 1 - \frac{3}{2}i}{1 - 5x + 8x^2 - 4x^3}, \quad (2.7)$$

$$f_{GC_n}(x) = \frac{(2 + 2i)x^2 - (2 + \frac{3}{2}i)x + 1 + \frac{1}{2}i}{1 - 5x + 8x^2 - 4x^3}. \quad (2.8)$$

### 3. Some Identities Connecting Gaussian modified Woodall, Gaussian modified Cullen, Gaussian Woodall and Gaussian Cullen Numbers

In this section, we obtain some identities on Gaussian modified Woodall, Gaussian modified Cullen, Gaussian Woodall and Gaussian Cullen numbers.

**THEOREM 3.1.** *The following equations hold for all integer  $n$ .*

$$GH_n = 2GG_{n+2} - 7GG_{n+1} + 6GG_n, \quad (3.1)$$

$$GH_n = 3GG_{n+1} - 10GG_n + 8GG_{n-1}, \quad (3.2)$$

$$GR_n = -2GC_{n+2} + 8GC_{n+1} - 7GC_n,$$

$$GG_n = -\frac{1}{2}GC_{n+2} + \frac{3}{2}GC_{n+1}, \quad (3.3)$$

$$GC_n = -\frac{7}{4}GR_{n+3} + \frac{27}{4}GR_{n+2} - 6GR_{n+1}, \quad (3.4)$$

$$GH_n = -\frac{1}{2}GR_{n+3} + \frac{5}{2}GR_{n+2} - 3GR_{n+1}, \quad (3.5)$$

$$GH_n = 5GG_n - 16GG_{n-1} + 12GG_{n-2}. \quad (3.6)$$

Proof. To proof identity (3.1), we can write

$$GH_n = aGG_{n+2} + bGG_{n+1} + cGG_n$$

and solving the system of equations

$$GH_0 = aGG_2 + bGG_1 + cGG_0,$$

$$GH_1 = aGG_3 + bGG_2 + cGG_1,$$

$$GH_2 = aGG_4 + bGG_3 + cGG_2.$$

We find that  $a = 2$ ,  $b = -7$ ,  $c = 6$ . Or using the relations  $GH_n = H_n + iH_{n-1}$ ,  $GG_n = G_n + iG_{n-1}$  and identity  $H_n = 2G_{n+2} - 7G_{n+1} + 6G_n$ , we obtain the identity (3.1). The others can be found similarly.  $\square$

LEMMA 3.2. Suppose that  $f(x) = \sum_{n=0}^{\infty} a_n x^n$  is the generating function of the sequence  $\{a_n\}_{n \geq 0}$ . Then the generating functions of the sequences  $\{a_{2n}\}_{n \geq 0}$  and  $\{a_{2n+1}\}_{n \geq 0}$  are given as

$$f_{a_{2n}}(x) = \sum_{n=0}^{\infty} a_{2n} x^n = \frac{f(\sqrt{x}) + f(-\sqrt{x})}{2}$$

and

$$f_{a_{2n+1}}(x) = \sum_{n=0}^{\infty} a_{2n+1} x^n = \frac{f(\sqrt{x}) - f(-\sqrt{x})}{2\sqrt{x}}$$

respectively.

The next Theorem presents the generating functions of even and odd-indexed Gaussian generalized Woodall sequences.

THEOREM 3.3. The generating functions of the sequences  $GW_{2n}$  and  $GW_{2n+1}$  are given by

$$f_{GW_{2n}}(x) = \frac{GW_0 - (9GW_0 - GW_2)x + (44GW_0 - 36GW_1 + 8GW_2)x^2}{1 - 9x + 24x^2 - 16x^3}$$

and

$$f_{GW_{2n+1}}(x) = \frac{GW_1 + (4GW_0 - 17GW_1 + 5GW_2)x + (32GW_0 - 20GW_1 + 4GW_2)x^2}{1 - 9x + 24x^2 - 16x^3}$$

respectively.

*Proof.* Both statements are consequences of Lemma (3.2) applied to (2.4) and some lengthy algebraic calculations.  $\square$

The previous theorem gives the following corollaries as particular examples.

COROLLARY 3.4. *We have the followings:*

$$\begin{aligned}
\text{(a): } f_{GR_{2n}}(x) &= -\frac{(24+22i)x^2-(16+\frac{29}{2}i)x+1+\frac{3}{2}i}{1-9x+24x^2-16x^3} \text{ and } f_{GR_{2n+1}}(x) = (x) \frac{-(24+24i)x^2+(14+16i)x+1-i}{1-9x+24x^2-16x^3}. \\
\text{(b): } f_{GC_{2n}}(x) &= \frac{(8+10i)x^2-\frac{3}{2}ix+1+\frac{1}{2}i}{1-9x+24x^2-16x^3} \text{ and } f_{GC_{2n+1}}(x) = \frac{(8+8i)x^2-2x+3+i}{1-9x+24x^2-16x^3}. \\
\text{(c): } f_{GG_{2n}}(x) &= \frac{(4+8i)x^2+(5+i)x}{1-9x+24x^2-16x^3} \text{ and } f_{GG_{2n+1}}(x) = \frac{4ix^2+(8+5i)x+1}{1-9x+24x^2-16x^3}. \\
\text{(d): } f_{GH_{2n}}(x) &= \frac{(24+20i)x^2-(18+13i)x+3+2i}{1-9x+24x^2-16x^3} \text{ and } f_{GH_{2n+1}}(x) = \frac{(32+24i)x^2-(28+18i)x+5+3i}{1-9x+24x^2-16x^3}.
\end{aligned}$$

From Corollary (3.4) we can obtain the following corollary which presents the identities on Gaussian Woodall sequences.

COROLLARY 3.5. *We have the following identities:*

$$\begin{aligned}
\text{(a): } (4+8i)GH_{2n-4}+(5+i)GH_{2n-2} &= (24+20i)GG_{2n-4}-(18+13i)GG_{2n-2}+(3+2i)GG_{2n}. \\
\text{(b): } (4+8i)GH_{2n-3}+(5+i)GH_{2n-1} &= (32+24i)GG_{2n-4}-(28+18i)GG_{2n-2}+(5+3i)GG_{2n}. \\
\text{(c): } -(24+24i)GG_{2n-4}+(14+16i)GG_{2n-2} &+ (1-i)GG_{2n} = (4+8i)GR_{2n-3}+(5+i)GR_{2n-1}. \\
\text{(d): } -(24+24i)GG_{2n-3}+(14+16i)GG_{2n-1} &+ (1-i)GG_{2n+1} = 4iGR_{2n-3}+(8+5i)GR_{2n-1}+ \\
&GR_{2n+1}. \\
\text{(e): } (8+10i)GG_{2n-4}-\frac{3}{2}iGG_{2n-2} &+ (1+\frac{1}{2}i)GG_{2n} = (4+8i)GC_{2n-4}+(5+i)GC_{2n-2}. \\
\text{(f): } (8+10i)GG_{2n-3}-\frac{3}{2}iGG_{2n-1} &+ (1+\frac{1}{2}i)GG_{2n+1} = 4iGC_{2n-4}+(8+5i)GC_{2n-2}+GC_{2n}. \\
\text{(g): } (8+8i)GG_{2n-4}-2GG_{2n-2} &+ (3+i)GG_{2n} = (4+8i)GC_{2n-3}+(5+i)GC_{2n-1}. \\
\text{(h): } (8+8i)GG_{2n-3}-2GG_{2n-1} &+ (3+i)GG_{2n+1} = 4iGC_{2n-3}+(8+5i)GC_{2n-1}+GC_{2n+1}. \\
\text{(i): } -(24+22i)GG_{2n-4}+(16+\frac{29}{2}i)GG_{2n-2} &- (1+\frac{3}{2}i)GG_{2n} = (4+8i)GR_{2n-4}+(5+i)GR_{2n-2}. \\
\text{(j): } -(24+22i)GG_{2n-3}+(16+\frac{29}{2}i)GG_{2n-1} &- (1+\frac{3}{2}i)GG_{2n+1} = 4iGR_{2n-4}+(8+5i)GR_{2n-2}+GR_{2n}. \\
\text{(k): } 4iGH_{2n-4}+(8+5i)GH_{2n-2}+GH_{2n} &= (24+20i)GG_{2n-3}-(18+13i)GG_{2n-1}+(3+2i)GG_{2n+1}. \\
\text{(l): } 4iGH_{2n-3}+(8+5i)GH_{2n-1}+GH_{2n+1} &= (32+24i)GG_{2n-3}-(28+18i)GG_{2n-1}+(5+3i)GG_{2n+1}. \\
\text{(m): } -(24+22i)C(2n-3)+(16+\frac{29}{2}i)C(2n-1) &- (1+\frac{3}{2}i)C(2n+1) = (8+8i)R(2n-4)-2R(2n- \\
&2)+(3+i)R(2n). \\
\text{(n): } -(24+24i)C(2n-4)+(14+16i)C(2n-2) &+ (1-i)C(2n) = (8+10i)R(2n-3)-\frac{3}{2}iR(2n- \\
&1)+(1+\frac{1}{2}i)R(2n+1).
\end{aligned}$$

*Proof.* From (3.4) we obtain

$$((4+8i)x^2+(5+i)x)f_{GH_{2n}} = ((24+20i)x^2-(18+13i)x+3+2i)f_{GG_{2n}}.$$

The LHS (left hand side) is equal to

$$\begin{aligned}
LHS &= ((5+i)x + (4+8i)x^2) \sum_{n=0}^{\infty} GH_{2n}x^n \\
&= (5+i)x \sum_{n=0}^{\infty} GH_{2n}x^n + (4+8i)x^2 \sum_{n=0}^{\infty} GH_{2n}x^n \\
&= (5+i) \sum_{n=0}^{\infty} GH_{2n}x^{n+1} + (4+8i) \sum_{n=0}^{\infty} GH_{2n}x^{n+2} \\
&= (5+i) \sum_{n=1}^{\infty} GH_{2n-2}x^n + (4+8i) \sum_{n=2}^{\infty} GH_{2n-4}x^n \\
&= (5+i)GH_0x \sum_{n=2}^{\infty} GH_{2n-2}x^n + (4+8i) \sum_{n=2}^{\infty} GH_{2n-4}x^n \\
&= (5+i)(3+2i)x + \sum_{n=2}^{\infty} ((4+8i)GH_{2n-4} + (5+i)GH_{2n-2})x^n
\end{aligned}$$

whereas the RHS is

$$\begin{aligned}
RHS &= (3+2i - (18+13i)x + (24+20i)x^2) \sum_{n=0}^{\infty} GG_{2n}x^n \\
&= (3+2i) \sum_{n=0}^{\infty} GG_{2n}x^n - (18+13i)x \sum_{n=0}^{\infty} GG_{2n}x^n + (24+20i)x^2 \sum_{n=0}^{\infty} GG_{2n}x^n \\
&= (3+2i) \sum_{n=0}^{\infty} GG_{2n}x^n - (18+13i) \sum_{n=0}^{\infty} GG_{2n}x^{n+1} + (24+20i) \sum_{n=0}^{\infty} GG_{2n}x^{n+2} \\
&= (3+2i) \sum_{n=0}^{\infty} GG_{2n}x^n - (18+13i) \sum_{n=1}^{\infty} GG_{2n-2}x^n + (24+20i) \sum_{n=2}^{\infty} GG_{2n-4}x^n \\
&= (3+2i)(GG_0 + GG_2x) \sum_{n=2}^{\infty} GG_{2n}x^n - (18+13i)(GG_0x) \sum_{n=2}^{\infty} GG_{2n-2}x^n \\
&\quad + (24+20i) \sum_{n=2}^{\infty} GG_{2n-4}x^n \\
&= (3+2i)(5+i)x + \sum_{n=2}^{\infty} ((24+20i)GG_{2n-4} - (18+13i)GG_{2n-2} + (3+2i)GG_{2n})x^n.
\end{aligned}$$

Compare the coefficients and the proof of the first identity (a) is done. The other identities can be proved similarly.  $\square$

We present an identity related with Gaussian general Woodall numbers and Woodall numbers.

**THEOREM 3.6.** *For all  $n, m \in \mathbb{Z}$ , the following identity holds:*

$$GW_{m+n} = G_{m+1}GW_n + (-8G_m + 4G_{m-1})GW_{n-1} + 4G_mGW_{n-2}. \quad (3.7)$$

*Proof.* First, we assume that  $m, n \geq 0$ . The other cases can be proved similarly. We prove the identity (3.7) by induction on  $m$ . If  $m = 0$  then

$$GW_n = G_1GW_n + (-8G_0 + 4G_{-1})GW_{n-1} + 4G_0GW_{n-2}$$

which is true because  $G_{-1} = 0, G_0 = 0, G_1 = 1$ . Assume that the equality holds for  $m \leq k$ . For  $m = k + 1$ , we have

$$\begin{aligned} GW_{(k+1)+n} &= 5GW_{n+k} - 8GW_{n+k-1} + 4GW_{n+k-2} \\ &= 5(G_{k+1}GW_n + (-8G_k + 4G_{k-1})GW_{n-1} + 4G_kGW_{n-2}) \\ &\quad - 8(G_kGW_n + (-8G_{k-1} + 4G_{k-2})GW_{n-1} + 4G_{k-1}GW_{n-2}) \\ &\quad + 4(G_{k-1}GW_n + (-8G_{k-2} + 4G_{k-3})GW_{n-1} + 4G_{k-2}GW_{n-2}) \\ &= (5G_{k+1} - 8G_k + 4G_{k-1})GW_n + (-8(G_k + G_{k-1} + G_{k-2}) \\ &\quad + 4(G_{k-1} + G_{k-2} + G_{k-3}))GW_{n-1} + 4(G_k + G_{k-1} + G_{k-2})GW_{n-2} \\ &= G_{k+2}GW_n + (-8G_{k+1} + 4G_k)GW_{n-1} + 4G_{k+1}GW_{n-2} \\ &= G_{(k+1)+1}GW_n + (-8G_{k+1} + 4G_{(k+1)-1})GW_{n-1} + 4G_{k+1}GW_{n-2}. \end{aligned}$$

By mathematical induction on  $m$ , this proves (3.6).  $\square$

The previous Theorem gives the following results as particular examples:

For all  $n, m \in \mathbb{Z}$ , we have ( taking  $GW_n = GG_n$  or  $GW_n = GH_n$  or  $GW_n = GR_n$  or  $GW_n = GC_n$  )

$$\begin{aligned} GG_{m+n} &= G_{m+1}GG_n + (-8G_m + 4G_{m-1})GG_{n-1} + 4G_mGG_{n-2}, \\ GH_{m+n} &= G_{m+1}GH_n + (-8G_m + 4G_{m-1})GH_{n-1} + 4G_mGH_{n-2}, \\ GR_{m+n} &= G_{m+1}GR_n + (-8G_m + 4G_{m-1})GR_{n-1} + 4G_mGR_{n-2}, \\ GC_{m+n} &= G_{m+1}GC_n + (-8G_m + 4G_{m-1})GC_{n-1} + 4G_mGC_{n-2}. \end{aligned}$$

#### 4. Simpson's Formula

In this chapter, we present Simpson's formula of generalized Gaussian Woodall numbers.

**THEOREM 4.1.** (*Simpson's formula of generalized Gaussian Woodall numbers*). For all integers  $n$ , we have

$$\begin{aligned} \begin{vmatrix} GW_{n+2} & GW_{n+1} & GW_n \\ GW_{n+1} & GW_n & GW_{n-1} \\ GW_n & GW_{n-1} & GW_{n-2} \end{vmatrix} &= 4^n \begin{vmatrix} GW_2 & GW_1 & GW_0 \\ GW_1 & GW_0 & GW_{-1} \\ GW_0 & GW_{-1} & GW_{-2} \end{vmatrix} \\ &= 4^n \left( \frac{1}{800} + \frac{7}{800}i \right) (W_0 - W_1 + \frac{1}{4}W_2) \left( (2 - 14i)W_0 - (3 - 21i)W_1 \right. \\ &\quad \left. + (1 - 7i)W_2 \right)^2. \end{aligned}$$

Proof. Use [46, Theorem 3.1].  $\square$

From the Theorem (4.1) we get the following corollary.

COROLLARY 4.2. *For all integer  $n$ , we get the following identities.*

$$\begin{aligned}
 \text{(a): } & \begin{vmatrix} GG_{n+2} & GG_{n+1} & GG_n \\ GG_{n+1} & GG_n & GG_{n-1} \\ GG_n & GG_{n-1} & GG_{n-2} \end{vmatrix} = (1-7i)2^{2n-4}. \\
 \text{(b): } & \begin{vmatrix} GH_{n+2} & GH_{n+1} & GH_n \\ GH_{n+1} & GH_n & GH_{n-1} \\ GH_n & GH_{n-1} & GH_{n-2} \end{vmatrix} = 0. \\
 \text{(c): } & \begin{vmatrix} GR_{n+2} & GR_{n+1} & GR_n \\ GR_{n+1} & GR_n & GR_{n-1} \\ GR_n & GR_{n-1} & GR_{n-2} \end{vmatrix} = -(1-7i)2^{2n-4}. \\
 \text{(d): } & \begin{vmatrix} GC_{n+2} & GC_{n+1} & GC_n \\ GC_{n+1} & GC_n & GC_{n-1} \\ GC_n & GC_{n-1} & GC_{n-2} \end{vmatrix} = (1-7i)2^{2n-4}.
 \end{aligned}$$

## 5. SUM FORMULAS

In this chapter, we give some sum formulas of generalized Gaussian Woodall numbers.

THEOREM 5.1. *For all integers  $n \geq 0$ , we have the following formulas:*

$$\begin{aligned}
 \text{(a): } & \sum_{k=0}^n GW_k = \frac{1}{2}W_2(2n-2^{n+1}(n-1)+2^{n+2}(n-2)+6) - \frac{1}{2}W_1(8n-2^{n+1}(3n-5)+2^{n+2}(3n-8)+22) \\
 & + W_0(4n-2^{n+1}(n-2)+2^{n+2}(n-3)+9) + i\left(\frac{1}{4}(28+16n-5 \times 2^{n+2}+2^{n+2}n)W_0 + (-33-16n+7 \times 2^{n+2}-3 \times 2^{n+1}n)W_1 + (9+4n-2^{n+3}+2^{n+1}n)W_2\right). \\
 \text{(b): } & \sum_{k=0}^n GW_{2k+1} = \frac{1}{18}W_2(18n-2^{2n+3}(2n+1)+2^{2n+5}(2n-1)+40) - \frac{1}{18}W_1(72n-2^{2n+3}(6n+1)+2^{2n+5}(6n-5)+150) \\
 & + \frac{1}{9}W_0(36n+2^{2n+5}(2n-2)-2 \times 2^{2n+3}n+64) + i\left(\frac{1}{9}W_0(36n-2^{2n+2}(2n-1)+2^{2n+4}(2n-3)+53) - \frac{1}{18}W_1(72n-2^{2n+2}(6n-2)+2^{2n+4}(6n-8)+120) + \frac{1}{18}W_2(18n+2^{2n+4}(2n-2)-2 \times 2^{2n+2}n+32)\right). \\
 \text{(c): } & \sum_{k=0}^n GW_{2k} = \frac{1}{9}W_0(36n-2^{2n+2}(2n-1)+2^{2n+4}(2n-3)+53) - \frac{1}{18}W_1(72n-2^{2n+2}(6n-2)+2^{2n+4}(6n-8)+120) \\
 & + \frac{1}{18}W_2(18n+2^{2n+4}(2n-2)-2 \times 2^{2n+2}n+32) + i\left(\frac{1}{9}W_0(36n-2^{2n+1}(2n-2)+2^{2n+3}(2n-4)+46) + \frac{1}{18}W_2((18n-2^{2n+1}(2n-1)+2^{2n+3}(2n-3)+\frac{53}{2}) - \frac{1}{18}W_1(72n-2^{2n+1}(6n-5)+2^{2n+3}(6n-11)+\frac{201}{2}))\right).
 \end{aligned}$$

*Proof.*

(a): When we use (2.2),

$$\sum_{k=0}^n GW_k = \sum_{k=0}^n W_k + i \sum_{k=0}^n W_{k-1}.$$

So, then we obtain

$$\begin{aligned} \sum_{k=0}^n W_k &= \frac{1}{2}W_2(2n - 2^{n+1}(n-1) + 2^{n+2}(n-2) + 6) - \frac{1}{2}W_1(8n - 2^{n+1}(3n-5) + 2^{n+2}(3n-8) + 22) \\ &\quad + W_0(4n - 2^{n+1}(n-2) + 2^{n+2}(n-3) + 9) \end{aligned}$$

and

$$\begin{aligned} \sum_{k=0}^n W_{k-1} &= \left(\frac{1}{4}((28 + 16n - 5 \times 2^{n+2} + 2^{n+2}n)W_0 + (-33 - 16n + 7 \times 2^{n+2} - 3 \times 2^{n+1}n)W_1 \right. \\ &\quad \left. + (9 + 4n - 2^{n+3} + 2^{n+1}n)W_2)\right) \end{aligned}$$

from sum formulas on the Generalized Woodall Sequence article. We get

$$\begin{aligned} \sum_{k=0}^n GW_k &= \frac{1}{2}W_2(2n - 2^{n+1}(n-1) + 2^{n+2}(n-2) + 6) - \frac{1}{2}W_1(8n - 2^{n+1}(3n-5) + 2^{n+2}(3n-8) + 22) \\ &\quad + W_0(4n - 2^{n+1}(n-2) + 2^{n+2}(n-3) + 9) + i\left(\frac{1}{4}((28 + 16n - 5 \times 2^{n+2} + 2^{n+2}n)W_0 \right. \\ &\quad \left. + (-33 - 16n + 7 \times 2^{n+2} - 3 \times 2^{n+1}n)W_1 + (9 + 4n - 2^{n+3} + 2^{n+1}n)W_2)\right). \end{aligned}$$

**(b):** When we use (2.1), we obtain the following equalities: If we rearrange the above equalities, we obtain. Now, if we add the above equations by side by, we get

$$\sum_{k=0}^n GW_{2k+1} = \sum_{k=0}^n W_{2k+1} + i \sum_{k=0}^n W_{2k}$$

and so we know

$$\begin{aligned} \sum_{k=0}^n W_{2k+1} &= \frac{1}{18}W_2(18n - 2^{2n+3}(2n+1) + 2^{2n+5}(2n-1) + 40) - \frac{1}{18}W_1(72n - 2^{2n+3}(6n+1) \\ &\quad + 2^{2n+5}(6n-5) + 150) + \frac{1}{9}W_0(36n + 2^{2n+5}(2n-2) - 2 \times 2^{2n+3}n + 64) \end{aligned}$$

and

$$\begin{aligned} \sum_{k=0}^n W_{2k} &= \left(\left(\frac{1}{9}W_0(36n - 2^{2n+2}(2n-1) + 2^{2n+4}(2n-3) + 53) - \frac{1}{18}W_1(72n - 2^{2n+2}(6n-2) \right. \right. \\ &\quad \left. \left. + 2^{2n+4}(6n-8) + 120) + \frac{1}{18}W_2(18n + 2^{2n+4}(2n-2) - 2 \times 2^{2n+2}n + 32)\right)\right). \end{aligned}$$

We get

$$\begin{aligned} \sum_{k=0}^n GW_{2k+1} &= \frac{1}{18}W_2(18n - 2^{2n+3}(2n+1) + 2^{2n+5}(2n-1) + 40) - \frac{1}{18}W_1(72n - 2^{2n+3}(6n+1) \\ &\quad + 2^{2n+5}(6n-5) + 150) + \frac{1}{9}W_0(36n + 2^{2n+5}(2n-2) - 2 \times 2^{2n+3}n + 64) \\ &\quad + i\left(\left(\frac{1}{9}W_0(36n - 2^{2n+2}(2n-1) + 2^{2n+4}(2n-3) + 53) - \frac{1}{18}W_1(72n - 2^{2n+2}(6n-2) \right. \right. \\ &\quad \left. \left. + 2^{2n+4}(6n-8) + 120) + \frac{1}{18}W_2(18n + 2^{2n+4}(2n-2) - 2 \times 2^{2n+2}n + 32)\right)\right). \end{aligned}$$

(c): We know

$$\begin{aligned} \sum_{k=0}^n W_{2k} &= \left( \frac{1}{9} W_0(36n - 2^{2n+2}((2n-1) + 2^{2n+4}((2n-3) + 53)) - \frac{1}{18} W_1(72n - 2^{2n+2}(6n-2) \right. \\ &\quad \left. + 2^{2n+4}(6n-8) + 120) + \frac{1}{18} W_2(18n + 2^{2n+4}(2n-2) - 2 \times 2^{2n+2}n + 32) \right) \end{aligned}$$

and

$$\begin{aligned} \sum_{k=0}^n W_{2k-1} &= \left( \frac{1}{9} W_0(36n - 2^{2n+1}(2n-2) + 2^{2n+3}(2n-4) + 46) + \frac{1}{18} W_2(18n - 2^{2n+1}(2n-1) \right. \\ &\quad \left. + 2^{2n+3}(2n-3) + \frac{53}{2}) - \frac{1}{18} W_1(72n - 2^{2n+1}(6n-5) + 2^{2n+3}(6n-11) + \frac{201}{2}) \right). \end{aligned}$$

So we know

$$\sum_{k=0}^n GW_{2k} = \sum_{k=0}^n W_{2k} + i \sum_{k=0}^n W_{2k-1}.$$

We get

$$\begin{aligned} \sum_{k=0}^n GW_{2k} &= \frac{1}{9} W_0(36n - 2^{2n+2}(2n-1) + 2^{2n+4}(2n-3) + 53) - \frac{1}{18} W_1(72n - 2^{2n+2}(6n-2) \\ &\quad + 2^{2n+4}(6n-8) + 120) + \frac{1}{18} W_2(18n + 2^{2n+4}(2n-2) - 2 \times 2^{2n+2}n + 32) \\ &\quad + i \left( \frac{1}{9} W_0(36n - 2^{2n+1}(2n-2) + 2^{2n+3}(2n-4) + 46) + \frac{1}{18} W_2((18n - 2^{2n+1}(2n-1) \right. \\ &\quad \left. + 2^{2n+3}(2n-3) + \frac{53}{2}) - \frac{1}{18} W_1(72n - 2^{2n+1}(6n-5) + 2^{2n+3}(6n-11) + \frac{201}{2}) \right). \end{aligned}$$

This completes the proof.  $\square$

As special cases of above Theorem, we have the following four Corollary, we get the following corollary:

First, taking  $GW_n = GG_n$  with  $GG_0 = 0, GG_1 = 1, GG_2 = 5 + i$ .

**COROLLARY 5.2.** (Sum of the Gaussian modified Woodall numbers). For  $n \geq 0$  we have the following formulas:

- (a):  $\sum_{k=0}^n GG_k = (1+i)n + (2+i)2^n n - (4+3i)2^n + (4+3i)$ .
- (b):  $\sum_{k=0}^n GG_{2k+1} = \frac{4}{9}((\frac{9}{4} + \frac{9}{4}i)n - (4+5i)2^{2n} + (12+6i)2^{2n}n + (\frac{25}{4} + 5i))$ .
- (c):  $\sum_{k=0}^n GG_{2k} = \frac{4}{9}((\frac{9}{4} + \frac{9}{4}i)n - (5+4i)2^{2n} + (6+3i)2^{2n}n + (5+4i))$ .

Second, taking  $GW_n = GH_n$  with  $GH_0 = 3 + 2i, GH_1 = 5 + 3i, GH_2 = 9 + 5i$ . We have the following corollary:

**COROLLARY 5.3.** (Sum of the Gaussian modified Cullen numbers). For  $n \geq 0$  we have the following formulas:

- (a):  $\sum_{k=0}^n GH_k = 2^{n+2} + n - 1 + i(n + 2^{n+1})$ .
- (b):  $\sum_{k=0}^n GH_{2k+1} = \frac{1}{3}(2^{2n+4} + 3n - 1) + i(\frac{1}{3}(2^{2n+3} + 3n + 1))$ .
- (c):  $\sum_{k=0}^n GH_{2k} = \frac{1}{3}(2^{2n+3} + 3n + 1) + i(n + \frac{1}{3}2^{2n+2} + \frac{2}{3})$ .

Third, taking  $GW_n = GR_n$  with  $GR_0 = -1 - \frac{3}{2}i, GR_1 = 1 - i, GR_2 = 7 + i$ . We get the following corollary:

COROLLARY 5.4. (*Sum of the Gaussian Woodall numbers*). For  $n \geq 0$  we have the following formulas:

- (a):  $\sum_{k=0}^n GR_k = (n-1)(2^{n+1}-1) + i(2^{n+1}(n-1) - n - 2^n n + \frac{1}{2})$ .
- (b):  $\sum_{k=0}^n GR_{2k+1} = \frac{1}{9}((6n+1)2^{2n+3} - 9n + 1) + i(\frac{1}{9}((3n-1)2^{2n+3} - 9n - 1))$ .
- (c):  $\sum_{k=0}^n GR_{2k} = \frac{1}{9}((3n-1)2^{2n+3} - 9n - 1) + i(\frac{1}{9}2^{2n+3}(2n-1) - \frac{1}{9}2^{2n+1}(2n+1) - n - \frac{7}{18})$ .

Fourth, taking  $GW_n = GC_n$  with  $GC_0 = 1 + \frac{1}{2}i, GC_1 = 3 + i, GC_2 = 9 + 3i$ . We have the following corollary:

COROLLARY 5.5. (*Sum of the Gaussian Cullen numbers*). For  $n \geq 0$  we have the following formulas:

- (a):  $\sum_{k=0}^n GC_k = (n-1)2^{n+1} + n + 3 + i(n + 2^{n+1}(n-1) - 2^n n + \frac{5}{2})$ .
- (b):  $\sum_{k=0}^n GC_{2k+1} = \frac{1}{9}((6n+1)2^{2n+3} + 9n + 19) + i(\frac{1}{9}((3n-1)2^{2n+3} + 9n + 17))$ .
- (c):  $\sum_{k=0}^n GC_{2k} = \frac{2}{9}((\frac{9}{2} + \frac{9}{2}i)n - (4 + 5i)2^{2n} + (12 + 6i)2^{2n}n + (\frac{17}{2} + \frac{29}{4}i))$ .

## 6. Matrix Formulation of $GW_n$

Consider the sequence  $\{G_n\}$  which is defined by the third-order recurrence relation

$$G_n = 5G_{n-1} - 8G_{n-2} + 4G_{n-3}$$

with the initial conditions

$$G_0 = 0, G_1 = 1, G_2 = 5.$$

We define the square matrix  $A$  of order 3 as

$$A = \begin{pmatrix} 5 & -8 & 4 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

such that  $\det A = 1$ . We give the following Lemma.

LEMMA 6.1. For  $n \geq 0$  the following identity is true

$$\begin{pmatrix} GW_{n+2} \\ GW_{n+1} \\ GW_n \end{pmatrix} = \begin{pmatrix} 5 & -8 & 4 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^n \begin{pmatrix} GW_2 \\ GW_1 \\ GW_0 \end{pmatrix}.$$

Proof. The Lemma (6.1) equality can be proved by strong induction on  $n$ . If  $n = 0$  we obtain

$$\begin{pmatrix} GW_{n+2} \\ GW_{n+1} \\ GW_n \end{pmatrix} = \begin{pmatrix} 5 & -8 & 4 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^0 \begin{pmatrix} GW_2 \\ GW_1 \\ GW_0 \end{pmatrix}$$

which is true. We assume that the identity given holds for  $n \leq k$ . So that the following identity is true.

$$\begin{pmatrix} GW_{n+2} \\ GW_{n+1} \\ GW_n \end{pmatrix} = \begin{pmatrix} 5 & -8 & 4 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^k \begin{pmatrix} GW_2 \\ GW_1 \\ GW_0 \end{pmatrix}.$$

For  $n = k + 1$ , we get

$$\begin{aligned} \begin{pmatrix} 5 & -8 & 4 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^{k+1} \begin{pmatrix} GW_2 \\ GW_1 \\ GW_0 \end{pmatrix} &= \begin{pmatrix} 5 & -8 & 4 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 5 & -8 & 4 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^k \begin{pmatrix} GW_2 \\ GW_1 \\ GW_0 \end{pmatrix} \\ &= \begin{pmatrix} 5 & -8 & 4 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} GW_{k+2} \\ GW_{k+1} \\ GW_k \end{pmatrix} \\ &= \begin{pmatrix} 5GW_{k+2} - 8GW_{k+1} + 4GW_k \\ GW_{k+2} \\ GW_{k+1} \end{pmatrix} \\ &= \begin{pmatrix} GW_{k+3} \\ GW_{k+2} \\ GW_{k+1} \end{pmatrix}. \end{aligned}$$

Consequently, by induction on  $n$ , the proof is finished.  $\square$

Note that

$$A^n = \begin{pmatrix} G_{n+1} & -8G_n + 4G_{n-1} & G_n \\ G_n & -8G_{n-1} + 4G_{n-2} & G_{n-1} \\ G_{n-1} & -8G_{n-2} + 4G_{n-3} & G_{n-2} \end{pmatrix}.$$

For the proof see [47].

**THEOREM 6.2.** *We assume that the matrices  $N_{GW}$  and  $E_{GW}$  are defined as follows*

$$N_{GW} = \begin{pmatrix} GW_2 & GW_1 & GW_0 \\ GW_1 & GW_0 & GW_{-1} \\ GW_0 & GW_{-1} & GW_{-2} \end{pmatrix},$$

$$E_{GW} = \begin{pmatrix} GG_{n+2} & GG_{n+1} & GG_n \\ GG_{n+1} & GG_n & GG_{n-1} \\ GG_n & GG_{n-1} & GG_{n-2} \end{pmatrix}.$$

*The following identity is true between  $A^n$ ,  $N_{GW}$  and  $E_{GW}$ .*

$$A^n N_{GW} = E_{GW}.$$

Proof. Note that one gets

$$\begin{aligned} A^n N_{GW} &= \begin{pmatrix} G_{n+1} & -8G_n + 4G_{n-1} & G_n \\ G_n & -8G_{n-1} + 4G_{n-2} & G_{n-1} \\ G_{n-1} & -8G_{n-2} + 4G_{n-3} & G_{n-2} \end{pmatrix} \begin{pmatrix} GW_2 & GW_1 & GW_0 \\ GW_1 & GW_0 & GW_{-1} \\ GW_0 & GW_{-1} & GW_{-2} \end{pmatrix} \\ &= \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \end{aligned}$$

such that

$$\begin{aligned} a_{11} &= GW_2 G_{n+1} + GW_1 (4G_{n-1} - 8G_n) + GW_0 G_n, \\ a_{12} &= GW_1 G_{n+1} + GW_0 (4G_{n-1} - 8G_n) + GW_{-1} G_n, \\ a_{13} &= GW_0 G_{n+1} + GW_{-1} (4G_{n-1} - 8G_n) + GW_{-2} G_n, \\ a_{21} &= GW_2 G_n + GW_1 (4G_{n-2} - 8G_{n-1}) + GW_0 G_{n-1}, \\ a_{22} &= GW_1 G_n + GW_0 (4G_{n-2} - 8G_{n-1}) + GW_{-1} G_{n-1}, \\ a_{23} &= GW_0 G_n + GW_{-1} (4G_{n-2} - 8G_{n-1}) + GW_{-2} G_{n-1}, \\ a_{31} &= GW_2 G_{n-1} + GW_1 (4G_{n-3} - 8G_{n-2}) + GW_0 G_{n-2}, \\ a_{32} &= GW_1 G_{n-1} + GW_0 (4G_{n-3} - 8G_{n-2}) + GW_{-1} G_{n-2}, \\ a_{33} &= GW_0 G_{n-1} + GW_{-1} (4G_{n-3} - 8G_{n-2}) + GW_{-2} G_{n-2}. \end{aligned}$$

Using the Theorem (3.6) the proof is completed.  $\square$

We have the following identities for  $N_{GW}$ ,  $E_{GW}$  :

$$\begin{aligned} N_{GG} &= \begin{pmatrix} 5+i & 1 & 0 \\ 1 & 0 & \frac{1}{4}i \\ 0 & \frac{1}{4}i & \frac{1}{4} + \frac{1}{2}i \end{pmatrix}, & N_{GH} &= \begin{pmatrix} 9+5i & 5+3i & 3+2i \\ 5+3i & 3+2i & 2+\frac{3}{2}i \\ 3+2i & 2+\frac{3}{2}i & \frac{3}{2} + \frac{5}{4}i \end{pmatrix}, \\ N_{GR} &= \begin{pmatrix} 7+i & 1-i & -1-\frac{3}{2}i \\ 1-i & -1-\frac{3}{2}i & -\frac{3}{2}-\frac{3}{2}i \\ -1-\frac{3}{2}i & -\frac{3}{2}-\frac{3}{2}i & -\frac{3}{2}-\frac{11}{8}i \end{pmatrix}, & N_{GC} &= \begin{pmatrix} 9+3i & 3+i & 1+\frac{1}{2}i \\ 3+i & 1+\frac{1}{2}i & \frac{1}{2} + \frac{1}{2}i \\ 1+\frac{1}{2}i & \frac{1}{2} + \frac{1}{2}i & \frac{1}{2} + \frac{5}{8}i \end{pmatrix}. \end{aligned}$$

and

$$\begin{aligned} E_{GG} &= \begin{pmatrix} GG_{n+2} & GG_{n+1} & GG_n \\ GG_{n+1} & GG_n & GG_{n-1} \\ GG_n & GG_{n-1} & GG_{n-2} \end{pmatrix}, & E_{GH} &= \begin{pmatrix} GH_{n+2} & GH_{n+1} & GH_n \\ GH_{n+1} & GH_n & GH_{n-1} \\ GH_n & GH_{n-1} & GH_{n-2} \end{pmatrix}, \\ E_{GR} &= \begin{pmatrix} GR_{n+2} & GR_{n+1} & GR_n \\ GR_{n+1} & GR_n & GR_{n-1} \\ GR_n & GR_{n-1} & GR_{n-2} \end{pmatrix}, & E_{GC} &= \begin{pmatrix} GC_{n+2} & GC_{n+1} & GC_n \\ GC_{n+1} & GC_n & GC_{n-1} \\ GC_n & GC_{n-1} & GC_{n-2} \end{pmatrix}. \end{aligned}$$

From the previous theorem presents, we have the following corollary.

COROLLARY 6.3. *The following identities are true:*

- (a):  $A^n N_{GG} = E_{GG}$ .
- (b):  $A^n N_{GH} = E_{GH}$ .
- (c):  $A^n N_{GR} = E_{GR}$ .
- (d):  $A^n N_{GC} = E_{GC}$ .

## 7. Conclusions

Recently, there have been so many studies of the sequences of numbers in the literature that concern about subsequences of the Horadam numbers and generalized third-order Pell numbers such as Fibonacci, Lucas, Pell and Jacobsthal numbers; third-order Pell, third-order Pell-Lucas, Padovan, Perrin, Padovan-Perrin, Narayana, third order Jacobsthal and third order Jacobsthal-Lucas numbers. The sequences of numbers were widely used in many research areas, such as physics, engineering, architecture, nature and art.

- In section 1, we present some background about the Gaussian generalized Woodall numbers and give some information about Gaussian sequences from literature.
- In section 2, we define Gaussian generalized Woodall numbers and give some properties such as Binet's formula and generating function.
- In section 3, we present some identities, using recurrence relation and generating function, on Gaussian modified Woodall, Gaussian modified Cullen, Gaussian Woodall, Gaussian Cullen numbers.
- In section 4, we give Simpson's formula of Gaussian generalized Woodall numbers.
- In section 5, we identify some sum formulas of Gaussian generalized Woodall numbers.
- In section 6, We give the square matrix  $A^n$  using modified Woodall sequence  $\{G_n\}$  and present some identities about Gaussian generalized Woodall numbers.

Linear recurrence relations (sequences) have many applications. Next, we list applications of sequences which are linear recurrence relations.

First, we present some applications of second order sequences.

- For the applications of Gaussian Fibonacci and Gaussian Lucas numbers to Pauli Fibonacci and Pauli Lucas quaternions, see [3].
- For the application of Pell Numbers to the solutions of three-dimensional difference equation systems, see [6].
- For the application of Jacobsthal numbers to special matrices, see [71].
- For the application of generalized k-order Fibonacci numbers to hybrid quaternions, see [22].
- For the applications of Fibonacci and Lucas numbers to Split Complex Bi-Periodic numbers, see [72].

- For the applications of generalized bivariate Fibonacci and Lucas polynomials to matrix polynomials, see [73].
- For the applications of generalized Fibonacci numbers to binomial sums, see [70].
- For the application of generalized Jacobsthal numbers to hyperbolic numbers, see [49].
- For the application of generalized Fibonacci numbers to dual hyperbolic numbers, see [50].
- For the application of Laplace transform and various matrix operations to the characteristic polynomial of the Fibonacci numbers, see [14].
- For the application of Generalized Fibonacci Matrices to Cryptography, see [37].
- For the application of higher order Jacobsthal numbers to quaternions, see [38].
- For the application of Fibonacci and Lucas Identities to Toeplitz-Hessenberg matrices, see [20].
- For the applications of Fibonacci numbers to lacunary statistical convergence, see [5].
- For the applications of Fibonacci numbers to lacunary statistical convergence in intuitionistic fuzzy normed linear spaces, see [29].
- For the applications of Fibonacci numbers to ideal convergence on intuitionistic fuzzy normed linear spaces, see [30].
- For the applications of  $k$ -Fibonacci and  $k$ -Lucas numbers to spinors, see [32].
- For the application of dual-generalized complex Fibonacci and Lucas numbers to Quaternions, see [40].
- For the application of special cases of Horadam numbers to Neutrosophic analysis see [19].
- For the application of Hyperbolic Fibonacci numbers to Quaternions, see [13].

We now present some applications of third order sequences.

- For the applications of third order Jacobsthal numbers and Tribonacci numbers to quaternions, see [11] and [10], respectively.
- For the application of Tribonacci numbers to special matrices, see [74].
- For the applications of Padovan numbers and Tribonacci numbers to coding theory, see [41] and [4], respectively.
- For the application of Pell-Padovan numbers to groups, see [15].
- For the application of adjusted Jacobsthal-Padovan numbers to the exact solutions of some difference equations, see [18].
- For the application of Gaussian Tribonacci numbers to various graphs, see [65].
- For the application of third-order Jacobsthal numbers to hyperbolic numbers, see [16].
- For the application of Narayan numbers to finite groups see [31].
- For the application of generalized third-order Jacobsthal sequence to binomial transform, see [51].
- For the application of generalized Generalized Padovan numbers to Binomial Transform, see [52].
- For the application of generalized Tribonacci numbers to Gaussian numbers, see [53].

- For the application of generalized Tribonacci numbers to Sedenions, see [54].
- For the application of Tribonacci and Tribonacci-Lucas numbers to matrices, see [55].
- For the application of generalized Tribonacci numbers to circulant matrix, see [56].
- For the application of Tribonacci and Tribonacci-Lucas numbers to hybrinomials, see [69].
- For the application of hyperbolic Leonardo and hyperbolic Francois numbers to quaternions, see [17].

Next, we now list some applications of fourth order sequences.

- For the application of Tetranacci and Tetranacci-Lucas numbers to quaternions, see [57].
- For the application of generalized Tetranacci numbers to Gaussian numbers, see [58].
- For the application of Tetranacci and Tetranacci-Lucas numbers to matrices, see [59].
- For the application of generalized Tetranacci numbers to binomial transform, see [60].

We now present some applications of fifth order sequences.

- For the application of Pentanacci numbers to matrices, see [42].
- For the application of generalized Pentanacci numbers to quaternions, see [44].
- For the application of generalized Pentanacci numbers to binomial transform, see [61].

We now present some applications of second order sequences of polynomials.

- For the application of generalized Fibonacci Polynomials to the summation formulas, see [62].
- For some applications of generalized Fibonacci Polynomials, see [63].

We now present some applications of third order sequences of polynomials.

- For some applications of generalized Tribonacci Polynomials, see [64].

## References

- [1] Aşçı M., Gürel E., Gaussian Jacobsthal and Gaussian Jacobsthal Lucas Numbers, *Ars Combinatoria, Soc.*, 111, 2013.
- [2] Ertaş A., Yılmaz F., On Quaternions with Gaussian Oresme Coefficients, *Turk. J. Math. Comput. Sci.*, 15(1), 192-202, 2023.
- [3] Azak, A.Z., Pauli Gaussian Fibonacci and Pauli Gaussian Lucas Quaternions. *Mathematics*, 2022, 10, 4655. <https://doi.org/10.3390/math10244655>
- [4] Basu, M., Das, M., Tribonacci Matrices and a New Coding Theory, *Discrete Mathematics, Algorithms and Applications*, 6 (1), 1450008, (17 pages), 2014.
- [5] Bilgin, N.G., Fibonacci Lacunary Statistical Convergence of Order  $\eta$  in IFNLS, *International Journal of Advances in Applied Mathematics and Mechanics*, 8(4), 28-36, 2021.
- [6] Büyük, H., Taşkara, N., On The Solutions of Three-Dimensional Difference Equation Systems Via Pell Numbers, *European Journal of Science and Technology, Special Issue 34*, 433-440, 2022.
- [7] P. Berrizbeitia, J. G. Fernandes, M. J. González, F. Luca, V. J. M. Hugueta, *On Cullen numbers which are Both Riesel and Sierpiński numbers*, *Journal of Number Theory*, **132** (2012), 2836-2841, <http://dx.doi.org/10.1016/j.jnt.2012.05.021>
- [8] Y. Bilu, D. Marques, A. Togbe, *Generalized Cullen numbers in linear recurrence sequences*, *Journal of Number Theory*, **202** (2019), 412-425, <https://doi.org/10.1016/j.jnt.2018.11.025>

- [9] Cerda-Morales, G., On Gauss third-order Jacobsthal numbers and their applications, , *Annals of the Alexandru Ioan Cuza University-Mathematics*, 67(2), 231-241, 2022.
- [10] Cerda-Morales, G., On a Generalization of Tribonacci Quaternions, *Mediterranean Journal of Mathematics* 14:239, 1-12, 2017.
- [11] Cerda-Morales, G., Identities for Third Order Jacobsthal Quaternions, *Advances in Applied Clifford Algebras* 27(2), 1043–1053, 2017.
- [12] A. J. C. Cunningham, H. J. Woodall, *Factorisation of  $Q = (2^q \mp q)$  and  $(q \cdot 2^q \mp 1)$* , *Messenger of Mathematics*, 47 (1917), 1–38.
- [13] Daşdemir, A., On Recursive Hyperbolic Fibonacci Quaternions, *Communications in Advanced Mathematical Sciences*, 4(4), 198-207, 2021. DOI:10.33434/cams.997824
- [14] Deveci, Ö., Shannon, A.G., On Recurrence Results From Matrix Transforms, *Notes on Number Theory and Discrete Mathematics*, 28(4), 589–592, 2022. DOI: 10.7546/nntdm.2022.28.4.589-592
- [15] Deveci, Ö., Shannon, A.G., Pell–Padovan–Circulant Sequences and Their Applications, *Notes on Number Theory and Discrete Mathematics*, 23(3), 100–114, 2017.
- [16] Dikmen, C.M., Altınoy, M., On Third Order Hyperbolic Jacobsthal Numbers, *Konuralp Journal of Mathematics*, 10(1), 118-126, 2022.
- [17] Dişkaya, O., Menken, H., Catarino, P.M.M.C., On the Hyperbolic Leonardo and Hyperbolic Francois Quaternions, *Journal of New Theory*, 42, 74-85, 2023.
- [18] Göcen, M., The Exact Solutions of Some Difference Equations Associated with Adjusted Jacobsthal-Padovan Numbers, *Kırklareli University Journal of Engineering and Science* 8(1), 1-14, 2022. DOI: 10.34186/klujes.1078836
- [19] Gökbaşı H., Topal, S., Smarandache, F., Neutrosophic Number Sequences: An Introductory Study, *International Journal of Neutrosophic Science (IJNS)*, 20(01), 27-48, 2023. <https://doi.org/10.54216/IJNS.200103>
- [20] Goy, T., Shattuck, M., Fibonacci and Lucas Identities from Toeplitz-Hessenberg Matrices, *Appl. Appl. Math.*, 14(2), 699–715, 2019.
- [21] J. Grantham, H. Graves, *The abc conjecture implies that only finitely many s-Cullen numbers are repunits*, (2021) <http://arxiv.org/abs/2009.04052v3>, MathNT.
- [22] Gül, K., Generalized k-Order Fibonacci Hybrid Quaternions, *Erzincan University Journal of Science and Technology*, 15(2), 670-683, 2022. DOI: 10.18185/erzifbed.1132164
- [23] R. Guy, *Unsolved Problems in Number Theory (2nd ed.)*, Springer-Verlag, New York, 1994.
- [24] Halıcı, S., Öz, S., On Some Gaussian Pell and Pell-Lucas Numbers, *Ordu Univ. J. Sci. Tech.*, 6(1), 8-18, 2016.
- [25] C. Hooley, *Applications of the sieve methods to the theory of numbers*, Cambridge University Press, Cambridge, 1976.
- [26] Horadam, A. F., Complex Fibonacci Numbers and Fibonacci quaternions, *Amer. Math. Monthly* 70, 289–291, 1963.
- [27] D. Kalman, *Generalized Fibonacci numbers by matrix methods*, *Fibonacci Quarterly*, 20(1) (1982), 73-76.
- [28] W. Keller, *New Cullen primes*, *Math. Comput.* 64 (1995), 1733-1741.
- [29] Kişi, Ö., Tuzcuoglu, I., Fibonacci Lacunary Statistical Convergence in Intuitionistic Fuzzy Normed Linear Spaces, *Journal of Progressive Research in Mathematics* 16(3), 3001-3007, 2020.
- [30] Kişi, Ö., Debnath, P., Fibonacci Ideal Convergence on Intuitionistic Fuzzy Normed Linear Spaces, *Fuzzy Information and Engineering*, 1-13, 2022. <https://doi.org/10.1080/16168658.2022.2160226>
- [31] Kuloğlu, B., Özkan, E., Shannon, A.G., The Narayana Sequence in Finite Groups, *Fibonacci Quarterly*. 60(5), 212–221, 2022.
- [32] Kumari, M., Prasad, K., Frontczak, R., On the  $k$ -Fibonacci and  $k$ -Lucas Spinors, *Notes on Number Theory and Discrete Mathematics*, 29(2), 322-335, 2023. DOI: 10.7546/nntdm.2023.29.2.322-335

- [33] F. Luca, P. Stanica, *Cullen numbers in binary recurrent sequences*, FT. Howard (ed.), Applications of Fibonacci Numbers : Proceedings of the Tenth International Research Conference on Fibonacci Numbers and their Applications, Kluwer Academic Publishers, **9** (2004) 167-175.
- [34] D. Marques, *On Generalized Cullen and Woodall numbers that are also Fibonacci numbers*, Journal of Integer Sequences, **17** (2014), Article 14.9.4
- [35] D. Marques, *Fibonacci s-Cullen and s-Woodall numbers*, , Journal of Integer Sequences, **18** (2015), Article 15.1.4
- [36] Pethe S., Horadam A.F., Generalised Gaussian Fibonacci numbers, Bull. Austral. Math. Soc., Vol.33, pp.37-48, 1986.
- [37] Prasad, K., Mahato, H., Cryptography Using Generalized Fibonacci Matrices with Affine-Hill Cipher, Journal of Discrete Mathematical Sciences & Cryptography, 25(8-A), 2341–2352, 2022. DOI : 10.1080/09720529.2020.1838744
- [38] Özkan, E., Uysal, M., On Quaternions with Higher Order Jacobsthal Numbers Components, Gazi University Journal of Science, 36(1), 336-347, 2023. DOI: 10.35378/gujs. 1002454
- [39] N. K. Meher, S. S. Rout, *Cullen numbers in sums of terms of recurrence sequence*, (2020), <http://arxiv.org/abs/2010.10014v1>, MathNT.
- [40] Şentürk, G.Y., Gürses, N., Yüce, S., Construction of Dual-Generalized Complex Fibonacci and Lucas Quaternions, Carpathian Math. Publ. 2022, 14 (2), 406-418, 2022. doi:10.15330/cmp.14.2.406-418
- [41] Shtayat, J., Al-Kateeb, A., An Encoding-Decoding algorithm based on Padovan numbers, arXiv:1907.02007, 2019.
- [42] Sivakumar, B., James, V., A Notes on Matrix Sequence of Pentanacci Numbers and Pentanacci Cubes, Communications in Mathematics and Applications, 13(2), 603–611, 2022. DOI: 10.26713/cma.v13i2.1725
- [43] Sloane, N.J.A., The on-line encyclopedia of integer sequences, <http://oeis.org/>
- [44] Soykan, Y., Özmen, N., Göcen, M., On Generalized Pentanacci Quaternions, Tbilisi Mathematical Journal, 13(4), 169-181, 2020.
- [45] Soykan, Y., Taşdemir, E., Okumuş, İ., Göcen, M., Gaussian Generalized Tribonacci Numbers, Journal of Progressive Research in Mathematics, 14(2) , 2018.
- [46] Soykan, Y., Simson Identity of Generalized m-step Fibonacci Numbers, Int. J. Adv. Appl. Math. and Mech. 7(2), 45-56, 2019.
- [47] Soykan, Y., A Study On Generalized (r,s,t)-Numbers, MathLAB Journal, 7, 101-129, 2020.
- [48] Soykan, Y., Generalized Woodall Numbers: An Investigation of Properties of Woodall and Cullen Numbers via Their Third Order Linear Recurrence Relations, Universal Journal of Mathematics and Applications, 5 : 2, 69-81, 2022.
- [49] Soykan, Y., Taşdemir, E., A Study On Hyperbolic Numbers With Generalized Jacobsthal Numbers Components, International Journal of Nonlinear Analysis and Applications, 13(2), 1965–1981, 2022. <http://dx.doi.org/10.22075/ijnaa.2021.22113.2328>
- [50] Soykan, Y., On Dual Hyperbolic Generalized Fibonacci Numbers, Indian J Pure Appl Math, 2021. <https://doi.org/10.1007/s13226-021-00128-2>
- [51] Soykan, Y., Taşdemir, E., Göcen, M., Binomial Transform of the Generalized Third-Order Jacobsthal Sequence, Asian-European Journal of Mathematics, 15(12), 2022. <https://doi.org/10.1142/S1793557122502242>.
- [52] Soykan, Y., Taşdemir, E., Okumuş, İ., A Study on Binomial Transform of the Generalized Padovan Sequence, Journal of Science and Arts, 22(1), 63-90, 2022. <https://doi.org/10.46939/J.Sci.Arts-22.1-a06>
- [53] Soykan, Y., Taşdemir, E., Okumuş, İ., Göcen, M., Gaussian Generalized Tribonacci Numbers, Journal of Progressive Research in Mathematics(JPRM), 14 (2), 2373-2387, 2018.
- [54] Soykan, Y., Okumuş, İ., Taşdemir, E., On Generalized Tribonacci Sedenions, Sarajevo Journal of Mathematics, 16(1), 103-122, 2020. ISSN 2233-1964, DOI: 10.5644/SJM.16.01.08
- [55] Soykan, Y., Matrix Sequences of Tribonacci and Tribonacci-Lucas Numbers, Communications in Mathematics and Applications, 11(2), 281-295, 2020. DOI: 10.26713/cma.v11i2.1102

- [56] Soykan, Y. Explicit Euclidean Norm, Eigenvalues, Spectral Norm and Determinant of Circulant Matrix with the Generalized Tribonacci Numbers, *Earthline Journal of Mathematical Sciences*, 6(1), 131-151, 2021. <https://doi.org/10.34198/ejms.6121.131151>
- [57] Soykan Y, Tetranacci and Tetranacci-Lucas Quaternions, *Asian Research Journal of Mathematics*, 15(1): 1-24, 2019; Article no.ARJOM.50749.
- [58] Soykan, Y., Gaussian Generalized Tetranacci Numbers, *Journal of Advances in Mathematics and Computer Science*, 31(3): 1-21, Article no.JAMCS.48063, 2019.
- [59] Soykan, Y., Matrix Sequences of Tetranacci and Tetranacci-Lucas Numbers, *Int. J. Adv. Appl. Math. and Mech.* 7(2), 57-69, 2019, (ISSN: 2347-2529).
- [60] Soykan, Y., On Binomial Transform of the Generalized Tetranacci Sequence, *International Journal of Advances in Applied Mathematics and Mechanics*, 9(2), 8-27, 2021.
- [61] Soykan, Y., Binomial Transform of the Generalized Pentanacci Sequence, *Asian Research Journal of Current Science*, 3(1), 209-231, 2021.
- [62] Soykan, Y., A Study on Generalized Fibonacci Polynomials: Sum Formulas, *International Journal of Advances in Applied Mathematics and Mechanics*, 10(1), 39-118, 2022. (ISSN: 2347-2529)
- [63] Soykan, Y., On Generalized Fibonacci Polynomials: Horadam Polynomials, *Earthline Journal of Mathematical Sciences*, 11(1), 23-114, 2023. E-ISSN: 2581-8147. <https://doi.org/10.34198/ejms.11123.23114>
- [64] Soykan, Y., Generalized Tribonacci Polynomials, *Earthline Journal of Mathematical Sciences*, 13(1), 1-120, 2023. <https://doi.org/10.34198/ejms.13123.1120>
- [65] Sunitha, K., Sheriba. M., Gaussian Tribonacci R-Graceful Labeling of Some Tree Related Graphs, *Ratio Mathematica*, 44, 188-196, 2022.
- [66] Taşcı, D., Gaussian Balancing And Gaussian Lucas Balancing Numbers, *Journal of Science and Arts*, 3(44),661-666,2018.
- [67] Taşcı, D., Gaussian Padovan and Gaussian Pell- Padovan numbers, *Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat.*, 67(2), 82-88, 2018.
- [68] Taşcı, D., On Gaussian Mersenne Numbers, *Journal of Science and Arts*, 4(57),1012-1028, 2021.
- [69] Taşyurdu, Y., Polat, Y.E., Tribonacci and Tribonacci-Lucas Hybrinomials, *Journal of Mathematics Research*; 13(5), 2021.
- [70] Ulutaş, Y.T., Toy, D., Some Equalities and Binomial Sums about the Generalized Fibonacci Number  $u_n$ , *Notes on Number Theory and Discrete Mathematics*, 28(2), 252–260, 2022. DOI: 10.7546/nntdm.2022.28.2.252-260
- [71] Vasanthi, S., Sivakumar, B., Jacobsthal Matrices and their Properties. *Indian Journal of Science and Technology* 15(5): 207-215, 2022, <https://doi.org/10.17485/IJST/v15i5.1948>
- [72] Yılmaz, N., Split Complex Bi-Periodic Fibonacci and Lucas Numbers, *Commun.Fac.Sci.Univ.Ank.Ser. A1 Math. Stat.* 71(1), 153–164, 2022. DOI:10.31801/cfsuasmas.704435
- [73] Yılmaz, N., The Generalized Bivariate Fibonacci and Lucas Matrix Polynomials, *Mathematica Montisnigri*, Vol LIII, 33-44, 2022. DOI: 10.20948/mathmontis-2022-53-5
- [74] Yılmaz, N, and N. Taskara, N., Tribonacci and Tribonacci-Lucas Numbers via the Determinants of Special Matrices, *Applied Mathematical Sciences*, 8, no. 39, 1947-1955, 2014.