

# Emerging Drinking Water Borne Diseases; A Review on Types, Sources and Health Precaution

## Abstract

The global demand for safe drinking water become increasingly important in the past few years due to the growing world population, civilization and increase the sources of contaminations. Millions of people every year suffer from different waterborne disease, which may include microbial or disinfectant related diseases. This review, presents an overview on drinking water and their health impact from different aspects, going through different types of physical, chemical, and biological contaminants and their potential effect on human body upon the consumption of their contaminated water. The review also furtherly discusses drinking water treatment techniques such as chlorination, ultraviolet radiation treatment, chlorination, ozonation, heating, and ultra-filtration and the efficiency of these techniques to eliminate different water microorganisms. Drinking waterborne diseases of bacteria, viruses, parasitic protozoa and even some fungi have been also discussed. Finally, health precaution regarding drinking water in term of monitoring of different indicators as well as following the hygienic behaviors was covered to ensure the sustainability and safety of water consumption.

**Keywords:** drinking water; water borne diseases; microbial contamination; water quality.

## 1. Introduction

The World Health Organization (WHO) reported in 2017 that more than 2.1 billion people cannot access to safe drinkable water free from all the sources of physical,

chemical and microbial contaminations, which equivalent to almost third the global population [1]. It has been estimated by WHO that the annual death caused by diarrheal diseases is about 1.8 million, which mainly due to the consumption of contaminated drinking water [2]. People who suffer from chronic diseases that weakening their immunity, such as diabetes, cancer, and AIDS, etc., and the young children as well as elder peoples are particularly more vulnerable to waterborne infections [3]. Environmental factors might also play a significant role, which influence more than 80% of the overall diseases. It was reported that in young children under the age of 5 years, the diseases with the highest environmental contribution including lower respiratory infections (32% of the overall cases), diarrhoeal and GIT diseases (22%), newborn associated conditions (15%), and parasitic-borne diseases (12%) [4]. Several millions of young children die every year from waterborne disease, particularly from acute diarrheal diseases due to the consumption or using of contaminated food or water [5]. These diseases can be transmitted to human through four basic routes including waterborne route by the consumption of water directly to the human body, water-washed route by using contaminated water in washing purposes causing contamination of hands or food-plates, water-based route during the recreational or other activities and insect vector route [6].

The global burden of infectious diseases associated with water is considerable, and even in developed and high-income countries, it continues to be a concern [7]. Countries all over the world are concerning with the potential effects of consuming unclean drinking water as it has been proved to be a main cause of numerous water-borne diseases leading to high range of morbidity and mortality especially in undeveloped and developing countries [8, 9]. As a daily consumed material, clean drinking water must be free from infectious pathogens as well as toxic and undesired physico-chemical pollutants [10, 11]. Clean drinking water is essential for overall health of the entire


human been and plays a substantial role especially in infants and young children health and survival as their immune system is weaker [12]. As such, extensive testing is required to understand the possible routes of the viral contagion, which is very expensive [13]. Therefore, monitoring the spread of COVID-19 early stages in communities through the wastewater-based epidemiology (WBE) approach is useful [14]. Additionally, this could possibly provide rapid results for effective and urgent interventions in the fight against COVID-19 [15].

Many review articles have been published discussing swimming pools associated viral waterborne diseases [16], transmission of protozoan parasites [17], and control of waterborne diseases [18]. Most of previous reviews discuss either location in the world, part from water-borne diseases factors or the outbreak of water borne diseases in particular location. This review delivers an overview on drinking water quality, the basic possible contaminant and their potential health impact upon consumption [19]. It also furtherly discusses waterborne diseases, their classification, drinking water disinfection and treatment techniques and highlight the health precaution regarding drinking water and water-borne diseases.

## **2. Drinking water quality**

Water quality is a general term used to describe the physical (such as turbidity, color, suspended solids, etc.), chemical (such as organic contaminant, inorganic contaminant, heavy metals, etc.), and biological (such as bacteria, plankton, algae, fungi, etc.) parameters may present in water [20, 21]. The consideration of water quality in most of cases corresponds to the origin and the purpose of water. However, the general instruction about drinking water standards have been published by many organizations such as United States environmental protection agency (USEPA) and the WHO, etc. [22]. The quantity of microorganisms have been reported to affect the taste and odor of water, variety of microbes produce certain pigments that cause changes in water color

[23, 24].The contamination of drinking water with the feces of hot blood organisms including animal and human is consider the main route of transmission of water borne diseases to human beings[25]. Abu-Amr and Yassin[26] concluded that self-reported diseases are well associated with insufficient chlorination of drinking water, intermittent water supply and sewage flooding. However, many developing countries have the issue of chronic shortages of fresh water, even costal country in Africa and southern Asia; the readily accessible resources are another major issue which been heavily polluted in most of times [27]. The treated drinking water pumped into the network of distribution system, which may contain different physical loads (dissolved or suspended), microbial loads (bacteria, yeasts, fungi or protozoa) and nutrient loads (different organic and inorganic nutrients), and thus many people use house hold purification system in order to maintain the desired quality of their water [28]. Figure 1 presents the main types of drinking water contamination.



**Figure 1.**Sources of drinking water contamination

## 2.1. Physico-chemical contamination of drinking water

The contamination of drinking water with different physical and chemical agents released from different anthropogenic contamination sources has become a global concern [29]. The contamination of drinking water resources may also occur due to the unsuccessful treatment or even after treatment, which has important repercussions for both environment and human health [30, 31]. Gul et al. [32] reported that different anions are present in drinking water including heavy metals such as Co, Cd, Cr, Ni, Zn, Hg, Pb, etc. These heavy metals have significant adverse effects on human health if they exceeded the quality guidelines either through deficiency or toxicity. Lu et al. [33] studied the potential for cancer risk related to the presence of carcinogenic metals when consumed with drinking water, and revealed high risk at certain levels. In a different study, Arshad and Imran [34] evaluated the effect of fluoride-contaminated drinking water consumption in a rural population of Pakistan on the prevalence of skeletal/dental fluorosis, and concluded that the consumption of high amounts of fluoride can lead to serious health issues. Nitrate (NO<sub>3</sub>) and nitrites (NO<sub>2</sub>) are another two parameters naturally found in water [35], the reduction of nitrate to nitrite is toxic to human health [36]. It has been reported that nitrite is involved in the oxidation of hemoglobin and transforms it to methemoglobin, a modified compound which is unable to transport oxygen to the cells and tissues [37]. Refer to table 1 for the health effect of different physico-chemical contaminants if they are present in drinking water in high levels.

**Table 1.** Potential health effect of physico-chemical contaminants upon consumed in high levels in drinking water

Contaminant	Potential health risk	Ref
Chlorine	Bladder, colon and rectum cancers in addition to adverse	[38]

	effects on reproduction	
Sodium	Excessive intake of sodium lead to raise inblood pressure, kidney damage, stroke, headacheand edema, etc.	[39]
Nitrates and nitrites	Major health problems such as Methemoglobinomia and increase the risk of abortion and even cancers.	[40]
Fluoride	Long-term consumption can lead to dental and skeletal fluorosis, Alzheimer and neurological problems.	[41]
Trihalomethanes	Increased risk of bladder cancer	[42]
Sulfate and calcium carbonate	Excessive intake of these element in water may lead to indigestion and severe diarrhea in consumers.	[43]
Heavy metals	Different heavy metals associated with different impact such as toxicity, stability, and carcinogenic and bioaccumulation.	[44]

## 2.2. Biological contamination of drinking water

Microorganisms may introduced into water distribution system, through the main source of raw water or by side contamination source, which all will undergo different changes in viability, density and diversity of the microorganisms due to the selective pressures of the system[45]. Phiri et al. [46] assessed the presence of drinking waterborne pathogens in New Zealand and tracked the potential sources of contamination. The authors reported the presence of *Escherichia coli* in more than half of the collected samples, which was used as an indicator organism. The same authors added that the use of water filters alone was not associated with the likelihood of compliance. In different study, Ward et al. [47] conducted a nine-month drinking water quality monitoring programme in rural Malawi and revealed highly abundance of *E. coli* as well as other fecal coliform bacterial in the tested samples. The application of

water treatment will lead to dramatic decrease in the microbial load [48]. The increase in microbial growth after water treatment has been termed as regrowth, it happened due to unsuccessful treatment process, which consider as a major problem and one of the main causes of many diseases[49]. Qin et al. [50] investigated the occurrence and distribution of water opportunistic pathogens in water storage tanks using species-specific qPCR assay. The authors determined the occurrence of many pathogens including *P. aeruginosa*, *Legionella spp*, *Mycobacterium spp*, and *Acanthamoeba spp*, etc., with 88% detection of *Mycobacterium spp* in the sediments and water samples, which was the most frequently detected pathogen. The same authors reported that microbial communities possibly influenced by constituents within the water in storage tank sediments, the compositions and diversity of microbial community and the types of elements may also influence the microbial water quality[51].

### **3. Microbial water quality**

The World Health Organization/the United Nations Children's Fund (WHO/UNICEF) report of 2017 regarding monitoring program of drinking water supplies and hygiene revealed that roughly 844 million people all over the world lack access to clean and drinkable water and 2.3 billion (about third of global population) lack basic sanitation services [1]. Two major indicators for microbial water quality monitoring usually used, total microbial counts and the presence of coliforms[52]. Coliform bacteria are group of rod-shaped gram-negative microorganisms, they are the most regular bacterial species ranged from harmless to pathogenic species. Coliforms have been widely isolated from soil and other environment in the defecation of creatures of warm blood [53]. They can be classified based on their origin to fecal and non-fecal coliforms, total and fecal coliforms assessment is the most used prominent method to measure and assess the drinking water bacteriological quality[54]. Wen et al. [55] furtherly discussed water quality indicator systems in variety of countries all over

the world, the review focused on different microbial indicators including bacteria, protozoa, and enteric virus, etc., that being tested and used in Asian, European and American countries to monitor water quality.

Various microbial indicator species such as, *Enterococcus spp*, coliforms, *Escherichia coli* and total counts of water bacterial are the most frequently monitored as detection or monitoring of many other water pathogens are expensive or impractical. Although the presence of *Escherichia coli* and fecal coliforms in drinking water indicates its contamination and its unsuitability for drinking, their absence does not particularly preclude the presence of one or more other harmful organisms [56]. Microbial biomass in water and liquid media is typically measured using heterotrophic plate counts technique, which detect and quantifies the viable bacteria in drinking water [57]. The actual bacterial species quantified by heterotrophic plate counts depends on the type of cultivation medium, as well as incubation time and temperature [58]. Potential concerns have been associated with the abundance of above 500 cfu/ml, which mainly due to the potential interference with the analytical detection of various pathogens such as total coliforms [59]. Various drawbacks have been associated with heterotrophic plate counts technique such as the requirement of a well-equipped laboratory as well as time consuming as the analyses typically take several days [60]. Faster methods such as flow cytometry cell counts method and adenosine tri-phosphate assessment method have been also developed but still relatively complex to interpret and implement, which prevent their widespread and usage for drinking water quality monitoring [61, 62].

Among 15000 species of protozoa present on earth, only four classes have been identified as they have significant impact on health including Flagellata, Sarcodina, Infusoria, and Sporozoa [63]. However, the most common drinking water-related parasitic infections among adults and children are cryptosporidiosis and giardiasis [64]. *Cryptosporidium* and *Giardia* are two zoonotic agents that mostly

identified during contaminated drinking water outbreaks, particularly causing diarrheal disease among young children in developing countries [65]. The cysts and oocysts of protozoa are found in different water sources including waste water, surface and groundwater sources, even after treatment of drinking water, it has been reported the presence of these parasitic forms in many drinking water samples after treatment with conventional methods [66, 67]. As these methods mainly focus on removing of pathogenic bacteria such as *Salmonella spp*, *Vibrio cholera* and *Escherichia coli* [68], parasitic protozoa such as *Giardia lamblia* and *Cryptosporidium parvum* are able to resist chlorine of particular concern [69].

### 3.1. Drinking water disinfection techniques

The quality of drinking water is a globally major concern and has a significant impact on human health [70]. Many plans have been developed for drinking water treatment, the suitable operation and design is normally determined based on various considerations such as the physical, chemical, and microbiological parameters of the water source [71]. It has been reported that weak sanitation and polluted water provisions are the main cause of more than 80% of the diseases in the poorest nations [72]. The maturation and developing of advanced municipal water purification approaches in the last few years has authorized developed countries to overcome most of water related disease. Many methods have been used for microbial disinfection of drinking water including chlorination, ultraviolet radiation, chlorination, ozonation, chlorine dioxide or ultra-filtration [73]. Refer to table 2 for the summary of these techniques.


**Table 2.** Drinking water disinfection techniques

Technique	Remark	Ref
Chlorination	Addition of chlorine or its products reacts with water to	[74]

	form hypochlorite ions and hypochlorous acid, which are able to disinfect many water microorganisms.	
Ultraviolet radiation	Exposing the water to shortwave radiation, which kills water microorganisms by directly affecting its DNA, it is non-residual in water and does not cause any harmful products in the drinking water.	[75]
Chloramination	It is the reaction of chlorine and ammonia under well-controlled conditions result in formation of monochloramine, which is able to reduce microorganisms and it doesn't form any harmful by-products.	[76]
Ozonation	Using ozone (O <sub>3</sub> ) as a powerful oxidizing agent for readily oxidizes various microbes, chemical residuals and organic matter in low concentrations and short contact times.	[77]
Heating	Heating up the water till boiling temperature for certain time to kill microorganisms, is a widely used method among the consumers of municipal water	[78]
Ultra-filtration	Filtration of water microorganisms present in water by passing the water through small pores do not permit passing of microbes	[79]

Chlorination has been reported as the most widely used all over the world, many countries follow their own guidelines for the added concentration of chlorine [80]. Gagnon et al. [81] reported that the presence of low concentration levels of chlorite in drinking water seemed not to be effective in elimination of heterotrophic bacteria, and indicated the need to maintain higher concentrations of chlorine dioxide residual in drinking water distribution systems for better control of microbial regrowth. A portion

of the chlorine at the point of its addition to water responds first with different natural and inorganic materials, as well as metals that may present in water, the rest of the remaining chlorine is known as total chlorine [82]. Mazhar et al. [38] reported that total chlorine present in drinking water is a sum of free chlorine and combined chlorine. From one side, the combined chlorine possesses weak disinfection properties and mostly consider inaccessible for disinfection especially when it combines with many inorganic and organic materials such as nitrates and urea, etc. From the other side, the free chlorine possess stronger disinfection properties and accessible to inactivate water pathogens, it is the left over chlorine, in which a measure of the potability of the water. For instance, if we are utilizing totally clean drinking water, which free from many contaminants, its chlorine demand will be zero, and it will have zero combined chlorine as well due to the absence of organic and inorganic material in that water [83]. Figure 2 summary the addition of chlorine to water as disinfectant [84].



**Figure 2.** The Chlorine addition to drinking water

Microbial communities in drinking water have been reported to be highly sensitive to the changes in the physiochemical and microbiological parameters (i.e., types and concentration of disinfectants, nutrients concentration, inorganic elements composition), which all could influence the types, numbers and growing rates of microbial population [85]. Wang et al. [86] reported reversible shifts in microbial communities in water upon the temporary switching between chlorination and chloramination. Yang et al. [87] concluded that in extreme case, discoloration of water could arise the increase of sulfate levels in feed water, which caused noteworthy increase of various types of sulfur-oxidizing and sulfate-reducing bacteria as well as iron-oxidizing bacteria. The authors explained this increased, as it was associated with the presence of iron erosion, which cause discoloration of the water [88].

#### **4. Water-borne diseases**

Water-borne or water related diseases encompass illnesses resulting from both direct and indirect exposure to water. It has been reported that the exposure of human to water-borne infections may occur by direct contact or consumption of contaminated drinking water, food containing contaminated water or even from recreational water [89]. Apart from microorganisms, water-borne diseases also include diseases associated with toxic substances presented in water [90]. Ghernaout et al. [91] reported that roughly, two billion people throughout the world don't reach protected potable water, leading to the death of 20 million babies annually, because of waterborne diarrheal diseases such as dysentery, cholera and typhoid fever. Maramraj et al. [92] estimated that around 1.7 billion cases of diarrheal diseases and 2.2 million deaths globally from acute diarrheal diseases are being recorded every year. World Health Organization reported that lack of access to clean and quality drinking water, in addition to inadequate sanitation and hygiene are the leading causes of acute diarrheal

disease [93]. Bivins et al. [94] estimated the median daily risk of infection of diarrheal disease attributable in people who consume fecally contaminated tap water without using suitable filter of good treatment approach. The authors reported 3 main reference pathogens including *Cryptosporidium*, *Campylobacter*, and rotavirus as a standard conservative risk proxies for infections for protozoa infection, bacterial infection, and viral infection respectively. The same authors concluded that around 17.20 million infections annually causing more than 4.52 million cases of diarrhea and almost 1560 deaths, underscoring the importance of water safety management specially in developing countries and the potential benefits of point-of-use treatment to mitigate risks[15]. A variety of drinking and daily used water reservoirs have been widely linked to nosocomial outbreaks including dialysis water, potable water, tub immersion, sinks, showers, faucet aerators, ice and ice machines, swimming pools, water baths, flower vases, and dental-unit water stations [95]. Decker et al. [96] reported that transmission of waterborne pathogens from contaminated water reservoir in direct or indirect contact and the ingestion of these pathogens with water, food or even inhalation of aerosols will lead to waterborne infections.

#### ***4.1. Water disinfectant associated diseases***

Although there are numerous ways have been used to sterilize drinking water such as filtration, heating and the addition of chemical and physical methods [97]. Many countries are still using chlorine in their water distribution systems and tanks to sterilize drinking water[98]. In the past few years, a strong correlation between residual free chlorine used in drinking water sterilization and the negative effects on health have been confirmed [82]. Water chlorine react with organic matters that may found in water producing chlorinated hydrocarbons, which have been linked with many water borne diseases [82]. Disinfection by-products formed when water chlorine react with organic matter have been reported to cause an adverse reproductive effect to pregnant women

including intrauterine growth retardation,spontaneous abortion and low birth weight [99]. Refer to table 3 for the effect water chlorine on the human body.

**Table 3.** Adverse health effect of chlorinated water drinking and/or exposure

<b>Method of contact</b>	<b>Body system</b>	<b>Remark</b>	<b>Ref</b>
Inhalation of high concentrations of chlorine	Respiratory system	Irritation and inflammation may occur of the throat and nose membranes in addition to possible occurrence of bronchitis and asthma.	[100, 101]
Swallowing high amount of chlorine	Digestive system	Vomiting, chest pain, irritation of the pharynx and esophagus, in addition to various types of cancer risk.	[102, 103]
Swallowing high amount of chlorine	Intestinal Microbiota	Development of food allergy in infancy, and increased risk of asthma.	[42]
Washing or during swimming	Eyes	Mild burns in the corneal membranes, but it may lead to an ulcer or even leave scars.	[104]
Bathing and ablution	Skin and hair	Potential scalp, skin redness and allergies in chlorine sensitive peoples. Destruction of hair proteins leading to dryness, fragile, losing color, and dandruff.	[38, 105]
Drinking and washing	Teeth	Decay and weakness of tooth, in addition to the possible loss of enamel by corrosive disintegration.	[106]
Swallowing high amount of chlorine	Reproductive system	Adverse reproductive effect in pregnant women, including, retardation of intrauterine growth, low birth weight and	[99, 107]

		even spontaneous abortion.	
--	--	----------------------------	--

#### ***4.2. Water microbes associated diseases***

Different microorganisms may present in drinking water and transmitted to human body either via drinking or upon consumption of food, if it washed or prepared with contaminated water [108]. Diverse range of drinking water pathogens have been isolated and studied, mainly include bacteria, viruses, parasites and in rare cases fungi. Some of these microbes has a serious health impact, which may be mortal, while other could be only mild [109].

##### **4.2.1. Bacteria**

Bacteria is the widest occurring pathogenic microbes affecting all water source types in almost all developed and developing countries [110]. *Escherichia coli* and *Enterococci* are two fecal coliforms commonly present in oceans as well as in fresh lakes and rivers water [111]. However, the presence of any amount of these bacterial in drinking water make it unsuitable for consumption. Fecal indicators have been used as an indicators for human or hot-blood animal wastes in drinking water, many scientists have linked the consumption of fecal coliform contaminated water with many diseases such as diarrhea, cramps, nausea, and headaches [112]. Odonkor et al. [113] used *Escherichia coli* as a tool for assessing disease risk for drinking water, and revealed that main causes of drinking water faecal contamination in all the studied samples were purported to be anthropogenic. Sometimes bacterial contamination of drinking water causes a serious kidney disease such as hemolytic uremic syndrome, and could be also cause potential lifelong complications [114]. Other water bacteria such as *Salmonella typhi*, *Vibrio cholera* and *Shigella* spp have been linked with water poisoning, cholera, typhoid fevers and bacillary dysentery [115]. Pedati et al. [116] investigated the outbreak resulted from *Campylobacteriosis* from municipal water supply, and reported

various symptoms of diarrhea, abdominal cramps, pain and fever. Numerous species of Mycobacterium have been isolated from hospital water distribution systems as well as from the potable water, which can cause nosocomial infections [117]. *Anabaena circinalis* another cyanobacterium that widely living in freshwater reservoirs in many countries all over the world [118]. The main health impact of this bacteria is the ability to produce neurotoxins, which cause respiratory arrest, and in most cases followed by death [119].

#### **4.2.2. Viruses**


Viruses are very small pathogens that can pass through all water filters, which retain bacteria. According to the reports of Centers for Disease Control and Prevention (CDC) viruses were responsible for more than 9% of all in the United States reported drinking water outbreaks from 1993 till 2012 [120]. Most drinking waterborne viruses have been reported to follow the typical fecal-oral route of infection [121]. Gall et al. [122] reported that more than 1.8 billion people worldwide rely on sewage-contaminated water, which is used in drinking, washing and/or cooking of food. Owing to their small size, viruses are persistent in the environment and able to significantly resist many of water treatment disinfection processes, making them difficult to remove or eliminate by particle separation [123]. Enteroviruses are types of non-enveloped viruses, which have been linked with number diseases as one of the drinking water borne pathogens [124]. Pearce-Walker et al. [125] reported the ability of these viruses to transmit mainly by fecal-oral routes, especially with drinking water and/or food, as well as ophthalmic secretions, dermal lesions, or soil. Many literatures have reported variety of diseases that caused by different types of enteroviruses; Echoviruses usually associated with aseptic meningitis diseases, encephalitis respiratory diseases, and common cold [126]. Coxsackieviruses have been reported to cause hand-foot-mouth disease, acute hemorrhagic conjunctivitis, aseptic meningitis, gastroenteritis, respiratory

system infections, and heart diseases[127]. In many of East Asian countries including China, Korea, Japan, and Taiwan, enteroviruses associated diseases are predominantly found during summer compared to the other seasons, suggesting that meteorological variables and the parameters of drinking water quality might influence the frequency and spread as well as distribution of enteroviruses [128-130].

#### **4.2.3. Protozoan parasites**

Protozoan parasites have been identified as the second most etiological responsible of the mortality among young and infant children in developing countries[131]. They are responsible for about 1.7 billion cases world wide of diarrhea, leading to annual of 842,000 deaths [17]. Bartram et al. [66] reported that drinking water transmitted parasitic diseases, particularly protozoa cause endemic and epidemic diseases not only in developing countries but all over the world. However, the former have better hygienic conditions, and thus parasitic protozoa in most of cases not considered as a major reason of these diseases [132]. Markedly high rates of protozoan associated diseases have been reported in the latter, where the consumption of both treated and untreated water occur [133]. Owing to their small size (1-17  $\mu\text{m}$ ), the cysts and oocysts of *Giardia* and *Cryptosporidium* can easily penetrate through the water treatment system, leading to purified drinking water outbreaks and epidemics [134]. The infection begins with drinking of cysts or oocysts with contaminated water, or by using such water in cold food preparation. Each single cyst produces two trophozoites inside the small intestine of the host, which undergo fast multiplication through longitudinal binary fission, and remain in the lumen (free or attached to the mucosa). Upon the transition of parasites toward the colon, encystation occurs and both trophozoites and cysts are suspended in the feces [135]. Cysts and oocysts in feces spread into the external environment and contaminate water, food or directly infect other hosts

[136]. Figure 3 present the life cycle of Giardia as one of typical drinking water pathogenic parasitic protozoa.



**Figure 3.** Schematic illustration of the life cycle of Giardia as one of typical drinking water pathogenic parasites. (Available online: [http://en.wikipedia.org/wiki/File:Giardia\\_life\\_cycle\\_en.svg](http://en.wikipedia.org/wiki/File:Giardia_life_cycle_en.svg)).


#### **4.2.4. Fungi**

Fungi have been reported to ubiquitously occur in different environments, including water, and many of fungal species are pathogenic and cause serious health problems[137]. De Toni et al. [138] investigated the effect of fungi on drinking water quality and reported that their presence can significantly change the smell and taste of drinking water, which makes it become unpleasant for drinking. However, many health issues have been reported in terms of the consumption of fungal contaminated water, Oliveira et al. [139] linked fungal contaminated water with respiratory illness, allergies and life-threatening meningitis, in addition to mycoses, invasive and contagious infections. Mycotoxins are one of many fungal secondary metabolites, which may cause food poisoning and be a life-threatening substance. Magwaza et al. [140] studied the occurrence and diversity of fungal waterborne in Africa and revealed their impact in many respiratory problems. Other studies have linked mycotoxins with more serious health issues such as reducing the immunity and even cancers [141, 142]. In most countries water microbial standards are based on bacterial indicators, such as fecal bacteria for the quality and safety of water for drinking and the presence of pathogenic bacteria and intestinal protozoa [143]. Thus, only the oral-fecal route is mostly taken in consideration, and as it is not the main path for most fungal entry into water systems, not much has been done regarding investigation and monitoring of drinking water fungi.

#### ***4.3. Health precaution***

Treatment of drinking water may decrease or eliminate most or even all the microorganisms present in water however, recontamination of treated water may occur during the distribution in the network on the way to consumers, or even if the consumers are not following the suitable hygienic behavior [144]. Obtaining rapid and affordable frequent assessments of the microbiological quality of drinking water at

the point of treatment as well as at the point where it reach the consumers is necessary to ensure the safety of consumed drinking water [145]. Indicator bacterial species such as, *Enterococcus spp*, coliforms,*Escherichia coli*and total bacterial countare the most frequently monitored, as the rest of pathogenic indicators tend to be more expensive or impractical. However, detection of other types of pathogens such as parasites and viruses is not performed in many countries, as it requires special labs and advanced skills [146]. Although the presence of *Escherichia coli*and fecal coliforms in drinking water indicates its contamination and its unsuitability for drinking, their absence does not particularly preclude the presence of one or more other harmful organisms [56]. Generalized monitoring system must be developed that can be affordable and easy to use among different technicians in developed and developing countries [147]. Omarova et al. [63]discussed the possibility to achieve drinking water free from bacteria, protozoan parasites, and even viruses as presented in figure 4. Following non hygienic personal behaviors will affect the person himself and the surrounding people who share the same water source. Effective drinking water treatment technique and hygienic behavior from all the consumers are necessary to maintain and achieve drinking water sustainable development goals.



**Figure 4.** Schematic illustration of systems approach to ensure improved drinking water, sanitation and hygiene. Adapted from [63]

## 5. Conclusion

The inaccessibility to safe and clean drinking water has been always a big challenge facing humanity in past and current the centuries. This paper reviewed different types of drinking water contaminants, including physico-chemical and biological contaminants and their effects on human health, and the most used water treatment technologies in term of microbial disinfection. In addition, this review presents water borne diseases resulted from both drinking water disinfectant and water microorganisms, and highlight health precaution regarding drinking water microorganisms. Many countries only uses bacterial indicators to judge the quality of drinking water, and ignore the other pathogenic microorganisms such as parasitic

protozoa, viruses and even fungi. However, this review highlighted the health impact of drinking water disinfectant that commonly used for water treatment and presented the health precaution to ensure the sustainability and safety of water consumption.

## 6. References

1. Organization, W.H., *Progress on drinking water, sanitation and hygiene: 2017 update and SDG baselines*. 2017.
2. Abodo, A., *Statistical analysis of the trend of diarrheal diseases in children under five years in Mulago National Referral Hospital, Kampala Uganda from 2015 to 2018*. 2019, Makerere University.
3. Bentham, R. and H. Whiley, *Quantitative microbial risk assessment and opportunist waterborne infections—are there too many gaps to fill?* International journal of environmental research and public health, 2018. **15**(6): p. 1150.
4. Organization, W.H., *Inheriting a sustainable world? Atlas on children's health and the environment*. 2017: World Health Organization.
5. Cissé, G., *Food-borne and water-borne diseases under climate change in low-and middle-income countries: Further efforts needed for reducing environmental health exposure risks*. Acta tropica, 2019. **194**: p. 181-188.
6. McClung, R.P., et al., *Waterborne disease outbreaks associated with environmental and undetermined exposures to water—United States, 2013–2014*. MMWR. Morbidity and mortality weekly report, 2017. **66**(44): p. 1222.
7. Murphy, H., et al., *A systematic review of waterborne disease burden methodologies from developed countries*. Journal of water and health, 2014. **12**(4): p. 634-655.
8. Clasen, T.F., et al., *Interventions to improve water quality for preventing diarrhoea*. Cochrane database of systematic reviews, 2015(10).
9. Organization, W.H., *UN-Water global annual assessment of sanitation and drinking-water (GLAAS) 2012 report: the challenge of extending and sustaining services*. 2012.
10. Yahya, E.B., et al., *The Effect of Natural and Petroleum Based Materials on the Growth Rate and Antibiotic Sensitivity of Pseudomonas aeruginosa*. International Journal for Research in Applied Sciences and Biotechnology, 2020. **7**(5): p. 295-298.
11. Rizal, S., et al., *Properties of Macroalgae Biopolymer Films Reinforcement with Polysaccharide Microfibre*. Polymers, 2020. **12**(11): p. 2554.
12. Allaire, M., H. Wu, and U. Lall, *National trends in drinking water quality violations*. Proceedings of the National Academy of Sciences, 2018. **115**(9): p. 2078-2083.
13. Ibrahim Mohamed Ali, S.M.A., Abdulmutalib Alabeed Allaq, Rasheed Abdulsalam and *Prevalence of Antibiotic Use and Factors Associated With Self-medication among University Students in Malaysia* Journal of Pharmaceutical Research International, 2021.
14. Abdulmutalib Alabeed Allaq, N.J.S., Aziyah Abdul-Aziz, Abdul Mutalib Alabeed Alkamil, Asita Elengoe, Esam Bashir Yahya, Muhanad Abdullah Abdulsamad, *Epidemiological studies of the novel Coronavirus (COVID-19) in Libya*. Pakistan Journal of Biotechnology, 2021. **VOL. 18 NO. (1-2) (2021): PAK. J. BIOTECHNOLOGY**(1-2): p. 7-16.
15. Noah Mohammed Saleh \*, , Ali Mohammed Saleh 2 , , Raed Abdulkareem Hasan1, Hadi Hamdi Mahdi3, *The Renewable, Sustainable, and Clean Energy in Iraq Between Reality and Ambition According to the Paris Agreement on Climate Change*. Mesopotamian Journal of Big Data, 2022: p. 36-43.

16. Bonadonna, L. and G. La Rosa, *A review and update on waterborne viral diseases associated with swimming pools*. International journal of environmental research and public health, 2019. **16**(2): p. 166.
17. Efstratiou, A., J.E. Ongerth, and P. Karanis, *Waterborne transmission of protozoan parasites: review of worldwide outbreaks—an update 2011–2016*. Water research, 2017. **114**: p. 14-22.
18. Isa, M.L.M., M.A. Jainul, and A.M. Yusof, *Review on the Control of Waterborne Diseases—The Islamic Approach*. IIUM Medical Journal Malaysia, 2018. **17**(2).
19. Hadi Hamdi Mahdi , A.M.S., Noah Mohammed Saleh , , *Production of Biofuels from Biomass as an Approach Towards Sustainable Development- A Short Review.pdf*. NTU JOURNAL FOR RENEWABLE ENERGY. **Vol. 3, No. 1**(P-ISSN: 2788-9912 E-ISSN: 2788-9920 ): p. 9-21.
20. Palansooriya, K.N., et al., *Occurrence of contaminants in drinking water sources and the potential of biochar for water quality improvement: A review*. Critical Reviews in Environmental Science and Technology, 2020. **50**(6): p. 549-611.
21. Yahya, E. and M.A. Abdulsamad, *In-vitro Antibacterial Activity of Carbopol-Essential Oils hydrogels*. Journal of Applied Science & Process Engineering, 2020. **7**(2): p. 564-571.
22. Shakya, A.K. and P.K. Ghosh, *Simultaneous removal of arsenic, iron and nitrate in an attached growth bioreactor to meet drinking water standards: Importance of sulphate and empty bed contact time*. Journal of Cleaner Production, 2018. **186**: p. 1011-1020.
23. Topalcengiz, Z., L.K. Strawn, and M.D. Danyluk, *Microbial quality of agricultural water in Central Florida*. PloS one, 2017. **12**(4): p. e0174889.
24. Yahya, E.B., et al., *Antibacterial cellulose-based aerogels for wound healing application: A review*. Biomedical Research and Therapy, 2020. **7**(10): p. 4032-4040.
25. Mina, S.A., et al., *Quality assessment of commercially supplied drinking jar water in Chittagong City, Bangladesh*. Applied water science, 2018. **8**(1): p. 24.
26. Amr, S.A. and M.M. Yassin, *Microbial contamination of the drinking water distribution system and its impact on human health in Khan Yunis Governorate, Gaza Strip: seven years of monitoring (2000–2006)*. Public health, 2008. **122**(11): p. 1275-1283.
27. Pant, N.D., N. Poudyal, and S.K. Bhattacharya, *Bacteriological quality of bottled drinking water versus municipal tap water in Dharan municipality, Nepal*. Journal of Health, Population and Nutrition, 2016. **35**(1): p. 17.
28. Prest, E.I., et al., *Biological stability of drinking water: controlling factors, methods, and challenges*. Frontiers in microbiology, 2016. **7**: p. 45.
29. Rapant, S. and K. Krčmová, *Health risk assessment maps for arsenic groundwater content: application of national geochemical databases*. Environmental Geochemistry and Health, 2007. **29**(2): p. 131-141.
30. Muhammad, S., M.T. Shah, and S. Khan, *Health risk assessment of heavy metals and their source apportionment in drinking water of Kohistan region, northern Pakistan*. Microchemical journal, 2011. **98**(2): p. 334-343.
31. Emmanuel, E., M.G. Pierre, and Y. Perrodin, *Groundwater contamination by microbiological and chemical substances released from hospital wastewater: Health risk assessment for drinking water consumers*. Environment international, 2009. **35**(4): p. 718-726.
32. Gul, N., et al., *Arsenic and heavy metals contamination, risk assessment and their source in drinking water of the Mardan District, Khyber Pakhtunkhwa, Pakistan*. Journal of Water and Health, 2015. **13**(4): p. 1073-1084.
33. Lu, S.-Y., et al., *Trace elements contamination and human health risk assessment in drinking water from Shenzhen, China*. Environmental monitoring and assessment, 2015. **187**(1): p. 4220.

34. Arshad, N. and S. Imran, *Assessment of arsenic, fluoride, bacteria, and other contaminants in drinking water sources for rural communities of Kasur and other districts in Punjab, Pakistan*. Environmental Science and Pollution Research, 2017. **24**(3): p. 2449-2463.
35. Jordao, C., et al., *Assessment of water systems for contaminants from domestic and industrial sewages*. Environmental Monitoring and Assessment, 2002. **79**(1): p. 75-100.
36. Yahya, E.B., et al., *Growth rate and antibiotic sensitivity effect of some natural and petroleum based materials on Staphylococcus aureus*. International Journal for Research in Applied Sciences and Biotechnology, 2020. **7**(5): p. 7-11.
37. Sapkota, S., et al., *Drinking water quality of Sagarmatha National Park, Nepal*. Agriculture and Society in Changing Climate, 2019: p. 146.
38. Mazhar, M.A., et al., *Chlorination disinfection by-products in Municipal drinking water—A review*. Journal of Cleaner Production, 2020: p. 123159.
39. Asghari, F.B., et al., *Data on assessment of groundwater quality with application of ArcGIS in Zanjan, Iran*. Data in brief, 2018. **18**: p. 375.
40. Chetty, A.A. and S. Prasad, *Flow injection analysis of nitrate and nitrite in commercial baby foods*. Food chemistry, 2016. **197**: p. 503-508.
41. Mohammadi, A.A., et al., *Skeletal fluorosis in relation to drinking water in rural areas of West Azerbaijan, Iran*. Scientific reports, 2017. **7**(1): p. 1-7.
42. Li, X.-F. and W.A. Mitch, *Drinking water disinfection byproducts (DBPs) and human health effects: multidisciplinary challenges and opportunities*. 2018, ACS Publications.
43. Lorenzen, J., R. Jacobsen, and A. Astrup. *Effect of short-term high dietary calcium intake on 24-h energy expenditure, fat oxidation, and fecal fat excretion*. in INTERNATIONAL JOURNAL OF OBESITY. 2004. NATURE PUBLISHING GROUP MACMILLAN BUILDING, 4 CRINAN ST, LONDON N1 9XW, ENGLAND.
44. Chakraborti, D., et al., *Arsenic groundwater contamination and its health effects in Patna district (capital of Bihar) in the middle Ganga plain, India*. Chemosphere, 2016. **152**: p. 520-529.
45. Power, K.N. and L.A. Nagy, *Relationship between bacterial regrowth and some physical and chemical parameters within Sydney's drinking water distribution system*. Water Research, 1999. **33**(3): p. 741-750.
46. Phiri, B.J., et al., *Microbial contamination in drinking water at public outdoor recreation facilities in New Zealand*. Journal of Applied Microbiology, 2020.
47. Ward, J.S., et al., *Tryptophan-like fluorescence as a high-level screening tool for detecting microbial contamination in drinking water*. Science of The Total Environment, 2020. **750**: p. 141284.
48. Khan, S., et al., *Drinking water quality and human health risk in Charsadda district, Pakistan*. Journal of cleaner production, 2013. **60**: p. 93-101.
49. Radfard, M., et al., *Drinking water quality and arsenic health risk assessment in Sistan and Baluchestan, Southeastern Province, Iran*. Human and Ecological Risk Assessment: An International Journal, 2018.
50. Qin, K., et al., *Opportunistic pathogens and microbial communities and their associations with sediment physical parameters in drinking water storage tank sediments*. Pathogens, 2017. **6**(4): p. 54.
51. Paramita, S., et al., *A Short Overview of Curcuma aeruginosa with Curative Potentials Against COVID-19*. Asian Journal of Chemistry, 2021. **33**(4): p. 789-792.
52. Samuel, O., N. Florence, and O. Ifeanyi, *Microbial quality assessment of commercial bottled water brands in major markets in Awka, Nigeria*. Universal Journal of Microbiology Research, 2016. **4**: p. 1-5.

53. Suzuki, Y., et al., *Bacterial flora analysis of coliforms in sewage, river water, and ground water using MALDI-TOF mass spectrometry*. Journal of Environmental Science and Health, Part A, 2018. **53**(2): p. 160-173.
54. Mishra, V.P., et al., *Assessment of bacteriological quality of drinking water in Belagavi City, India*. MicroMedicine, 2019. **7**(1): p. 19-25.
55. Wen, X., et al., *Microbial Indicators and Their Use for Monitoring Drinking Water Quality—A Review*. Sustainability, 2020. **12**(6): p. 2249.
56. Pascual-Benito, M., et al., *Assessment of dead-end ultrafiltration for the detection and quantification of microbial indicators and pathogens in the drinking water treatment processes*. International Journal of Hygiene and Environmental Health, 2020. **230**: p. 113628.
57. Van Nevel, S., et al., *Flow cytometric bacterial cell counts challenge conventional heterotrophic plate counts for routine microbiological drinking water monitoring*. Water Research, 2017. **113**: p. 191-206.
58. Rygala, A., J. Berlowska, and D. Kregiel, *Heterotrophic Plate Count for Bottled Water Safety Management*. Processes, 2020. **8**(6): p. 739.
59. Horn, S., R. Pieters, and C. Bezuidenhout, *Pathogenic features of heterotrophic plate count bacteria from drinking-water boreholes*. Journal of water and health, 2016. **14**(6): p. 890-900.
60. Manickum, T., *Total Colony Counts (TCC) By Flow Cytometry (FCM) Should Replace the Heterotrophic Plate Count (HPC) Test for Bacteriological Enumeration of Water—Some Recent Developments in Flow Cytometry: A Review*.
61. van der Wielen, P.W. and D. van der Kooij, *Effect of water composition, distance and season on the adenosine triphosphate concentration in unchlorinated drinking water in the Netherlands*. Water Research, 2010. **44**(17): p. 4860-4867.
62. Ighalo, J.O. and A.G. Adeniyi, *A comprehensive review of water quality monitoring and assessment in Nigeria*. Chemosphere, 2020: p. 127569.
63. Omarova, A., et al., *Protozoan parasites in drinking water: A system approach for improved water, sanitation and hygiene in developing countries*. International journal of environmental research and public health, 2018. **15**(3): p. 495.
64. Saheb, E.J., *The prevalence of parasitic protozoan diseases in Iraq, 2016*. Karbala International Journal of Modern Science, 2018. **4**(1): p. 21-25.
65. Huang, D.B. and A.C. White, *An updated review on Cryptosporidium and Giardia*. Gastroenterology Clinics, 2006. **35**(2): p. 291-314.
66. Bartram, J. and G. Craun, *Waterborne zoonoses: identification, causes, and control*. 2004: World Health Organization.
67. Jain, S., et al., *Current and emerging tools for detecting protozoan cysts and oocysts in water*. TrAC Trends in Analytical Chemistry, 2019. **121**: p. 115695.
68. Ding, S., et al., *Disinfection byproduct formation during drinking water treatment and distribution: A review of unintended effects of engineering agents and materials*. Water research, 2019. **160**: p. 313-329.
69. Adeyemo, F.E., et al., *Efficiency of chlorine and UV in the inactivation of Cryptosporidium and Giardia in wastewater*. PloS one, 2019. **14**(5): p. e0216040.
70. Abdulmutalib Alabeed Allaq1, Norrizah Jaafar Sidik1, Aziyah Abdul-Aziz1, Idris Adewale Ahmed2, *Cumin (Cuminum cyminum L.): A review of its ethnopharmacology, phytochemistry*. Biomedical Research and Therapy,, 2020. **7**(9):4016-4021: p. 4016-4021.
71. Price, J.I. and M.T. Heberling, *The effects of source water quality on drinking water treatment costs: a review and synthesis of empirical literature*. Ecological Economics, 2018. **151**: p. 195-209.

72. Novotna, K., et al., *Microplastics in drinking water treatment—Current knowledge and research needs*. Science of The Total Environment, 2019. **667**: p. 730-740.
73. Gelete, G., et al., *Evaluating disinfection techniques of water treatment*. Desalination and Water Treatment, 2020. **177**: p. 408-415.
74. Pichel, N., M. Vivar, and M. Fuentes, *The problem of drinking water access: A review of disinfection technologies with an emphasis on solar treatment methods*. Chemosphere, 2019. **218**: p. 1014-1030.
75. Chatzisymeon, E., et al., *Disinfection of water and wastewater by UV-A and UV-C irradiation: application of real-time PCR method*. Photochemical & Photobiological Sciences, 2011. **10**(3): p. 389-395.
76. Yang, F., et al., *Haloacetonamides versus haloacetonitriles formation and toxicity in chloraminated drinking water*. Journal of hazardous materials, 2014. **274**: p. 156-163.
77. Ding, W., et al., *Ozone disinfection of chlorine-resistant bacteria in drinking water*. Water research, 2019. **160**: p. 339-349.
78. De Pasquale, A., et al., *District heating by drinking water heat pump: Modelling and energy analysis of a case study in the city of Milan*. Energy, 2017. **118**: p. 246-263.
79. Karakurt, S., et al., *Dynamics of wastewater effluent contributions in streams and impacts on drinking water supply via riverbank filtration in germany—a national reconnaissance*. Environmental science & technology, 2019. **53**(11): p. 6154-6161.
80. Issa, H.M., *Comparative Analysis of Different Disinfection Techniques Performances in Drinking Water Treatment Plant Using a Process Simulation Software*. Zanco Journal of Pure and Applied Sciences, 2019. **31**(s3): p. 1-8.
81. Gagnon, G., et al., *Disinfectant efficacy of chlorite and chlorine dioxide in drinking water biofilms*. Water research, 2005. **39**(9): p. 1809-1817.
82. Mohsen, I.H., A.H. Mohsen, and H.K. Zaidan, *Health effects of chlorinated water: a review article*. Pak. J. Biotechnol, 2019. **16**: p. 163-167.
83. Pickering, A.J., et al., *Effect of in-line drinking water chlorination at the point of collection on child diarrhoea in urban Bangladesh: a double-blind, cluster-randomised controlled trial*. The Lancet Global Health, 2019. **7**(9): p. e1247-e1256.
84. Ali Mohammed Saleh 1, a), Noah Mohammed Saleh 2,b), Hadi Hamdi Mahdi 1c, *Production of Biofuels from Biomass as an Approach Towards Sustainable Development: A Short Review*. NTU JOURNAL FOR RENEWABLE ENERGY, 2022. **Vol. 3, No. 1**(P-ISSN: 2788-9912 E-ISSN: 2788-9920 ): p. 9-20.
85. Gomez-Alvarez, V., et al., *Resilience of microbial communities in a simulated drinking water distribution system subjected to disturbances: role of conditionally rare taxa and potential implications for antibiotic-resistant bacteria*. Environmental Science: Water Research & Technology, 2016. **2**(4): p. 645-657.
86. Wang, H., et al., *Microbial community response to chlorine conversion in a chloraminated drinking water distribution system*. Environmental science & technology, 2014. **48**(18): p. 10624-10633.
87. Yang, F., et al., *Effect of sulfate on the transformation of corrosion scale composition and bacterial community in cast iron water distribution pipes*. Water Research, 2014. **59**: p. 46-57.
88. Hasan Mohammed Hamid Agha<sup>1</sup>, A.M.S., Hadi Hamdi Mahdi<sup>3</sup>, Amjad Abdulhadi Mohammed<sup>4</sup>, Abdulmutalib Alabeed Alkamil Allaq<sup>5</sup>, *OVERVIEW OF EFFECT OF PLASTIC WASTE POLLUTION ON MARINE ENVIRONMENT*. Journal of Asian Scientific Research, 2022. **Vol. 12, No. 4, 260-268**.(226-5724): p. 260.
89. Spagnolo, A.M., et al., *Hospital water and prevention of waterborne infections*. Reviews in Medical Microbiology, 2016. **27**(1): p. 25-32.

90. Allou, N., et al., *Waterborne Infections in Reunion Island, 2010–2017*. The American Journal of Tropical Medicine and Hygiene, 2018. **99**(3): p. 578-583.
91. Ghernaout, D., *Disinfection and DBPs removal in drinking water treatment: A perspective for a green technology*. International Journal of Advances in Applied Sciences, 2018. **5**: p. 108-117.
92. Maramraj, K.K., et al., *A community-wide acute diarrheal disease outbreak associated with drinking contaminated water from shallow bore-wells in a tribal village, India, 2017*. BMC Public Health, 2020. **20**(1): p. 1-8.
93. Organization, W.H., *Preventing diarrhoea through better water, sanitation and hygiene: exposures and impacts in low-and middle-income countries*. 2014: World Health Organization.
94. Bivins, A.W., et al., *Estimating infection risks and the global burden of diarrheal disease attributable to intermittent water supply using QMRA*. Environmental science & technology, 2017. **51**(13): p. 7542-7551.
95. Rutala, W.A. and D.J. Weber, *Water as a reservoir of nosocomial pathogens*. Infection Control & Hospital Epidemiology, 1997. **18**(9): p. 609-616.
96. Decker, B.K. and T.N. Palmore, *The role of water in healthcare-associated infections*. Current opinion in infectious diseases, 2013. **26**(4): p. 345.
97. Hossain, S., et al., *Spectrophotometric Online Detection of Drinking Water Disinfectant: A Machine Learning Approach*. Sensors, 2020. **20**(22): p. 6671.
98. Van Haute, S., et al., *Chlorine dioxide as water disinfectant during fresh-cut iceberg lettuce washing: disinfectant demand, disinfection efficiency, and chlorite formation*. LWT, 2017. **75**: p. 301-304.
99. Richardson, S.D., J.E. Simmons, and G. Rice, *Peer reviewed: disinfection byproducts: the next generation*. 2002, ACS Publications.
100. Bougault, V., et al., *Airway remodeling and inflammation in competitive swimmers training in indoor chlorinated swimming pools*. Journal of allergy and clinical immunology, 2012. **129**(2): p. 351-358. e1.
101. Bougault, V., et al., *The respiratory health of swimmers*. Sports Medicine, 2009. **39**(4): p. 295-312.
102. Villanueva, C.M., et al., *Bladder cancer and exposure to water disinfection by-products through ingestion, bathing, showering, and swimming in pools*. American journal of epidemiology, 2007. **165**(2): p. 148-156.
103. Evlampidou, I., et al., *Trihalomethanes in drinking water and bladder cancer burden in the European Union*. Environmental health perspectives, 2020. **128**(1): p. 017001.
104. Bernard, A., *Chlorination products: emerging links with allergic diseases*. Current medicinal chemistry, 2007. **14**(16): p. 1771-1782.
105. Wu, A., *The Effects of Chlorine on Different Ethnicities of Hair*. 2018.
106. Dawes, C. and C.L. Boroditsky, *Rapid and severe tooth erosion from swimming in an improperly chlorinated pool: case report*. Journal of the Canadian Dental Association, 2008. **74**(4).
107. Mashau, F., E.J. Ncube, and K. Voyi, *Drinking water disinfection by-products exposure and health effects on pregnancy outcomes: a systematic review*. Journal of water and health, 2018. **16**(2): p. 181-196.
108. Gomes, F.M.d.S., et al., *Access to drinking water and sewage treatment in Brazil: a challenge for the control of waterborne infectious diseases*. Revista do Instituto de Medicina Tropical de São Paulo, 2020. **62**.
109. Pal, M., et al., *Public health hazards due to unsafe drinking water*. Air Water Borne Dis, 2018. **7**(1000138): p. 2.
110. Bain, R., et al., *Global assessment of exposure to faecal contamination through drinking water based on a systematic review*. Tropical Medicine & International Health, 2014. **19**(8): p. 917-927.

111. Stocker, M.D., et al., *Seasonality of E. coli and Enterococci Concentrations in Creek Water, Sediment, and Periphyton*. Water, Air, & Soil Pollution, 2019. **230**(9): p. 223.
112. Niroumand, A., et al., *A Survey on Drinking Water Contamination to Indicator Bacteria in Dairy Farms of Mashhad Suburb*. Iranian Journal of Veterinary Medicine, 2020. **14**(3): p. 251-259.
113. Odonkor, S.T. and T. Mahami, *Escherichia coli as a Tool for Disease Risk Assessment of Drinking Water Sources*. International Journal of Microbiology, 2020. **2020**.
114. Crude, S.E. and R. Walker, *Tragedy could have been prevented*. Opflow, 2005. **31**(6): p. 1-7.
115. Hess-Erga, O.-K., et al., *Microorganisms in ballast water: disinfection, community dynamics, and implications for management*. Science of the Total Environment, 2019. **657**: p. 704-716.
116. Pedati, C., et al., *Campylobacteriosis Outbreak Associated with Contaminated Municipal Water Supply—Nebraska, 2017*. Morbidity and Mortality Weekly Report, 2019. **68**(7): p. 169.
117. Loret, J.-F. and N. Dumoutier, *Non-tuberculous mycobacteria in drinking water systems: A review of prevalence data and control means*. International Journal of Hygiene and Environmental Health, 2019. **222**(4): p. 628-634.
118. Abobaker, H.M. and H.E. Elsalhin, *Antibacterial Activity of Anabaena circinalis Isolated from Fresh Water*. Journal of Advances in Microbiology, 2019: p. 1-7.
119. Li, J. and K.M. Persson, *Paralytic Shellfish toxins (PST toxins) monitoring in freshwaters for drinking water supply-A review*. Chemosphere, 2020: p. 128591.
120. WHO, M.S.F.M., et al., *Essential medicines and health products*. 2015.
121. Xagorarakis, I., Z. Yin, and Z. Svambayev, *Fate of viruses in water systems*. Journal of Environmental Engineering, 2014. **140**(7): p. 04014020.
122. Gall, A.M., et al., *Waterborne viruses: a barrier to safe drinking water*. PLoS Pathog, 2015. **11**(6): p. e1004867.
123. Tanneru, C.T., J.D. Rimer, and S. Chellam, *Sweep flocculation and adsorption of viruses on aluminum flocs during electrochemical treatment prior to surface water microfiltration*. Environmental science & technology, 2013. **47**(9): p. 4612-4618.
124. Moazeni, M., et al., *Estimation of health risks caused by exposure to enteroviruses from agricultural application of wastewater effluents*. Water research, 2017. **125**: p. 104-113.
125. Pearce-Walker, J., et al., *Managing leafy green safety from adenoviruses and enteroviruses in irrigation water*. Agricultural Water Management, 2020. **240**: p. 106272.
126. Shirasaki, N., et al., *Elimination of representative contaminant candidate list viruses, coxsackievirus, echovirus, hepatitis A virus, and norovirus, from water by coagulation processes*. Journal of hazardous materials, 2017. **326**: p. 110-119.
127. El-Senousy, W.M., et al., *Coxsackievirus B4 as a causative agent of diabetes mellitus type 1: is there a role of inefficiently treated drinking water and sewage in virus spreading?* Food and environmental virology, 2018. **10**(1): p. 89-98.
128. Kim, B.I., et al., *Effect of climatic factors on hand, foot, and mouth disease in South Korea, 2010-2013*. PLoS One, 2016. **11**(6): p. e0157500.
129. Bukhari, N., et al., *Prevalence of Human Papilloma Virus Sub Genotypes following Head and Neck Squamous Cell Carcinomas in Asian Continent, A Systematic Review Article*. Asian Pacific Journal of Cancer Prevention: APJCP, 2019. **20**(11): p. 3269.
130. Saadoun, I., et al., *Molecular Surveillance of Enteroviruses in Al-Zarqa River, Jordan*. Jordan Journal of Biological Sciences, 2020. **13**(3).
131. Ahmed, S.A., M. Guerrero Flórez, and P. Karanis, *The impact of water crises and climate changes on the transmission of protozoan parasites in Africa*. Pathogens and global health, 2018. **112**(6): p. 281-293.
132. Fletcher, S.M., et al., *Enteric protozoa in the developed world: a public health perspective*. Clinical microbiology reviews, 2012. **25**(3): p. 420-449.

133. Mtapuri-Zinyowera, S., et al., *Human parasitic protozoa in drinking water sources in rural Zimbabwe and their link to HIV infection*. *Germs*, 2014. **4**(4): p. 86.
134. Smith, H., *Detection of parasites in the environment*. *Parasitology*, 1999. **117**(7): p. 113-141.
135. Koehler, A.V., et al., *Cryptosporidium and Giardia taxa in faecal samples from animals in catchments supplying the city of Melbourne with drinking water (2011 to 2015)*. *Parasites & vectors*, 2016. **9**(1): p. 315.
136. Borowski, H., P.L. Clode, and R.A. Thompson, *Active invasion and/or encapsulation? A reappraisal of host-cell parasitism by Cryptosporidium*. *Trends in parasitology*, 2008. **24**(11): p. 509-516.
137. Arroyo, M.G., et al., *Broad Diversity of Fungi in Hospital Water*. *The Scientific World Journal*, 2020. **2020**.
138. De Toni, P.S.-A. and K. Reilly, *A REVIEW OF FUNGI IN DRINKING WATER AND THE IMPLICATIONS FOR HUMAN HEALTH*. 2011.
139. Oliveira, H., et al., *Fungi from a groundwater-fed drinking water supply system in Brazil*. *International Journal of Environmental Research and Public Health*, 2016. **13**(3): p. 304.
140. Magwaza, N.M., et al., *The occurrence and diversity of waterborne fungi in African aquatic systems: their impact on water quality and human health*. *International journal of environmental research and public health*, 2017. **14**(5): p. 546.
141. Pitt, J., et al., *Mycotoxins and toxigenic fungi*. *Medical mycology*, 2000. **38**(sup1): p. 41-46.
142. Bhat, R., R.V. Rai, and A.A. Karim, *Mycotoxins in food and feed: present status and future concerns*. *Comprehensive reviews in food science and food safety*, 2010. **9**(1): p. 57-81.
143. Besner, M.-C., M. Prévost, and S. Regli, *Assessing the public health risk of microbial intrusion events in distribution systems: conceptual model, available data, and challenges*. *Water research*, 2011. **45**(3): p. 961-979.
144. Tambekar, D., S. Dhote, and G. Shinde, *Household drinking water: Assessment of microbiological contamination between source and point of use*. *Bioscience Discovery*, 2016. **7**(2): p. 152-7.
145. Wu, J., et al., *Are microbial indicators and pathogens correlated? A statistical analysis of 40 years of research*. *Journal of water and health*, 2011. **9**(2): p. 265-278.
146. García-Aljaro, C., et al., *Mobilisation of microbial indicators, microbial source tracking markers and pathogens after rainfall events*. *Water research*, 2017. **112**: p. 248-253.
147. Hazarika, R. and R. Jandl, *The Nexus between the Austrian Forestry Sector and the Sustainable Development Goals: A Review of the Interlinkages*. *Forests*, 2019. **10**(3): p. 205.