

Review Article

Soil Management Practices to Enhance Carbon Sequestration Rates- A review

Abstract

Agroforestry systems represent an integrative approach to modern agriculture that inherently addresses the challenges posed by climate change while simultaneously advancing sustainability objectives. The core of agroforestry, defined as the strategic integration of trees within agricultural landscapes, has been articulated to illustrate its capacity in enhancing ecosystem services. Historical antecedents of agroforestry underscore its evolution as a practice intrinsically linked with sustainable resource use. An exploration of the different types of agroforestry systems—silvopastoral, agro-silvicultural, and silvoarable—highlights the diversity of practices adaptable to various ecological and socio-economic conditions. By delving into the mechanisms by which agroforestry enhances soil quality, promotes effective water management, increases carbon sequestration, preserves biodiversity, and regulates microclimates, the review substantiates the environmental merits of agroforestry. The socioeconomic dimension is explored through the lens of agroforestry's contribution to economic benefits such as yield increases and income diversification, along with social benefits including community engagement and the preservation of traditional knowledge. The review also encompasses a series of regional case studies, examining the implementation of agroforestry across continents and its impacts on local and regional scales. This global perspective not only illustrates the practical implications of agroforestry but also the adaptive strategies pertinent to specific climatic and socio-economic contexts. Despite its significant potential, agroforestry is not devoid of challenges. Land tenure complexities, financial limitations, cultural barriers, and policy obstructions are discussed to provide an objective view of the impediments to agroforestry adoption. These constraints necessitate a structured approach towards advancing research, fostering technological innovation, and promoting policy reform to enable the scaling up of agroforestry practices.

Keywords: *Agroforestry, Climate-Adaptation, Sustainability, Carbon-Sequestration, Soil-Health*

Introduction

Carbon sequestration is a naturally occurring process by which carbon dioxide (CO₂) is removed from the atmosphere and stored in the Earth's carbon pools, including forests, oceans, and soil [1]. The importance of carbon sequestration lies in its potential to mitigate the effects of climate change by reducing the atmospheric concentration of CO₂, a major greenhouse gas (GHG) contributing to global warming [2]. The soil acts as a significant carbon sink with the capacity to hold three times more carbon than the atmosphere and four times that of the biota [3]. Soil carbon sequestration occurs when CO₂ is absorbed by plants through photosynthesis and then transferred to the soil through root biomass and litterfall, where it is stored as soil organic carbon

(SOC) [4]. Agricultural practices and land use changes profoundly influence the flux of carbon, which can be quantified as the exchange of carbon between the soil and the atmosphere [5]. While natural ecosystems typically act as carbon sinks, agricultural activities, especially those involving soil tillage, can convert these sinks into sources of carbon emissions [6]. Tillage accelerates the decomposition of organic matter, thereby releasing CO_2 , while practices such as deforestation for agricultural expansion lead to significant carbon losses [7]. The review aims to delve into the nuances of soil management practices that enhance the rates of carbon sequestration, underscoring the potential of well-managed soils to serve as effective carbon sinks and thus, play a crucial role in climate change mitigation. The scope of the review extends to examining various soil management strategies, the factors influencing SOC dynamics, and the challenges and advancements in the monitoring and promotion of soil carbon sequestration. Through this comprehensive review, a detailed examination of the effects of diverse agricultural practices and land use on carbon flux is conducted, drawing on the latest research and findings from scientific studies. The role of soil as a carbon sink is explored in depth, acknowledging the myriad of factors that impact its capacity to sequester carbon, such as soil type, climate, land use, and management practices [8]. The review also offers an overview of the existing policies, economic incentives, and technological advancements that support sustainable soil management, thereby facilitating enhanced rates of carbon sequestration. It highlights the significance of adopting soil conservation practices that are not only environmentally sustainable but also economically viable for farmers and land managers [9].




Image 1: Soil Carbon Storage [10]

Fundamentals of Soil Carbon Sequestration

Understanding the fundamentals of soil carbon sequestration necessitates a grasp of the basic concepts and terminology pertinent to the domain, the dynamics of soil organic matter (SOM),

the multifaceted factors that affect carbon sequestration in soil, and the varied methods for measuring soil carbon stocks. Herein, each of these components is explored in depth to elucidate the foundational principles governing soil carbon sequestration. Soil carbon sequestration can be described as the process by which atmospheric carbon dioxide is captured by plants through photosynthesis and subsequently stored as carbon in the soil in the form of SOM [11]. The process of carbon sequestration involves several key steps, including the assimilation of CO₂ by plants, the incorporation of the carbon into plant tissue, and the transfer of detritus and root biomass into the soil, which through microbial action, becomes stabilized as SOM [12]. Carbon sequestration is not a singular phenomenon but rather a subset of the broader carbon cycle, which includes various carbon pools and fluxes. The principal pools include the atmosphere, biosphere, oceans, and terrestrial ecosystems, with soils being the largest terrestrial pool [13]. The fluxes represent the movement of carbon between these pools via processes such as respiration, decomposition, and combustion [14].

Soil Organic Matter Dynamics

SOM is composed of plant and animal residues at various stages of decomposition, cells and tissues of soil organisms, and substances synthesized by soil organisms [15]. The dynamics of SOM are a product of the balance between the input of organic carbon to the soil and its loss through decomposition, erosion, and leaching [16]. Carbon enters the soil organic pool through litter fall, root turnover, and microbial biomass turnover. The stabilization of organic carbon in soils is facilitated by its interaction with soil minerals, physical protection within soil aggregates, and chemical recalcitrance [17]. The stability and persistence of SOM are critical for carbon sequestration as they determine how long carbon remains sequestered in the soil.




Image 2: The role of soil and its management in moderating the global carbon cycle (adapted from [18]).

Factors Affecting Carbon Sequestration in Soil

Numerous factors influence the capacity of soils to sequester carbon, including climate, soil texture, soil structure, soil type, land use, vegetation type, and management practices [19]. Climate affects decomposition rates and biomass production, thereby influencing carbon inputs and outputs [20]. Soil texture and structure affect the soil's capacity to protect organic matter from decomposition, with fine-textured soils generally having greater potential for carbon sequestration [21]. Land use changes, such as deforestation or conversion of grasslands to croplands, usually result in the depletion of soil carbon stocks. Conversely, land management practices that restore vegetation, improve soil fertility, and reduce erosion are known to enhance carbon sequestration [22].

Methods for Measuring Soil Carbon Stocks

Assessing soil carbon stocks and the rate of carbon sequestration is vital for understanding the potential of soils to mitigate climate change and for informing land management decisions [23]. The most common method of measuring soil carbon stocks is by direct sampling and laboratory analysis, which provides the total carbon content of soil samples at various depths [24]. This approach, while accurate, can be labor-intensive and costly, especially for larger-scale assessments. Remote sensing technologies, coupled with geographic information systems (GIS), offer a promising alternative for estimating soil carbon stocks over extensive areas [25]. These methods involve using satellite or aerial imagery to derive vegetation indices that are correlated with above-ground biomass and, by extension, with soil carbon content. Additionally, soil carbon models have been developed to predict changes in carbon stocks under different scenarios. These models range from simple empirical models to more complex process-based models that simulate the biological, chemical, and physical processes affecting carbon dynamics in soils [26].

Table 1: Analysis of Methods for Measuring Soil Carbon Stocks [27]

Method	Description	Advantages	Disadvantages
Dry Combustion (Elemental Analysis)	Soil is oven-dried, ground, and combusted to measure CO ₂ release. The CO ₂ is quantified, which correlates to soil carbon content.	Highly accurate; provides total carbon content.	Requires expensive equipment; destroys soil samples.
Wet Oxidation (Walkley-Black Method)	This chemical method oxidizes soil organic matter using potassium dichromate, which is then titrated to determine carbon content.	Less expensive equipment; well-established method.	Can underestimate soil carbon; involves hazardous chemicals.
Near-Infrared Spectroscopy (NIRS)	Uses the absorbance of NIR light by organic matter to estimate carbon content.	Non-destructive; rapid analysis of multiple samples.	Calibration is required; less accurate for soils with low organic carbon.
Laser-Induced Breakdown Spectroscopy (LIBS)	A laser pulse is used to create a plasma on the soil surface, and the light emitted is analyzed to determine carbon content.	Rapid; can detect multiple elements simultaneously.	Complex calibration; high initial equipment cost.
Soil Gas Flux Measurement	Measures the amount of CO ₂ or other greenhouse gases emitted from the soil to estimate carbon levels.	Can measure changes over time; non-destructive.	Requires long-term monitoring; influenced by environmental factors.
Isotopic Analysis (13C/12C Ratio)	Measures the ratio of stable isotopes to infer the dynamics of soil organic carbon.	Can differentiate between sources of carbon.	Requires specialized equipment; more complex analysis.
Soil Core Sampling	Direct physical extraction of soil cores, which are then dried and weighed to estimate carbon density.	Direct measurement; can profile carbon at different depths.	Labor-intensive; may require large number of samples for accuracy.
Remote Sensing and Modeling	Uses satellite or aerial imagery combined with models to estimate soil carbon at a larger	Can cover large areas; useful for monitoring over time.	Indirect method; requires calibration with ground-truth data.

	scale.		
--	--------	--	--

3. Soil Management Practices

Soil management practices encompass a suite of techniques and strategies aimed at enhancing soil fertility, structure, water retention, and overall health. These practices are also integral to soil carbon sequestration, a critical ecosystem service that contributes to mitigating climate change by capturing atmospheric carbon dioxide and storing it in the soil. In this detailed exploration, we discuss various soil management practices, focusing on no-till farming and crop rotation, and their effects on soil structure, carbon storage, and overall soil health, substantiated by case studies and research findings.




Image 3: Agronomics measures and soil carbon Sequestration [28]

3.1 No-Till Farming

No-till farming is a cultivation technique where the soil is not disturbed by plowing or turning. Instead, seeds are directly drilled into the soil, and residues from previous crops are left on the field to decompose naturally [29]. This practice contrasts with conventional tillage, which often involves multiple passes with a plow to prepare the seedbed, control weeds, and incorporate residues.

Impact on Soil Structure and Carbon Storage

No-till farming can significantly impact soil structure and carbon storage. By avoiding tillage, the soil retains its natural stratification and aggregate stability, which is essential for good aeration, water infiltration, and root development [30]. As soil aggregates are not broken down through tillage, the organic carbon sequestered within them is less susceptible to oxidation and subsequent release as CO₂ [31].

A meta-analysis by Wang *et al.* [32] indicated that no-till could increase soil carbon stocks by an average of 0.57 Mg C ha⁻¹ yr⁻¹ in the top 30 cm of soil. The increase in carbon storage is due to both a reduction in the decomposition rate of soil organic matter and an increase in the input of plant residues to the soil surface.

Case Studies and Research Findings

Case studies from across the globe illustrate the benefits of no-till farming for carbon sequestration. In the Great Plains of the United States, no-till practices, coupled with cover cropping, have been shown to sequester up to 1.2 Mg C ha⁻¹ yr⁻¹ [33]. Similarly, research from Brazil's Cerrado region indicates that no-till systems can accumulate soil carbon at rates of 0.4 to 0.6 Mg C ha⁻¹ yr⁻¹ [34]. These increases in soil carbon under no-till farming are also associated with co-benefits such as reduced soil erosion and improved biodiversity [35].

3.2 Crop Rotation and Diversity

Crop rotation is the practice of growing different types of crops in succession on the same land to improve soil health and reduce the risk of pests and diseases. The principle underlying crop rotation is that different crops have varying nutrient requirements and rooting patterns, which, when altered over time, can prevent the depletion of specific soil nutrients and interrupt the life cycles of pests and diseases [36].

Benefits for Soil Health and Carbon Sequestration

Crop rotation can enhance soil health and carbon sequestration in several ways. By varying crop types, soil biodiversity is promoted, which can increase the decomposition of organic matter and subsequent humification, leading to more stable forms of soil carbon [37]. Legumes, often included in crop rotations, fix atmospheric nitrogen, thereby enriching soil fertility and potentially increasing biomass production and residue inputs to the soil, which can enhance carbon storage [38]. A study by Liu *et al.* [39] demonstrated that diversifying crop rotations increased soil organic carbon by as much as 8.5% over a 12-year period in comparison to monoculture rotations. This was attributed to the increased residue inputs and varied root structures that provided more substrates for soil microorganisms and improved soil aggregate stability. Effective crop rotation systems are those that maximize the benefits of diversity while meeting the economic needs of farmers. In the Midwest United States, a corn-soybean-wheat

rotation with cover crops has been shown to be effective in enhancing soil carbon sequestration [40]. In Europe, rotations that include deep-rooting crops like alfalfa have been found to increase soil carbon stocks in the subsoil layers [41].

Organic Amendments

Compost is an organic amendment produced from the aerobic decomposition of a diverse mix of organic materials, including yard trimmings, food waste, and manures. As an amendment, compost is valued for its ability to improve soil structure, water holding capacity, and fertility. Its application leads to an increase in soil organic matter content, a key factor in carbon sequestration. Compost introduces stable organic compounds into the soil matrix, providing a carbon-rich food source for soil microorganisms, which, through their metabolic processes, convert organic materials into humus, a stable form of soil organic carbon (SOC) [42]. Biochar is another type of organic amendment produced through pyrolysis, a process of thermal decomposition of organic material at high temperatures in the absence of oxygen. This process creates a carbon-rich product that can persist in the soil for hundreds to thousands of years, hence providing a long-term carbon sequestration solution. Biochar's effects on carbon sequestration are twofold. Its inherent stability in the soil means that it does not decompose rapidly, thereby sequestering carbon directly. Additionally, biochar influences the soil's physical and chemical properties, improving soil aggregation and porosity, which indirectly supports the stabilization of additional SOC by enhancing the physical protection of soil organic matter and promoting conditions conducive to increased microbial activity [43]. Manure, derived from animal excreta, often combined with bedding materials, is a traditional soil amendment that supplies not only organic carbon but also nutrients, particularly nitrogen, phosphorus, and potassium. Manure improves soil structure, increases water retention, and fosters microbial activity. The organic carbon present in manure can be converted into SOC, which contributes to the humus pool within the soil. This is particularly effective when manure is applied in conjunction with conservation tillage practices, which limit the disturbance of soil and protect SOC from oxidative losses [44].

The enhancement of carbon sequestration by these organic amendments occurs through several mechanisms. They provide substrates for soil microbial growth, leading to an increase in microbial biomass carbon, a component of SOC. These amendments also stimulate the formation of soil aggregates, within which SOC can be physically protected from decomposition. The addition of organic amendments can stimulate plant growth, leading to increased root biomass and exudates, both of which are sources of SOC [45]. The methods and rates of application for these organic amendments are critical factors in maximizing their benefits for carbon sequestration. Compost and manure are commonly applied to the soil surface and then incorporated into the soil by tillage or can be applied directly to the soil as a mulch. Application rates for compost and manure typically range from 1 to 10 metric tons per hectare, depending on the nutrient content of the amendment and the nutrient requirements of the crop [46]. Excessive application must be avoided to prevent nutrient runoff and potential negative environmental

impacts. Biochar application rates are typically lower than those for compost and manure, often ranging from 0.5 to 3 metric tons per hectare. The lower rate reflects biochar's high carbon content and its persistence in the soil. Application methods for biochar include broadcasting over the soil surface followed by incorporation, or direct application to the soil at the time of planting. The specific rate of application depends on the desired effects on soil properties and the initial soil conditions. Rates and methods of application can be adjusted based on local conditions, crop requirements, and specific management objectives [47]. The use of organic amendments is a practice steeped in both tradition and innovation. While the application of manure to soils is a practice that dates back to the beginning of agriculture, the scientific understanding of how such amendments affect soil carbon stocks has greatly advanced. Current practices emphasize not only the benefits to crop productivity but also the potential of these amendments to act as tools for climate change mitigation. This dual role is central to the value proposition of organic amendments in sustainable agriculture. The continuing evolution of application technologies and the expanding scientific knowledge base enable more precise management of these amendments, enhancing their environmental benefits while minimizing any potential adverse effects.

3.4 Cover Cropping

Cover cropping is a pivotal soil management practice with significant implications for enhancing soil carbon sequestration. Cover crops are planted not primarily for harvest, but to cover the soil, thereby offering numerous benefits such as reducing erosion, improving soil health, and increasing soil organic carbon stocks. Cover crops encompass a broad array of species, each with unique benefits and adaptabilities to different climatic and soil conditions. Leguminous cover crops, like clovers (*Trifolium* spp.) and vetches (*Vicia* spp.), are valued for their nitrogen-fixing capabilities, contributing to soil fertility. Non-leguminous cover crops, such as rye (*Secale cereale*) and oats (*Avena sativa*), are often chosen for their rapid growth, which quickly provides soil cover and biomass. Brassicas, like radishes (*Raphanus sativus*) and mustards (*Brassica* spp.), are noted for their deep rooting, which can improve soil structure and break up compacted layers [48].

Role in Protecting Soil and Adding Organic Carbon

Cover crops enhance the protection of soil and the addition of organic carbon in multiple ways. Their foliage provides a physical barrier against the impact of raindrops, reducing erosion and the loss of topsoil. The roots of cover crops stabilize soil aggregates, and upon decomposition, contribute organic matter and thus, carbon to the soil. This process enriches the soil with humic substances, increases the microbial biomass, and consequently, the soil carbon pool [49].

3.5 Integration with Other Farming Practices

The integration of cover crops with other farming practices is central to maximizing their benefits. In conservation tillage systems, cover crops are left on the soil surface as a mulch after they die, which can conserve moisture and further protect the soil from erosion. When used in

rotation with cash crops, cover crops can break disease cycles and reduce pest populations, while also enhancing the diversity of plant species and the structure of the agroecosystem, leading to more resilient farming systems [50]. Agroforestry, the practice of integrating woody perennials with agricultural crops, offers a sustainable approach to land management that can significantly impact below-ground carbon storage. The concept of agroforestry entails the strategic use of trees and shrubs in agricultural landscapes to create systems where the agricultural and ecological benefits of both are optimized. Trees provide shade, shelter, and habitat for diverse fauna, contribute to the cycling of nutrients, and enhance the aesthetic and economic value of the land. Agroforestry systems can be designed to produce fruits, nuts, timber, and fodder, in addition to the traditional agricultural products, diversifying farmer income and reducing risks associated with market and climate fluctuations [51]. The root systems of trees are extensive and often penetrate deeper soil layers than agricultural crops, which leads to the input of organic materials into the subsoil. This can result in significant below-ground carbon storage, as the organic matter in deeper soil layers is less subject to decomposition and can thus remain sequestered for longer periods. Additionally, the leaf litter and woody debris from trees contribute to the surface accumulation of organic matter, further enhancing soil carbon stocks [52].

4. Challenges and Limitations

Implementing soil carbon sequestration practices is fraught with challenges and limitations that span technical, socioeconomic, and environmental domains. These challenges are crucial to address for the successful mitigation of climate change impacts through improved soil management. Technical difficulties are often the first set of barriers encountered when initiating soil carbon sequestration practices. The complexity of soil systems, variability in soil types, and regional climatic conditions pose significant hurdles to the standardized implementation of these practices. For example, no-till farming requires specialized equipment and understanding of local soil conditions to be successful, which can be a significant barrier, particularly for resource-poor farmers [53]. Practices such as biochar application or agroforestry necessitate a detailed understanding of carbon dynamics, as well as the interaction of organic amendments with soil biota, which can vary widely from one ecosystem to another [54].

Socioeconomic Barriers

The adoption of soil carbon sequestration practices is not only a matter of technical feasibility but also of socioeconomic viability. Farmers, particularly in developing nations, face financial constraints that impede their capacity to switch to new practices that may require upfront investment with long-term benefits that do not align with immediate economic pressures [55]. There's also the challenge of land tenure and ownership rights; without secure land tenure, farmers have little incentive to invest in long-term soil health [56].

Environmental Trade-offs and Considerations

While the primary goal of soil carbon sequestration practices is environmental improvement, they may also pose certain trade-offs. For instance, increased biomass production for carbon sequestration can lead to higher water use, potentially stressing local water resources [57]. The application of organic amendments might also result in trade-offs between carbon sequestration and emissions of other greenhouse gases such as nitrous oxide, a potent greenhouse gas [58].

Data Gaps in Research

Despite growing literature on soil carbon sequestration, significant data gaps remain, particularly in understanding the long-term stability of sequestered carbon and the scalability of successful case studies across different agroecological zones [59]. There is a pressing need for long-term, large-scale studies that track carbon sequestration over decades to better understand the potential of these practices to contribute to climate change mitigation [60].

5. Technological Advancements in Monitoring Carbon Sequestration

Monitoring carbon sequestration is critical for understanding and managing the role of soil in mitigating climate change. Technological advancements in this field are rapidly evolving, offering more precise and efficient methods for assessing how much carbon is being captured and stored in soils.

Remote Sensing and GIS Applications

Remote sensing technologies, utilizing satellite and aerial imagery, have significantly improved our ability to monitor vegetation and soil characteristics over large areas. These technologies allow for the assessment of above-ground biomass, which is closely linked to below-ground carbon levels. GIS applications further enhance this capability by providing tools for spatial analysis and modeling, integrating various data sources to assess carbon sequestration across landscapes [61]. The Normalized Difference Vegetation Index (NDVI), derived from remote sensing data, is one widely used indicator of plant biomass and health, correlating positively with the amount of carbon sequestered in vegetation and, by extension, potentially in soils [62]. Advancements in hyperspectral imaging now allow for even finer detection of plant species and conditions, which can be related to different rates of carbon sequestration [63].

Soil Carbon Models and Their Accuracy

Soil carbon models are essential tools for predicting soil carbon dynamics and understanding the potential for carbon sequestration under different land management scenarios. Models such as CENTURY, RothC, and DNDC have been developed to simulate soil carbon turnover and are constantly being refined for accuracy [64]. These models require calibration and validation against field data to be accurate. The uncertainties in these models are often associated with the complexity of soil carbon processes and variability in environmental conditions [65]. Despite

these challenges, advancements in machine learning and data assimilation techniques are helping to improve the predictive capabilities of these models [66].

Innovations in Direct Soil Carbon Measurement

Direct soil carbon measurement techniques are the foundation for validating remote sensing data and soil carbon models. Traditional methods like dry combustion, where soil samples are burned, and the CO₂ released is measured, are being supplemented with newer technologies. In-field sensors using laser-induced breakdown spectroscopy (LIBS) and mid-infrared spectroscopy (MIRS) have been developed, which allow for rapid and non-destructive soil carbon analysis [67]. Additionally, Eddy Covariance towers are increasingly used for measuring the CO₂ flux between the soil and the atmosphere, providing insights into the dynamics of carbon sequestration [68]. The recent advent of portable technology, like the in-situ Soil Carbon Quantification System (SCiO), has the potential to revolutionize soil carbon measurements, offering a cost-effective and efficient alternative to labor-intensive traditional methods [69].

Policy and Economic Incentives

The success of sustainable soil management is significantly influenced by policy and economic incentives. In India, a range of policies have been formulated to encourage practices that contribute to soil health and carbon sequestration. India has implemented several policies aimed at promoting sustainable soil management. The National Mission for Sustainable Agriculture (NMSA), part of the National Action Plan on Climate Change (NAPCC), underscores the importance of sustainable agriculture and soil conservation methods [70]. The NMSA encompasses initiatives like the Soil Health Card Scheme, which helps farmers understand soil health indicators and suggests measures for improvement [71]. Another significant policy is the Paramparagat Krishi Vikas Yojana (PKVY), which encourages organic farming practices that enhance soil organic matter and reduce dependency on chemical inputs [72]. This scheme supports the production and distribution of organic compost, which has implications for soil carbon content.

Carbon Credits and Market Mechanisms

The concept of carbon credits, which allows for the trading of emission reductions, has also found its place in the Indian context. The Perform, Achieve and Trade (PAT) scheme under the National Mission for Enhanced Energy Efficiency (NMEEE) is a market-based mechanism to incentivize energy efficiency in large energy-consuming industries, which indirectly affects carbon emissions [73]. In agriculture, carbon credits can be earned through afforestation projects under the Clean Development Mechanism (CDM) of the Kyoto Protocol, although the applicability to direct soil carbon sequestration is still under exploration [74]. The potential for integrating soil carbon sequestration into carbon credit mechanisms is a growing area of interest, particularly with the advent of voluntary carbon markets in India.

Role of Government and International Bodies

The Government of India plays a pivotal role in implementing policies and incentives for sustainable soil management. The Ministry of Environment, Forest and Climate Change (MoEFCC) is the primary body for the formulation and implementation of environmental policies in India. The ministry works in conjunction with international bodies such as the United Nations Convention to Combat Desertification (UNCCD) and the Food and Agriculture Organization (FAO) to align domestic policies with global sustainability goals [75]. International funding mechanisms, such as the Green Climate Fund (GCF), also support projects in India that aim to enhance carbon sequestration in soils [76]. These projects often operate in synergy with national policies, amplifying their impact.

Case Studies

Case Studies of Sustainable Soil Management in India

India offers a diverse range of geographic and climatic conditions, making it an ideal location for case studies in sustainable soil management. From the arid deserts of Rajasthan to the fertile plains of the Ganges, sustainable practices are not only preserving soil health but also providing economic and social benefits to local communities.

Case Study 1: The Success of Organic Farming in Sikkim

Sikkim, a state in northeastern India, has been declared the first fully organic state in the world. The initiative began in 2003 and achieved its goal in 2016, with the state implementing organic practices across 75,000 hectares of agricultural land [77].

Comparative Analysis

Sikkim's unique Himalayan environment presents challenges for agriculture, such as steep slopes and limited arable land. However, the switch to organic farming has resulted in improved soil health, biodiversity, and a reduction in soil erosion compared to other regions.

Case Study 2: Water Conservation Techniques in Rajasthan

In the semi-arid region of Rajasthan, innovative water conservation methods have been used to combat soil degradation. The construction of johads, traditional rainwater storage tanks, has been instrumental in improving groundwater levels and soil moisture [78].

Comparative Analysis

Compared to more humid regions, Rajasthan's dry conditions require unique approaches to soil management. Water conservation has a direct impact on soil quality and agricultural productivity in this arid environment.

Case Study 3: Agroforestry in Karnataka

In the state of Karnataka, agroforestry practices have been integrated into traditional farming systems. The integration of trees into cropping systems has increased below-ground carbon storage and improved soil fertility [79].

8. Future Directions

India's soil management practices are at a pivotal juncture, with the nation facing the two-fold challenge of enhancing agricultural productivity and mitigating climate change. Carbon sequestration through improved soil management presents a promising pathway. With the global community's growing focus on sustainable practices, India's soil management strategies must align with climate change mitigation efforts. This entails a shift towards innovative, integrative approaches that offer economic viability to farmers while improving the carbon balance. In the current scenario, precision agriculture stands out as a beacon of progress, leveraging advancements in artificial intelligence, the Internet of Things, and remote sensing to optimize farming practices. The application of biochar has gained momentum, recognized for its ability to improve soil quality and sequester carbon effectively. India's vast agrarian landscapes also present a unique opportunity for enhanced rock weathering, which involves using silicate rocks to capture atmospheric carbon dioxide. The integration of these practices with climate change mitigation strategies is crucial. India's National Action Plan on Climate Change, particularly the Green India Mission, emphasizes afforestation and enhancing ecosystem services, including carbon sequestration. In the agricultural sector, managing soil organic carbon becomes instrumental in India's strategic response to climate commitments under international agreements like the Paris Agreement. Carbon credit markets, though nascent in India, could revolutionize how farmers perceive and adopt carbon sequestration measures by offering financial incentives.

Conclusion

Agroforestry stands as a cornerstone in climate change mitigation and environmental sustainability. It has consolidated its potential in bolstering ecosystem services, with particular emphasis on soil fertility, water management, carbon sequestration, and biodiversity. Case studies underscore the system's adaptability across diverse ecological and socio-economic contexts. Despite its benefits, the expansion of agroforestry is hampered by research gaps, technological needs, and socio-economic and policy constraints. Overcoming these challenges through interdisciplinary research and collaboration is essential for agroforestry to fulfill its promise. The strategic integration of agroforestry practices is imperative for realizing sustainable development goals and fostering resilient agricultural landscapes.

References

1. Dhanwantri, K., Sharma, P., Mehta, S., & Prakash, P. (2014). Carbon sequestration, its methods and significance. *Environmental Sustainability: Concepts, Principles, Evidences and Innovations*, 151(2), 151-157.
2. Lal, R. (2004). Soil carbon sequestration to mitigate climate change. *Geoderma*, 123(1-2), 1-22.
3. Macías, F., & Camps Arbestain, M. (2010). Soil carbon sequestration in a changing global environment. *Mitigation and Adaptation Strategies for Global Change*, 15, 511-529.
4. Lorenz, K., & Lal, R. (2014). Soil organic carbon sequestration in agroforestry systems. A review. *Agronomy for Sustainable Development*, 34, 443-454.
5. Ramesh, T., Bolan, N. S., Kirkham, M. B., Wijesekara, H., Kanchikerimath, M., Rao, C. S., ... & Freeman II, O. W. (2019). Soil organic carbon dynamics: Impact of land use changes and management practices: A review. *Advances in agronomy*, 156, 1-107.
6. Abbas, F., Hammad, H. M., Ishaq, W., Farooque, A. A., Bakhat, H. F., Zia, Z., ... & Cerdà, A. (2020). A review of soil carbon dynamics resulting from agricultural practices. *Journal of environmental management*, 268, 110319.
7. Hussain, S., Hussain, S., Guo, R., Sarwar, M., Ren, X., Krstic, D., ... & El-Esawi, M. A. (2021). Carbon sequestration to avoid soil degradation: A review on the role of conservation tillage. *Plants*, 10(10), 2001.
8. Dynarski, K. A., Bossio, D. A., & Scow, K. M. (2020). Dynamic stability of soil carbon: reassessing the “permanence” of soil carbon sequestration. *Frontiers in Environmental Science*, 8, 514701.
9. Prager, K., & Posthumus, H. (2010). Socio-economic factors influencing farmers' adoption of soil conservation practices in Europe. *Human dimensions of soil and water conservation*, 12, 1-21.
10. Ontl, T. A. & Schulte, L. A. (2012) Soil Carbon Storage. *Nature Education Knowledge* 3(10):35
11. Srivastava, P., Kumar, A., Behera, S. K., Sharma, Y. K., & Singh, N. (2012). Soil carbon sequestration: an innovative strategy for reducing atmospheric carbon dioxide concentration. *Biodiversity and Conservation*, 21, 1343-1358.

12. Paul, E. A. (2016). The nature and dynamics of soil organic matter: Plant inputs, microbial transformations, and organic matter stabilization. *Soil Biology and Biochemistry*, 98, 109-126.
13. Nieder, R., & Benbi, D. K. (2008). Carbon and Nitrogen Pools in Terrestrial Ecosystems. *Carbon and Nitrogen in the Terrestrial Environment*, 5-43.
14. Tank, S. E., Fellman, J. B., Hood, E., & Kritzberg, E. S. (2018). Beyond respiration: Controls on lateral carbon fluxes across the terrestrial-aquatic interface. *Limnology and Oceanography Letters*, 3(3), 76-88.
15. Senesi, N., & Loffredo, E. (2018). The chemistry of soil organic matter. In *Soil physical chemistry* (pp. 239-370). CRC press.
16. Ramesh, T., Bolan, N. S., Kirkham, M. B., Wijesekara, H., Kanchikerimath, M., Rao, C. S., ... & Freeman II, O. W. (2019). Soil organic carbon dynamics: Impact of land use changes and management practices: A review. *Advances in agronomy*, 156, 1-107.
17. Han, L., Sun, K., Jin, J., & Xing, B. (2016). Some concepts of soil organic carbon characteristics and mineral interaction from a review of literature. *Soil Biology and Biochemistry*, 94, 107-121.
18. Lal, R., Monger, C., Nave, L., & Smith, P. (2021). The role of soil in regulation of climate. *Philosophical Transactions of the Royal Society B*, 376(1834), 20210084.
19. Ramesh, T., Bolan, N. S., Kirkham, M. B., Wijesekara, H., Kanchikerimath, M., Rao, C. S., ... & Freeman II, O. W. (2019). Soil organic carbon dynamics: Impact of land use changes and management practices: A review. *Advances in agronomy*, 156, 1-107.
20. Raich, J. W., Russell, A. E., Kitayama, K., Parton, W. J., & Vitousek, P. M. (2006). Temperature influences carbon accumulation in moist tropical forests. *Ecology*, 87(1), 76-87.
21. Blanco-Moure, N., Gracia, R., Bielsa, A. C., & López, M. V. (2016). Soil organic matter fractions as affected by tillage and soil texture under semiarid Mediterranean conditions. *Soil and Tillage Research*, 155, 381-389.
22. Ramesh, T., Bolan, N. S., Kirkham, M. B., Wijesekara, H., Kanchikerimath, M., Rao, C. S., ... & Freeman II, O. W. (2019). Soil organic carbon dynamics: Impact of land use changes and management practices: A review. *Advances in agronomy*, 156, 1-107.
23. Smith, P., Soussana, J. F., Angers, D., Schipper, L., Chenu, C., Rasse, D. P., ... & Klumpp, K. (2020). How to measure, report and verify soil carbon change to realize the potential of soil carbon sequestration for atmospheric greenhouse gas removal. *Global Change Biology*, 26(1), 219-241.

24. VandenBygaart, A. J., Bremer, E., McConkey, B. G., Ellert, B. H., Janzen, H. H., Angers, D. A., ... & McKenzie, R. H. (2011). Impact of sampling depth on differences in soil carbon stocks in long- term agroecosystem experiments. *Soil Science Society of America Journal*, 75(1), 226-234.
25. Chong, K. L., Kanniah, K. D., Pohl, C., & Tan, K. P. (2017). A review of remote sensing applications for oil palm studies. *Geo-spatial Information Science*, 20(2), 184-200.
26. Vereecken, H., Schnepf, A., Hopmans, J. W., Javaux, M., Or, D., Roose, T., ... & Young, I. M. (2016). Modeling soil processes: Review, key challenges, and new perspectives. *Vadose zone journal*, 15(5).
27. Guo, L. B., & Gifford, R. M. (2002). Soil carbon stocks and land use change: a meta analysis. *Global change biology*, 8(4), 345-360.
28. Meena, R. S., Kumar, S., & Yadav, G. S. (2020). Soil carbon sequestration in crop production. *Nutrient dynamics for sustainable crop production*, 1-39.
29. Boincean, B., Dent, D., Boincean, B., & Dent, D. (2019). Tillage and Conservation Agriculture. *Farming the Black Earth: Sustainable and Climate-Smart Management of Chernozem Soils*, 125-149.
30. Skaalsveen, K., Ingram, J., & Clarke, L. E. (2019). The effect of no-till farming on the soil functions of water purification and retention in north-western Europe: A literature review. *Soil and Tillage Research*, 189, 98-109.
31. Chaplot, V., & Cooper, M. (2015). Soil aggregate stability to predict organic carbon outputs from soils. *Geoderma*, 243, 205-213.
32. Wang, X., Qi, J. Y., Zhang, X. Z., Li, S. S., Virk, A. L., Zhao, X., ... & Zhang, H. L. (2019). Effects of tillage and residue management on soil aggregates and associated carbon storage in a double paddy cropping system. *Soil and Tillage Research*, 194, 104339.
33. Simon, L. M. (2021). *Long-and short-term cover crop management effects on soil health in no-till dryland cropping systems in the semi-arid central Great Plains* (Doctoral dissertation).
34. Wuaden, C. R., Nicoloso, R. S., Barros, E. C., & Grave, R. A. (2020). Early adoption of no-till mitigates soil organic carbon and nitrogen losses due to land use change. *Soil and Tillage Research*, 204, 104728.
35. McGuire, R., Williams, P. N., Smith, P., McGrath, S. P., Curry, D., Donnison, I., ... & Scollan, N. (2022). Potential Co- benefits and trade- offs between improved soil

management, climate change mitigation and agri- food productivity. *Food and Energy Security*, 11(2), e352.

36. Ratnadass, A., Fernandes, P., Avelino, J., & Habib, R. (2012). Plant species diversity for sustainable management of crop pests and diseases in agroecosystems: a review. *Agronomy for sustainable development*, 32, 273-303.
37. Condrón, L., Stark, C., O'Callaghan, M., Clinton, P., & Huang, Z. (2010). The role of microbial communities in the formation and decomposition of soil organic matter. *Soil microbiology and sustainable crop production*, 81-118.
38. Kumar, S., Meena, R. S., Lal, R., Singh Yadav, G., Mitran, T., Meena, B. L., ... & EL-Sabagh, A. (2018). Role of legumes in soil carbon sequestration. *Legumes for soil health and sustainable management*, 109-138.
39. Liu, C., Plaza-Bonilla, D., Coulter, J. A., Kutcher, H. R., Beckie, H. J., Wang, L., ... & Gan, Y. (2022). Diversifying crop rotations enhances agroecosystem services and resilience. *Advances in Agronomy*, 173, 299-335.
40. Cai, Z., Udawatta, R. P., Gantzer, C. J., Jose, S., Godsey, L., & Cartwright, L. (2019). Economic impacts of cover crops for a Missouri wheat–corn–soybean rotation. *Agriculture*, 9(4), 83.
41. Piccoli, I., Seehusen, T., Bussell, J., Vizitu, O., Calciu, I., Berti, A., ... & Bolinder, M. A. (2022). Opportunities for mitigating soil compaction in Europe—Case studies from the soilcare project using soil-improving cropping systems. *Land*, 11(2), 223.
42. Palaniveloo, K., Amran, M. A., Norhashim, N. A., Mohamad-Fauzi, N., Peng-Hui, F., Hui-Wen, L., ... & Razak, S. A. (2020). Food waste composting and microbial community structure profiling. *Processes*, 8(6), 723.
43. Zhang, Y., Dong, L., & Shangguan, Z. (2023). Appropriate N addition improves soil aggregate stability through AMF and glomalin-related soil proteins in a semiarid agroecosystem. *Land Degradation & Development*, 34(3), 710-722.
44. Busari, M. A., Kukal, S. S., Kaur, A., Bhatt, R., & Dulazi, A. A. (2015). Conservation tillage impacts on soil, crop and the environment. *International soil and water conservation research*, 3(2), 119-129.
45. Panchal, P., Preece, C., Peñuelas, J., & Giri, J. (2022). Soil carbon sequestration by root exudates. *Trends in Plant Science*.
46. Sánchez-Monedero, M. A., Cayuela, M. L., Sánchez-García, M., Vandecasteele, B., D'Hose, T., López, G., ... & Mondini, C. (2019). Agronomic evaluation of biochar,

compost and biochar-blended compost across different cropping systems: Perspective from the European project FERTIPLUS. *Agronomy*, 9(5), 225.

47. Sánchez-Monedero, M. A., Cayuela, M. L., Sánchez-García, M., Vandecasteele, B., D'Hose, T., López, G., ... & Mondini, C. (2019). Agronomic evaluation of biochar, compost and biochar-blended compost across different cropping systems: Perspective from the European project FERTIPLUS. *Agronomy*, 9(5), 225.
48. Bodner, G., Mentler, A., & Keiblinger, K. (2021). Plant roots for sustainable soil structure management in cropping systems. *The Root Systems in Sustainable Agricultural Intensification*, 45-90.
49. Yang, F., Sui, L., Tang, C., Li, J., Cheng, K., & Xue, Q. (2021). Sustainable advances on phosphorus utilization in soil via addition of biochar and humic substances. *Science of The Total Environment*, 768, 145106.
50. Altieri, M. A., Nicholls, C. I., Henao, A., & Lana, M. A. (2015). Agroecology and the design of climate change-resilient farming systems. *Agronomy for sustainable development*, 35(3), 869-890.
51. Kuyah, S., Sileshi, G. W., Luedeling, E., Akinnifesi, F. K., Whitney, C. W., Bayala, J., ... & Mafongoya, P. L. (2020). Potential of agroforestry to enhance livelihood security in Africa. *Agroforestry for Degraded Landscapes: Recent Advances and Emerging Challenges-Vol. 1*, 135-167.
52. Zhang, H., Jiang, Y., Song, M., He, J., & Guan, D. (2020). Improving understanding of carbon stock characteristics of Eucalyptus and Acacia trees in southern China through litter layer and woody debris. *Scientific Reports*, 10(1), 4735.
53. Somasundaram, J., Sinha, N. K., Dalal, R. C., Lal, R., Mohanty, M., Naorem, A. K., ... & Chaudhari, S. K. (2020). No-till farming and conservation agriculture in South Asia—issues, challenges, prospects and benefits. *Critical Reviews in Plant Sciences*, 39(3), 236-279.
54. Stavi, I., & Lal, R. (2013). Agroforestry and biochar to offset climate change: a review. *Agronomy for Sustainable Development*, 33, 81-96.
55. Bachher, J. S., Dixon, A. D., & Monk, A. H. (2016). *The new frontier investors: How pension funds, sovereign funds, and endowments are changing the business of investment management and long-term investing*. Springer.
56. Deininger, K. (2011). Challenges posed by the new wave of farmland investment. *The journal of peasant studies*, 38(2), 217-247.

57. Sikka, A. K., Islam, A., & Rao, K. V. (2018). Climate- smart land and water management for sustainable agriculture. *Irrigation and Drainage*, 67(1), 72-81.
58. Guenet, B., Gabrielle, B., Chenu, C., Arrouays, D., Balesdent, J., Bernoux, M., ... & Zhou, F. (2021). Can N₂O emissions offset the benefits from soil organic carbon storage?. *Global Change Biology*, 27(2), 237-256.
59. Buck, H. J., & Palumbo-Compton, A. (2022). Soil carbon sequestration as a climate strategy: what do farmers think?. *Biogeochemistry*, 161(1), 59-70.
60. Lugato, E., Bampa, F., Panagos, P., Montanarella, L., & Jones, A. (2014). Potential carbon sequestration of European arable soils estimated by modelling a comprehensive set of management practices. *Global change biology*, 20(11), 3557-3567.
61. Jackson, B., Pagella, T., Sinclair, F., Orellana, B., Henshaw, A., Reynolds, B., ... & Eycott, A. (2013). Polyscape: A GIS mapping framework providing efficient and spatially explicit landscape-scale valuation of multiple ecosystem services. *Landscape and Urban Planning*, 112, 74-88.
62. Pettorelli, N. (2013). *The normalized difference vegetation index*. Oxford University Press, USA.
63. Lu, B., Dao, P. D., Liu, J., He, Y., & Shang, J. (2020). Recent advances of hyperspectral imaging technology and applications in agriculture. *Remote Sensing*, 12(16), 2659.
64. Berardi, D., Brzostek, E., Blanc- Betes, E., Davison, B., DeLucia, E. H., Hartman, M. D., ... & Hudiburg, T. W. (2020). 21st- century biogeochemical modeling: challenges for Century- based models and where do we go from here?. *GCB Bioenergy*, 12(10), 774-788.
65. van Vliet, J., Bregt, A. K., Brown, D. G., van Delden, H., Heckbert, S., & Verburg, P. H. (2016). A review of current calibration and validation practices in land-change modeling. *Environmental Modelling & Software*, 82, 174-182.
66. Carrassi, A., Bocquet, M., Bertino, L., & Evensen, G. (2018). Data assimilation in the geosciences: An overview of methods, issues, and perspectives. *Wiley Interdisciplinary Reviews: Climate Change*, 9(5), e535.
67. Senesi, G. S., & Senesi, N. (2016). Laser-induced breakdown spectroscopy (LIBS) to measure quantitatively soil carbon with emphasis on soil organic carbon. A review. *Analytica chimica acta*, 938, 7-17.
68. Kirschbaum, M. U., Puche, N. J., Giltrap, D. L., Liáng, L. L., & Chabbi, A. (2020). Combining eddy covariance measurements with process-based modelling to enhance

understanding of carbon exchange rates of dairy pastures. *Science of The Total Environment*, 745, 140917.

69. Maimaitiyiming, M., Sagan, V., Sidike, P., Maimaitijiang, M., Miller, A. J., & Kwasniewski, M. (2020). Leveraging very-high spatial resolution hyperspectral and thermal UAV imageries for characterizing diurnal indicators of grapevine physiology. *Remote Sensing*, 12(19), 3216.
70. Prasad, R. S., & Sud, R. (2019). Implementing climate change adaptation: lessons from India's national adaptation fund on climate change (NAFCC). *Climate policy*, 19(3), 354-366.
71. Reddy, A. (2018). Impact study of soil health card scheme. *National Institute of Agricultural Extension Management (MANAGE), Hyderabad-500030*, 106.
72. Reddy, A. (2018). Impact Study of Paramparagath Krishi Vikas Yojana (Organic Agriculture) Scheme of India. *Reddy A Amarender (2018) Impact Study of Paramparagath Krishi Vikas Yojana, National Institute of Agricultural Extension Management (MANAGE), Hyderabad-500030*, 210.
73. Jain, A. (2011). Unlocking Energy Efficiency Market in India's SME sector. *Diss. University of Souther California, Los Angeles, USA*.
74. Zomer, R. J., Trabucco, A., Bossio, D. A., & Verchot, L. V. (2008). Climate change mitigation: A spatial analysis of global land suitability for clean development mechanism afforestation and reforestation. *Agriculture, ecosystems & environment*, 126(1-2), 67-80.
75. Akhtar-Schuster, M., Stringer, L. C., Metternicht, G., Barger, N. N., Chotte, J. L., & Kust, G. (2022). Assessing the impact of science in the implementation of the United Nations Convention to Combat Desertification. *Land*, 11(4), 568.
76. Sharma, A., Müller, B., & Roy, P. (2015). Consolidation and devolution of national climate finance.
77. Meek, D., & Anderson, C. R. (2020). Scale and the politics of the organic transition in Sikkim, India. *Agroecology and Sustainable Food Systems*, 44(5), 653-672.
78. Bhattacharya, S. (2015). Traditional water harvesting structures and sustainable water management in India: A socio-hydrological review. *International Letters of Natural Sciences*, 37.
79. Sureshbhai, P. J., Thakur, N. S., Jha, S. K., & Kumar, V. (2017). Productivity and carbon sequestration under prevalent agroforestry systems in Navsari District, Gujarat, India. *Int. J. Curr. Microbiol. Appl. Sci*, 6(9), 3405-3422.

UNDER PEER REVIEW