

Original Research Article
**Bacteriological Quality of Ready to Eat
Vegetable Salads Vended in Ilala District
Markets and Antibiotic Sensitivity Profiles of
Isolated Contaminant Bacteria**

ABSTRACT

Aims: The consumption of ready-to-eat salads (RTES) in fast food outlets has increased as a result of heightened awareness about the nutritional status of these foods and their appeal to consumers looking for healthy and convenient meals. However, microbial contamination during the preparation, transportation, and storage of fresh ingredients has become a public health concern. People who consume RTES are at high risk of contracting food-borne diseases since they are eaten uncooked. Some microbial contaminants in RTES are resistant to antibiotics, increasing the chances of hard-to treat food-borne infections.

Study design: An experimental cross-sectional study was conducted.

Methodology: Twenty-four RTES samples were bought from randomly chosen fast food centers ~~that were chosen at random~~. The RTE salads were analyzed at the Pharmaceutical Microbiology Laboratory. The total viable counts (TVC) were determined, and the standard procedures for microbial identification were performed and confirmed by physiological tests. The identified microbial contaminants were subjected to antibiotic sensitivity testing (AST) using the Kirby-Bauer disc diffusion method. Six widely used antibiotics: amoxicillin (AX30), sulfamethoxazole/ trimethoprim-(SXT-25), amoxicillin/clavulanic acid (AMC30), gentamicin (CN5), ciprofloxacin (C5), and chloramphenicol (C30) were used for the AST.

Results: The TVC of isolated microbial contaminants ranged from 10^6 to 10^8 cfu/g, which was above the acceptable limit and unbounded and hence unfit for human consumption. Five bacterial species comprised of *Escherichia coli*, *Klebsiella pneumoniae*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, and *Klebsiella oxytoca* were isolated and subjected to the AST. All bacteria were resistant to AX30 and AMC30. *Escherichia coli* was isolated from 10% of the RTES. Association between bio-burden and antibiotic resistance was observed.

Conclusion: The RTES harbored contaminant bacteria beyond acceptable limits. The predominant contaminants were *P. aeruginosa* and *E. coli*. One-fifth of the samples contained *E. coli*, indication of poor sanitation. All the isolated bacteria were resistant to AX30 and AMC30. Prompt measures are required to curb the spread of antibiotic-resistant microorganisms.

Comment [SM1]: The aim of the study is not clearly mentioned, please correct

Comment [SM2]: It would be great to extend on this. Maybe mention the area as well

Formatted: Strikethrough

Comment [SM3]: Check again

Keywords: [Bacteriological quality, contaminant bacteria, antibiotic sensitivity testing]

1. INTRODUCTION

Ready-to-eat vegetable salads (RTES) are commonly prepared at home and food vendor outlets in several localities of Dar es Salaam, Tanzania. The most common ingredients for RTES include tomatoes, onions, or cucumbers, carrots, cabbages, salts, and sometimes lemon juice or vinegar [1]. Generally, salads are considered a source of vitamins, minerals, proteins, and other nutrients that are important for the proper body functioning of the body and skin cancer and disease fighting [2, 3]. Since RTES are consumed without cooking uncooked, there is no step to ensure the removal of microbes as compared to cooked food [4]. Hence, RTES they are also considered potential sources of enteropathogenic microbes and other food-borne diseases. Bacteria are regarded as considered the second to none as the leading source of food-borne diseases accounting for a significant (66% of food-borne diseases), followed by chemicals (26%) [5, 6]. The most common enteropathogenic bacterial contaminants in fresh produce include *Campylobacteria* spp., *Salmonella* spp., *Escherichia coli*, and *Shigella* spp. [7]. of which some are, These Such microorganisms are some of the common bacteria that causes of food-related illnesses.

On the other hand, the spread of antibiotic resistance through the food chain is another also a global health concern [8]. Not only because antibiotics are used in aquaculture, agriculture, and veterinary medicine, but also because antibiotic-resistant bacteria and genes can simply spread through the food chain [8-10]. Currently, different types of antibiotic-resistant microorganisms have been found in both food products and humans [11, 12]. However, basic safety measures such as proper food handling of such RTES, and effective cooking with adequate temperature, and proper food handling may significantly control the dissemination of antibiotic-resistant food-borne pathogens [13-15]. Utilization of pesticides and manures derived from animals treated with antibiotics has greatly contributed to the further spread of antibiotic resistance, as some traces of antibiotics remain intact [16, 17].

We therefore aimed to assess the microbiological quality of RTES by isolating the microbial contaminants and determining their antibiotic sensitivity patterns. The findings, thus, can provide an insight into the microbial quality of RTES and their potential role in spreading antibiotic resistance emanating from the contaminant bacteria.

2. MATERIAL AND METHODS

2.1 Study design, areas and samples collection

This was a cross-sectional experimental based study involving the collection of RTES from various fast-food vendors in three localities (Kariakoo, Muhimbili and Buguruni) of Ilala District, Dar es Salaam was conducted Region. The district is one of the populous areas in Dar es Salaam City with over 1.3 (17.6%) million habitants of Dar es Salaam Region. Samples were randomly bought from randomly selected fast-food vendors between March 2021 and April from three randomly selected localities namely Kariakoo, Muhimbili and Buguruni. Eight (8) samples were collected from each locality and aseptically transported in cool box to Pharmaceutical Microbiology Laboratory at School of Pharmacy, MUHAS. The collected samples were processed within two hours upon arrival at the laboratory.

2.2 Microbial quantification, isolation and identification

About ten (10) grams of each sample was weighed, and transferred into 90 mL normal saline and gently agitated for 5 minutes. Supernatant of each content was ten-fold serially diluted, from which one aliquot (1 mL) of it was deposited onto selective and non-selective agar plates, and incubated at 37°C for 24-48 hours. After 24 hours' incubation, all discrete colonies were enumerated and designated as TVC and expressed in-term of colony forming unit/gram (cfu/gr). Pure colonies were identified through conventional methods such as colony morphology, Gram staining and biochemical tests [18].

2.3 Antibacterial sensitivity testing

The identified microbial contaminants (bacteria) from RTES were overnight sub-cultured in freshly prepared nutrient broth, and the resultant turbidity was compared to the McFarland 0.5 standard (equivalent to 1.5×10^8 cfu/ml) prior to performing antibiotic sensitivity tests (AST). Then, the bacteria were subjected to AST profiling against six commonly used antibiotics: amoxicillin (AX30µg), ciprofloxacin (CIP5µg), gentamicin (CN10µg), sulfamethoxazole/trimethoprim (SXT25µg), chloramphenicol (C30µg), and amoxicillin/clavulanic acid (AMC30µg) (Oxoid, Hampshire, UK). All assays were performed on Mueller-Hinton agar plates (Roth, Germany) using the Kirby-Bauer disk-diffusion method. Following an overnight incubation at 37 °C, the diameters of inhibition zones (IZ) were determined in millimeters. Each of the above tests was

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Comment [SM4]: Please rephrase: the proper use of antibiotics cannot facilitate the spread of resistance

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Comment [SM5]: Which samples were used for this study?

Comment [SM6]: Which year?

Formatted: Strikethrough

Comment [SM7]: It would be great to include a map showing the study area

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Comment [SM8]: What are these media?

Comment [SM9]: The method is not clear. What guided you to use this method?

Comment [SM10]: Which tests were used in this case. Did you use sugars or APIs? And which API?

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Comment [SM11]: Commonly used antibiotics in which sector? What are they used for?

performed in triplicate for statistical purposes and reproducibility. Therefore, the resultant IZ was expressed as means and interpreted as per the Clinical Laboratory Standards Institute [19] as sensitive (S), intermediate (I), or resistance (R).

3. RESULTS AND DISCUSSION

The presence of microbial contaminants in RTEs is a major health concern. Since salads are consumed raw, they can act as major vehicles for food-borne illness outbreaks, as has been observed worldwide [7, 11-12, 20]. Generally, raw vegetables, salads included, are capable of supporting bacterial growth due to their high water, neutral pH, and nutrient content. Hence, cautious and minimal handling of such produce and cleanliness of equipment and premises are important [11, 13, 16].

In this study, a total of 24 RTEs samples were analyzed, of which all revealed the presence of contaminant bacteria that exceeded the allowable limits (Tables 1 and 2). Our study shows a higher incidence of *Pseudomonas aeruginosa* and *Escherichia coli*, which differs from other previous studies showing variability in incidences depending on socio-geographic factors [20, 22]. Cross-contamination is one of the main causes of the microbiological contamination of RTEs at various points, from farming through preparation to the distribution chain [8, 9]. During farming or irrigation with microbiologically contaminated water, field-grown vegetables may also become microbiologically contaminated, resulting in food-borne disease outbreaks [22, 23]. Factors such as reduced or enhanced microbial transfer efficiency, cells' adhesion, and the chemical and physical properties of the vegetables and abiotic surfaces have been attributed to the facilitation of the contamination process [23].

Comment [SM12]: This section needs a proper makeover
 1. there is a lot of repetition
 2. there is statistical analysis, which was not mentioned in the methods and materials.
 3. The section sounds more like a literature review
 4. Content not clear enough e.g. 24 samples were tested. It is not clear to how many samples were positive. Furthermore how many bacteria were isolated from each sample. The results in the text are different to the one in the figures
 5. The section is too long, maybe it would be great to include important things

Formatted: Strikethrough

Comment [SM13]: Acceptable standard

Comment [SM14]: I think its better to use 'acceptable'

Comment [SM15]: showed

Comment [SM16]: I think 'detection' would be a better word

Formatted: Strikethrough

Formatted: Strikethrough

Comment [SM17]: I think it is better to include values so that a reader doesnt go around looking for what you talking about.

Table 1. International Microbiological Criteria [24].

Microbes	Microbiological Criteria (cfu/gr)	Microbiological Quality
<i>Pseudomonas</i>	$< 10^2$	Satisfactory
	30 to 3×10^2	Acceptable
	$> 3 \times 10^2$	Unsatisfactory
<i>E. coli</i>	Absence	Satisfactory
	Presence (< 20)	Acceptable
	$20 < 10^2$	Unsatisfactory
<i>S. aureus</i>	$< 10^2$	Satisfactory
	$10^2 < 10^4$	Acceptable
	$> 10^4$	Unsatisfactory
Enteric coliforms	$< 10^2$	Satisfactory
	$10^2 < 10^4$	Acceptable
	$> 10^4$	Unsatisfactory

Table 2. Bacterial contaminants obtained from RTEs (expressed as colony forming unit per each tested gram of sample)

bacteria	Isolated	cfu/gr (Minimum-maximum)	Number Isolates/samples

Formatted: Strikethrough


Comment [SM18]: This is not clear and change it in the text as well. Are you referring to number of isolates/ sample? How many samples were positive for this bacteria

<i>Pseudomonas aeruginosa</i>	6.0×10^6 - 2.08×10^7	11
<i>Escherichia coli</i>	4.4×10^6 - 1.44×10^7	5
<i>Staphylococcus aureus</i>	6.0×10^6 - 2.0×10^7	4
<i>Klebsiella pneumoniae</i>	4.0×10^6 - 8.6×10^6	3
<i>Klebsiella oxytoca</i>	7.6×10^6	1

There were statistically significant differences between the established microbiological limits and TVC revealed from the tested RTES ($p < .01$). The most frequently isolated contaminant bacteria were *P. aeruginosa* (45.83%; $n = 11$), while only one isolate of *Klebsiella oxytoca* (4.17%) was found (Table 2). All tested bacterial isolates were sensitive to CIP5 and CN10. Except for one isolate of *E. coli* and another of *P. aeruginosa* that exhibited an intermediate susceptibility to CIP5 and CN10, respectively (Fig. 1). This observation was confirmed statistically by both the RTES-derived bacteria and the reference bacteria/control, yielding no significant differences in ZI ($p = .51$).

The T-test (independent samples) analysis of the antibiotic sensitivity of the contaminant bacteria with respect to their respective control organisms revealed different patterns: Significant differences in susceptibility against AMC30, CN10 ($p < .001$), and SXT25 ($p = .014$) when *S. aureus* was compared to control organisms. *Klebsiella oxytoca* exhibited a significant difference ($p < .001$) against CIP5, C30, AMC30, and AX30, in respect to the control organism. Significant differences in sensitivity between control organisms and *P. aeruginosa* were also observed against AMC30, AX30, and CIP4 ($p < .001$). For *K. pneumoniae*, there were significant differences against C30, AMC30, and AX30. While *E. coli* demonstrated a significant difference against AMC30 and AX30 ($p < .001$), C30 ($p = .038$), and CN10 ($p = .023$).

Fig. 1. Antibiotic sensitivity patterns of isolated contaminant bacteria to ACM30, AX30 and C30.



All isolated bacteria were resistant to AX30 and AMC30 (Fig. 2). Significant differences between reference microorganisms and the isolated contaminant bacteria against AX30 ($p < .01$; $df = 21$; 2-tailed) were observed. Six (25%) of 24 isolated bacteria were resistant to C30 and five (20.8%) to SXT25, as shown in Table 3.

Comment [SM19]: There is no mention of statistical analysis in your methods. How was it determined?

Comment [SM20]: Contradiction: even on the table, all are sensitive. Please be clear

Comment [SM21]: Figure 2 depicts something else

Comment [SM22]: 24 isolates from 24 samples, does it mean one bacteria was isolated from each sample?


Table 3: Proportions of antibiotic resistant bacteria isolated from RTES

Isolated microbes	Antibiotics (%)					
	AX30	AMC30	C30	SXT25	CIP5	CN10
<i>P. aeruginosa</i> (11)	3(27.3)	11(100.0)	3(27.3)	4(36.4)	-	-
<i>E. coli</i> (5)	5(100.0)	5(100.0)	3(60)	1(20)	-	-
<i>S. aureus</i> (4)	4(100.0)	4(100.0)	-	-	-	-
<i>K. pneumoniae</i> (3)	3(100.0)	3(100.0)	-	-	-	-
<i>K. oxytoca</i> (1)	1(100.0)	1(100.0)	-	-	-	-
Total	24 (100.0)	24(100.0)	6(25.0)	5(20.8)	-	-

Key: (-) means none was resistant to the antibiotic

A multi-resistant *P. aeruginosa* was isolated in one sample that exerted resistance against three antibiotics, namely SXT25, as depicted in Fig. 1 above, as well as against AX30 and AMC30 (Fig. 2). About 21% of the isolated contaminants were resistant to SXT25. Of these, four (36.4%) are *P. aeruginosa* and one (*E. coli*) (26%), as shown in Table 3.

Fig. 2. Antibiotic susceptibility patterns of the isolated contaminant bacteria against CIP5, CN10 and SXT25.



Antibiotic resistance is largely caused by the usage of antibiotics [25]. According to certain research on AST patterns, there is a correlation between the community's widespread use of antibiotics and irrational antibiotic usage, which could influence an individual's antibiograms [26, 27]. Our study shows that about 49% of the tested RTES-derived bacteria were antibiotic-resistant *P. aeruginosa*, followed by *E. coli* (23.7%), as shown in Fig. 3. The incidence of *E. coli* obtained from RTES is slightly lower than what was previously reported (34%) in Pakistan [28] and (8.7%) in Nigeria [21]. *Escherichia coli* is a commensal bacterium; however, some strains have acquired virulence factors and therefore have evolved to pathogenic *E. coli*. When in food, they may cause gastroenteritis and diarrhea [29]. The presence of *E. coli* in RTES samples could suggest poor hygienic practices among the street food vendors, and it is a fecal contamination indicator.

Comment [SM23]: You said all in the above paragraph

Comment [SM24]: Normal proper usage cannot confer resistance




Fig. 3. Proportions of antibiotic resistant bacteria isolated from RTEs grouped by species

The association between bio-burden (TVC) and prevalence rates of antibiotic resistance was relatively higher among the Gram-negative bacteria (*P. aeruginosa*, *E. coli*, *K. pneumoniae*, and *K. oxytoca*) (Pearson's Chi-square = 231.94; $p < .01$). While such an association was observed among isolates of *S. aureus* (Gram-positive bacterial contaminants) against AX30 and AMC30 (Pearson's Chi-square = 299.05; $p < .01$) to which they were resistant to the two antibiotics.

Staphylococcus aureus is a pathogenic bacterium that causes staphylococcal food poisoning along with other illnesses. Human-origin isolates of *Staphylococcus* are the main source of food poisoning outbreaks in the country [30, 31], where about 1% of deaths annually are attributed to food-related contamination by *S. aureus* [31]. As little as 100 thousand *S. aureus* in food may have detrimental effects. *Pseudomonas aeruginosa* is an opportunistic pathogen widely distributed in soil, water, and other moist environments. The presence of *P. aeruginosa* in the food sample indicates infective decontamination during preparation [32]. Consumption of contaminated RTEs with such a pathogen may have serious human health consequences, particularly for children, elders, and immunologically compromised persons [5, 33].

Klebsiella genus is a common opportunistic pathogen for humans and other animals, as well as being transient flora (especially in the gastrointestinal tract). Other habitats include sewage, drinking water, soils, and surface waters [33]. *Klebsiella pneumoniae* is one of the most important members of the *Klebsiella* genus in the Enterobacteriaceae family, known to cause several infections in the upper respiratory and gastrointestinal tracts. The infections are more serious among immunocompromised individuals [34]. It has been reported that clinically relevant features of *K. pneumoniae* may be preserved in wastewater, even after treatment. This evidence highlights the potential of *K. pneumoniae* for spreading through wastewater, enhancing the risks of transmission back to humans [35].

Klebsiella oxytoca is rising as a significant opportunistic pathogen, causing health care facility-acquired infections in neonates as well as adults. The bacterium is responsible for a wide range of ailments, from colitis to infective endocarditis, other than the common urinary and respiratory tract infections [36]. Our study shows that *Klebsiella oxytoca* was resistant to AX30 and AMC30 (Table 3). This implies that these antibiotics may not be useful to individuals infected with *Klebsiella oxytoca*, particularly those with weakened immune systems. More importantly, it is also clear now that RTEs harbors microorganisms that may carry various antibiotic resistance genes that can be vertically or horizontally transmitted [37].

4. CONCLUSION

The microbial quality of RTEs available in the surveyed areas raises health concerns. All tested samples had bacterial loads beyond permissive levels. *Pseudomonas aeruginosa* and *E. coli* were the most abundantly isolated contaminants. The presence of *E. coli* is an indication of poor hygienic practices and insanitary conditions. All the isolated bacteria were resistant to AX30 and AMC30. An association between bacterial load and antibiotic resistance among microbial contaminants was observed.

The authors hypothesize that improper handling of the RTE ingredients and cross-contamination of vegetables during pre- and post-harvesting processes could have been attributed to the poor microbial quality. They recommend that

Comment [SM25]: Reference?

Comment [SM26]: Ineffective?

Comment [SM27]: Check you figure and correct accordingly

responsible authorities establish more stringent measures to ensure RTEs are not the cause of food-borne outbreaks in our community and vehicles of dissemination of antibiotic-resistant pathogens. Further studies employing a larger sample size and covering wider study areas need to be conducted to ascertain the microbial quality of RTEs and the magnitude of resistant pathogens emanating from RTEs.

Comment [SM28]: What measures do you suggest?

Comment [SM29]: Include this when you write 'limitation of this study'

CONSENT (WHERE EVER APPLICABLE)

The study was approved by both the School of Pharmacy Research Project Task force and the University Research Ethical Committee prior obtaining permission from the local authorities to conduct the study.

No manuscripts will be peer-reviewed if a statement of patient consent is not presented during submission (wherever applicable).

REFERENCES

- Mbae KM, Ndwiga MK, Kiruki FG. Microbiological Quality of Kachumbari, a Raw Vegetable Salad Popularly Served Alongside Roast Meat in Kenya. *J. Food Qual.* 2018;1–5. DOI: <https://doi.org/10.1155/2018/8539029>
- Sarkar T, Salauddin M, Roy S, Chakraborty R, Rebezov M, Shariati MA, Thiruvengadam M, Rengasamy KR. Underutilized green leafy vegetables: Frontier in fortified food development and nutrition. *Rev. Food Sci. Nutr.* 2022 Jun 29;1-55. DOI: 10.1080/10408398.2022.2095555
- Sharma RK, Coniglio MA, Laganà P. *Natural Inflammatory Molecules in Fruits and Vegetables.* Berlin/Heidelberg, Germany: Springer; 2022. ISBN: 978-3-030-88472-7
- Arienzo A, Murgia L, Fraudentali I, Gallo V, Angelini R, Antonini G. Salads during Shelf-Life and Home-Refrigeration. *Foods.* 2020 Oct 8;9(10):1421. DOI: <https://doi.org/10.3390/foods9101421>
- Addis M., Sisay D. A review on major food borne bacterial illnesses. *J. Trop. Dis.* 2015;3(4):1–7. DOI: 10.4176/2329-891X.1000176
- Guptaa AK, Chaudharyb A. Food Poisoning: causes, its effects and control. *INWASCON Technology Magazine (i-TECH MAG).* 2022; 4:42-8. DOI: <http://doi.org/10.26480/itechmag.04.2022.59.61>
- Łepecka A, Zielińska D, Szymański P, Buras I, Kolożyn-Krajewska D. Assessment of the Microbiological Quality of Ready-to-Eat Salads—Are There Any Reasons for Concern about Public Health? *Int. J. Environ. Res. Public Health.* 2022 Jan 29;19(3):1582. DOI: <https://doi.org/10.3390/ijerph19031582>
- Bester LA, Essack SY. Antibiotic resistance via the food chain: Fact or fiction? *S. Afr. J. Sci.* 2010 Sep 1;106(9):1-5. DOI:10.4102/sajs.v106i9/10.281
- Sagar P, Azeem A, Banjara SK, Veleri S. The role of food chain in antimicrobial resistance spread and One Health approach to reduce risks. *Int. J. Food Microbiol.* 2023 Feb 24:110148. DOI: <https://doi.org/10.1016/j.ijfoodmicro.2023.110148>
- Riwu KH, Effendi MH, Rantam FA, Khairullah AR, Widodo A. A review: Virulence factors of *Klebsiella pneumoniae* as emerging infection on the food chain. *Vet. World.* 2022 Sep 1;15(9). DOI: 10.14202/vetworld.2022.2172-2179
- Ma F, Xu S, Tang Z, Li Z, Zhang L. Use of antimicrobials in food animals and impact of transmission of antimicrobial resistance on humans. *Bio-saf. Health.* 2021, 1;3(1):32-8. DOI: <https://doi.org/10.1016/j.bshealth.2020.09.004>
- Abebe E, Gugsu G, Ahmed M. Review on major food-borne zoonotic bacterial pathogens. *J. Trop. Med.* 2020 Oct; 2020. DOI: <https://doi.org/10.1155/2020/4674235>
- Caniça M, Manageiro V, Abriouel H, Moran-Gilad J, Franz CM. Antibiotic resistance in foodborne bacteria. *Trends Food Sci Technol.* 2019 Feb 1; 84:41-4. DOI: <https://doi.org/10.1016/j.tifs.2018.08.001>
- Abakari G, Cobbina SJ, Yeleliere E. Microbial quality of ready-to-eat vegetable salads vended in the central business district of Tamale, Ghana. *Int J food Contam.* 2018;1–8. DOI: <https://doi.org/10.1186/s40550-018-0065-2>
- Kapaya F, Mwansa FD, Sakubita P, Gama A, Langa N, Chewe O, et al. A foodborne disease outbreak investigation experience in a college in Lusaka, Zambia, 2017. *Pan Afr Med J.* 2018;29 (February). DOI:10.11604/pamj.2018.29.100.14737
- Iwu CD, Korsten L, Okoh AI. The incidence of antibiotic resistance within and beyond the agricultural ecosystem: A concern for public health. *MicrobiologyOpen.* 2020;(January):1–28. doi: 10.1002/mbo3.1035
- Mwambete KD, Semvua EK. Antimicrobial resistance profiles of bacteria isolated from chicken droppings in Dar es Salaam. *Int J Pharm PharmSci*, Vol 7, Issue 9, 268-271. ISSN- 0975-1491
- Cheesbrough M. *District laboratory practice in tropical countries*, part 2. Cambridge university press; 2005.
- Clinical and Laboratory Standards Institute. *Performance standards for antimicrobial susceptibility testing*, 30th ed. CLSI supplement M100. Clinical and Laboratory Standards Institute, Wayne, PA. 2020.
- Calonico C, Delfino V, Pesavento G, Mundo M, Nostro AL. Microbiological quality of ready-to-eat salads from processing plant to the consumers. *J. Food Nutr. Res.* 2019; 7:427-34. DOI:10.12691/jfnr-7-6-3

Comment [SM30]: The journal name as well as the species name must be in italics. Do the same with all references
The heading of the study: put caps where necessary and be consistent

21. Ezemba, C. C., C. J. Mmaduekwe, C. P. Udeze, F. C. Ogumuo, N. Ozoekwe, B. N. Duru, F. K. Ndulue et al. "Antibiogram of *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* and *Escherichia coli* isolates recovered from ready to eat food samples in ULI campus." *Int. J. Front. Res. Sci.* 1 (2022): 016-021. DOI: 10.56355/ijfrms.2022.1.1.0027
22. Kayombo MC, Mayo AW. Assessment of microbial quality of vegetables irrigated with polluted waters in Dar es Salaam City, Tanzania. *Environ. Ecol. Res.* 2018;6(4):229-39. DOI: 10.13189/eer.2018.060403
23. Possas A, Pérez-Rodríguez F. New insights into Cross-contamination of Fresh-Produce. *Curr. Opin. Food Sci.* 2023 Feb 1; 49:100954. <https://doi.org/10.1016/j.cofs.2022.100954>
24. Institute of Medicine (US) and National Research Council (US) Committee on the Review of the Use of Scientific Criteria and Performance Standards for Safe Food. *Scientific Criteria to Ensure Safe Food*. Washington (DC): National Academies Press (US); 2003. Appendix E, International Microbiological Criteria. Available from: [https://www.ncbi.nlm.nih.gov/books/NBK221566/Accessed 23 October 2023](https://www.ncbi.nlm.nih.gov/books/NBK221566/Accessed%2023%20October%202023)
25. Olesen SW, Barnett ML, MacFadden DR, Brownstein JS, Hernández-Díaz S, Lipsitch M, Grad YH. The distribution of antibiotic use and its association with antibiotic resistance. *Elife.* 2018 Dec 18;7: e39435. DOI: <https://doi.org/10.7554/eLife.39435.001>
26. Truong WR, Hidayat L, Bolaris MA, Nguyen L, Yamaki J. The antibiogram: key considerations for its development and utilization. *JAC-Antimicrob Resist.* 2021 Jun 1;3(2): dlab060. doi:10.1093/jacamr/dlab060
27. Joshi S. Hospital antibiogram: a necessity. *Indian J. Med. Microbiol.* 2010 Oct 1;28(4):277-80. DOI: 10.4103/0255-0857.71802
28. Shah MS, Eppinger M, Ahmed S, Shah AA, Hameed A, Hasan F. Multidrug-resistant diarrheagenic *E. coli* pathotypes are associated with ready-to-eat salad and vegetables in Pakistan. *J. Korean Soc. Appl. Biol. Chem.* 2015 Apr;58(2):267-73.
29. Uzeh RE, Adepoju A. Incidence and survival of *Escherichia coli* O157: H7 and *Listeria monocytogenes* on salad vegetables. *Int. Food Res. J.* 2013 Jul 1;20(4):1921.
30. Saifullah S, Abbas F, Samad A, Rizwan M, Bugti FS, Saima R, Yousaf M, Mykhalto T, Raziq A. 31. *Staphylococcus aureus* prevalence in the fresh salad and vegetables of the Quetta city. *Pure Appl. Biol.* 2018 Mar 2;7(1):255-62. <http://dx.doi.org/10.19045/bspab.2018.70031>
31. Le HH, Dalsgaard A, Andersen PS, Nguyen HM, Ta YT, Nguyen TT. Large-scale *Staphylococcus aureus* foodborne disease poisoning outbreak among primary school children. *Microbiol. Res.* 2021 Feb 11;12(1):43-52. <https://doi.org/10.3390/microbiolres12010005>
32. Alldice-Francis K, Brown PD. Diversity of antimicrobial resistance and virulence determinants in *Pseudomonas aeruginosa* associated with fresh vegetables. *International journal of microbiology.* 2012 Oct;2012. <https://doi.org/10.1155/2012/426241>
33. Laborda P, Sanz-García F, Hernando-Amado S, Martínez JL. *Pseudomonas aeruginosa*: An antibiotic resilient pathogen with environmental origin. *Curr. Opin. Microbiol.* 2021 Dec 1; 64:125-32. <https://doi.org/10.1016/j.mib.2021.09.010>
34. Monteiro A, Cardoso J, Guerra N, Ribeiro E, Viegas C, Cabo Verde S, Sousa-Uva A. Exposure and health effects of bacteria in healthcare units: An overview. *Appl. Sci.* 2022 Feb 13;12(4):1958. <https://doi.org/10.3390/app12041958>
35. Rocha J, Ferreira C, Mil-Homens D, Busquets A, Fialho AM, Henriques I, Gomila M, Manaia CM. Third generation cephalosporin-resistant *Klebsiella pneumoniae* thriving in patients and in wastewater: what do they have in common? *BMC Genom.* 2022 Dec;23(1):1-4. DOI: <https://doi.org/10.1186/s12864-021-08279-6>
36. Neog N, Phukan U, Puzari M, Sharma M, Chetia P. *Klebsiella oxytoca* and emerging nosocomial infections. *Curr Microbiol.* 2021 Apr;78(4):1115-23. DOI: <https://doi.org/10.1007/s00284-021-02402-2>
37. Zhou SY, Wei MY, Giles M, Neilson R, Zheng F, Zhang Q, Zhu YG, Yang XR. Prevalence of antibiotic resistance in ready-to-eat salad. *Front Public Health.* 2020 Mar 25; 8:92. <https://doi.org/10.3389/fpubh.2020.00092>