

ANTIBACTERIAL ACTIVITIES OF CLOVE EXTRACTS ON VEGETABLE

SPOILAGE BACTERIA.

Abstract

The present study was carried out to analyze the antibacterial and phytochemical properties of clove extracts (aqueous and methanol) on vegetable spoilage bacteria. The study seeks to determine its potential use as bio-preservatives. Eight bacterial isolates were obtained from infected leaves of bitter leaf₁, pumpkin₂ and cocoyam₃ using pour plate technique. The isolates include *Bacillus* sp₁, *Bacillus* sp₂, *Bacillus* sp₃, *Micrococcus* sp₁, *Staphylococcus* sp₁, *Staphylococcus* sp₂, *Actinomyces* sp₁ and *Actinomyces* sp₂. The pumpkin leaf had the highest average number of bacterial colonies of 2.5×10^2 followed by bitter leaf which had 1.5×10^2 while cocoyam leaf had the least with 7.0×10^1 . *Actinomyces* sp₂ has the highest zone of inhibition of 22.0 mm followed by *Bacillus* sp₁ which has 20.00 mm, *Bacillus* sp₂ has 18.00 mm, *Micrococcus* sp₁ has 17.00 mm, *Actinomyces* sp₁ has 19.00 mm, *Staphylococcus* sp₁ has 15 mm, *Bacillus* sp₃ has 14.00 mm while *Staphylococcus* sp₂ has the least zone of inhibition of 13.00 mm all at 500mg/ml. The organisms were more susceptible at higher concentrations than the lower concentrations for both the aqueous and methanol extracts of clove. For aqueous extracts, *Bacillus* sp₃, *Staphylococcus* sp₁ had the highest zone of inhibition of 16.00 mm respectively followed by *Bacillus* sp₁ and *Bacillus* sp₂ which has 12.00 mm respectively at concentrations of 500 mg/ml. The least zones were observed in *Staphylococcus* sp₂ and *Bacillus* sp₃ and concentration of 250 mg/ml and 125 mg/ml respectively. *Bacillus* sp₁, *Bacillus* sp₂, *Micrococcus* sp₁ and *Actinomyces* sp₁ has MIC of 125 mg/ml while *Staphylococcus* sp₁ and *Staphylococcus* sp₂ has 250mg/ml respectively for methanol extract while *Bacillus* sp₂, *Micrococcus* sp₁ and *Actinomyces* sp₁ has MIC of 250mg/ml respectively and *Staphylococcus* sp₁ and *Staphylococcus* sp₂ has MIC of 500 mg/ml for aqueous extracts. The Minimum Bactericidal Concentration (MBC) of the Clove extract of methanol was lower than the aqueous extracts across all the bacterial isolates. The MBC for the methanol extracts was 125 mg/ml for *Actinomyces* sp₁ while *Bacillus* sp₂, *Bacillus* sp₃, *Micrococcus* sp₁ and *Actinomyces* sp₂ were 250 mg/ml respectively while *Bacillus* sp₁, *Bacillus* sp₂, *Micrococcus* sp₂, *Staphylococcus* sp₂ and *Actinomyces* sp₁ all has MBC of 500mg/ml respectively on the aqueous extracts. The phytochemical analysis of the aqueous and methanol extracts of Clove shows that saponins, tannins, amino acids, proteins, reducing sugar, glycosides, alkaloids, anthraquinone and phenols are present though in varying amounts.

Key words:Antimicrobial, Phytochemical, Inhibition, Aqueous, Concentration and Dilution

1.0 INTRODUCTION

Environmental and health concern of the carcinogenic effect of chemical preservatives, food safety and huge economic losses of agricultural products due to spoilage by microorganisms, has

aroused the thought for adoption of natural preservatives for inhibition of bacterial activity and growth over chemical preservatives. Microorganisms such as bacteria and fungi, are the major contributors of food spoilage (Liu *et al.*, 2017). Spices possess antimicrobial activities (Corles – Rojas *et al.*, 2014; Martinez- Gracia *et al.*, 2015) and is one of the most used natural antimicrobial agents traditionally for preservation of food. Similarly, spices have been reported as rich sources of phytochemicals which are responsible for the inhibition of bacterial growth (Li *et al.*, 2019). Phytochemicals such as saponins, alkaloids, tannins, steroids etc. are partially responsible for the antimicrobial activities of plants (Nethatheet *et al.*, 2011). Cloves are brown, dried, unopened flower buds of *Syzygium aromaticum*, an evergreen tree native to Indonesia (Salvador, 2014) and of height, 8 – 30m, with many semi- erect branches (Nadem *et al.*, 2023). It possess a pleasant smell and is commonly and traditionally used as a spice for seasoning of meat and a wide variety of food and drinks. Additionally, some rural dwellers prepares aqueous extract of cloves, although with no standard concentration and drink same as herbal remedy for bacterial infection and to boost general health as they believed on its antibacterial activity

The antibacterial activity of clove has been tested and reported by many researchers. Saeed *et al.*, (2013) reported that solvent extract of clove has a great potential for the inhibition of microbial load. Likewise, Gupta and Prakash (2021) reported clove extract having the potential to inhibit the growth of bacterial species and be used as a natural antibacterial agent for pathogens.

Likewise, the phytochemical analysis of clove extract revealed the presence of carbohydrates, lipids, alkaloids, flavonoids, tannins, sterols and triterpenes (Lone and Jain, 2022). Khatri *et al.*, (2014) reported an essential ingredient of clove oil to include tannins, vanillin, flavonoids etc. and tannins was found to be in high amounts in fresh clove (Batiha *et al.*, 2020).

This study therefore seeks to evaluate the antibacterial and phytochemical properties of extracts of clove and to determine the concentrations of the extracts at which the antibacterial activity could be achieved. The study also seeks to assess its use as a natural preservative of vegetables.



Fig. 1. Clove (flower bud of *Syzygium aromaticum*)

Scientific Classification of *Syzygium aromaticum*

Kingdom: Plantae

Division: Magnoliophyta

Class: Magnoliopsida

Order: Myrtales

Family: Myrtaceae

Genus: *Syzygium*

Species: *S. aromaticum*

2.0 MATERIALS AND METHODS

2.1 Collection of Samples

Cloves were purchased from a local trader at the Oron main market, Akwa Ibom State of Nigeria and identified by a botanist at the Department of Biological Sciences, Clifford University, Owerri, Abia State where the research was carried out.

2.2 Sample Preparation

The Cloves were washed in tap water, dried under shade and then homogenized to fine powder using a disinfected manual blender and stored in airtight bottles. Fifty (50) grams of leaf powder was placed in 450 ml of methanol and water respectively in conical flasks and stirred on a rotary shaker at 200 rpm for 48 h. The suspension was filtered through muslin cloth and the extracts were centrifuged at 5000 rpm for 10 min. Supernatants were evaporated at 45 °C to dry and stored at 4 °C in air tight bottles. The dried extract was re-dissolved at 10% (100 mg/ml) in DMSO for the antifungal testing, when not in use, it was stored in a refrigerator in 4 °C (Mahesh *et al.*, 2008)

2.3 Determination of percentage yield

The percentage yield of the extract was calculated using the following formula:

$$\% \text{ yield} = \text{mass of extract} / \text{mass of leaves} \times 100/1$$

2.4 Isolation of Bacterial Plant Pathogen

The infected parts of vegetable (pumpkin), bitter leaf and cocoyam leaves were aseptically collected from the farm with clean nylon bag. After washing the leaves thoroughly in sterile water, the infected leaves along with adjacent small uninfected leaves were cut into small pieces (2-5 mm squares) and flamed – sterilized forceps was used to transfer them into 0.1% mercuric chloride solution used for surface sterilization of plant tissues, and rinsed. A sterile mortar and pestle was used to squash the various samples into a fine homogenized form. A clean filter paper

and digital weighing balance was used to weigh one gram of each of the samples into 9 ml of sterile water, they were shaken to homogenize respectively. Double fold serial dilution was carried out by transferring 1 ml from the stock into the first test containing 9 ml of sterile water, it was equally shake to homogenize. The dilution was continued till the last tube at which 1ml aliquot was discarded. The samples (10^{-3} , 10^{-4} and 10^{-5}) were inoculated on a nutrient agar using spread plate technique (Cheesbrough, 2005). The plates were incubated at 37°C for 24hrs for the complete growth of bacteria.

2.5 Purification of Isolates

The colonial bacterial growths were purified by sub culturing them on sterile nutrient agar plates using the streak method. Thereafter the pure cultures were stored in agar slants.

2.6 Identification of Isolates

The bacterial isolates were identified by Gram staining and biochemical tests which include: Catalase, coagulase, Indole, Citrate utilization, Urease, Hyrogensulphide, sugar fermentation and Oxidaze tests.

2.6.1 Gram staining

Gram staining reaction was used to identify gram status of bacteria in the culture. Heat-fixed smears of young cultures were prepared by placing a drop of water on a grease free microscopic slide. Using a sterilized and cooled inoculation loop, a very small sample of the bacterial colony was obtained and gently immersipated into the drop of water on the slide. This was then placed on the staining rack and allowed to air-dry. The smears were flooded with crystal violet and allowed to stand for 30 seconds after which the stains were rinsed with a slow running tap water for 5 seconds. Again, the smears were flooded with Gram's iodine mordant and allowed to stand for 60 seconds. The iodine was rinsed with a slow running tap water for 5 seconds. The

smears were decolorized with 95% ethanol until stain no longer washes off the slides; the decolorizer was rinsed with slow running tap water for 5 seconds. The smear was counterstained with safranin and left for 60 seconds and was rinsed with a slow running tap water for 5 seconds. The slides were air-dried and examined under $\times 100$ with the use of oil immersion (Cheersbrough, 2000).

2.6.2 Biochemical test

1. Catalase test

3ml of 3% hydrogen peroxide solution was placed in a test tube and distilled water was dropped on another test tube as the control. Colonies of the test organism was picked using a sterile wire loop and was immersed in the hydrogen peroxide solution and the test tube containing water. It was kept for reaction for 2 seconds. Active bubble indicates catalase positive, No bubbles indicates catalase negative.

2. Coagulase test

A drop of distilled water was placed on two glass slides separately. A colony of the test organism was emulsified on each of the drops to make thick suspensions. A loop full of human plasma was added to one of the suspension and was gently mixed. It was kept for 10 seconds for reaction. Clumping of the bacterial cell indicates positive result.

3. Oxidase test

A piece of filter paper was placed on a clean test tube and three drops of the freshly prepared oxidase reagent was dropped. A colony of the test organism was removed using a sterile wire loop and was dropped on the filter paper. The development of a deep purple color indicates the production of oxidase by the organism which oxidizes the phenylenediamine in the reagent to the deep purple color.

4. Citrate utilization test

Simmons Citrate agar was prepared and sterilized according to the manufacturer's instruction. Test tubes were slanted and were allowed to solidify. A sterile wire loop was used to pick a colony of the test organism and inoculated in the slant by first stabbing the butt and finally streaking the slant. It was incubated at 37°C for 24 hours and was checked for color change. Medium changing from green to blue color indicates positive result.

5. Indole production test

Trypton broth was prepared and the test organism was inoculated in it and incubated for 24 hours at 37°C. Ten drops of Kovac's reagent was added to the trypton broth. The test tube was allowed to stand for 5 minutes, and then observation for the appearance of a dark-red upper layer which indicates the presence of Indole and straw which indicates absence of indole. Dark red upper layer indicates presence of indole, but straw or yellow color indicates absence of indole.

6. Hydrogen Sulphide (H₂S) production and sugar fermentation test using TSI agar

TSI agar was prepared according to the manufacturer's instructions. It was sterilized in an autoclave and slanted to solidify. A sterile wire loop was used to pick a colony of the test organism and inoculated in the slant by first stabbing the butt and then streaking the slope. It was incubated at 37°C for 24 hours and observations were taken. Formation of black precipitate in the

butt of the tube indicates H₂S positive, if acid is produced by the breakdown of dextrose to acid, the slant will change from red (alkaline) to yellow (acid). Also when dextrose is fermented by the organism, gas is released and the gas appears in bubbles in the butt of the tube or split the media or part of the slant will rise upwards (Cheersbrough, 2005)

2.7 Phytochemical analysis

Phytochemical analysis of the extracts (Cloves) was carried out according to the general method of Harbone (1998). Basic phytochemical screening was carried out using simple chemical tests to detect the presence of secondary plant constituents such as alkaloids, tannins, flavonoids, saponins, phenols, glycoside, reducing sugar, soluble carbohydrate etc. in the sample. The methods used were those outlined by Harbone (1998) except otherwise stated.

2.7.1 Saponin (frothing test)

Extract of (0.5 g) was suspended in 5 ml of distilled water in a test tube. The contents were shaken vigorously and observed for a stable persistent froth. The frothing was mixed with 3 drops of olive oil and again shaken vigorously. The formation of an emulsion, if any, was observed.

2.7.2 Test for Alkaloids

The extract (0.1mg) was added to 6ml of dilute hydrochloric acid and boiled. After boiling, it was cooled and filtered. Few drops of Mayer's reagent were added. The formation of turbidity indicated the presence of alkaloids.

2.7.3 Glycoside

Five milligram (5 mg) extract was suspended in 5 ml ethanol and two drops each of 1% ferric chloride and concentrated sulphuric acid were added. The development of reddish-brown coloration at the junction of two layers and the bluish green color in the upper layer indicated the presence of glycoside.

2.7.4 Test for Terpenoids

Nine milliliters (9 ml) of ethanol was added to 1 g of the extract and refluxed for a five min and filtered. The filtrate was concentrated down to 2.5 ml in a boiling water bath. Hot distilled water (5 ml) was added to the concentrated solution; the mixture was allowed to stand for 1 hour and the waxy mater was filtered off. The filtrate was extracted with 2.5 ml of chloroform using a separating funnel. The chloroform extract was evaporated to dryness in a water bath and dissolved in 3 ml of concentrated sulphuric acid and then heated for 10 min in a water bath. A grey color indicated the presence of terpenoids.

2.7.5 Test for Steroids

A known quantity of the test sample was extracted in the chloroform and filtered. The filtrate was mixed with 2 ml of conc. H_2SO_4 carefully so that the sulphuric acid formed a lower layer. A reddish-brown color at the interphase indicated the presence of steroidal ring.

2.7.6 Reducing sugar

Extract of 0.5 mg was dissolved in 1 ml of water. To it 5 – 8 drops of Fehling's solution were added and solution was heated. Appearance of brick red precipitate indicated the presence of reducing sugar.

2.7.7 Amino acid and Protein

Extract of 0.1 mg was boiled with 1ml of 0.2% Ninhydrin solution. The appearance of violet blue color indicates the presence of protein in the extract.

2.7.8 Phenols

Extract of 0.2 mg was taken in 3 ml of 5% ferric chloride and then add 1 ml each of methyl alcohol and Folin – Ciocalteu reagent and shaken gently. Formation of deep blue or black color indicated the presence of phenol.

2.7.9 Tannins

Extract of 0.5 g of extract was boiled in 10 ml of water in a test tube and then filtered. Few drops of 0.1% ferric chloride were added. Production of brownish green or blue-black color indicated the presence of tannins.

2.7.10 Test for Flavonoids

A quantity of the extract was boiled in ethyl acetate (10 ml) for 3 mins, filtered and cooled. Then the filtrate (4 ml) was shaken with 1 ml of dilute ammonia solution. An intense yellow coloration indicated the presence of flavonoids.

2.7.11 Test for Soluble Carbohydrate (Molisch test)

The extract (0.1 g) was boiled with 2 ml of distilled water and filtered. To the filtrate, few drops of naphthol solution in ethanol (molisch's reagent) were added. Concentrated sulphuric acid in a Pasteur pipette was then gently poured down the side of the test tube to form a lower layer. A purple interfacial ring indicated the presence of carbohydrate.

2.7.12 Test for Anthraquinone

Diluted H_2SO_4 was added to 0.5 g of the extract; the solution was then boiled and filtered. The filtrate was cooled and to it equal volume of benzene was added. The solution was shaken well

and the organic layer was separated. Equal volume of dilute ammonia solution was added to the organic layer. If the ammonia layer turns pink, then it indicates the presence of anthraquinone.

2.8 Preliminary antifungal screening of the extract

The Mueller Hinton agar was prepared according to the manufacturer's instruction and poured aseptically into Petri-dishes and was allowed to gel. The surface of the plates was then streaked with inoculums of the test organisms using an inoculating loop. Thereafter, a sterilized 6mm long cork-borer with depth 3-4 mm, was used to create holes on the agar plates. The extract was then serially diluted in test-tubes containing 1ml of the solvents. In the first test-tube containing the weighed plant extract, 2 ml of 10% Dimethyl sulphoxide (DMSO) was used to dissolve the extract (the stock solution of 500 mg/4ml), then 1 ml of the solvent was added into the other test-tubes. From the first test-tube, 1000ul was pipette and put in the second test-tube until it got to the last one (500 mg/4ml, 125 mg/ml, 62.5 mg/ml, 31.25 mg/ml, 15.625 mg/ml, 7.8 mg/ml). This was done for the methanol and aqueous extracts respectively. After the dilution, 20 µl of the extracts were pipette according to their concentration and put in the hole. The Petri-dishes were allowed to sit for 30 min for pre-diffusion of the plant extracts and the plates were incubated at 37 °C for 24hours. The inhibition zones were then measured after incubation by using a transparent meter rule in millimeters (mm). This was also done for the test drug (Chloramphenicol 200mg) which served as the positive control while 10% DMSO alone served as the negative control.

2.8.1 Minimum Inhibitory Concentration (MIC) Determination

MIC is the lowest concentration of extracts that inhibited noticeable growth of the organisms after 24 hours using the tube dilution method as described by the Clinical and laboratory standards institute one (1 ml) milliliter of different concentrates (500 mg/ml, 250mg/ml, 125

mg/ml, 62.5 ml/ml and 31.25 mg/ml, 15.625 mg/ml) of the various extracts. Sterile pipette was used to transfer 20 µl into each well, with the bacterial colonies streaked on the media surface. The MHA plates were incubated at 37°C 24hours. After the incubation the visible zone of inhibition was measured (CLSI, 2016).

2.8.2 Minimum Bactericidal Concentration (MBC) Determination

The MBC was determined by inoculating from the MIC plate showing a visible inhibition and was incubated at 37°C for 24 hours.

3.0 RESULT

Table 1 below recorded the percentage yield of methanol and aqueous extracts of cloves. Result shows higher percentage yield of methanol extract to the aqueous extracts.

Table 1: Percentage (%) Yield of Extracts

Extracts	Percentage (%) Yield
Methanol	44.68
Aqueous	14.32

Table 2: Phytochemical Analysis of Clove Extracts

S/n	Phytochemical	Aqueous	Methanol
1.	Saponins	++	+++
2.	Tannins	+	++
3.	Amino acid and proteins	++	++
4.	Reducing sugar	+	++

5.	Glycosides	++++	
6.	Alkaloids	+	++
7.	Anthraquinone	+	+
8.	Phenol	+	++

Key: + = Positive test

Table 2 above records the phytochemical analysis of the aqueous and methanol extracts of clove.

Result shows the presence of saponins, tannins, Amino acid and proteins, reducing sugar, glycosides, alkaloids, anthraquinone and phenol in both aqueous and methanol extracts. Methanol seems to extract more of the phytochemicals than water as indicated in the table.

Table 3: Microbiological and Biochemical Characteristics of Bacteria Isolates and Their Possible Identity.

Organism code	Cultural characteristics	Gram rxn	C at	C oa	In d	C it	M	O x	B	S	G	H 2s	Possible organisms
PLM1	Milky, round, flat, smooth and shiny	+ve cocci	+	-	-	+	-	+	+	-	-	-	<i>Micrococcus</i> sp
PLM2	Milky, rough, flat, sticky and shiny	+ve rod	+	-	-	+	-	+	+	-	-	-	<i>Acetomyctes</i> sp ₁
PLY1	Yellow, circular, smooth, shinny and non sticky.	+ve cocci	+	+	-	+	-	-	-	-	-	-	<i>Staphylococcus</i> sp ₁
PIO1	Orange, sticky, ,round, smooth and	+ve rod	+	+	-	+	-	+	+	-	-	-	<i>Bacillus</i> sp ₁

	shiny														
BLO2	Orange, sticky, round, smooth and shiny	+ve	+	-	-	+	+	-	+	+	+	-	<i>Bacillus sp₂</i>		
BLM3	Milky, filamentous, sticky and shiny	+ve	+	-	-	+	+	+	+	+	+	+	<i>Bacillus sp₃</i>		
BLY2	Yellow, round, non sticky, raised and shinny	+ve	+	+	-	+	+	-	+	+	+	-	<i>Staphylococcus sp₂</i>		
CLM4	Milky, rhizoid shiny and non sticky	+ve	+	-	-	-	-	+	+	+	-	-	<i>Acetomyces sp₂</i>		
CLW1	White, non sticky and flat	+ve	-	+	-	+	-	-	+	+	+	+	<i>Bacillus sp₄</i>		

Key: Cat: catalase test, Coa: coagulase test, Ind: indole test, Cit: Citrate utilization test, MR: Methyl red test, OX: Oxidase test, SFT: Sugar fermentation test, B=butt, S= slope, G= gas production, H₂S: Hydrogen Production.

Table 3 records the possible bacterial isolates from the infected leaves sample used for the study. Eight bacterial isolates were obtained from the infected leaves of bitter leaf₁, pumpkin₂ and cocoyam₃ using pour plate technique. The isolates include *Bacillus sp₁*, *Bacillus sp₂*, *Bacillus sp₃*,

Micrococcus sp₁, *Staphylococcus* sp₁, *Staphylococcus* sp₂, *Actinomyces* sp₁ and *Actinomyces* sp₂.




Fig. 2. Colonial Distribution of Isolates

Table 4: Zone of Inhibition of Bacterial Isolates by Methanol Extract of Clove

Conc. (Mg/ml)	<i>Bacillus</i> s sp ₁	<i>Bacillus</i> s sp ₂	<i>Micrococcus</i> ssp	<i>Bacillus</i> s sp ₃	<i>Staph.</i> sp ₁	<i>Actinomyces</i> es sp ₁	<i>Actinomyces</i> es sp ₂	<i>Staph.</i> . sp ₂
Zone of Inhibition (mm)								
500	20.00	18.00	17.00	14.00	15.00	19.00	22.00	13.00
250	13.00	14.00	12.00	12.00	12.00	11.00	14.00	10.00
125	12.00	0.00	0.00	11.00	0.00	0.00	0.00	0.00

Table 6: Zone of Inhibition of Test Drug (Chloramphenicol)

Conc. (250mg/ ml).	<i>Bacillu</i> <i>s sp</i> ₁	<i>Bacillu</i> <i>s sp</i> ₂	<i>Micrococcu</i> <i>ssp</i>	<i>Bacillu</i> <i>s sp</i> ₃	<i>Staph</i> <i>. sp</i> ₁	<i>Actinomycet</i> <i>es sp</i> ₁	<i>Actinomycet</i> <i>es sp</i> ₂	<i>Staph</i> <i>. sp</i> ₂
Zone of Inhibition (mm)								
250	28.00	28.20	30.00	27.00	24.00	30.00	27.00	30.00
125	20.00	23.30	28.00	25.00	18.00	25.00	25.00	27.00
62.25	17.00	20.10	26.00	23.00	15.00	23.00	23.00	25.00
31.125	13.50	18.00	24.00	20.00	12.00	20.00	20.00	22.00
15.625	11.10	15.00	20.00	16.00	10.00	15.00	18.00	20.00
7.813	8.00	7.00	15.00	10.00	8.00	10.00	14.00	15.00

Table 7: Minimum Inhibitory Concentration (MIC) of the Clove Leaf Extracts and the Test Drug (Chloramphenicol).

Extract and Test drug	<i>Bacill</i> <i>us sp</i> ₁	<i>Bacill</i> <i>us sp</i> ₂	<i>Bacill</i> <i>us sp</i> ₃	<i>Micrococ</i> <i>cus sp</i>	<i>Stap</i> <i>h.</i>	<i>Stap</i> <i>h sp</i> ₂	<i>Actinomyc</i> <i>etes sp</i> ₁	<i>Actinomyc</i> <i>etes sp</i> ₂
Methanol	125	125	-	125	250	250	125	-

Aqueous	-	250	-	250	500	500	250	-
Chloramphenicol	15.62	15.62	15.62	15.625	15.6	15.6	15.625	15.625
	5	5	5		25	25		

Table: 8. Minimum Bactericidal Concentration (MBC) of the Cloves Extract

Extract and Test drug	<i>Bacillus</i> sp ₁	<i>Bacillus</i> sp ₂	<i>Bacillus</i> sp ₃	<i>Micrococcus</i> sp	<i>Staphylococcus</i> sp ₁	<i>Staphylococcus</i> sp ₂	<i>Actinomyces</i> sp ₁	<i>Actinomyces</i> sp ₂
Methanol	-	250	250	250	250		125	250
Aqueous	500	500	-	500	-	500	500	-
Chloramphenicol	7.813	7.813	7.813	7.813	7.81	7.81	7.813	7.813
					3	3		




Fig. 3. Minimum Inhibitory Concentration of Clove Extracts




Fig. 4. Minimum Bactericidal Concentration of Clove Extracts

DISCUSSION

The phytochemical analysis of aqueous and methanol extracts of Clove shows that saponins, tannins, amino acids, proteins, reducing sugar, glycosides, alkaloids, anthraquinone and phenols were present though in varying amounts. This is in agreement with the previous report of Sanusi *et al.*, (2023), who equally has all the phytochemicals present. The phytochemical result is contrary to the report of Ajobiewee *et al.*, (2022), where tannins, glycosides, alkaloids and saponins were absent in ethanol and aqueous extracts respectively. Also, the result suggests the ability of methanol to extract more of the bioactive compounds in clove than water. Therefore, alcohol could be recommended as a better option for extraction of active and chemical compounds in cloves than water. To achieve best result in its use as bio preservatives as well as herbal remedy for bacterial infections, alcohol extract will be more active and preferable to aqueous extracts.

Eight bacterial isolates were obtained from the infected leaves of Bitter leaf, pumpkin and Coco yam leaf. The isolates include *Bacillus* sp₁, *Bacillus* sp₂, *Bacillus* sp₃, *Micrococcus* sp, *Staphylococcus* sp₁, *Staphylococcus* sp₂, *Actinomycetes* sp₁ and *Actinomycetes* sp₂. These are common infections causing bacteria in humans. There is tendency that high cases of infections in rural areas may also be due to consumption of infected or spoiled vegetables. The application of cloves extracts on these vegetables could inhibit the activities of these bacteria on the vegetables and thus preserve them from spoilage easily. Therefore, it could save a lot of economic waste and combat cases of infectious diseases among the rural dwellers.

The pumpkin leaf has the highest average number of bacterial colonies of 2.5×10^2 followed by bitter leaf which had 1.5×10^2 while Coco yam has the least with 7.0×10^1 . The result of the bacteriological load agrees with Patience N. Okafor (2022) whose bacterial load on nutrient agar ranged from 6.7×10^4 to 1.02×10^5 and 7.2×10^4 to 1.64×10^5 from pumpkin, bitter leaf and

scent leaf surfaces. The organisms isolated were *Bacillus* sp, *Erwiniaspp*, *Pseudomonas* spp and *Streptococcus* spp. It also agrees with the work of Muhammad *et al.*, (2021) who reported on the isolation and identification of bacteria responsible for the spoilage of fluted pumpkin and bitter leaf. The isolates includes *E. coli*, *Klebsiella* and *Proteus mirabilis* from the fluted pumpkin and *Proteus mirabilis*, *E. coli* and *Bacillus cereus* from bitter leaf.

Actinomyces sp₂ has the highest zone of inhibition of 22.0 mm followed by *Bacillus* sp₁ which has 20.00 mm, *Bacillus* sp₂ has 18.00 mm *Micrococcus* sp has 17.00 mm, *Actinomyces* sp₁ has 19.00 mm, *Staphylococcus* sp₁ has 15 mm, *Bacillus* sp₃ has 14.00 mm while *Staphylococcus* sp₂ has the least zone of inhibition of 13.00 mm all at 500mg/ml. The organisms were more susceptible at higher concentrations than the lower concentrations for both the methanol and aqueous extracts of clove flower bud. For aqueous extracts *Bacillus* sp₃, *Staphylococcus* sp₁ has the highest zone of inhibition of 16.00 mm respectively followed by *Bacillus* sp₁ and *Bacillus* sp₂ which has 12.00 mm respectively at concentrations of 500 mg/ml. The least zones were observed in *Staphylococcus* sp₂ and *Bacillus* sp₃ and concentration of 250 mg/ml and 125 mg/ml respectively. This work is in line with the previous report of Sanusi *et al.*, (2023) whose inhibition zone diameter of clove bud extracts for *S. aureus* was 21.00 mm and 22.00 mm for the aqueous and ethanol extracts. Ajobiewe *et al.*, (2022) had zone of inhibitions of 27, 22 to 9 mm on cold water extracts of clove plant extracts on *E. coli* and 28, 25 to 8mm respectively on ethanol extract at concentrations of 500mg/ml to 15.63 mg/ml.

Bacillus sp₁, *Bacillus* sp₂, *Micrococcus* sp and *Actinomyces* sp₁ has MIC of 125 mg/ml while *Staphylococcus* sp₁ and *Staphylococcus* sp₂ has 250mg/ml respectively for methanol extract while *Bacillus* sp₂, *Micrococcus* sp and *Actinomyces* sp₁ has MIC of 250mg/ml respectively and *Staphylococcus* sp₁ and *Staphylococcus* sp₂ has MIC of 500 mg/ml for aqueous extracts.

The Minimum Bactericidal Concentration (MBC) of the Clove extract of methanol is lower than the aqueous extracts across all the bacterial isolates. The MBC for the methanol extracts is 125 mg/ml for *Actinomyces* sp₁ while *Bacillus* sp₂, *Bacillus* sp₃, *Micrococcus* sp and *Actinomyces* sp₂ are 250 mg/ml respectively while *Bacillus* sp₁, *Bacillus* sp₂, *Micrococcus* sp, *Staphylococcus* sp₂ and *Actinomyces* sp₁ all has MBC of 500mg/ml respectively on the aqueous extracts. The MIC results above is contrary to the MIC result obtained by Sanusi *et al.*, (2023) were 6.25 mg/ml was obtained for the isolates on ethanol extract. The isolates includes *E. coli*, *Salmonella* sp, *Staphylococcus aureus* and *Pseudomonas* sp while for aqueous extracts *E. coli* has the highest MIC of 12.5mg/ml followed by *Salmonella* sp, *S. aureus* and *Pseudomonas* sp which has 6.25 mg/ml respectively.

CONCLUSION

This study has revealed that the Clove extracts possess significant amount of phytochemicals such as saponins, tannins, reducing sugar, amino acids, proteins, phenols and glycosides. The extracts also shows antimicrobial properties against *Bacillus* spp, *Micrococcus* sp, *Staphylococcus* spp, *Actinomyces* sp and *Actinomyces* sp. These findings are interesting and essential at the present time where the problems of emerging and re - emerging resistant strains of microorganisms are becoming the order of the day and scarcity of food due to bacterial infection on plants.

RECOMMENDATION

This study is recommending Clove extract as an alternative source for bio- preservation in food and allied products.

References

- Ajobiewe, H. F., Elisha E, I., Ajobiewe J. O., Salami, A. O., Umeji, L. C, Udefuna, P.A., Yashim, A. N, Alau, K. K (2022). Antimicrobial (*Syzygium aromaticum*) on *Escherichia coli*. *ScholarJournal of Applied Medical Sciences*. 10 (5): 673 – 684.
- Batiha, G. E., Alkazmi, L. M., Wasef, L. G., Beshbishy, A. M., Nadwa, E. H., Rashwan, E. K. (2020). *Syzygium aromaticum* L (*Myrtaceae*): Traditional uses, bioactive chemical constituents, Pharmacological and Toxicological Activities. *Biomolecules*, 10(2): 202
- Cheesbrough, (2005). *District Laboratory Practice in Tropical Countries*: Cambridge University press. Pp 70 – 62.
- CLSI (Clinical and Laboratory Standards Institute), (2016). Performance standard for antimicrobial susceptibility testing. CLSI approved standard M100S. Clinical and Laboratory Standards Institute, Wayne, PA.
- Cortes-Rojas, D. F., Femandes de Souza, C. R and Oliveira, W. P (2014). Clove (*Syzygium aromaticum*): A Precious Spice. *Asian Pacific Journal of Tropical Biomedicine*, 4(2): 90 - 96
- Cupta, C and Prakash, D (2021). Comparative study of the Antimicrobial Activity of Clove oil and Clove extract on oral pathogens. *Dentistry Open Journal*, 7(1): 12 – 15
- Harbone J. B (1998) “phytochemical methods, A Guide to Modern Techniques of Plant Analysis,” 2nd ed. Chapman and Hill Ltd, New York. 100 – 104.
- Khatri, M., Mittal M., Cupta, N., Parashar, P., Mehra, V. (2014). Phytochemical evaluation and pharmacological activity of *Syzygium aromaticum*: A Comprehensive Review. *International Journal of Pharmacy and Pharmaceutical Sciences*, 6(8): 67 - 72

- Li, G. H., Li, Y. R., Li, V. Y., Wu, D. M., Ren, H. X., Lou, X. N., Wang, T. S. (2019). Lignin and Flavonoid support the prevention of cinnamon against oxidative stress related diseases. *Phytomedicine*, 53: 143 - 153
- Lone, Z. A and Jain, N. K. (2022). Phytochemical Analysis of Clove (*Syzygium aromaticum*) Dried Flower buds Extract and it's Therapeutic Importance. *Journal of Drug Delivery and Therapeutics*, 12(4-S): 87 – 92
- Mahesha B. and Satish s., Antimicrobial activity of some important medicinal plant against plant and human pathogens, *World Journal of Agricultural Science*, 4, 839 – 843 (2008).
- Martinez-Gracia, C., Gonzalez-Bermueles, A. M., Cabellero-Valcareel, M., Santaella-Paschal, C., Frontela-Saseta (2015). Use of herbs and spices for food preservatives: Advantages and Limitations. *Current Opinion of Food Science*, 6: 35 - 43
- Muhammad A.S, Ani S.A and Nasiru A.M. (2021). Isolation and Identification of Bacteria Responsible for The Spoilage of Fluted Pumpkin and Bitter leaf In Sokoto Metropolis. *International Journal of Multidisciplinary Research*. 7 (1): 372 – 378.
- Nadeem M., Imran, M., Din, A and Khan, A. (2023). Essential oils, Extraction characterization and applications. Academic Press, Pp 325 -344
- Nethathe, B. B., Ndip, R. N. (2011). Bioactivity of *Hydnora africana* on selected bacterial pathogens: Preliminary Phytochemical Screening. *African Journal of Microbiology Research*, 5(18): 2820 - 2826

Patience N. Okafor (2022). Isolation and Identification of Bacteriological Constituents of Some Leavy Vegetables Sold in Rural Markets In Anambra State, Nigeria. *International Journal of Advance Research*, 10(2) : 845 – 851.

Saeed, M., Nadeem, M., Khan, M. R., Shabbir, M. A., Shehzad, A and Amir, R. M. (2013) . Antimicrobial activity of *Syzygium aromaticum* extracts against food spoilage bacteria. *African Journal of Microbiology Research*, 7(4): 4848 – 4856

Salvador, M. J. (2014). Clove (*Syzygium aromaticum*): A precious Spice. *Asian Pacific Journal of Tropical Biomedicine*, 4(2): 90 – 96

Sansi, S. B., Shehu I. and Saka, H. K. (2023). Study on Antimicrobial Activity of Clove (*Syzygium aromaticum*) Crude Extract Against *Staphylococcus aureus*, *Escherichia coli*, *Salmonella* and *Pseudomonas* sp. *Science World Journal* 18 (1): 97 – 100.