

THERAPEUTIC POTENTIALS OF *Azadirachta indica* AND *Ocimum gratissimum*, AND THEIR SYNERGISTIC EFFECTS IN TREATMENT OF *Plasmodium berghei* INFECTED ALBINO MICE

ABSTRACT

Antimalarial resistance is a major challenge for effective control of malaria. This triggered the need to monitor the efficacy of *Azadirachta indica* and *Ocimum gratissimum* in the disease management. The present study compared antimalarial activities of the leaf extracts of the individual plants and evaluated their combined effects on malaria infected mice. The leaves of the different plants were extracted with absolute ethanol (BDL 95%) for the test. Clean albino mice were experimentally infected intraperitoneally with chloroquine-sensitive *Plasmodium berghei* Nk65 strain. Parasitaemia level was determined before parasite inoculation and at 24 hours post treatment period. Efficacy of the leaf extracts was tested on the infected mice using Peter's 4-days suppressive and curative tests, and secondary biological assessment procedures. The lethal median dose (LD₅₀) recorded for neem and clove basil leaf extracts were 31.62 and 1246.9 mg/kg body weight, respectively. Infected mice treated with leaf extracts of the plants and their combinations produced significant dose-dependent activity against the parasite (P<0.05). Highest reduction of parasitaemia was observed on day 4. Maximum parasitaemia reduction (78.65%) was attained with 30mg/kg of the combination of the extracts on the 7th day. These observations indicate better anti-malarial activity of the combination therapy as compared with the individual extracts of the plants, and also show their good antimalarial potentials.

Key Words: Malaria Treatment, Drug resistance, Phytochemicals, Leaf extraction, Synergism of phytochemicals.

INTRODUCTION

Malaria is an infectious and life-threatening disease caused by *Plasmodium* species and vectored by the female *Anopheles* mosquitoes [1, 2]. Of over forty recognized species of the *Plasmodium* parasites, five species namely: *Plasmodium vivax*, *P. ovale*, *P. malariae*, *P. falciparum*, and *P. knowlesi* have been reported to infect humans [3, 4, 5]. A few other species such as *P. Berghei*, *P. yoelii*, *P. chabaudi*, and *P. vinckei*, have also been reported to be infective in other mammals like rodents, and monkeys [6].

According to the World Health Organization (WHO), approximately 219 million cases and 440 thousand deaths were reported due to malaria in 2018 [7]. Nigeria accounts for more malaria cases and deaths than any other country in the world and there were an estimated 68 million malaria cases with over 194,000 deaths in 2021[8]. *Plasmodium falciparum*, the most virulent of the malaria parasites, is responsible for the vast majority of the mortality and morbidity associated with malaria infections in Nigeria. Pregnant women and children less than five years are most vulnerable to malaria attack [3, 9].

The widespread resistance of malaria parasites against many antimalaria drugs is a factor in the economic constraints of malaria control. There is evidence of resistance against most antimalarial drugs which is worsened by the emergence of chloroquine-resistant strains of *P. falciparum*, the malaria parasite responsible for most of the death cases every year. This led to the use of artemisinin combination therapy (ACT) and further search for new phytotherapeutic agents [10, 11].

If not promptly treated with effective medicines, malaria can cause severe illness that is often fatal [3]. Hence, treatment of malaria with combination drug regimens have become the practice of choice because of their increased therapeutic efficacy over monotherapy and the other benefits including decreased cytotoxicity, delay or prevention of the development of drug resistance [12].

But affordability and accessibility limit the use of artemisinin combination therapy and these can lead to poor treatment practices, production of substandard forms of the drug, inadequate patient adherence to prescribed antimalaria regimen, which may in turn lead to treatment failures. Following the problems associated with the implementation of ACT, majority of the populations depend on traditional medical remedies [13], mainly from plants. The traditional herbal medicines are often more available, affordable, and sometimes perceived as being more effective than conventional anti-malarial drugs including artemisinin combination therapy (ACT) [14].

Traditional herbs have been used to treat malaria for thousands of years. Most of these plants are used in the form of monotherapies and only a few plants are taken in combined therapies [15]. Antimalarial plants used in combination may promote the effectiveness of each plant, with efficacy being achieved by the use of a lower dose of each plant extract. This would confer pharmacological benefits, as one extract of the plants clears infection from one body system, and the other clears it from a different site of the same body [14]. Also, synergism with antimalarial agents could be utilized to prevent or delay the emergence *in vivo* of resistant populations of the parasite [16].

Azadirachta indica (Neem) plants from the Meliaceae family are extensively used as traditional medicinal remedies against malaria in the tropics [17, 18, 19]. Several studies demonstrated that *A. indica* leaf, seed and stem bark extract possess *in vivo* inhibitory activity on *P. falciparum* asexual stages [18, 11, 20]. Antimalarial activity of *Ocimum gratissimum* (African basil/clove basil) has also been demonstrated [12]. Although these plants are used in the traditional treatment of malaria, there is the

need to scientifically assess the efficacy of the individual plant extract and their combinations in the treatment of malaria.

The main aim of the present study was to compare the antimalarial effects of *Azadirachta indica* and *Ocimum gratissimum* extracts with chloroquine in *Plasmodium berghei*-infected mice. The specific objectives were to assess the effects of the ethanolic leaf extracts of *A. indica*, *O. gratissimum*, and their combined extracts in reducing parasitaemia in the albino mice; and the synergistic or inhibitory effects of the combined extracts in treating rodent malaria.

MATERIALS AND METHODS

Study Area

This study was undertaken in Owerri at the Federal Medical Centre Laboratory. Owerri is the capital city of Imo State of Nigeria, located between latitudes $5^{\circ} 31'$ and $6^{\circ} 27'$ N and Longitudes $7^{\circ} 00'$ and $7^{\circ} 05'$, and approximately 100 square kilometers (40sq miles) in area, and consists of three Local Government Areas including Owerri Municipal, Owerri North, and Owerri West (Figure 1). Owerri has an estimated population of 1,401,873 according to the 2006 census. It is bordered by the Otamiri River to the East and the Nworie River to the South [21].

Owerri has an annual temperature of 20°C - 30°C , relative humidity of 71%, rainfall distribution of 113.5m. Tropical rainforest is the predominant vegetation in Owerri although its density has drastically reduced due to anthropogenic activities such as urbanization, deforestation and agricultural activities.




Figure 1: Showing the Map of Owerri

Ethical Considerations

Approval for the study was obtained from the Ethical Committee, Public Health Unit of the Owerri Municipal Council (OWMC), Owerri, Imo State, Nigeria. The caring and experimental use of the mice was in accordance with the National Institute of Health Guidelines for care of Laboratory Animals.

Study Design

The work was an experimental study aimed at investigating the *in vivo* antimalarial effects of ethanolic leaf extracts of *A. indica* and *O. gratissimum* in *Plasmodium berghei*-infected Swiss albino mice (*Mus musculus* L: Muridae), using chloroquine as the standard drug. The parasites and the mice were sourced from the National Institute of Medical Research (NIMR), Yaba, Lagos, Nigeria. The plant materials (*A. indica* and *O. gratissimum*) were collected from the Botanical Garden of the Ministry of Agriculture and Natural Resources, Owerri, Imo State, Nigeria. Assessment of parasitaemia, and antimalarial effects of the plant extracts were done using standard laboratory analytical procedures at the Federal Medical Centre, Owerri.

Preparation of Plant Materials

Two kilogrammes (2kg) of each plant material was harvested, washed twice in clean water, and allowed to drip dry. The plants leaves were air-dried separately in the laboratory at room temperature for three weeks to a constant weight before

pulverizing. A 100g each, of the pulverized leaves of the plants was measured and soaked in 500ml of absolute ethanol (BDL 95%) for 24 hours. The mixture was then filtered using Watman 2.0 filter paper. The filtrate was evaporated to dryness in a Rotary evaporator at a temperature of 40°C. The extracts were kept in a tightly closed bottle in a refrigerator until needed for anti-malaria testing.

The stock solution of each plant filtrate was prepared by dissolving 10g of the extract in 100ml of distilled water to give a stock concentration of 0.10g/ml [22].

Maintenance of Study Animals.

One hundred and seventeen mature Swiss albino mice (*Mus musculus*) (25-30g) of either sex free from infection, were used. The animals were housed in mosquito screened-cages lined with wood chip beddings and were stabilized for ten days in the laboratory before being used for the experiments. The mice were maintained on a standard rat diet (Pfizer) and water *ad libitum*. They were kept at ambient room temperature of 12hour light and dark exposure cycles.

Maintenance of Parasites

Plasmodium berghei (NK65 strain) was used as the rodent malaria parasite in the study. The chloroquine sensitive cryopreserved parasites stored at -80°C were revived, stabilized and maintained by serial passage of blood from infected mice to clean mice that served as donor mice in the study.

Collection, Preparation of Inoculum, Parasite Inoculation and Establishment of Infection

The blood of a donor mouse was collected in heparinized syringe and diluted in phosphate buffered saline to 10⁸ parasitized erythrocytes per ml.

The infection of mice was initiated by needle passage of the *P.berghei* parasite preparation from the donor mouse to healthy test mice via an intraperitoneal route [23]. Parasitaemia was monitored by microscopic examination of Giemsa-stained thin blood smears. The number of parasitized erythrocytes in each of the ten such fields was counted thrice and the average count was computed to give the level of parasitaemia of each mouse [23, 24].

The drug chloroquine was administered intramuscularly while the plant extracts and distilled water were administered orally with the aid of an intravenous medicut/mediflon cannula used as an improvised oral cannula [22].

Acute Toxicity Studies

Albino mice (25-30g) of either sex were used. The median lethal dose (LD₅₀) was determined for each of the neem leaf and love leaf extracts using the method previously described by Lorke [25]. In the first phase, the mice were divided into three groups with three mice in each group and were administered with the ethanol leaf extracts at doses of 10, 100, and 1000 mg/kg body weight respectively via the oral route. The mice were then observed for signs of toxicity and death for 24 hrs. In

the second phase, groups of one mouse each were treated with more specific doses of the extract respectively depending on the result obtained from the first phase and observed for signs of toxicity and death in 24hrs. The final LD₅₀ was calculated as the geometric mean of the lowest dose that caused death and the highest dose for which the animal survived.

LD₅₀ = $\sqrt{\text{Mean of the lowest dose that caused death and the highest dose for which the animal survived.}}$

Matsumura [26] and Corbett *et al.* [27] classified chemicals based on their LD₅₀ values as follows:

Extremely toxic	LD ₅₀ ≤ 1 mg/kg
Highly toxic	LD ₅₀ 1-50 mg/kg
Moderately toxic	LD ₅₀ 50-500 mg/kg
Slightly toxic	LD ₅₀ 500-5000 mg/kg
Practically non-toxic	LD ₅₀ 5000-15000 mg/kg
Harmless	LD ₅₀ > 15g/kg

Deriving from the outcome of the acute toxicity studies, dosage ranging from 20-60 mg/kg body weight for both extracts of *Azadirachtaindica* and *Ocimumgratissimum* leaves were screened for therapeutic activities from which the dose(s) that gave optimum and consistent results were selected. The graded dose used in this study was 50mg/kg body weight for both extracts.

Experimental Set up and Treatments.

The mice were divided into three experimental groups for the assessment of the efficacy of the plant materials. These were
 Group 1 or Primary Biological Assessment Group
 Group 2 or Secondary Biological Assessment Group and
 Group 3 or Curative / Therapeutic Group

In Group 1, fifty four mice were used and they were divided into six (6) groups (A,B,C,D,E and F) of nine (9) mice each. The nine (9) mice in each group were sub- grouped into three (1,2,3), of three (3) mice in each subgroup to represent the experimental mice and control mice (i.e positive and negative controls). The mice were infected by method described by Peter and Anatoli [23], and David *et al* [28], with 0.2ml of diluted (10⁸) parasitised erythrocytes/ml infected blood each intraperitoneally (i.p), except those in group F that formed control group. Treatment of mice in experimental groups A to E commenced 2 hours after infection at a dose of 10mg/kg body weight for the individual/single plant extracts, and 10ml/kg for chloroquine and each of combined plant extracts, and treatment was repeated daily with the same dose for 72 hours. Percentage parasitaemia was assessed beginning

from 24 hours post treatment, and was repeated on daily basis throughout the treatment period.

- Group A - Control (Infected with no treatment).
- Group B - Infected and treated with chloroquine (CHQ) (10ml/kg)
- Group C - Infected and treated with *A. indica* leaf extract (10mg/kg).
- Group D - Infected and treated with *O. gratissimum* leaf extract (10mg/kg).
- Group E - Infected and treated with combined extract each of *A. indica* and *O. gratissimum leaf* extracts (10mg/kg).
- Group F - Not infected (Normal Control).

Twenty four hours after the last treatment (96 hours post-infection) blood smears from all the mice were prepared and stained with Giemsa stain for assessment of parasitaemia.

In Group 2, forty five mice were used and they were divided into fifteen groups of 3 mice per group as follows:

- Group A - Infected with no treatment (negative control)
- Group B - Infected and treated with 10ml/kg chloroquine (standard control)
- Group C₁ - Infected and treated with 20mg/kg *A. indica* leaf extract
- Group C₂ - Infected and treated with 30mg/kg *A. indica* leaf extract
- Group C₃ - Infected and treated with 40mg/kg *A. indica* leaf extract
- Group C₄ - Infected and treated with 50mg/kg *A. indica* leaf extract
- Group D₁ - Infected and treated with 20mg/kg *O. gratissimum* leaf extract
- Group D₂ - Infected and treated with 30mg/kg *O. gratissimum* leaf extract
- Group D₃ - Infected and treated with 40mg/kg *O. gratissimum* leaf extract
- Group D₄ - Infected and treated with 50mg/kg *O. gratissimum* leaf extract
- Group E₁ - Infected and treated with 15mg/kg each of *A. indica* and *O. gratissimum* leaf extracts
- Group E₂ - Infected and treated with 20mg/kg each of *A. indica* and *O. gratissimum* leaf extracts
- Group E₃ - Infected and treated with 25mg/kg each of *A. indica* and *O. gratissimum* leaf extracts
- Group E₄ - Infected and treated with 30mg/kg each of *A. indica* and *O. gratissimum* leaf extracts
- Group F - Not infected, but administered with 10ml/kg distilled water (normal control).

Treatment of mice in experimental groups A to E commenced 2 hours after infection at varying doses for the individual/single plant extracts and each of the combined plant extracts, and 10ml/kg for chloroquine . Treatment was repeated daily with the same dose for 72 hours. Twenty four hours after the treatment, blood smears were

made from blood bled from the tails of the treated and untreated mice, and stained with Giemsa to assess parasitaemia. Parasite reduction was computed.

In Group 3, eighteen mice were used for the experiment. Mice maintenance, infection, drug preparation and application were same as for the basic 4-day treatment protocol. Treatment commenced on day four when parasitaemia was well observed in the infected mice and percentage parasitaemia was assessed beginning from 24h post treatment. Fifty milligram per kilogram body weight of *A. indica* leaf extract, 50mg/kg body weight of *O. gratissimum* leaf extract, 10ml/kg body weight of chloroquine, and 30mg/kg body weight each of *O. gratissimum* leaf extract and *A. indica* leaf extract were used for the treatment.

Assessment of Parasitaemia

Before parasite inoculation and at 24 hours post treatment period, the experimental mice were bled from their tails, thin blood smears were made by fixing for 5 minutes using methanol stained with 10% Giemsa stain in phosphate buffer, P^H 7.2, and examined microscopically under oil immersion at x100. Parasitaemia level was determined by counting the parasites observed in 10 fields of approximately 100 erythrocytes per field. The difference between the mean values of the experimental group was calculated and expressed as percentage parasitaemia inhibition using the following equation:

% Inhibition =

$$\frac{\text{ParasitisedRBCinNegativeControl} - \text{ParasitisedRBCinstudygroup} \times 100}{\text{ParasitisedRBCinNegativeControl}}$$

Data Analysis

Data was analysed using the statistical package for social sciences (SPSS) version 2.0.

Tabulations, percentages and graphs were used to present results. Statistically significant differences were analyzed using Analysis of variance (ANOVA). Graphs were plotted using Microsoft Excel, version 2007.

RESULTS

Acute Toxicity Study and Behavioural Effects of the Plants' Leaf Extracts

Acute toxicity test of the *A. indica* leaf extract at concentrations of 100 and 1000 mg/kg produced mortality after 24hrs of observation. The median lethal dosage (LD₅₀) of this extract was 31.80 mg/kg. However, the animals were observed to experience slow movement within the first six hours of the extracts. The *O. gratissimum* leaf extract did not show any mortality for the 100mg/kg dose and it recorded a higher lethal dosage of 1264.9 mg/kg (Table 1). According to Matsumura [26] and Corbett *et al.* [27], the *A. indica* leaf is highly toxic while the *O. gratissimum* is classified as slightly toxic.

Table 1: Lethal Dose (LD₅₀) of the Ethanol Extracts of *Azadirachta indica* and *Ocimum gratissimum*

Extract	LD ₅₀ (mg/kg)	Verdict
<i>A. indica</i>	31.62	Highly toxic
<i>O. gratissimum</i>	1264.9	Slightly toxic

Intensity of Parasitaemia

Few minutes prior to infection of the mice, no malaria parasite was observed in the peripheral blood of all the experimental animals. Detectable parasitaemia was recorded from day 3 post infection (p.i). On the 4th day of post infection, parasite intensity (Table 2), was between 12030±61.83 and 12292±101.61 in all the infected mice. Parasite intensity was not significantly different in the treatment groups B,C,D and E (P>0.05; P= 0.958, 1.00, 0.958, and 0.99).

After treatment of mice with 10mg/kg body weight on the day 4 post infection, the parasite intensity in group A (infected but not treated) rose significantly from 12174±1.00 to 13689±1.00 (P<0.05; P=0.00). The parasite intensity of mice infected and treated with chloroquinein (group B, standard control) significantly decreased to 3.00 ± 0.58 (P<0.05; P=0.00). There were no significant reductions in parasite intensity among those treated with 10mg/kg body weight of mice with the extracts of *A. indica* and *O. gratissimum* separately or combined.

Table 2: Changes in Parasite Intensity before Infection, after Infection, and after Administration of Ethanolic Leaf Extracts of *A. indica*, and *O.gratissimum* in *P.berghei*-infected Mice (Primary Biological Assessment of *A.indica* and *Ogratissimum* Extracts Efficacy)

Experimental groups	Parasite intensity					
	Before infection(x10 ³)		After infection (x10 ³)		After treatment (x10 ³)	
A - Infected, No Treatment	0.000± 0.00	0.000 Sig P<0.05	12174.00 ± 1.00	0.000 Sig P<0.05	136891.00 ± 1.00	0.000 Sig P<0.05
B - 10ml/kg of Chloroquine	0.000± 0.00	0.000 Sig P<0.05	12030.00 ± 61.83	0.958 NS P>0.05	3.00 ± 0.58	0.000 Sig P<0.05
C - 10mg/kg of <i>A. indica</i>	0.000± 0.00	0.000 Sig P<0.05	12267.00 ± 274.64	0.754 NS P>0.05	3601.00 ± 330.77	0.157 NS P>0.05
D - 10mg/kg of <i>O. gratissimum</i>	0.000± 0.00	0.000 Sig P<0.05	12292.00± 101.61	1.000 NS P>0.05	4367.00 ± 275.99	0.992 NS P>0.05
E - 10mg/kg each of <i>A.i</i> & <i>O.g</i>	0.000± 0.00	0.000 Sig P<0.05	12224.00 ± 22.61	0.999 NS P>0.05	1710.00± 438.25	0.416 NS P>0.05
F-Not Infected, Normal Control	0.000± 0.00	0.000 Sig P<0.05	0000.00 ± 0.00	0.000 Sig P<0.05	0.00 ± 0.00	0.000 Sig P<0.05

Key: (*A.i*)=*Azadirachta indica*, (*O.g*)=*Ocimum gratissimum*.

.Table 3 indicates that few minutes before infection of the mice, there were no significant differences in parasite intensity in both the control groups A, B, and F, and treatment groups C₁-C₄,D₁-D₄ and E₁-E₄ (P<0.05; P=0.00). No parasite was observed in the peripheral blood of all the experimental mice.

Two hours after infection, parasite intensity was between 12316±656.81 and 12705±261.40 in all the infected mice. Parasite intensity was not significantly different in treatment groups B, C₁-C₄,D₁-D₄ and E₁-E₄ (P>0.05; P=1.00), that were infected with the parasites.

Ninety-six hours after treatment with varying doses (10ml/kg, 15mg/kg, 20mg/kg, 25mg/kg, 30mg/kg, 40mg/kg, and 50mg/kg) body weight of the mice, parasite intensity in group A (infected but not treated) rose significantly from 12467±611.01 to 312526 (P>0.05; P=1.00). The parasite intensity in group B (infected and treated with chloroquine) significantly decreased to 3.00±0.58 (P>0.05; P=1.00). There were non-significant reductions in parasite intensity among those treated with the plant extracts of *A. indica* and *O. gratissimum* and their combined regimens (P>0.05; P=1.00).

Table 3: Changes in Parasite Intensity before Infection, after Infection, and after Administration of Ethanolic Leaf Extracts of *A. indica* and *O. gratissimum* in *P. berghei*-infected Mice at different Concentrations (Secondary Biological Assessment of *A. indica* and *O. gratissimum* Extracts Efficacy).

Experimental groups	Parasite intensity					
	Before infection(x10 ³)		After infection(x10 ³)		After treatment (x10 ³)	
A - Infected, No Treatment	0.000± 0.00	0.000Sig P<0.05	12467.00 ± 611.01	1.000 NS P>0.05	312526.00 ± 149404.59	0.000Sig P<0.05
B - 10ml/kg of chloroquine	0.000± 0.00	0.000Sig P<0.05	12316.00 ± 656.81	1.000 NS P>0.05	3.00 ± 0.58	1.00. NS P>0.05
C ₁ - 20mg/kg of <i>A. indica</i>	0.000± 0.00	0.000Sig P<0.05	12439.00 ± 356.31	1.000 NS P>0.05	3929.00 ± 53.15	1.00. NS P>0.05
C ₂ - 30mg/kg of <i>A. indica</i>	0.000± 0.00	0.000Sig P<0.05	12504.00 ± 517.04	1.000 NS P>0.05	3614.00 ± 16.70	1.00. NS P>0.05
C ₃ - 40mg/kg of <i>A. indica</i>	0.000± 0.00	0.000Sig P<0.05	12450.00 ± 671.38	1.000 NS P>0.05	3372.00 ± 25.51	1.00. NS P>0.05
C ₄ - 50mg/kg of <i>A. indica</i>	0.000± 0.00	0.000Sig P<0.05	12363.00 ± 469.67	1.000 NS P>0.05	3115.00 ± 13.01	1.00. NS P>0.05
D ₁ - 20mg/kg of <i>A. indica</i>	0.000± 0.00	0.000Sig P<0.05	12472.00 ± 408.24	1.000 NS P>0.05	4872.00 ± 37.02	1.00. NS P>0.05
D ₂ - 30mg/kg of <i>O. gratissimum</i>	0.000± 0.00	0.000Sig P<0.05	12658.00 ± 415.44	1.000 NS P>0.05	4646.00 ± 45.13	1.00. NS P>0.05
D ₃ - 40mg/kg of <i>O. gratissimum</i>	0.000± 0.00	0.000Sig P<0.05	12508.00 ± 370.66	1.000 NS P>0.05	4422.00 ± 10.50	1.00. NS P>0.05
D ₄ - 50mg/kg of <i>O. gratissimum</i>	0.000± 0.00	0.000Sig P<0.05	12625.00 ± 341.15	1.000 NS P>0.05	4129.00 ± 91.35	1.00. NS P>0.05
E ₁ - 15mg/kg each of <i>A.i</i> & <i>O.g.</i>	0.000± 0.00	0.000Sig P<0.05	12551.00 ± 155.38	1.000 NS P>0.05	2785.00 ± 20.59	1.00. NS P>0.05
E ₂ - 20mg/kg each of <i>A.i</i> & <i>O.g.</i>	0.000± 0.00	0.000Sig P<0.05	12705.00 ± 261.40	1.000 NS P>0.05	2576.00 ± 12.50	1.00. NS P>0.05
E ₃ -25mg/kg each of <i>A.i.</i> & <i>O.g.</i>	0.000± 0.00	0.000Sig P<0.05	12578.00 ± 241.33	1.000 NS P>0.05	2321.00 ± 16.56	1.00. NS P>0.05
E ₄ - 30mg/kg each of <i>A.i.</i> & <i>O.g.</i>	0.000± 0.00	0.000Sig P<0.05	12396.00 ± 221.74	1.000 NS P>0.05	2106.00 ± 31.76	1.00. NS P>0.05
F-Not Infected, Normal Control	0.000± 0.00	0.000Sig P<0.05	0.000 Sig P<0.05	0.000 Sig P<0.05	0000.00± 0.000	0.00 Sig P<0.05

Table 4 shows that few minutes prior to infection, there were no significant differences in parasite intensity in both the control groups A,B,.and F, and treatment groups C, D, and E, ($P<0.05$; $P=0.00$). No parasite was observed in the peripheral blood of all the experimental mice.

Twenty-four hours after infection, parasite intensity was between 136853 ± 1022.01 and 138633 ± 852.31 in all the infected mice. Parasite intensity was not significantly different in treatment groups B,C,D, and E ($P>0.05$; $P=0.995$, 0.599 , 0.9902 , and 1.00), that were infected with the parasites.

Twenty-four hours after last treatment with 30mg/kg and 50mg/kg body weight of the mice, parasite intensity in groups A (infected but not treated) rose significantly from 138083 ± 1.00 to 181358.00 ± 1.00 ($P<0.05$; $P=0.00$). The parasite intensity in groups B (infected and treated with chloroquine) significantly decreased to 3.00 ± 0.58 ($P<0.05$; $P=0.00$). There were non-significant reductions in parasite intensity among those treated with plant extracts of *A.indica*,and*O. gratissimum* and their combined regimens ($P>0.05$; $P=1.00$).

Table 4: Changes in Parasite Intensity before Infection, after Infection, and after Administration of Ethanolic Leaf Extracts of *A. indica*, and *O. gratissimum* in *P. berghei*-infected Mice (Therapeutic Effects of *A. indica* and *O. gratissimum*).

Experimental groups	Parasite intensity					
	Before infection(x10 ³)		After infection (x10 ³)		After treatment (x10 ³)	
A - Infected, No Treatment	0.000 ± 0.00	0.000 Sig P<0.00	138083.00 ± 1.000	0.000 Sig P<0.05	181358.00 ± 1.00	0.000 Sig P<0.00
B - 10ml/kg of Chloroquine	0.000 ± 0.00	0.000 Sig P<0.00	138633.00 ± 852.31	0.995 NS P> 0.05	3.00 ± 0.58	0.000 Sig P<0.05
C - 50/mg/kg <i>A. indica</i>	0.000 ± 0.00	0.000 Sig P<0.00	136853.00 ± 1022.01	0.599 NS P> 0.05	1957.00 ± 694.54	0.137 NS P>0.05
D -50mg/kg of <i>O. gratissimum</i>	0.000 ± 0.00	0.000 Sig P<0.00	137971.00 ± 903.51	0.9902NSP> 0.05	3083.00 ± 93.54	0.625 NS P>0.05
E - 30mg/kg each of <i>A.i</i> & <i>O.g</i>	0.000 ± 0.00	0.000 Sig P<0.00	137634.00 ± 2301.11	1.000 NS P> 0.05	1235.00 ± 111.53	0.172 NS P>0.05
F-Not Infected,Normal Control	0.000 ± 0.00	0.000 Sig P<0.00	0.00± 0.00	0.000 Sig P< 0.05	0.00 ± 0.00	0.000 Sig P<0.05

Key: (*A.i*)=*Azadirachta indica*, (*O.g*)=*Ocimum gratissimum*

Antimalarial Effects of Different Dosages of the Plants Leaf Extracts

The treatment of animals with different dosages of ethanolic leaf extracts of the plants showed a different response to increased dosage in line with protection against infection (Figure 2). The parasitaemia in the infected but untreated group increased with days of infection while the parasite-infected and treated groups had significant reduction ($P < 0.05$; $P = 0.00, 0.009, 0.002, \text{ and } 0.00$) in parasitaemia level from day one to seven. From day 1 to 7 of the treatment period, all groups showed significant difference in parasitaemia levels as compared to the untreated control group ($P > 0.05$; $P = 0.122, 0.060, \text{ and } 0.088$).

The extract of *A. indica* at 20mg/kg, 30mg/kg, and 40mg/kg showed parasitaemia reduction of 57.80%, 59.0%, and 60.10%; and 62.50%, 63.60%, and 64.50% on days 4 and 7 respectively, while the 50mg/kg of *A. indica* treated group recorded 61.40% parasitaemia reduction on day 4 and 66.00% on day 7. Values observed with 20mg/kg, 30mg/kg and 40mg/kg of *O. gratissimum* were 49.08%, 52.70%, and 55.10%; and 54.10%, 57.00%, and 59.88% on days 4 and 7, respectively, while 50mg/kg of *O. gratissimum* gave 56.88% and 61.10% reduction on days 4 and 7, respectively (Figure 4). Combination doses of 15mg/kg, 20mg/kg, 25mg/kg and 30mg/kg of each of the plant leaf extracts gave a significantly higher reduction in parasitaemia as compared to combined treatments with 40 and 50mg/kg. Maximum reduction of 78.65% parasitaemia was attained for the extracts combinations of 30mg/kg of each plant extract on the seventh day, indicating synergistic action of the extracts. By the twenty-eight day post treatment, the parasitaemia had gone below detectable levels in all treated groups. The plant extracts showed increase in parasitaemia inhibition with increase in dose administered on the infected animals, i.e. dose-dependent inhibition of parasitaemia, and even at the higher dosages, the animals survived longer than the chloroquine treated ones.

The therapeutic effect of the leaf extracts of the plants on the infected animals (Figure 3) showed that the animals responded to the extracts within the first seven days of treatment with 69.89% reduction of parasitaemia for *O. gratissimum* treated group, 71.11% for *A. indica* treated group, and 82.70% for combined extracts of *A. indica* and *O. gratissimum* treated groups before the parasitized red blood cells started to increase which later led to the deterioration of the health condition of the animals. The antimalarial action of the combined leaf extracts of *A. indica* and *O. gratissimum* (82.70%) was close to that of chloroquine (97.98%).




Figure 2a: Effects of Different Doses of *A. indica* Extract on Parasitized RBC of *P.berghei* Infected Mice




Figure 2b: Effects of Different Doses of *O. gratissimum* Extracts on Parasitized RBC of *P. berghei* Infected Mice




Figure 2c: Effects of Different Doses of Combined Extracts of *A. indica* and *O. gratissimum* on Parasitized RBC of *P. berghei* Infected Mice




Figure 2d: Effects of chloroquine on Parasitized RBC of *P. berghei* Infected Mice




Figure 3: Therapeutic Effect of Ethanolic Extracts of the Plants Leaves on albino Mice Infected with *P. berghei*

DISCUSSION

The *in vivo* antimalarial activities of *A. indica* and *O. gratissimum* leaf extracts were studied in *P. berghei*-infected mice to determine the treatment outcomes of the extracts from January to March, 2021.

Before parasite inoculation and at 24 hours post treatment period, parasitaemia was monitored by microscopic examination of Giemsa-stained thin blood smears in all the experimental mice. The experimental mice were placed into various groups for the assessment of the efficacy of the plant materials used in the study. The significant increase in the level of parasitaemia in the infected untreated group recorded from day 1 to day 7, as symptomised by the enlargement of the red blood cells, anaemia, loss of appetite, and loss of weight as major clinical signs observed during the course of treatment is in tune with the view that parasitaemia increases progressively after infection until the point of death if no suitable treatment is administered [29, 9]. There were significant increases in parasitaemia reduction with increase in doses of the extracts administered in the treated groups. This indicates that as the dose increases, the strength of the antiplasmodial activity of the plant extracts also increases.

The reduction in parasitaemia was highest in the group administered with the combined extracts of *A. indica* and *O. gratissimum* (Figure 3, [82.70%]) than those administered with single plant extracts. This may be attributed to the antiplasmodial activities of the various ethanolic extracts, on one hand, and result of the synergistic effect of the compounds or their metabolites on the other hand. These findings agree with the reports of Alehegn *et al* [30], Ceravolo *et al* [31], Maximus *et al* [32], Oladeji *et al* [33], Singh *et al* [34], Ekpo and Ekanemesang [35], and Cissy *et al* [36], which indicate that phytochemicals are responsible for the antimalarial activities of many plants. Also, these metabolites may be acting individually or in synergy with one another to produce antimalarial activity. This is attributable to the ability of the plant extracts to reduce the intensity of parasitaemia in the

infected and treated groups, which is in tune with studies reported by Chaniad *et al* [37], Joseph & Samson [38], and Osonwaet *al* [10]. The activity of the combined leaf extracts of *A. indica* and *O. gratissimum* was also observed to be better than the individual/separate extracts of *A. indica* and *O. gratissimum*, which corroborates the results observed in their antiplasmodial activity [39, 10, 40].

CONCLUUSION

From observations, the individual leaf extracts of *A. indica* and *O. gratissimum* have been shown to be effective in the treatment of malaria parasite in mice by reducing the parasite intensity. All the therapeutic observations were more pronounced in the combined extracts treatments indicating synergistic action of the extracts. It is therefore recommended that the coadministration of these locally used plants, especially at the concentrations investigated in this study be taken for the treatment and management of malaria fever as their action is very close to that of the antimalarial drug, chloroquine which was the standard drug used in this study.

CONFLICT OF INTERESTS

The authors declare no conflict of interest.

AUTHORS' CONTRIBUTIONS

The study was designed by Anyasodor Charles C., who also performed the statistical analysis, wrote the protocol as well as the first draft of the manuscript. Prof. Onyido Angus E, Dr. Anyasodor Anayochukwu E., and Dr. Ogbuefi Emmanuel O, managed the analysis of the study. Prof. Onyido managed the literature searches, among others. All authors read and approved the final manuscript.

The authors declare no conflict of interest.

REFERENCES

1. Sitali, L, Miller, JM, Mwenda, MC, Bridges, DJ, Hawela, MB, Hamainza, B. Distribution of *Plasmodium* species and assessment of performance of diagnostic tools used during a malaria survey in southern and western provinces of Zambia. *Malaria Journal*. 2019; 18,130-133.
2. Sato, S. Plasmodium – a brief introduction to the parasites causing human malaria and their basic biology. *Journal of physiology/Anthropology*. 2021; 40: 1-5.
3. World Health Organization/RBM. Documentation of the socio-economic impact of malaria epidemics in Africa. Final report. 2019; [www.rollbackmalaria.org/seim2019/\(WHO/HTM/MAL/2019.1102\)](http://www.rollbackmalaria.org/seim2019/(WHO/HTM/MAL/2019.1102)).
4. Woldearegai, TG, Lalremruata, A, Nguyen, IT, Gmeiner, M, Veletzky, L, Tazenda-kuitsouc, G.B. Characterisation of *Plasmodium* infections among inhabitants of rural areas in Gabon. *Journal of Scientific Reports*. 2019; 9(1): 41597-41599.
5. Kotepui, M, Kotepui, KU, Millanez, GD, Masangkau, FR. Severity and mortality of severe *Plasmodium ovale* infection: a systematic review and meta-analysis. *PLoS ONE*. . 2020. 15:e0235014.
6. Larson, B, Barthelemy, N, Stephan, N, Nancy-Diamela, M, Francois, R, Virginie, R, Frank, P. Rodent Malaria in Gabon: Diversity and Host range. *International Journal of Parasitology*. 2019; 10:117-124.
7. World Health Organisation. World malaria report. 2018. <https://www.who.int/publications/report>.
8. World Health Organisation.. Report on malaria in Nigeria 2022. [https://www.afro.who.int/publications/report-malaria-nigeria-2022#:~:text=Malaria%20is%20a%20major%20public,the%20country%2C%20all%20year%20round](https://www.afro.who.int/publications/report-malaria-nigeria-2022#:~:text=Malaria%20is%20a%20major%20public,the%20country%2C%20all%20year%20round.). 2023.
9. Iwuchukwu, IC, Vincent CN. Studies on prevalence of malaria and its adverse foetal outcomes in FMC, Owerri, Imo State, Nigeria. *Archives of Community Medicine and Public Health*. 2021; 7(2): 151-163.
10. Osonwa, EU, Mbonu, DO, Eluu, CS, Oli, AN. Antiplasmodial and biochemical effects of combination of ethanolic leave-extract of *Azadirachta indica* and *Ocimum gratissimum* on *Plasmodium berghei*-infected mice. *African Journal of Pharmaceutical Sciences and Pharmacy*. 2017; 5:1-6.

11. Afolabi, OJ, Simon-oke, IA, Oladokun, OI. Antiplasmodial Activity of Ethanolic Extract of Neem Leaf (*Azadirachta indica*) in Albino Mice Infected with *Plasmodium berghei*. *International Archives of Clinical Pharmacology*.2021; **7**:2572-2578.
12. Nwankwo, EN, Ezekwesili, OO, Chude, CM. In vitro antiplasmodial and in vivo toxicity potentials of *Menthapiperita* and *Ocimum gratissimum* essential oils and their synergistic effect with conventional antimalarial drugs against *Plasmodium falciparum*. *International Journal of Mosquito Research*.2022;9(1): 114-122.
13. Omagha, R, Idowu, ET, Adeneye, AK. Survey of ethnobotanical cocktails commonly used in the treatment of malaria in southwestern Nigeria. *Future Journal of Pharmaceutical Sciences*.2021;7:152-158.
14. Nigussie, G, Wale, M. Medicinal plants used in traditional treatment of malaria in Ethiopia: a review of ethnomedicine, anti-malarial and toxicity studies. *Malaria Journal*.2022; 21: 262-266.
15. Million, E, Mulugeta, T, Umeta, B. Traditional Medicine Practice and its Role in the Management of Malaria in Jimma Town, Oromia, Ethiopia. *Journal of Infection and Drug Resistance*.2022; 15:2187-2198.
16. Erhirhie, EO, Ikegbune, C, Okonkwo, OB. Antimalarial herbal drugs: a review of their interactions with conventional antimalarial drugs. *International Journal of Phytomedicine and Physiotherapy*.2021; 7:4-10.
17. Yarmohammadi, F, Mehri, S, Najan, N, Salar, AS, Hosseinzadeh, H. The Protective Effect of *Azadirachta indica* (Neem) against Metabolic Syndrome: A Review. *Iran Journal of Basic Medical Sciences*.2021;24(3): 280-292.
18. Achi, NK., Onyeabo C, Nnate, DA, Ekeleme-Egedigwe, CA, Kalu, IK, Chibundu, IC, Wokoma, GC. Therapeutic effects of *Azadirachta indica* A. juss. leaves in malaria-induced male wistar rats. *Journal of Pharmacy and Pharmacognosy Research*.2018 6(3): 191-204.
19. Mbugi, EV, Sife, AS, Mboni, R., Grace, EPM, Bestina, D, Edda, TL. Effectiveness of *Azadirachta indica* (neem tree) on prevention and treatment of clinical human malaria: A systematic review. *Journal of East Africa Science*. 2021;3:1-4.
20. Annette, H, Barbara, P, Sofia, T, Judith, N, Fabrizio, B. Effects of *Azadirachta indica* seed kernel extracts on early erythrocytic schizogony of *Plasmodium berghei* and proinflammatory response in inbred mice. *Malaria Journal*.2019;18: 35-43.
21. Alex, DWA, Okorie, PU. "Water quality studies of Nworie Rivers in Owerri, Nigeria." *Journal of Mississippi Academy of Sciences*. 2008; 20(1): 55-60.

22. Ejebe, DE, Emudianohwo, JO, Ozoko, T, Simialayi, IM, Esume, CO, Maduodi, UV. An Investigation into the antiplasmodial effect of the ethanol extract of the leaves of *Helliantusannus* in Swiss Albino mice. *Global Journal of Pharmacology*.2011; 5(2): 92-96.
23. Peter, IT, Anatoli, VK. The current global malarial situation. *Malaria parasite biology, pathogenesis, and protection*. American Society for Microbiology, WashingtonDC.1998;pp.11-22.
24. Dawit D, Eyassu, M, Asfaw, D, Dawit,A, Kelbessa, U, Wallelign, M, Daniel, M, Ashenafi, A,yared, M. *In vivo* antimalaria activity of hydrocoholic extracts from *Asparagus africanus* Lam. In mice infected with *Plasmodium berghei*. *Ethiopian Journal of Health Development*. 2006; 20(2): 112-118.
25. Lorke, DA. New Approach to Practical Acute Toxicity Testing. *Achives of Toxicology*. 1983; 54: 275-287.
26. Matsumura, F. *Toxicology of insecticides*. Plenum press, York and London. 1975; pp. 24.
27. Corbett, JR, Wright, K, Baillie, AC. *The Biochemical mode of Action of pesticides*, 2nd Academic press, London and New York. 1984.
28. David, AF, Lip, JR, Simon, LC, Rato, B, Solomon N. Antimalarial drug discovery: efficacy models for compound screening. *Nature Review*. 2004;16: 522-528.
29. White, NJ. Malaria parasite clearance. *Malaria journal*. 2017; 16(1), 88-92.
30. Alehegn, AA., Yesuf, JS, Birru, EM. Antimalarial activity of crude extract and solvent fractions of the leaves of *Bersamaabyssinica*fresen.(*Meliantaceae*) against *Plasmodium berghei* infection in Swiss albino mice. *Evidence-based Complementary and Alternative Medicine*. 2020; 202-212.
31. Ceravolo, IP, Aguiar, AC, Adebayo, JO, Krettli, AU. Studies on activities and chemical characterization of medicinal plants in search for new Antimalarials: a ten year review on Ethnopharmacology. *Frontiers in Pharmacology*. 2021; 12: 734-763.
32. Maximus, MT, Tukan, GD, Prajogo, BE,Mangestuti, A. Antiplasmodial activity and phytochemical constituents of selected antimalarial plants used by native people in west timor Indonesia. *Turkish Journal of Pharmaceutical Sciences*. . 2021; 18(1): 80-86.

33. Oladeji, OS, Oluyori, AP, Dada, AO. Genus *Morinda*: An insight to its ethnopharmacology, phytochemistry, pharmacology and Industrial Applications. *Arabian Journal of Chemistry*. 2022; 15(9): 104-124.
34. Singh, N, Chatterjee, A, Chanu, WK, Vaishalli, PM, Singh, CB, Nagaraj, VA. Antimalarial activity of *Toonaciliata* MJ Roem aqueous methanolic leaf extract and its antioxidant and phytochemical properties. *Journal of Traditional and Complementary Medicine*. 2023;
35. Ekpo, DE, Ekanemesang, MU. Antiplasmodial/antimalarial effect of ethanol extracts of leaves of *Vernonia amygdalina* and *Gongronema latifolium* on the activity of catalase in *Plasmodium berghei*-parasitized mice. *International Journal of BCRR*. 2016; 10(4): 1-9.
36. Cissy, N, Engeu, OP, Berna, O, Norbert, A, Esther, M. *Artemisia annua* L. and *Vernonia amygdalina* Del: A potential Herbal Artemisinin Combination Treatment against malaria. *British Journal of Pharmaceutic Resources*. 2016; 14(3):1-7.
37. Chaniad, P, Techarang, T, Phuwaroanpong, A, Horata, N, Septama, AW, Punsawad, C. Exploring potential antimalarial candidate from medicinal plants of Kheaw Hom remedy. *Tropical Medicine and Infectious Disease*. 2022; 7(11): 368- 377.
38. Joseph, AO, Samson, OT. Antiplasmodial Efficacy of *Anacardium occidentale* in Albino Mice Infected with *Plasmodium berghei*. *Journal of Family Medicine and Disease Prevention*. 2020. DOI: 10.23937/2469-5793/1510123.
39. Ekasari, W, Mahardiani, A, Putri, NT, Wahyuni, TS, Arwati, H. Toxicological evaluation and protective effects of ethanolic leaf extract of *Cassia spectabilis* DC on liver and kidney function of *Plasmodium berghei*-infected mice. *Veterinary Medicine International*, 2022. <https://doi.org/10.1155/2022/6770828>.
40. Anigboro, AA, Onakurhefe, P, Tonukari, NJ, Avwioroko, OJ, Egbeme, E. Quantitative determination of some phytochemicals (phenol, flavonoid, saponin and alkaloid) in twenty-two Nigeria medicinal plants. *Nigerian Journal of Science and Environment*. 2018; 13(1): 86-93.