

Review Article

Neisseria gonorrhoeae: Its Dominant Properties to Establish Contact and Attachment In order to Facilitate Epithelial Invasion and Colonization From a Biochemist Perspective

ABSTRACT

Aim: to discuss about recent epidemiology of gonorrhea and updates regarding its pathogenesis with focus on biochemical aspects of contact and adhesion that precede its epithelial invasion.

Discussion: *Neisseria gonorrhoeae*, a Gram-negative obligate human pathogenic bacterium, infects human epithelial cells and causes sexually transmitted disease in both male and female. Gonorrhea rates are rising in many countries. It could lead to long-term health problems and even infertility. Vulnerable group including men who have sex with men and sex workers that appear to bear disproportionate burdens of gonorrhea. As *N. gonorrhoeae* advances through the steps of disease formation and pathogenesis (transmission, adherence, colonization and invasion, and also immune evasion), the bacterium expresses some virulence factors to facilitate its survival and replication; while on the same time keeping their existence barely invasive and almost undiscoverable by active immune cells. Adherence to epithelial cell becomes the first event that precedes invasion.

Conclusion: Adhesion of gonococci to mucosal epithelial cells appears to be a critical step in the pathogenesis of gonococcal infection. Gonococci have the ability to adhere to a variety of human cells. Gonorrhea has multiple surface proteins that facilitate adhesion. *N. gonorrhoeae* utilizes type IV pili and Opa, opacity-associated proteins, a surface protein involved in cellular attachment that precedes its invasion, Lipooligosaccharides (LOS) and also the outer membrane protein porin (POrB).

Keywords: *mucosal, microcolonies, type IV pili, conserved hydrophobic N terminus, Opa protein, lipooligosaccharide, porin*

1. INTRODUCTION

Neisseria gonorrhoeae is responsible for causing gonorrhea, a sexually transmitted infection which has been reported to infect humans since ancient times [1]. It is a well-adapted organism that is able to overcome the already existing normal microbiome [2], having an awesome ability to accommodate the harsh internal condition of their human host [3] and has maintained ways to evade the host immune properties [4]. Its initial predilection is mainly shallow or restricted near the surface, namely the epithelial mucosa of the female endocervix [5]. Colonization and disruption of the epithelium is actually an important route of infection used by this mucosal pathogen [6]. The normal epithelium actually neutralizes invasion by managing to exfoliate damaged portions of the affected cells while keeping the best mucosal barrier in business [7].

Over the past decade, extensive studies of *N. gonorrhoeae* have shown that in media containing glucose [8], the compound lactate (a classical byproduct of glucose metabolism) actually encourages the bacteria's metabolism [9], and even ensuring its survival inside the human polymorphonuclear leukocytes [10]. These properties could improve its colonization [11], increased lipopolysaccharide synthesis [12], increase gonococcal resistance to complement-mediated killing by human serum [13] and also contribute to antibiotic resistance [14]; or in other words lactate play a central role in *N. gonorrhoeae* pathogenicity. Lactate and glucose are present together in most sites where infection and active inflammation occurs in vivo [15]. This mini-review describes the gonococci dominant properties to establish contact, attachment, epithelial invasion and colonization from a biochemist perspective.

2. EPIDEMIOLOGY AND PATHOGENESIS OF *N. GONORRHOEAE*

By the year 2020, the WHO roughly estimated 82.4 million new STIs due to *N. gonorrhoeae* among adults aged 15 - 49 years old, globally [17]. Prevalence of gonorrhoea is highest among specific populations such as men who have sex with men [18], sex workers [19], transgender individuals [20] and adolescents and young people in high burden countries [17]. Gonorrhoea prevalence in infertile sub populations is several folds higher than in the general sub population, with even higher prevalence in sub population of women suffer from tubal factor infertility and in people with secondary infertility [21]. These allegations contribute the potential role of gonorrhoea in causing infertility and indicate that some type of infertility is very likely preventable by means of preventing gonorrhoea transmission [22].

The main obstacles for portraying its pathogenesis are include its all-in exclusive human host [23] and in combination with its active and continuously modifying superficial sector [24]. Its primary niche is in the surface portion of human internal genital organ [3]. While *N. gonorrhoeae* infects both sexes (men and also women), most commonly manifests as urethritis in men and cervicitis in women [25]. Unfortunately for female patients, silent but active infection tends to develop more intricated consequences, ranging from simple colonization without any distinct clinical appearance [26] to a more severe clinical conditions such as pelvic inflammatory disease [27], ectopic pregnancy [28] and tubal infertility [22,28]. Unfortunately, due to the larger part of infections among female group are asymptomatic [26], this condition give the bacteria the privilege to escalate silently, until then these microorganisms reach deeper anatomical structures and cause more serious problems with more prominent symptoms; a condition called complications. Eventhough a study conducted by Teker et al [29] revealed a significant proportion of asymptomatic female patients somehow able to unloaded their infections spontaneously.

The integrity of the vaginal epithelium is crucial for women's reproductive health and for providing protection against sexually transmitted infections and even HIV [30]. The surface of entire female reproductive tract coated with a variety of epithelial cells [31]. *Neisseria gonorrhoeae* adheres for the first time to the urogenital tract surface by attaching to surface structures namely epithelium. This microorganism using at least four structure to facilitate their attachment, namely: (1) Type IV pili (Tfp), (2) opacity (Opa) proteins, (3) Lipooligosaccharide (LOS), or (4) outer membrane protein porin (PorB) and all of these four, together and simultaneously, arbitrate the complex interactions between *N. gonorhea* and epithelial cells or phagocytes subsequent to adhesion [32].

3. BIOCHEMICAL ASPECTS OF CONTACT AND ATTACHMENT

Major infection mechanism of any mucosal pathogens actually involved the combination of colonization and perturbation of the epithelium at the surface of organ [46]. The epithelium

neutralizes infection by suddenly shedding all broken epithelial cells while keeping the most important function of mucosal, named barrier [47].

Vaginal epithelium is actually inhabited by diverse strains and species of microorganisms considered as normal microbiota [2]. The bacterial composition of vaginal microbiota controls the balance between bacteria and yeast, or in simplified word the equilibrium between health and disease. There must be at least relative endowment of the epithelial cell combine with bacterial cell surfaces regarding to its adhesion, that according to Ferrari et al [48] there is also influence of surface properties, and in addition to that whether and how adhesion is controlled due to certain cell membrane portions [49]. The forces that facilitate Bacterial adhesion combine with the portion affected (cell membrane regions) which was not located above the nucleus are actually bind better compare to regions located above the nucleus; this condition apply to all inhabited organism whether pathogens or commensal and probiotic organism such as Lactobacillus strains that actually implicated in maintaining health condition.

One must kept in mind that, adhesion force ratios over certain membrane regions abroad from and upward the nucleus happen at the same time with the ratios between the amount of attaching microorganism over both portion of the infected regions [50]. Bacterial adhesion forces were dramatically decreased by depleting the epithelial cell membrane of cholesterol [51] or sub-membrane cortical actin [52]. Pathogenic obligate intracellular organism use principally host-derived structures to facilitate its movement through (and also between) cells [53]. Epithelial cells actually have the authority to regulate affected membrane regions to which bacterial adhesion is prevented, very likely to secure the nucleus and maintain its function [47].

Infection of the genital mucosa initiate by *N. gonorrhoeae* entail contact, adherence to and invasion of epithelial cells which preceded colonization [34]. Early bonding of gonococci to the columnar epithelial cells is helped by type IV pili [54] constructed from pilin subunit PilE proteins [55] and pilus tip-associated PilC proteins [56]. Type IV pili are composite structure in the form of long fibers that are assembled by polymerization of a major pilin protein in the periplasm of a wide range of bacteria and archaea. Type IV pili are distinctive dynamic filaments at the surface of many bacteria that can rapidly extend and retract and withstand strong forces. They play crucial roles in pathogenesis, DNA transformation, self-agglutination, proficiency for natural transformation, and twitching/spasm type motility and also are capable of rapid retraction, generating powerful motor forces [53].

In the context of adherence properties of *N. gonorrhoeae*, type IV pili consists of multifunctional polymers of the major pilin protein, which share a conserved hydrophobic N terminus that is a curved extended α -helix, $\alpha 1$, in X-ray crystal structures. Type IV pili plays an important role, that as a Gram-negative causative agent of gonorrhea, to attach to the specific epithelial cells and actively determine the colonization process [55,56]. Type IV pili-mediated gonococcal attachment [59] not directly related to pilin subunit protein PilE but by choice is prefer to be compatible with the expression of the PilC protein, which actually less profuse; and this copurifies with type IV pili. An almost identical system seems to work for the highly related species that come from the same ancestor named *N. meningitidis* [60].

Gonococcal attachment is also facilitate and strengthen by the utterance of phase-variable opacity-associated (Opa) proteins [6,7,57,58]. Opa proteins are specific Gonococcal's variable outer membrane proteins that facilitate closed host-pathogen interaction; it account for the cell tropisms demonstrated by *N. gonorrhoeae*, e.g., for human leukocytes and epithelial cells [63]. Its extensive exploration have successfully allowing researchers to reveal recognition of different host receptors responsible for attachment and also organizing

the certain cell pattern tropism advertised by this bacteria, including epithelial and neutrophils [57,58] and this action being defined by precise adhesin-receptor interactions [62].

Lipooligosaccharide (LOS) also contribute to adherence ability [12,63,64]. This sugar is the main constituent within the outer membrane. Beside adherence, LOS also contributes in pathogenesis by arousing the host inflammatory reflex and also permitting evasion of the initially aroused host's innate immunity by way of sialylation [65]. A naturally occurring variation of the terminal carbohydrates on the lipooligosaccharide (LOS) molecule correlates with altered disease states [12, 63-65].

Biochemically, Neisserial LOS molecules constructed from hexa-acylated lipid A, two keto-deoxyoctulosonate, a carbohydrate (KDO) molecules, and one or more carbohydrate chains of 8–12 saccharide units, the core oligosaccharide; with all the three oligosaccharide chains attached to a lipid A core [66]. The oligosaccharide chains of LOS are almost identical both in sequence and linkage with the oligosaccharides expressed on the surface of human cells [63]. The synthesis of these compounds requires a series of glycosyl transferases. Besides their highly conserved core structures, the terminal oligosaccharides of LOS molecules undergo rapid phase variation [64]. LOS variation is mediated by a change in the number of guanines in the middle of the coding sequences of several key enzymes, which results in alterations in the expression of these glycosyl transferases and the surface expression of various LOS isoforms. At any given time, several LOS structures with varying terminal oligosaccharides may also be pronounced on the part of *N. gonorrhoeae* outer membrane. Due to its strain based variability, it has been difficult to study the structure and function of LOS. The asialoglycoprotein receptor is also expressed in human urethral epithelial cells. Neisserial LOS has a relative paucity of glycosylation when compared to lipopolysaccharide of enteric Gram-negative species. Variations in gonococcal LOS are largely due to phase-variable expression of genes involved in LOS biosynthesis, which determine core oligosaccharide structure and Ist, which determines LOS sialylation.

Last but not least, the outer membrane protein porin (POrB) of *N. gonorrhoeae*. The major outer membrane porin (PorB) expressed by *N. gonorrhoeae* plays multiple roles during infection, in addition to its function as an outer membrane pore [67]. PorB is a voltage-gated pore that mediates ion exchange between *N. gonorrhoeae* and the environment and is essential for bacterial viability, likely due to its ability to allow nutrients access to the periplasm [68]. Porin is found in the outer membrane as a homotrimeric β -pleated barrel; each monomer is 32 to 35 kDa and consists of 16 transmembrane-spanning segments and 8 extracellular loops. *N. gonorrhoeae* strains contain a single porB gene in one of two allelic forms (P.IA or P.IB), and the two alleles have been associated with different biological phenotypes [67,68]. P.IA-expressing strains tend to be associated with disseminated disease, whereas P.IB-expressing isolates typically cause localized urogenital infections. There is ~80% nucleotide sequence similarity between P.IA and P.IB, and the majority of the variation between alleles and between different isolates expressing the same allele occurs within the extracellular loops

In the context of *N. gonorrhoeae*, the known target receptors are being explored and the potential role of receptor modulation in increasing host susceptibility to infection is always be considered and perhaps become the major focus in future study. Most bacteria, commensal or pathogenic, actually while associating with their hosts produce certain specific adhesive molecules superficially. This molecules promotes reciprocation with surface receptors of their host or with available soluble macromolecules. Eventhough bacterial contact and then followed by attachment to superficial lining made by epithelial cells may be beneficial for bacterial colonization, adhesion may also causing potential harm because attachment to

waken immune cells can simplify the process of phagocytosis and elimination. Certain pathogenic microorganisms have deciphered this difficulty by way of releasing an antiphagocytic surface substance that is usually always made from polysaccharide and alternatively by expressing their adhesins on polymeric structures that extend out from the cell surface. This is certainly an interesting topic, related to the virulence of *N. gonorrhoea*, to be explored in the future.

4. CONCLUSION

Adhesion of gonococci to mucosal epithelial cells appear to be a critical step in the pathogenesis of gonococcal infection. Gonococci have the ability to adhere to a variety of human cells. Gonorrhoea has multiple surface proteins that facilitate adhesion. *N. gonorrhoeae* utilize type IV pili and Opa, opacity-associated proteins, a surface proteins involved in cellular attachment that precede its invasion.

CONSENT

Not needed

ETHICAL APPROVAL

Not needed

REFERENCES

1. Vicentini CB, Manfredini S, Maritati M, Di Nuzzo M, Contini C. Gonorrhoea, a current disease with ancient roots: from the remedies of the past to future perspectives. *Infez Med.* 2019;27(2):212-221.
2. Sunarti LS. Microbial Normal Flora: Its Existence And Their Contribution To Homeostasis. *Journal of Advances in Microbiology*, 2022;22 (9): 1-15 <https://doi.org/10.9734/JAMB/2022/v22i930483>.
3. Song W, Yu Q, Wang L-C, Stein DC. Adaptation of *Neisseria gonorrhoeae* to the Female Reproductive Tract. *Microbiology Insights.* 2020;13. <https://doi.org/10.1177/1178636120947077>
4. Płaczkiewicz J. Avoidance Of Mechanisms Of Innate Immune Response By *Neisseria Gonorrhoeae*. *Postępy Mikrobiologii - Advancements of Microbiology.* 2019;58: 367-73. <https://doi.org/10.21307/PM-2019.58.4.367>.
5. Wang LC, Yu Q, Edwards V, Lin B, Qiu J, Turner JR, Stein DC, Song W. *Neisseria gonorrhoeae* infects the human endocervix by activating non-muscle myosin II-mediated epithelial exfoliation. *PLoS Pathog.* 2017;13(4):e1006269. <https://doi.org/10.1371/journal.ppat.1006269>.
6. Yu Q, Wang LC, Di Benigno S, Stein DC, Song W. Gonococcal invasion into epithelial cells depends on both cell polarity and ezrin. *PLoS Pathog.* 2021;17(12):e1009592. doi: <https://doi.org/10.1371/journal.ppat.1009592>.
7. Dong M, Dong Y, Bai J, Li H, Ma X, Li B, et al. Interactions between microbiota and cervical epithelial, immune, and mucus barrier. *Front Cell Infect Microbiol.* 2023;13:1124591. <https://doi.org/10.3389/fcimb.2023.1124591>.

8. Takei M, Yamaguchi Y, Fukuda H, Yasuda M, Deguchi T. Cultivation of *Neisseria gonorrhoeae* in liquid media and determination of its in vitro susceptibilities to quinolones. *J Clin Microbiol.* 2005;43(9):4321-7. <https://doi.org/10.1128/JCM.43.9.4321-4327.2005>.
9. Smith H, Yates EA, Cole JA, Parsons NJ. Lactate stimulation of gonococcal metabolism in media containing glucose: mechanism, impact on pathogenicity, and wider implications for other pathogens. *Infect Immun.* 2001;69(11):6565-72. doi: <https://doi.org/10.1128/IAI.69.11.6565-6572.2001>.
10. Atack JM, Ibranic I, Ong CL, Djoko KY, Chen NH, Vanden Hoven R, Jennings MP, Edwards JL, McEwan AG. A role for lactate dehydrogenases in the survival of *Neisseria gonorrhoeae* in human polymorphonuclear leukocytes and cervical epithelial cells. *J Infect Dis.* 2014;210(8):1311-8. <https://doi.org/10.1093/infdis/jiu230>.
11. Exley RM, Wu H, Shaw J, Schneider MC, Smith H, Jerse AE, Tang CM. Lactate acquisition promotes successful colonization of the murine genital tract by *Neisseria gonorrhoeae*. *Infect Immun.* 2007;75(3):1318-24. <https://doi.org/10.1128/IAI.01530-06>.
12. Gao L, Parsons NJ, Curry A, Cole JA, Smith H. Lactate causes changes in gonococci including increased lipopolysaccharide synthesis during short-term incubation in media containing glucose. *FEMS Microbiol Lett.* 1998;169(2):309-16. doi: <https://doi.org/10.1111/j.1574-6968.1998.tb13334.x>.
13. Exley RM, Shaw J, Mowe E, Sun YH, West NP, Williamson M, Botto M, Smith H, Tang CM. Available carbon source influences the resistance of *Neisseria meningitidis* against complement. *J Exp Med.* 2005 May 16;201(10):1637-45. <https://doi.org/10.1084/jem.20041548>.
14. Ayala JC, Shafer WM. Transcriptional regulation of a gonococcal gene encoding a virulence factor (L-lactate permease). *PLoS Pathog.* 2019;15(12):e1008233. <https://doi.org/10.1371/journal.ppat.1008233>.
15. Ratter JM, Rooijackers HMM, Hooiveld GJ, Hijmans AGM, de Galan BE, Tack CJ, et al. In vitro and in vivo Effects of Lactate on Metabolism and Cytokine Production of Human Primary PBMCs and Monocytes. *Front Immunol.* 2018;9:2564. <https://doi.org/10.3389/fimmu.2018.02564>.
16. Quillin S, Seifert H. *Neisseria gonorrhoeae* host adaptation and pathogenesis. *Nat Rev Microbiol* 2018;16: 226–40. <https://doi.org/10.1038/nrmicro.2017.169>
17. World Health Organization. Gonorrhoea (*Neisseria gonorrhoeae* infection). 18 July 2023. Downloaded from [https://www.who.int/news-room/fact-sheets/detail/gonorrhoea-\(neisseria-gonorrhoeae-infection\)](https://www.who.int/news-room/fact-sheets/detail/gonorrhoea-(neisseria-gonorrhoeae-infection))
18. Budkaew J, Chumworathayi B, Pientong C, Ekalaksananan T. Prevalence and factors associated with gonorrhea infection with respect to anatomic distributions among men who have sex with men. *PLoS One.* 2019 Apr 3;14(4):e0211682. doi: <https://doi.org/10.1371/journal.pone.0211682>.
19. Whelan J, Abbing-Karahagopian V, Serino L, Unemo M. Gonorrhoea: a systematic review of prevalence reporting globally. *BMC Infect Dis.* 2021;21(1):1152. <https://doi.org/10.1186/s12879-021-06381-4>.
20. Van Gerwen OT, Jani A, Long DM, Austin EL, Musgrove K, Muzny CA. Prevalence of Sexually Transmitted Infections and Human Immunodeficiency Virus in Transgender Persons: A Systematic Review. *Transgend Health.* 2020;5(2):90-103. <https://doi.org/10.1089/trgh.2019.0053>.
21. Chemaitelly H, Majed A, Abu-Hijleh F, Blondeel K, Matsaseng TC, Kiarie J, et al. Global epidemiology of *Neisseria gonorrhoeae* in infertile populations: systematic

- review, meta-analysis and metaregression. *Sex Transm Infect.* 2021;97(2):157-169. <https://doi.org/10.1136/sextrans-2020-054515>.
22. Tsevat DG, Wiesenfeld HC, Parks C, Peipert JF. Sexually transmitted diseases and infertility. *Am J Obstet Gynecol.* 2017; 216(1): 1-9. <https://doi.org/10.1016/j.ajog.2016.08.008>.
 23. Ngampasutadol J, Tran C, Gulati S, Blom AM, Jerse EA, Ram S, Rice PA. Species-specificity of *Neisseria gonorrhoeae* infection: do human complement regulators contribute? *Vaccine.* 2008;26 Suppl 8:162-6. <https://doi.org/10.1016/j.vaccine.2008.11.051>.
 24. Farage MA, Maibach HI. Morphology and physiological changes of genital skin and mucosa. *Curr Probl Dermatol.* 2011;40:9-19. <https://doi.org/10.1159/000321042>.
 25. Springer C, Salen P. Gonorrhea. [Updated 2023 Apr 17]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK558903/>
 26. Martín-Sánchez M, Fairley CK, Ong JJ, Maddaford K, Chen MY, Williamson DA, et al. Clinical presentation of asymptomatic and symptomatic women who tested positive for genital gonorrhoea at a sexual health service in Melbourne, Australia. *Epidemiol Infect.* 2020;148:e240. doi: <https://doi.org/10.1017/S0950268820002265>.
 27. Darville T. Pelvic Inflammatory Disease Due to *Neisseria gonorrhoeae* and *Chlamydia trachomatis*: Immune Evasion Mechanisms and Pathogenic Disease Pathways. *J Infect Dis.* 2021;224(12 Suppl 2):S39-S46. doi: <https://doi.org/10.1093/infdis/jiab031>.
 28. Reekie J, Donovan B, Guy R, Hocking JS, Kaldor JM, Mak D, Preen D, Ward J, Liu B; Chlamydia and Reproductive Health Outcome Investigators. Risk of Ectopic Pregnancy and Tubal Infertility Following Gonorrhea and Chlamydia Infections. *Clin Infect Dis.* 2019;69(9):1621-3. <https://doi.org/10.1093/cid/ciz145>.
 29. Teker B, de Vries H, Heijman T, van Dam A, van der Loeff MS, Jongen VW. Spontaneous clearance of asymptomatic anogenital and pharyngeal *Neisseria gonorrhoeae*: a secondary analysis from the NABOGO trial. *Sexually Transmitted Infections* 2023; 99: 219-25. <https://doi.org/10.1136/sextrans-2022-055488>
 30. Liang C, Zhao Z, Liu S, Zhang T, Zuo W. Single-cell transcriptome profiling of the vaginal epithelium reveals the heterogeneity of suprabasal cells. *Precis Clin Med.* 2023;6(1):pbad006. <https://doi.org/10.1093/pcmedi/pbad006>.
 31. Anderson DJ, Marathe J, Pudney J. The structure of the human vaginal stratum corneum and its role in immune defense. *Am J Reprod Immunol.* 2014;71(6):618-23. doi: <https://doi.org/10.1111/aji.12230>.
 32. Mendes AC, Ciccone M, Gazolla B, Bahia D. Epithelial Haven and Autophagy Breakout in Gonococci Infection. *Front Cell Dev Biol.* 2020;8:439. doi: <https://doi.org/10.3389/fcell.2020.00439>.
 33. Taktikos J, Lin YT, Stark H, Biais N, Zaburdaev V. Pili-Induced Clustering of *N. gonorrhoeae* Bacteria. *PLoS One.* 2015; 10(9): e0137661. <https://doi.org/10.1371/journal.pone.0137661>.
 34. Sunarti LS. Gonorrhoeae Must Compete With The Naturally Inhabitant of Microbial Community Right At The Outer Mucosal Surface To Authorize Infection. [Preprint]. 2023. Downloaded from https://www.researchgate.net/publication/374695474_Gonorrhoeae_Must_Compete_With_The_Naturally_Inhabitant_of_Microbial_Community_Right_At_The_Outer_Mucosal_Surface_To_Authorize_Infection

35. Pönisch W, Weber CA, Juckeland G, Biais N, Zaburdaev V. Multiscale modeling of bacterial colonies: how pili mediate the dynamics of single cells and cellular aggregates. *New J Phys.* 2017;19(1):015003. <https://doi.org/10.1088/1367-2630/aa5483>.
36. Kouzel N, Oldewurtel ER, Maier B. Gene Transfer Efficiency in Gonococcal Biofilms: Role of Biofilm Age, Architecture, and Pilin Antigenic Variation. *J Bacteriol.* 2015;197(14):2422-31. <https://doi.org/10.1128/JB.00171-15>.
37. Kurzyp K, Harrison OB. Bacterium of one thousand and one variants: genetic diversity of *Neisseria gonorrhoeae* pathogenicity. *Microb Genom.* 2023;9(6):mgen001040. <https://doi.org/10.1099/mgen.0.001040>.
38. Stevens JS, Criss AK. Pathogenesis of *Neisseria gonorrhoeae* in the female reproductive tract: neutrophilic host response, sustained infection, and clinical sequelae. *Curr Opin Hematol.* 2018;25(1):13-21. <https://doi.org/10.1097/MOH.0000000000000394>.
39. Lacroix G, Gouyer V, Gottrand F, Desseyn JL. The Cervicovaginal Mucus Barrier. *Int J Mol Sci.* 2020;21(21):8266. <https://doi.org/10.3390/ijms21218266>.
40. Shukair SA, Allen SA, Cianci GC, Stieh DJ, Anderson MR, Baig SM, Gioia CJ, Sponberg EJ, Kauffman SM, McRaven MD, Lakouagna HY, Hammond C, Kiser PF, Hope TJ. Human cervicovaginal mucus contains an activity that hinders HIV-1 movement. *Mucosal Immunol.* 2013;6(2):427-34. <https://doi.org/10.1038/mi.2012.87>.
41. Jennings LK, Krywko DM. Pelvic Inflammatory Disease. [Updated 2023 Mar 13]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK499959/>
42. Prendiville W, Sankaranarayanan R. Colposcopy and Treatment of Cervical Precancer. Lyon (FR): International Agency for Research on Cancer; 2017. (IARC Technical Report, No. 45.) Chapter 2., Anatomy of the uterine cervix and the transformation zone. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK568392/>
43. Dash S, Sethy PK, Behera SK. Cervical Transformation Zone Segmentation and Classification based on Improved Inception-ResNet-V2 Using Colposcopy Images. *Cancer Inform.* 2023; 22: 11769351231161477. <https://doi.org/10.1177/11769351231161477>.
44. Walker E, van Niekerk S, Hanning K, Kelton W, Hicks J. Mechanisms of host manipulation by *Neisseria gonorrhoeae*. *Front Microbiol.* 2023;14:1119834. <https://doi.org/10.3389/fmicb.2023.1119834>.
45. Zhu W, Cardenas-Alvarez MX, Tomberg J, Little MB, Duncan JA, Nicholas RA. Commensal *Neisseria* species share immune suppressive mechanisms with *Neisseria gonorrhoeae*. *PLoS One.* 2023; 18(4): e0284062. <https://doi.org/10.1371/journal.pone.0284062>.
46. Peric A, Weiss J, Vulliamoz N, Baud D, Stojanov M. Bacterial Colonization of the Female Upper Genital Tract. *Int J Mol Sci.* 2019;20(14):3405. <https://doi.org/10.3390/ijms20143405>.
47. Younes JA, Klappe K, Kok JW, Busscher HJ, Reid G, van der Mei HC. Vaginal epithelial cells regulate membrane adhesiveness to co-ordinate bacterial adhesion. *Cell Microbiol.* 2016;18(4):605-14. <https://doi.org/10.1111/cmi.12537>.
48. Ferrari M, Cirisano F, Morán MC. Mammalian Cell Behavior on Hydrophobic Substrates: Influence of Surface Properties. *Colloids and Interfaces.* 2019; 3(2):48. <https://doi.org/10.3390/colloids3020048>

49. Kreve S, Reis ACD. Bacterial adhesion to biomaterials: What regulates this attachment? A review. *Jpn Dent Sci Rev.* 2021;57:85-96. <https://doi.org/10.1016/j.jdsr.2021.05.003>.
50. Busscher H, Norde W, Sharma P, van der Mei H. (2010). Interfacial re-arrangement in initial microbial adhesion to surfaces. *Current Opinion in Colloid & Interface Science.* 2010, 15. 510–7. <https://doi.org/10.1016/j.cocis.2010.05.014>.
51. Bukrinsky MI, Mukhamedova N, Sviridov D. Lipid rafts and pathogens: the art of deception and exploitation. *J Lipid Res.* 2020;61(5):601-10. <https://doi.org/10.1194/jlr.TR119000391>.
52. Giardina PC, Williams R, Lubaroff D, Apicella MA. *Neisseria gonorrhoeae* induces focal polymerization of actin in primary human urethral epithelium. *Infect Immun.* 1998;66(7):3416-9. <https://doi.org/10.1128/IAI.66.7.3416-3419.1998>.
53. Karuppiah V, Collins R, Thistlethwaite A, Gao Y, Derrick J. Structure and assembly of an inner membrane platform for initiation of type IV pilus biogenesis. *Proceedings of the National Academy of Sciences of the United States of America.* 2013. <https://doi.org/10.1073/pnas.1312313110>.
54. Hu LI, Stohl EA, Seifert HS. The *Neisseria gonorrhoeae* type IV pilus promotes resistance to hydrogen peroxide- and LL-37-mediated killing by modulating the availability of intracellular, labile iron. *PLoS Pathog.* 2022;18(6):e1010561. <https://doi.org/10.1371/journal.ppat.1010561>.
55. Garnett JA, Atherton J. Structure Determination of Microtubules and Pili: Past, Present, and Future Directions. *Front Mol Biosci.* 2022; 8: 830304. <https://doi.org/10.3389/fmolb.2021.830304>.
56. Hu LI, Yin S, Ozer EA, Sewell L, Rehman S, Garnett JA, Seifert HS. Discovery of a New *Neisseria gonorrhoeae* Type IV Pilus Assembly Factor, TfpC. *mBio.* 2020;11(5):e02528-20. <https://doi.org/10.1128/mBio.02528-20>.
57. Ball LM, Criss AK. Constitutively Opa-expressing and Opa-deficient *neisseria gonorrhoeae* strains differentially stimulate and survive exposure to human neutrophils. *J Bacteriol.* 2013;195(13):2982-90. <https://doi.org/10.1128/JB.00171-13>.
58. Alcott AM, Werner LM, Baiocco CM, Belcher Dufresne M, Columbus L, Criss AK. Variable Expression of Opa Proteins by *Neisseria gonorrhoeae* Influences Bacterial Association and Phagocytic Killing by Human Neutrophils. *J Bacteriol.* 2022;204(4):e0003522. <https://doi.org/10.1128/jb.00035-22>.
59. Winther-Larsen HC, Hegge FT, Wolfgang M, Hayes SF, van Putten JP, Koomey M. *Neisseria gonorrhoeae* PilV, a type IV pilus-associated protein essential to human epithelial cell adherence. *Proc Natl Acad Sci U S A.*;98(26):15276-81. <https://doi.org/10.1073/pnas.261574998>.
60. Kolappan S, Coureuil M, Yu X, Nassif X, Egelman EH, Craig L. Structure of the *Neisseria meningitidis* Type IV pilus. *Nat Commun.* 2016;7:13015. <https://doi.org/10.1038/ncomms13015>
61. Kupsch EM, Knepper B, Kuroki T, Heuer I, Meyer TF. Variable opacity (Opa) outer membrane proteins account for the cell tropisms displayed by *Neisseria gonorrhoeae* for human leukocytes and epithelial cells. *EMBO J.* 1993;12(2):641-50. <https://doi.org/10.1002/j.1460-2075.1993.tb05697.x>.
62. Hung MC, Christodoulides M. The biology of *Neisseria* adhesins. *Biology (Basel).* 2013;2(3):1054-109. <https://doi.org/10.3390/biology2031054>.
63. Song W, Ma L, Chen R, Stein DC. Role of lipooligosaccharide in Opa-independent invasion of *Neisseria gonorrhoeae* into human epithelial cells. *J Exp Med.* 2000;191(6):949-60. <https://doi.org/10.1084/jem.191.6.949>.

64. van Vliet SJ, Steeghs L, Bruijns SC, Vaezirad MM, Snijders Blok C, Arenas Busto JA, Deken M, van Putten JP, van Kooyk Y. Variation of *Neisseria gonorrhoeae* lipooligosaccharide directs dendritic cell-induced T helper responses. *PLoS Pathog.* 2009;5(10):e1000625. <https://doi.org/10.1371/journal.ppat.1000625>.
65. Ram S, Shaughnessy J, de Oliveira RB, Lewis LA, Gulati S, Rice PA. Gonococcal lipooligosaccharide sialylation: virulence factor and target for novel immunotherapeutics. *Pathog Dis.* 2017;75(4):ftx049. <https://doi.org/10.1093/femspd/ftx049>.
66. Duncan JA, Brown LB, Leone PA. CHAPTER 25 - Gonococcal and Other Neisserial Infections in Guerrant RL, Walker DH, Weller PF. *Tropical Infectious Diseases: Principles, Pathogens and Practice (Third Edition)*, W.B. Saunders, 2011, Pages 184-190, <https://doi.org/10.1016/B978-0-7020-3935-5.00025-2>.
67. Chen A, Seifert HS. Structure-function studies of the *Neisseria gonorrhoeae* major outer membrane porin. *Infect Immun.* 2013;81(12):4383-91. <https://doi.org/10.1128/IAI.00367-13>.
68. Zeth K, Kozjak-Pavlovic V, Faulstich M, Fraunholz M, Hurwitz R, Kepp O, Rudel T. Structure and function of the PorB porin from disseminating *Neisseria gonorrhoeae*. *Biochem J.* 2013;449(3):631-42. <https://doi.org/10.1042/BJ20121025>. Erratum in: *Biochem J.* 2013;454(2):359.