

Original Research Article

**Age-dependent Radiological Risk Assessment of Radon (^{222}Rn) in Samples of
Commercial Bottled Water from Benin City, Nigeria**

ABSTRACT

Radon inhalation as well as ingestion through the use of water has a high potential of causing serious harm to sensitive cells and organs of the body when absorbed into the bloodstream. Therefore, this study aimed to assess the radiological health damage caused by radon in drinking bottled water available in Benin City, Edo State, Nigeria. Radon concentration was measured using a RAD7 electronic radon detector. The mean activity concentration radon is 137.18 ± 0.25 mBq/L. This mean value is slightly higher than 0.1 Bq/L recommended by the Standard Organisation of Nigeria but lower than the maximum contaminant level (MCL) of 11.1 Bq/L set by the United States Environmental Protection Agency and 100 Bq/L set by the World Health Organisation (WHO) for consumption of radon in drinking water. The mean total annual effective dose is 1.07 ± 1.91 ($\mu\text{Sv}/\text{y}$) for infants, 0.61 ± 1.09 ($\mu\text{Sv}/\text{y}$) for children, and 0.70 ± 1.24 ($\mu\text{Sv}/\text{y}$) for adults. The computed annual effective dose to the public by inhalation and ingestion radon through the use of bottled water in the study areas are lower than the 0.1 mSv/y limit recommended by WHO. The estimated average for the age group excess lifetime cancer risk are 3.21 ± 5.71 , 1.836 ± 3.26 , 2.086 ± 3.71 ($\times 10^{-6}$) for infants, children and adults respectively. These values are below the world average permissible levels. Hence, consumption of the water examined poses no serious health risk to consumers.

Keywords: Bottled water, radon (^{222}Rn), Benin City, cancer risk, effective dose, RAD7 detector

INTRODUCTION

Water is incredibly an important aspect of lives (Jain and Singh 2019). Water makes up about 60% of body weight in men and 55% in women and for infants it is about 70%–80% (Jain and Singh 2019). Unrefined water includes a range of pollutants, including biological, chemical, and radioactive contaminants (Oni et al. 2021). Most people depend profoundly on groundwater sources as a result of population growth, the contamination of streams, and the inadequacy of the government's potable water supply, numerous individuals have resorted to consuming treated water, such as bottled water and sachet water. In Nigeria, sachet water and bottled water have become increasingly embraced as a substitute for the inadequate safe drinking water from the government's pipe-borne water system. Bottled drinking water is water derived from any source of potable water (well, bore well, groundwater) which may be subjected to different treatment processes such as **decantation**, filtration, aeration, reverse osmosis, **ozonation**, ultraviolet treatment, silver ionization, other similar methods (Jain and Singh 2019)

Radioactivity is a natural phenomenon and natural sources of radiation are features of the environment. Exposure of human tissues or organs to radiation can induce the death of cells on a scale that can be extensive enough to impair the function of the exposed tissue or organ (IAEA 2014). Radon is a radioactive, colourless, odourless, and tasteless gas that is formed through the breakdown of uranium in soil and rocks (Opoku-Ntim et al. 2019). Radon is acknowledged as a cancer-causing substance. It is the most dangerous radioisotope to human health of all the radioactive elements that contribute to natural background radiation (NRC 1999). The most abundant form of radon is Rn-222, with a half-life of 3.82 days. Rn-222 is a decay product of radium (Ra-226), and a member of the U-238 decay chain (the most common isotope of

uranium). It occurs at low levels in most rocks, soil, water body, plants, and animals (Kamba and Okunade 2016).

Many uses of water release radon into the indoor air because radon is easily released by agitation in water. This contributes to the total indoor airborne radon concentrations. Ingestion of radon in water is also thought to pose a direct health risk through irradiation of sensitive cells in the gastrointestinal tract and in other organs once it is absorbed into the bloodstream. Thus, radon in drinking water could potentially produce adverse health effects in addition to lung cancer (NRC 1999). Once radon has entered the blood, through either the stomach or the small intestine, it is distributed among the organs according to the blood flow to them and the relative solubility of radon in the organs and in blood. Radon dissolved in blood that enters the lung will equilibrate with air in the gas-exchange region and be removed from the body (NRC 1999).

Radon levels and radiological safety of bottled and sachet water have been investigated by several researchers in Nigeria and other countries of the world. The sachet water consumed in Damaturu, Yobe State, Nigeria, as reported by Abba and Umaru (2020), has some level of radiological implications due to radon contamination. Yong et al. (2020) measured the radon concentrations of 15 different brands of commercial bottled drinking water produced in China and assessed the impacts on human health from commercial bottled water. The relationship between different mineral content and radon concentration in drinking mineral-water was also explored. Despite bottled mineral water not posing a significant health risk to the public, study results showed radon levels were strongly correlated with its total dissolved solids and calcium concentration. Recently, with the RAD7 Radon monitoring system, Al Mahmud et al. (2023) measured radon levels in commercial bottled water and deep well water samples, finding results that were significantly lower than USEPA and WHO recommendations. Oni et al.

(2021) determined the radon concentrations present in seventeen brands of sachet and bottled water packaged in Ile-Ife, South West Nigeria. Kadhim et al. (2021) assessed the radon concentrations from samples of bottled water available in the Iraqi markets. A study conducted by Bayaya et al. (2020) measured the levels of activity in radon-222 in water samples collected from different sachet water companies in **Dutsin-Ma** local government area of **Katsina** State, Nigeria. According to the authors, water consumption does not pose significant health risks from a radiological standpoint. In spite of this, radiation levels in the environment need to be continuously monitored.

Thorough assessments and monitoring of drinking water quality, as well as concentrations of radon activity, are paramount to public health and safety. However, no study of this sort has been carried out in Edo State. Benin City which is the commercial hub of Edo State was chosen for this study. The study, therefore, aimed to determine the activity concentrations of radon (^{222}Rn) in bottled water in Benin City, Nigeria, and assess the health hazards associated with ingestion and inhalation of radon.

MATERIALS AND METHODS

Study area

Benin City is the capital of Edo State, Nigeria and is located in the south-south geopolitical zone of Nigeria. The city has an area of 1, 204 km² and a population of approximately 1.84 million people (2022 population projection). The area is bounded by longitudes 4°36'– 4°39' E and latitudes 8°27' – 8°30' N and area of about 500 square kilometres. Benin City is located within a tropical equatorial zone where there are two distinct seasons, one wet season from April to October and one dry season from November to March (Erah et al. 2002).

Sampling and preparation

Ten brands of bottled water produced and sold or vended in Benin City were examined. Three bottles (75 cl each) of each of the brands were purchased from diverse stores, markets, and street hawkers in Benin city. A 1.5 L bottle, previously rinsed and washed with distilled water was used for the collection of water samples. In the process of putting the water into the collection bottles, water bubbling was carefully prevented to stop dissolved radon from escaping. The experimental procedures used in this investigation were those published by Osahon *et al.* (2018). The prepared samples were sealed and taken to the Ladoké Akintola University's Radiation and Health Physics Laboratory, Department of Physics, for radon analysis.

Radon measurement using RAD7

RAD7 (Figure 1) is a radon detector that is well-calibrated, quick, and accurate. The electronic detector employs the alpha spectrometry method with a bubbling kit. Alpha spectrometry method using RAD7 has widely been used by (Abba and Umaru 2020; Al Mahmud *et al.* 2023; Kadhim *et al.* 2021; Oni *et al.* 2021; Osahon *et al.* 2018). It can precisely assess the radon levels in a water sample in 20 minutes. The time required is quite short when compared to the radon half-life of 3.8 days, and as a result, the RAD7 detector is a good choice for measuring radon in water (DURRIDGE 2020; Oni *et al.* 2014).



Figure 1: Setup of RAD-7 for radon measurement in water(DURRIDGE, 2020)

Having obtained the values for the radon activity concentrations of each of the natural radionuclides in the bottled water, a calculation of the yearly effective dosage of radon from ingestion ()based on equation (1) was made(UNSCEAR 2000; Oni *et al.* 2016; Yong *et al.* 2020; Kadhim *et al.* 2021; and Sukanya *et al.* 2021)

$$AED_{Ing} = C_{RnW} \times C_f \times A_{WI} \quad (1)$$

Where C_{RnW} is the activity concentration of ^{222}Rn in (Bq/L), A_{WI} is the yearly consumption of drinking water (L/y), C_f is the age-dependent ingested dose conversion factor for ^{222}Rn .

As per UNSCEAR (2000), the mean effective doses of radon due to inhalation (AED_{inh}) in drinking water samples were calculated (Yong *et al.* 2020; Opoku-Ntim *et al.* 2019):

$$AED_{Inh} (\mu\text{Sv/y}) = C_{RnW} \times R_{a.W} \times F \times O \times D_{CF} \quad (2)$$

where AED_{Inh} is the effective dose for inhalation, C_{RnW} is the radon concentration in water, $R_{a.W}$ is the ratio of radon in air to radon in tap water (1×10^{-4}), F is the equilibrium factor between radon and its decay products (0.4), O is the average indoor occupancy time per person (7000 h/y)

and D_{CF} is the Dose Conversion Factor for radon exposure. Infants, children, and adults consume approximately 260, 350, and 730 litres of water per year, respectively (Opoku-Ntim *et al.* 2019; Orosun *et al.* 2022) and the ingested dose conversion factor per age group are 2.3×10^{-8} , 5.9×10^{-9} , and 3.5×10^{-9} (Sv/Bq) respectively (UNSCEAR, 2000; Yong *et al.* 2020; Kadhim *et al.* 2021; Sukanya *et al.* 2021).

The total Annual Effective Dose () as a result of ingestion and inhalation of radon through the use of the bottled water in the study area was obtained using equation (3)

$$AED_{Tot} = AED_{Ing} + AED_{Inh} \quad (3)$$

The lifetime cancer risk associated with radon concentrations in drinking water was calculated using equation (4) (Kadhim *et al.* 2021; Dankawu *et al.* 2022).

$$ELCR = \text{Annual effective dose} \times \text{Average lifespan} \times \text{Risk coefficient} \quad (4)$$

where, the nominal risk coefficient for the whole population is based on the International Commission on Radiological Protection public (ICRP 2007). In Nigeria, the average life expectancy at birth is 54.5 years, according to the WHO (2015) report.

RESULTS AND DISCUSSION

Radon concentration

The radon activity concentrations, as presented in Table 1, are from a minimum value of 20.83 ± 0.18 mBq/L in BTW 2 to a maximum of 340.28 ± 0.33 mBq/L in BTW 1 with a mean of 137.18 ± 0.25 mBq/L. The radon concentration value of BTW 8 is slightly closer to the maximum

Table 1: Radon concentrations (mBq/L) in bottled water in Benin City.

Sample ID	Radon Concentrations (mBq/L)
BTW 1	340.28 ± 0.33

BTW 2	20.83 ± 0.18
BTW 3	21.10 ± 0.19
BTW 4	20.83 ± 0.19
BTW 5	20.94 ± 0.19
BTW 6	41.65 ± 0.20
BTW 7	41.65 ± 0.20
BTW 8	337.52 ± 0.34
BTW 9	232.01 ± 0.30
BTW 10	294.95 ± 0.32
Minimum	20.83 ± 0.18
Maximum	340.28 ± 0.33
MEAN	137.18 ± 0.25

concentration recorded. There is a slight increase in the mean radon activity concentrations in this study over the 0.1 Bq/L recommended by the Standard Organisation of Nigeria (SON 2015) for human consumption of ^{222}Rn concentrations in drinking water, but it is significantly below the maximum contaminant level (MCL) of 11.1 Bq/L recommended by the USEPA (USEPA 1999) and 100 Bq/L suggested by the WHO (WHO 2011).

Table 2: Comparison of range/mean of radon activity concentrations in other part of the world with this study.

Location	Water type	^{222}Rn Conc. (mBq/L)	References
Turkey	Bottled water	30.6–57.6 /43.1	Seid et al. 2020
Dhaka/Bangladesh	Bottle water	105–1300/590	Al Mahmud et al. 2023
Urumqi/China	Bottled water	13.3–300/21.6	Yong et al. 2020
Ile-Ife	Bottled water	0–9449.3/2442.8	Oni et al. 2021

Iraq	Bottle water	28.4–283/104.14	Kadhim et al. 2021
Damaturu/Yobe	Sachet water	4420–20210	Abba and Umaru 2020
Dutsin-Ma/Katsina	Sachet water	11690–13040/12037	Baraya et al. 2020
Benin City	Bottled water	20.83–340.28/137.18	This study

The mean radon concentrations obtained in this study are slightly higher than those obtained by Yong *et al.* (2020) in China, Seid *et al.* (2020) in Turkey, Kadhim *et al.* (2021) in Iraq and Baraya *et al.* (2020) in Katsina, Nigeria, as shown in Table 2. Osahon *et al.* (2018), Al Mahmud *et al.* (2023) and Oni *et al.* (2021) reported radon mean concentrations greater than the obtained value in this study. The explanation for this low amount of radon in bottled water can be linked to the processing method employed by the bottled water factories, because radon gas is easily emitted when groundwater is agitated. In addition, the geological formation, soil porosity, meteorological parameters of the study area can also be a contributory factor (Opoku-Ntim *et al.* 2019).

Annual Mean Effective Dose of Radon

The results of the analysis of bottled water sold in Benin City indicate that the AED have minimum, maximum and mean values of 0.11 ± 0.95 , 1.80 ± 1.80 and 0.73 ± 1.29 ($\mu\text{Sv/y}$) for infants; 0.04 ± 0.35 , 0.66 ± 0.66 and 0.27 ± 0.48 ($\mu\text{Sv/y}$) for children; 0.05 ± 0.46 , 0.87 ± 0.87 and 0.35 ± 0.62 ($\mu\text{Sv/y}$) for adults respectively. The inhalation for all ages has minimum, maximum and mean values of 0.05 ± 0.45 , 0.86 ± 0.86 and 0.35 ± 0.61 ($\mu\text{Sv/y}$). These findings are summarized in Table 3 and

Table 3: Annual effective dose of radon ($\mu\text{Sv/y}$) in bottled water in Benin City.

Sample ID	Ingestion ($\mu\text{Sv/y}$)			Inhalation ($\mu\text{Sv/y}$)	Total AED ($\mu\text{Sv/y}$)		
	Infant	Children	Adults		Infant	Children	Adults
BTW 1	1.80 ± 1.75	0.66 ± 0.64	0.87 ± 0.84	0.86 ± 0.83	2.66 ± 2.58	1.52 ± 1.47	1.73 ± 1.67
BTW 2	0.11 ± 0.95	0.04 ± 0.35	0.05 ± 0.46	0.05 ± 0.45	0.16 ± 1.41	0.09 ± 0.80	0.11 ± 0.91
BTW 3	0.11 ± 1.01	0.04 ± 0.37	0.05 ± 0.49	0.05 ± 0.48	0.16 ± 1.48	0.09 ± 0.85	0.11 ± 0.96
BTW 4	0.11 ± 1.01	0.04 ± 0.37	0.05 ± 0.49	0.05 ± 0.48	0.16 ± 1.48	0.09 ± 0.85	0.11 ± 0.96
BTW 5	0.11 ± 1.01	0.04 ± 0.37	0.05 ± 0.49	0.05 ± 0.48	0.16 ± 1.48	0.09 ± 0.85	0.11 ± 0.96
BTW 6	0.22 ± 1.06	0.08 ± 0.39	0.11 ± 0.51	0.11 ± 0.50	0.33 ± 1.56	0.19 ± 0.89	0.21 ± 1.02
BTW 7	0.22 ± 1.06	0.08 ± 0.39	0.11 ± 0.51	0.11 ± 0.50	0.33 ± 1.56	0.19 ± 0.89	0.21 ± 1.02
BTW 8	1.79 ± 1.80	0.66 ± 0.66	0.86 ± 0.87	0.85 ± 0.86	2.64 ± 2.66	1.51 ± 1.52	1.72 ± 1.73
BTW 9	1.23 ± 1.59	0.45 ± 0.58	0.59 ± 0.77	0.59 ± 0.76	1.81 ± 2.34	1.04 ± 1.34	1.18 ± 1.52
BTW 10	1.56 ± 1.69	0.57 ± 0.62	0.75 ± 0.82	0.74 ± 0.81	2.30 ± 2.50	1.32 ± 1.43	1.50 ± 1.62
Minimum	0.11 ± 0.95	0.04 ± 0.35	0.05 ± 0.46	0.05 ± 0.45	0.16 ± 1.41	0.09 ± 0.80	0.11 ± 0.91
Maximum	1.80 ± 1.80	0.66 ± 0.66	0.87 ± 0.87	0.86 ± 0.86	2.66 ± 2.66	1.52 ± 1.52	1.73 ± 1.73
MEAN	0.73 ± 1.29	0.27 ± 0.48	0.35 ± 0.62	0.35 ± 0.61	1.07 ± 1.91	0.61 ± 1.09	0.70 ± 1.24




Fig. 2: Total annual effective dose of radon ($\mu\text{Sv/y}$) of bottled water in Benin City, Edo State.

depicted in Figure 2. The mean total AED are 1.07 ± 1.91 , 0.61 ± 1.09 and 0.70 ± 1.24 ($\mu\text{Sv/y}$) for infants, children and adults respectively. The infants recorded the highest value contributing 45% of the mean total AED, whereas the children and adults contributed 29% and 26% respectively. All the sampled bottled waters are below 0.1 mSv/y (WHO 2009).

The infants age category recorded the highest radiation dose contributing 45% of the mean total AED. This implies that infants are the most susceptible to radiation exposure from radon intake. The probable cause can be connected to the high quantity of radon that they absorb per kilogram of body mass as a result of low organ mass and rapidly dividing cells (Sukanya *et al.* 2021), despite the lower volume of water intake compared to children and adults. Adults are the second most vulnerable group to radon exposure, probably due to increased water use. However, there is just a little variation between adult and children category dosages, which is consistent with (Sukanya *et al.* 2021). The computed annual effective dose to the general population by

inhalation and ingestion in the study areas are less than the 0.1 mSv/y level recommended by WHO (2009) and 1 mSv/y ICRP, (2000). Comparatively, this study's annual effective dose (Sv/y) of radon is consistent with the findings of Seid *et al.* (2020) and Yong *et al.* (2020) and lower than 0.023-0.148 mSv for adults and 0.046-0.295 mSv for children reported by Abba and Umaru (2020) and 25.56 ± 2.69 , 19.48 ± 2.05 and 42.56 ± 5.58 Sv/y for adults, children, and infants respectively reported by Osahon *et al.* (2018).

Table 4: Excess Lifetime Cancer Risk from consumption of bottled water in Benin City, Edo State.

Sample ID	ELCR ($\times 10^{-3}$)		
	Infant	Children	Adults
BTW 1	7.913 ± 7.73	4.526 ± 4.41	5.141 ± 5.01
BTW 2	0.492 ± 4.23	0.282 ± 2.40	0.321 ± 2.73
BTW 3	0.492 ± 4.44	0.282 ± 2.55	0.321 ± 2.88
BTW 4	0.492 ± 4.44	0.282 ± 2.55	0.321 ± 2.88
BTW 5	0.492 ± 4.44	0.282 ± 2.55	0.321 ± 2.88
BTW 6	0.983 ± 4.68	0.564 ± 2.67	0.638 ± 3.06
BTW 7	0.983 ± 4.68	0.564 ± 2.67	0.638 ± 3.06
BTW 8	7.913 ± 7.97	4.526 ± 4.56	5.141 ± 5.19
BTW 9	5.431 ± 7.01	3.105 ± 4.02	3.528 ± 4.56
BTW 10	6.906 ± 7.49	3.951 ± 4.29	4.487 ± 4.86
Minimum	0.492 ± 4.23	0.282 ± 2.40	0.321 ± 2.73
Maximum	7.913 ± 7.97	4.526 ± 4.56	5.141 ± 5.19
MEAN	3.21 ± 5.71	1.836 ± 3.26	2.086 ± 3.71

Evaluation of Excess Lifetime Cancer Risk

The results of the excess lifetime cancer risk presented in Table 4 indicate that the mean value is 3.21 ± 5.71 , 1.836 ± 3.26 , $2.086 \pm 3.71 (\times 10^{-6})$ for infants, children and adults respectively.

The infants recorded higher value than adults and children. Internal radon exposure is extremely dangerous and can result in respiratory disorders such as asthma (Avwiri et al. 2014). However, the result obtained in the study, for all age categories, were below the world average permissible level of 0.2×10^{-3} (UNSCEAR 2000). Hence, consumption of the water examined poses no health threat to anyone that takes it.

CONCLUSION

This study examined the radiological evaluation of ^{222}Rn in several random samples of bottled water available in Benin City. The results demonstrate that the mean radon concentration is somewhat greater than the 0.1 Bq/L suggested by the Standard Organization of Nigeria (SON 2015) but still relatively low when compared to the standard levels recommended by the USEPA (1999), UNSCEAR (2000), and WHO (2011). The computed annual effective dose to the general public via inhalation and ingestion in the study areas were lower than the 0.1 mSv/y limit recommended by (WHO 2009) and 1 mSv/y (ICRP 2000). The infants are mostly irradiated by consumption of these bottled waters. The age group evaluation of the lifetime cancer risk, were below the set limit of 0.2×10^{-3} (UNSCEAR 2000). By implication, consumption of the water examined poses no significant health hazard to consumers. Nevertheless, the water factories should improve on the purification process by employing a more effective radon reduction methods to ensure radon free water for consumption. Subsequently, the findings of this study might be used to establish a baseline for a radiological implication of radon in water in Edo State.

REFERENCES

- Abba HT and Umaru D 2020 Evaluation of radon concentration and annual effective dose in sachet drinking water in Damaturu, Yobe State, Nigeria. *Dutse J. Pure and Appl. Sci.*6(2): 85–94.
- Al Mahmud J, Siraz MMM., Alam MS, Das SC, Bradley DA, Khandaker MU, Tokonami S, Shelley A and Yeasmin S 2023 A study into the long-overlooked carcinogenic radon in bottled water and deep well water in Dhaka, Bangladesh. *Int. J. Environ. Analyt. Chem.* 1–13
- Amakom CM and Jibiri NN 2010 Chemical and radiological risk assessment of uranium in borehole and well waters in the Odeda area, Ogun state, Nigeria. *Int. J. Phys. Sci.*5(7): 1009–1014.
- Avwiri GO, Ononugbo CP and Nwokeoji IE 2014 Radiation hazard indices and excess lifetime cancer risk in soil, sediment and water around Mini-Okoro/Oginigba Creek, Port Harcourt, Rivers State, Nigeria. *Comprehen. J. of Environm. and Earth Sci.* 3(1): 38–50.
- Baraya TJ, Sani MH and Joshua J 2020 Measurement of radon-222 concentration levels in brands of sachet drinking-water produce in Dutsin-Ma Local Government Area (LGA) of Katsina State, Nigeria. *J. Asian Sci. Res.*10(1): 33–42.
- Dankawu UM, Shuaibu HY, Maharaz MN, Zangina T, Lariski FM, Ahmadu M, Zarma SS, Benedict JN, Uzair M, Adamu GD and Yakubu A 2022 Estimation of excess life cancer risk and annual effective dose for boreholes and well water in Dutse, Jigawa State Nigeria. *Dutse J. Pure and Appl. Sci.*7(4a): 209–218.
- DURRIDGE 2020 *RAD H₂O user manual. DURRIDGE Company Inc.*
- Erah P, Akujieze C and Oteze G 2002 The Quality of Groundwater in Benin City: A baseline study on inorganic chemicals and microbial contaminants of health importance in boreholes and open wells. *Trop.l J. Pharmac. Res.*1(2): 75.
- IAEA 2014 Radiation Protection and Safety of Radiation Sources : International Basic Safety Standards. *General Safety Requirements Part 3*, 1-4.
- International Commission on Radiological Protection 2000 Protection of the Public in situations of prolonged radiation exposure 2000. ICRP Publication 82. *Ann. ICRP*, 29(1-2).
- International Commission on Radiological Protection ICRP 2007 The 2007 Recommendations of the International Commission on Radiological Protection. ICRP Publication 103. *Ann ICRP* 37 (2-4).
- Jain B and Singh AK (2019). The world around bottled water. In AMH Alexandru Mihai Grumezescu (Ed.), *Bottled and Packaged Water* 39–61
- Kadhim NA., Mdekil ASI, Abbas HH and Abojassim AA 2021 Age-dependent health risk assessment for radon concentrations from drinking water available in the Iraqi markets. *Egypt. J. Chem.*64(4): 1889–1895.
- Kamba AM and Okunade IO 2016 Radon measurement in commercial borehole water from some selected areas of Kaduna metropolis using Liquid Scintillation Counter. *Int. J.*

*Mathem. and Phys. Sci. Res.*3(2): 71–81.

National Research Council 1999 Risk Assessment of Radon in Drinking Water. Washington, DC: The National Academies Press.

Oni EA, Adagunodo TA, Adegbite AA and Omeje M 2021 Determination of radon gas in bottled and sachet water in Ile-Ife , Nigeria. *IOP Conferen. Series: Earth and Environ. Sci.*655: 012092.

Oni EA, Oni OM, Oladapo OO, Olatunde ID and Adediwura FE 2016 Measurement of Radon Concentration in Drinking Water of Ado-Ekiti , Nigeria. *J. Academ. Ind. Res.*4(8): 190–192.

Oni OM, Oladapo OO, Amuda DB, Oni EA., Olive-adelodun AO, Adewale KY and Fasina MO 2014 Radon Concentration in Groundwater of Areas of High Background Radiation Level in Southwestern Nigeria. 64–67.

Opoku-Ntim I, Andam AB, Akiti TT, Flectcher JJ and Roca V 2019 Annual effective dose of radon in groundwater samples for different age groups in Obuasi and Offinso in the Ashanti region, Ghana. *Environ. Res. Comm.*1: 105002.

Orosun MM, Ajibola BT, Ehinlafa EO, Issah AK, Salawu BN, Ishaya SD, Ochommadu KK and Adewuyi AD 2022 Annual effective dose assessment of radon in drinking water from abandoned tin and cassiterite mining site in Oyun, Kwara State, Nigeria. *Pollution*, 8(1): 181–192.

Osahon OD, Popoola FA, Igumbor E and Achurefe ME 2018 Radon activity concentration in water samples around a tourmaline mining site in Budo-Are, Oyo State, Nigeria. *J. Sci. Eng. and Tech.* 65–70.

Seid AMA, Turhan Ş, Kurnaz A, Bakır TK and Hançerlioğulları A. 2020 Radon concentration of different brands of bottled natural mineral water commercially sold in Turkey and radiological risk assessment. *Int. J. Environ. Analy. Chem.* 1–13.

SON 2015 Nigerian Standard for Drinking Water Quality. Standard Organisation of Nigeria. *Nigeria Industrial Standard, NIS-554*, 20.

Sukanya S, Joseph S and Noble J 2021 Evaluation of radiation dose from radon ingestion and inhalation in groundwater of a small tropical river basin, Kerala, India. *Isotopes in Environ. Health Stud.*57(2): 204–215.

UNSCEAR 2000 Sources and effects of ionizing radiation Volume I New York (NY): United Nations Scientific Committee on the Effects of Atomic Radiation.

United State Environmental Protection Agency USEPA 1999 *Cancer Risk Coefficients for environmental exposure to radionuclides*. 13: 1–335.

WHO 2009 *Indoor Radon a Public Health Perspective*. 110.

World Health Organization WHO 2011 Guidelines for drinking water quality 1st Ed. World Health Organization Geneva ISBN-10: 9241541687, pp: 465.

World Health Organization WHO 2015 WHO statistical profile. Country Statistics and Global Health Estimates by WHO and UN Partners Global Health Observatory World Health Organization.

Yong J, Feng G, Liu Q, Tang C, Wu B, Hu Y, Cai C and Mao P 2020 Radon concentration measurement and effective dose assessment in different brands of commercial bottled water produced in China. *Water Sci. and Tech.: Water Supply*, 20(5): 1581–1591.

UNDER PEER REVIEW