

Growth Response of Wheat as Affected by Different Nutrient Sources and NPK levels Under Vertisols

ABSTRACT

Growth analysis such as crop growth rate, relative growth rate, leaf area index and net assimilation rate are the most important traits in prediction of yield. Keeping in this view a field experiment was conducted during winter (*Rabi*) 2019 growing season at the experimental field of JNKVV, Jabalpur to estimate growth response of wheat as affected by different nutrient sources and NPK levels under Vertisols. The treatment comprised of 3 sources of nutrient M₁- Inorganic sources (NPK fertilizers), M₂- organic sources (FYM, vermicompost, biofertilizers *Azotobacter* and PSB) and M₃- Integrated sources (50% Inorganic + 50% organic) as main treatments and 5 NPK levels S₁- control, S₂- 100% RDF, S₃- 150% RDF, S₄- 200% RDF, S₅- Based on soil test value (STV) for target yield of 6t ha⁻¹ as sub treatments were replicated thrice in a split plot design. The results showed that the among the different sources of nutrients maximum growth responses was found with inorganic sources of nutrient which was significant over organic sources of nutrient followed by integrated sources of nutrient. All the growth characters (LAI, NAR, RGR and NAR) were higher at high fertility (200% RDF) which was found significant over 100% RDF and STV based value and at par with 150% RDF. Thus, from the present investigation concluded that the adequate supply of plant nutrient in inorganic sources with 150% RDF levels helps in better growth of wheat which ultimately helps in higher yield.

Keywords: leaf area index, net assimilation rate, relative growth rate and relative growth rate.

1. INTRODUCTION

Progressive growth in the human population is a fact, and it will reach at least nine billion by 2050 [1]. The food gap can be covered by several actions, but the most important and effective are both the increase in yields of crops, and area of arable soils [2]. In the past, the first factor was responsible for 55%-60% increase in the food production [3]. Wheat (*Triticum aestivum* L.) is the most important cereal crop in world and is the staple food for humans [2]. Wheat is one of the most important crops in the world that can cover the food gap [4]. The yielding potential of this crop is high. But the world average yield of wheat is drastically lower and amounts to only 3.2 t/ha. Nutrient management has played an important role in increasing the productivity of this crop. Fertilizers have an axial role in enhancing the growth and production in developing countries especially after the introduction of high yielding and fertilizer responsive crop cultivars [5]. In commercial production, quantity's of NPK fertilizers is very important factor affecting plant growth and production, especially the balance among N, P and K in soil [6]. Application of optimum dose of nutrients with appropriate method is considered as a key to success in increasing yield of any crop [7]. Therefore, application of fertilizers might be a successful tool for improving the agrochemical conditions of the soil. It could induce simulative effect on plant growth and productivity, especially with applying

chemical fertilizer [6]. The 'Green Revolution' in India has increased the yields tremendously, however, it served as a mixed blessing, as on one hand ambitious use of agrochemicals boosted the food grain production and on the other hand, it destroyed the agricultural ecosystem [8]. The farmers use CF to increase crop production as they are more economical, affordable, easy to use and quick in response [9]. No doubt, the use of chemical fertilizers is the quickest way of boosting crop production, but their increasing prices, soil health deterioration, sustainability and pollution considerations in general have led to renewed interest in the use of organic manures [8]. Use of organics alone does not result in spectacular increase in crop yields due to their low nutrient status but the supplementary and complementary use of such sources is known to enhance the utilization efficiency of fertilizers [10]. So, there is a need to draw a mid-way between organic and inorganic extremities that may sustain crop yields without deteriorating soil fertility and/or productivity [9]. This can only be maintained at sustainable level by nutrients via integrated approach [11]. Combined application of organic and inorganic nutrient sources improved synergism and synchronization between nutrient release and plant recovery thus resulted in better crop growth and yield [12]. Growth is a vital function of plants and is an indication of a gradual increase in number and size of cells [13]. Growth analysis such as crop growth rate, relative growth rate and leaf area index are the most important traits in prediction of yield [14]. Growth analysis is a suitable method for plant response to different environmental conditions during plant life also observed significant positive interaction between fertilizer treatments and physiological stages of wheat growth [15, 16]. The allocation and partition of photosynthates in plants can be better understood by using the technique of growth analysis of plants throughout their entire cycle or even part of it. Growth analysis is a laborious technique, but relatively simple and easy to perform using a few pieces of equipment. It basically consists of determining the leaf area of plants and the dry mass of each organ of the entire plant, normally every week [17]. With the obtention of the leaf area and the dry mass of the different plant organs, the instantaneous or average physiological indicators of growth such as crop growth rate, relative growth rate, net assimilation rate, specific leaf area, leaf area ratio and leaf area duration can be determined [18]. Many researchers reported that the use of balanced fertilizers have a promising role in growth and development of crop plants which resulted in improved quality and quantity of the agricultural produce [15]. Therefore, this study was conducted to identify the most effective source of nutrient and NPK levels for wheat to enhance physiological growth and ultimately higher productivity.

2. MATERIAL AND METHODS

2.1 Experiment details

A field experiment was carried out in the research field of Department of Soil Science and Agricultural Chemistry Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur during the 2018-2019. The soil of the experimental site was Typic Haplustert, clay in texture neutral in reaction, non calcareous, medium in organic carbon content, medium in available nitrogen, phosphorus, and potassium and low in DTPA extractable Zn. The treatment comprised of 3 sources of nutrient M₁- chemical, M₂- organic and M₃- integrated (chemical and organic) as main treatments and 5 fertility levels S₁- control, S₂- 100% RDF, S₃- 150% RDF, S₄- 200% RDF, S₅- Based on soil test value for target yield of 6t ha⁻¹ as sub treatments were replicated thrice in a split plot design. The wheat crop (GW-366) sown on with spacing of 22.5 cm row to row. The observations were recorded from each plot at 21, 45, 65 and 90 DAS of wheat.

2.2 Growth analysis

For computing LAI, leaves from 5 plants were collected and cleaned with water and then wiped with tissue paper. Using a leaf area meter (Model LICOR 3000, USA), the area of

fresh green leaves was measured and expressed in cm²/plant. While placing the leaf on the roller utmost care was taken to avoid overlapping of the leaf. Plants from one-meter row length were cut near the ground surface for dry-matter estimation from each treatment, at 21, 45, 60 and 90 DAS. The whole plants along with leaf in 1 m² area were collected and shade dried for 7 days and then oven dried at 65±5°C for one week. The dry weight was recorded with the help of electronic balance and expressed in g/m². The recorded dry weight data at 21, 45, 60 and 90 DAS was used to calculate the mean leaf area index (LAI), crop growth rate (CGR), relative growth rate (RGR), net assimilation rate (NAR). The mean LAI, CGR, RGR and NAR worked out with the following formulas:

1. Leaf area index (LAI)

$$\text{Leaf area index (LAI)} = \frac{\text{Total leaf area}}{\text{Unit land area}}$$

2. Crop growth rate (CGR) g m² day

$$\text{CGR} = \frac{W_2 - W_1}{P(t_2 - t_1)}$$

Where, P = Ground area, W₁ = Dry weight of plant⁻¹ m² recorded at time t₁, W₂ = Dry weight of plant⁻¹ m² recorded at time t₂, t₁ and t₂ were the interval of time, respectively and it is expressed in g m² day

3. Relative growth rate (RGR) (g⁻¹ g⁻¹ day)

$$\text{RGR} = \frac{(\ln W_2 - \ln W_1)}{(t_2 - t_1)}$$

Where, ln = Natural log, W₁ = Dry weight of plant⁻¹ m² recorded at time t₁, W₂ = Dry weight of plant⁻¹ m² recorded at time t₂, t₁ and t₂ were the interval of time, respectively and is expressed as g⁻¹g⁻¹ day

4. Net Assimilation Rate (NAR) (g m⁻² day⁻¹)

$$\text{NAR} = \frac{(W_2 - W_1)(\text{Log } L_2 - \text{Log } L_1)}{(t_2 - t_1)(L_1 - L_2)}$$

Where L₁ and L₂ are total leaf are at time t₁ and t₂ respectively. W₁ and W₂ are total dry wt. time t₁ and t₂ respectively.

2.3 Statistical analysis:

Data were analysed using SPSS for analysis of variance and Fisher's LSD multiple range test was employed for the means comparisons.

3. RESULTS AND DISCUSSION

3.1 Leaf area index (LAI)

The perusal of the results in table 1 suggested that the highest LAI of 0.75, 0.88, 1.22 and 1.87, respectively observed in treatment where inorganic source of nutrient (M₁) was applied, it was significantly superior over integrated sources of nutrient (M₃) and sources of nutrient (M₂) at 21, 45, 60 and 90 DAS. Application of M₃ with the LAI of 0.67, 0.86, 0.95 and 1.67, respectively was also found significantly superior over organic sources of nutrient with the lowest LAI of 0.39, 0.47, 0.55 and 1.61 at each stage. The increase was 92.30, 87.23, 121.81 and 16.14 per cent & 71.79, 57.37, 94.54 and 3.72 per cent respectively, due to M₁ and M₃ over M₂ at all the respective stages.

Among the nutrient levels, LAI increased from 0.50, 0.65, 0.74 and 1.58 in S₁ (control) to 0.71, 0.96, 1.23 and 2.06 in S₄ (200% RDF) at 21, 45, 60 and 90 DAS, respectively. Application of S₄ (200% RDF) obtained significantly higher LAI over S₂ (100% RDF) and S₅ (STV based RDF) and it was on par with S₃ (150% RDF) at each growth stage. However, the application of S₃ (150% RDF) significantly increased LAI over S₂ (100% RDF)

and S₅ (STV based RDF) at each growth stages except S₃ at milking stage. Application of S₂(100% RDF) was also found significant over S₅ (STV based RDF) 21, 45 and 60 DAS. The magnitude of increase was 42.0, 57.37, 89.23 and 66.12 per cent, respectively & 38.0, 45.90, 70.76 and 37.09 per cent, respectively due to application of S₄ (200% RDF) and S₃ (150% RDF), respectively over control at all growth stages.

Interaction effect of different sources of nutrient and NPK levels were found to be significant overall growth stages. At 21 DAS, maximum value found in combination of M₁S₄ (0.92) which was found significant over all combinations but found at par with M₁S₃ (0.87) and M₃S₄ (0.87). All levels of different sources of nutrient were found significant over their control. At 45 DAS, highest LAI found with M₁S₄ (1.22) which was found significant over all treatments followed by M₁S₃ (1.06), M₃S₄ (1.09) and M₃S₃ (1.09) which were found at par among themselves. At 60 DAS, maximum values found in M₃S₄ (1.86) which was found significant over all combinations. However, the application of M₁S₃ (1.58) also found significant over all combinations. At 90 DAS, maximum values found under M₁S₄ (2.47) which was at par with M₁S₅ (2.16), M₃S₄ (1.98), M₁S₅ (1.90) and M₁S₃ (1.86) followed by remaining treatments.

The leaf area (LA) is a central factor for plant growth studies because it represents the surface that receives radiation, triggering the photosynthetic process on which the production of plant biomass depends and consequently, agricultural production [19, 20]. Increase leaf area in inorganic fertilizers and different NPK levels may be due to nitrogen application boosted the photosynthetic rate, leaf expansion and leaf persistency. As leaf area directly relates to the rate as well as duration of leaf expansion, so LAI finds to be sensitive to N availability to crop plants [21]. Soldati et al. [22] reported that an increase in dry matter accumulation leads to an increase in leaf area because proportion of dry matter allocated to leaves remain constant while an increase in leaf area leads to an increase in rate of dry matter accumulation because light interception is directly related to leaf area during this phase of development. N at highest rates boosted up the tissue formation with better plant growth which increases its concentration in leaves and results in higher LAI. As leaf area directly relates to the rate as well as duration of leaf expansion, so LAI finds to be sensitive to N availability to crop plants [21].

Crop growth rate (CGR)

Data has been presented in table 2 indicated that the treatment where inorganic source of nutrient (M₁) was applied registered the highest crop growth rate (CGR) of 1.35, 3.13 and 21.45 g⁻¹ m⁻² day⁻¹, respectively which was significantly better than integrated sources of nutrient (M₃) and organic sources of nutrient(M₂) at 21-45, 45-60 and 60-90 DAS. The CGR of 1.24, 3.07 and 20.52 g⁻¹ m⁻² day⁻¹ recorded in M₃ was also found significant superior over M₂ which registered the lowest CGR of 1.17, 2.74 and 20.01 g⁻¹ m⁻² day⁻¹, respectively at each stage. The response of increase was 15.38, 14.23 and 7.19 per cent respectively and 5.98, 12.04 and 2.54 per cent, respectively due to M₁ and M₃ over M₂ at each stage.

The crop growth rate (CGR) increased from 1.07, 2.38 and 19.43 g⁻¹ m⁻² day⁻¹, respectively in S₁ (control) to 1.38, 3.25 and 21.70 g⁻¹ m⁻² day⁻¹ in S₄ (200% RDF) at each stage.

However, the application of S₄ (200% RDF) significantly increased over S₂ (100% RDF) and S₅ (STV based RDF) at all growth stages and found at par with S₃ (150% RDF) which registered the CGR of 1.34, 3.17 and 21.03 g⁻¹ m⁻² day⁻¹. Effect of S₂ (100% RDF) and S₅ (STV based RDF) for CGR was significantly similar with each other at each stage. The increase was up to 28.97, 36.55 and 11.68 percent respectively and 25.32, 33.19 and 8.23 per cent, respectively due to S₄ (200% RDF) and S₃ (150% RDF) over control at each stage. The lowest crop growth rate (CGR) values were recorded during early vegetative growth stages but increased to maximum during the flowering stage which is like the reports of other workers [23, 24, 5]. The significant increase in CGR due to combined NPK fertilizer nutrition

might be owing to better availability of nutrients and effective conversion of macronutrients at the site of photosynthesis into pigments [25]. In fact, the combined function of NPK nutrients might have maximum photosynthate accumulation towards the leaf biomass, because in the initial stage, leaf is the more powerful sink than any other plant parts in most of the crops [26]. Kumar et [27] reported that increasing the fertilizer doses 150% RDF have brought increment in growth indices (plant height, tillers, DM, LAI, CGR, RGR, NAR, photosynthetic rate). Thus, the number of leaves plant⁻¹ justified the ultimate final expression of growth parameters of the growing plants.

Relative growth rate (RGR)

An appraisal of data presented in the table (3) revealed that the significantly maximum value of relative growth rate (RGR) of 0.0683, 0.0236 and 0.0178 g⁻¹ m⁻² day⁻¹ was recorded with the application of inorganic sources of nutrient (M₁) which proved its significant superiority over integrated sources of nutrient (M₃) and organic sources of nutrient (M₂) at CRI, tillering and jointing. The addition of M₃ also gave significantly higher value of RGR which was 0.0225, 0.0631 and 0.0169 g⁻¹ m⁻² day⁻¹, respectively over M₂ with the lowest RGR of 0.0603, 0.0214 and 0.0159 g⁻¹ m⁻² day⁻¹, respectively at each stage. Application of M₁ and M₃ accounted 13.26, 10.28, 11.94 per cent, respectively and 4.64, 5.14 and 6.28 per cent, respectively increase over M₂ at each stage.

The RGR increases from 0.0588, 0.0183 and 0.0153 g⁻¹ m⁻² day⁻¹ in S₁ (control) to 0.0673, 0.0247 and 0.0181 g⁻¹ m⁻² day⁻¹ in S₄ (200% RDF) at each stage. However, the application of S₃ (150% RDF) and S₄ (200% RDF) significantly increased RGR over S₂ (100% RDF) and S₅ (STV based RDF) at each growth stages. Highest RGR was obtained in S₄ (200% RDF), which was significantly similar with S₃ (150% RDF) with the RGR of 0.0668, 0.0242 and 0.0177 g⁻¹ m⁻² day⁻¹ respectively. Effect of S₂ (100% RDF) and S₅ (STV based RDF) was found at par with each other at each stage. The corresponding increase was 14.45, 34.97 and 13.83 per cent, respectively and 13.60, 32.24 and 11.32 per cent, respectively due to S₄ (200% RDF) and S₃ (150% RDF) at each stage.

Relative growth rate (RGR) expresses the dry weight increase in time interval in relation to the initial weight N fertilizers, which might be due to the high concentrations of nutrients in fairly fertilized plots causing production of more leaves which in turn contributes for the production of high dry matter which resulted in more RGR [28]. The phenomena of RGR tend to be low again during later stage and negative towards maturity considerably due the amount of CO₂ lost by respiration, which occurs uninterruptedly in all metabolically active tissues, was greater than that assimilated by photosynthesis which occurs only in cells with chlorophyll when exposed to light [29]. These results are in agreement with those obtained by [30, 31, 32]

Net Assimilation Rate (NAR)

Data presented in table 4 indicated that the among the different sources, inorganic sources of nutrient showed significant superiority with the NAR of 0.0176, 0.0141 and 0.0138 g⁻¹ m⁻² day⁻¹ over integrated sources of nutrient (M₃) and organic source of nutrients (M₂) at CRI, tillering and flowering. Application of M₃ registered comparatively higher NAR of 0.0168, 0.0132 and 0.0130 g⁻¹ m⁻² day⁻¹ over M₂, which have lowest NAR of 0.0158, 0.0129 and 0.0122 g⁻¹ m⁻² day⁻¹ at each stage. Increase in NAR of 11.39, 2.32 and 13.11 per cent respectively and 6.32, 9.30 and 6.55 respectively in response to M₁ and M₃ over M₂ at each stage.

The NAR increased from 0.0153, 0.0118 and 0.0113 g⁻¹ m⁻² day⁻¹ in S₁ (control) to 0.0178, 0.0147 and 0.0143 g⁻¹ m⁻² day⁻¹ in S₄ (200% RDF) at each stage. However, the application of S₃ (150% RDF) and S₄ (200% RDF) significantly increased NAR over S₂ (100% RDF) and S₅ at each growth stages. The treatment receiving S₄ (200% RDF) obtained the maximum NAR which was significantly similar with S₃ (150% RDF), with the resultant value of 0.0172, and 0.0143 and 0.0139 g⁻¹ m⁻² day⁻¹. The NAR of S₂ (100% RDF)

and S₅ were found at par with each other. The increase in NAR over control was 16.33, 24.57 and 26.54 per cent respectively and 12.41, 23.00 and 21.18 per cent, respectively in response to S₄ (200% RDF) and S₃ (150% RDF) at each stage.

Interaction effect was nonsignificant at 21-45 DAS but found significant at 45-60 DAS and 60-90 DAS. At 45-60 DAS, maximum NAR found with the application of M₁S₄ (0.0160g⁻¹ m⁻² day⁻¹) which was found at par with M₁S₃ (0.0157g⁻¹ m⁻² day⁻¹) and M₃S₄ (0.0146g⁻¹ m⁻² day⁻¹) followed by remaining treatments. At 60-90 DAS maximum values found under M₁S₄ (0.0158) which was found over all treatment but found at par with M₁S₃ (0.0158g⁻¹ m⁻² day⁻¹) and M₃S₄ (0.0144) followed by remaining treatments. NAR is a measure of the rate of dry matter accumulation per unit leaf area. An increase in NAR during the growing season is indicative of response of the photosynthetic apparatus to an increase in demand for assimilates to afford rapid growth of the grain fraction [24]. Portes reported that [29] NAR decrease with maturity because the loss of CO₂ by respiration exceeded the amount assimilated for photosynthesis. Elevated nitrogen supply can boost dry matter content through production of photo-assimilates via leaves which is the centre of plant growth during vegetative stage and later distribution of assimilates to the reproductive organs [24]. The low net assimilation rate might be due to restricted availability of essential nutrients and decreased photosynthetic efficiency [33].

4. CONCLUSION

A result obtained from present investigation indicates that application of different sources of nutrients and varied levels of NPK fertilizers significantly influences the growth and growth attributes of wheat. Adequate supply of plant nutrient in inorganic sources with 150% RDF levels helps in better photosynthesis and growth of wheat which helps in higher yield.

REFERENCES

1. Hunter MC, Smith RG, Schipanski ME, Atwood LW, Mortensen DA. Agriculture in 2050: Recalibrating targets for sustainable intensification. *Bioscience*. 2017;67(4):386-391.
2. Alexandre C, Mamadou B, Rebouh NY, Valentinovich VV, Polityko PM. Nitrogen Fertilizer Influence on Winter Wheat (*Triticum aestivum* L.) Grain Yield and Grain Quality Production under Non-Chernozem Soils of Central Russia: A Review. *Agrotechnology*. 2022; 11:288.
3. Ray DK, Ramankutty N, Mueller ND, West PC, Foley JA. Recent patterns of crop yield growth and stagnation. *Nat Commun*. 2012; 3(1):1-7.
4. Mandic V, Krnjaja V, Tomic Z, Bijelic Z, Simic A, Ruzic Muslic D, et al. Nitrogen fertilizer influence on wheat yield and use efficiency under different environmental conditions. *Chilean journal of agricultural research*. 2015; 75(1): 92-97.
5. Kumar N, Mankotia BS, Manuja S, Pareek B, Kumar P, Sharma R and Mandian IS. Effect of varying fertility levels on physiological aspects of rice cultivars in the Northwestern Himalayas. *Journal of Crop and Weed*, 2020; 16(3): 143-147.
6. Abdelaziz ME, Hannfy Ahmed AH, Bekhit RS, Pokluda R. Response of growth patterns in sweet pepper to different NPK levels. *Acta univ. agric. et silvic. Mendel. Brun*. 2008; 1: 241-244.
7. Pandey D, Bhatnagar A, Singh G and Samartha. Growth response of intercropped maize (*Zea mays* L.) and urdbean (*Vigna mungo* L.) under different planting patterns and nutrient management practices. *Pantnagar Journal of Research*. 2019; 17(2): 99-105

8. Verma RK, Shivay YS and Ghasal PS. Effect of different cropping systems and nutrient sources on growth, productivity and economics of direct seeded basmati rice (*Oryza sativa*), Indian Journal of Agricultural Sciences. 2017; 87 (10): 1377–83.
9. Faisal M, Khan I, Ashraf U, Shahzad T, Hussain S, Shahid M, Abid M, Ullah S. Effects of organic and inorganic manures on maize and their residual impact on soil physico-chemical properties. Journal of Soil Science and Plant Nutrition. 2017; 17 (1): 22-32.
10. Yadav RL. Factor productivity trends in a ricewheat cropping system under long-term use of chemical fertilizers. Exptl. Agric. 1998; 34 : 1-8.
11. Priya S, Kaushik MK, Sharma SK, Kumawat P. Impact of Integrated Nutrient Management on Growth and Productivity of Hybrid Maize (*Zea mays L.*). Annals of Biology. 2014; 30 (1) : 106-108.
12. Huang, S., Weijian, Z.W., Yu, X., Huang, Q. Effects of long-term fertilization on corn productivity and its sustainability in an Ultisol of southern China. Agri. Ecosyst. Environ. 2010; 138: 44–50.
13. Khan MA, Abid M, Hussain N, Imran T. Growth and analysis of wheat (*Triticum aestivum L.*) cultivars under saline conditions. International Journal of Agriculture and Biology. 2005; 7 (3): 508–510.
14. Tafes B, Alemayehu Y. Physiological growth indices of durum wheat (*Triticum turgidum L. Var. durum*) as affected by rates of blended and nitrogen fertilizers, American Journal of Life Sciences. 2020; 8(4): 52-59.
15. Tesar MB. Physiological basis of crop growth and development. American Society of Agronomy. Madison. Wisconsin. 1984: 291-321.
16. Mandal A, Patra AK, Singh D, Swarup A and Masto RE. Effect of long-term application of manure and fertilizer on biological and biochemical activities in soil during crop development stages. Bioresource Technology. 2007; 98: 3585-3592.
17. Portes TA, Carvalho SIC. Crescimento e alocação de fitomassa de cincogramíneasforrageirasem condições de Cerrado. Revista de Biologia Neotropical. 2009; 6 (2): 1-14.
18. Portes TA, Castro JLG. Análise de crescimento de plantas: um programacomputacional auxiliar. Revista Brasileira Fisiologia Vegetal. 1991; 3: 53-56.
19. Srinivasan V. Decreasing, not increasing, leaf area will raise crop yields under global atmospheric change. Global Change Biology. 2017; 23 (4) :1626–1635.
20. Reis EM. Relationship between soybean plant defoliation and Asian soybean rust severity. Summa Phytopathologica. 2019; 45(3): 252-254.
21. Wajid A, Ahmad A, Awais M, Habib-ur-Rahman M, Sammar A, Bashir U, Arshad MN, Sana Ullah, Irfan M, Gull U. Nitrogen Requirements of Promising Cotton Cultivars in Arid Climate of Multan. Sarhad Journal of Agriculture. 2017; 33(3): 397-405.
22. Soldati A, Stehli A, Stamp P. Temperature adaptation of tropical highland maize (*Zea mays L.*) during early growth and in controlled conditions. Crop Science. 1984; 24:28-32.
23. Awal MA, Khan MAH. Mulch Induced Eco-physiological growth and Yield of maize. Pakistan Journal of Biological Sciences. 20003; (1): 61-64.
24. Valero A, Juan D, Maturano M, Ramírez AA. Tarjuelo Martín-Benito JM, Ortega Álvarez JF. Growth and nitrogen use efficiency of irrigated maize in a semiarid region as affected by nitrogen fertilization. Spanish Journal of Agricultural Research. 2005; 3(1): 134-144.
25. Prajapati N and Rawat GS. Effect of tillage practices and fertility levels on growth analysis, yield of clusterbean and residual soil nutrients. International Journal of Chemical Studies. 2019; 7(5): 3255-3258.

26. Singh BV, Singh Rakesh. Effect of sources of nutrients on physiological growth parameters and yield of onion varieties. *Annals of Plant and Soil Research*. 2014; 16(4):362-366.
27. Kumar R, Singh A, Yadav RB, Kumar A, Kumar S, Shahi UP and Singh AP. Growth, development and yield response of rice (*Oryza sativa L.*) as influenced by efficient nitrogen management under subtropical climatic condition. *Journal of Pharmacognosy and Phytochemistry*. 2017; SP1: 791-797.
28. Nataraja TH, Halepyati A S, Pujari B T and Desai B K. Influence of phosphorus levels and micronutrients on physiological parameters of wheat (*Triticum durum Dcsf.*). *Karnataka Journal Agricultural Science*. 2006; 19 (3): 685-687.
29. Portes ATD, Araújo BRBD, Melo HCD. Growth analysis, photosynthate partition and nodulation in bean and soybean. *Ciência Rural*, 2022; 52 (10): 1-14.
30. Azarpour E, Moraditochae M, Bozorgi HR. Effect of nitrogen fertilizer management on growth analysis of rice cultivars. *International Journal of Biosciences*. 2014; 4:35-47.
31. Paul SK, Isam MS, Sarkar MAR, Das KR, Islam SMM. Impact of variety and levels of nitrogen on the growth performance of HYV transplant aman rice. *Progressive Agriculture*. 2016; 27:32-38.
32. Salem AKM, Elkhoby WM, Khalifa Abou, Ceesay M. Effect of nitrogen fertilizer and seedling age on inbred and hybrid rice varieties. *American-Eurasian Journal Agricultural and Environmental Science*. 2011; (5):640- 646.
33. Datta SC. *Plant Physiology*. Wiley Eastern Ltd. New Age Intl. Ltd. New Delhi, India. 1994.

Table 1. Effect of different sources and NPK levels on LAI at different growth stages

M/S	LAI															
	21 DAS				45 DAS				60 DAS				90 DAS			
	M1	M2	M3	Mean	M1	M2	M3	Mean	M1	M2	M3	Mean	M1	M2	M3	Mean
S1	0.60	0.37	0.54	0.50	0.68	0.42	0.74	0.61	0.74	0.45	0.77	0.65	1.35	1.35	1.03	1.24
S2	0.71	0.45	0.68	0.61	0.75	0.49	0.78	0.67	1.41	0.52	0.83	0.92	1.79	1.48	1.53	1.60
S3	0.87	0.46	0.73	0.69	1.06	0.54	1.09	0.89	1.58	0.55	1.20	1.11	1.86	1.58	1.66	1.70
S4	0.92	0.40	0.81	0.71	1.22	0.56	1.09	0.96	1.25	0.59	1.86	1.23	2.47	1.72	1.98	2.06
S5	0.65	0.27	0.56	0.49	0.70	0.33	0.59	0.54	1.10	0.64	0.70	0.81	1.90	1.90	2.16	1.99
Mean	0.75	0.39	0.67	0.60	0.88	0.47	0.86	0.74	1.22	0.55	1.07	0.95	1.87	1.61	1.67	1.72
SEm±		0.028				0.027				0.023				0.088		
CD(p=0.05)		0.110				0.106				0.089				NS		
SEm±		0.023				0.025				0.026				0.069		
CD(p=0.05)		0.066				0.073				0.075				0.202		
Int I		0.039				0.043				0.045				0.120		
		0.114				0.126				0.130				0.349		
Int II		0.066				0.066				0.060				0.206		
		0.193				0.194				0.176				0.601		

*M*₁- Inorganic sources (NPK fertilizers), *M*₂- organic sources (FYM, vermicompost, biofertilizers Azotobacter and PSB) and *M*₃- Integrated sources (50% Inorganic + 50% organic) as main treatments and 5 NPK levels *S*₁- control, *S*₂- 100% RDF, *S*₃- 150% RDF, *S*₄- 200% RDF, *S*₅- Based on soil test value (STV) for target yield of 6t ha⁻¹

Table 2- Effect of different sources and NPK levels on CGR at different growth stages

M/S	CGR ($\text{g}^{-1} \text{m}^{-2} \text{day}^{-1}$)											
	21-45 DAS				45- 60 DAS				60- 90 DAS			
	M ₁	M ₂	M ₃	Mean	M ₁	M ₂	M ₃	Mean	M ₁	M ₂	M ₃	Mean
S ₁	1.05	1.07	1.09	1.07	2.38	2.41	2.36	2.38	19.10	19.96	19.24	19.43
S ₂	1.35	1.16	1.22	1.25	3.18	2.67	3.17	3.01	21.66	19.70	20.55	20.64
S ₃	1.49	1.23	1.30	1.34	3.41	2.80	3.30	3.17	22.01	20.25	20.82	21.03
S ₄	1.54	1.24	1.36	1.38	3.45	2.90	3.40	3.25	22.76	20.61	21.71	21.70
S ₅	1.32	1.14	1.22	1.22	3.24	2.90	3.10	3.08	21.71	19.55	20.26	20.51
Mean	1.35	1.17	1.24	1.25	3.13	2.74	3.07	2.98	21.45	20.01	20.52	20.66
SEm±		0.035				0.058				0.253		
CD(p=0.05)		0.136				0.228				0.994		
SEm±		0.042				0.068				0.249		
CD(p=0.05)		0.124				0.199				0.726		
Int I		0.074				0.118				0.431		
		NS				NS				NS		
Int II		0.095				0.157				0.636		
		NS				NS				NS		

M₁- Inorganic sources (NPK fertilizers), M₂- organic sources (FYM, vermicompost, biofertilizers Azotobacter and PSB) and M₃- Integrated sources (50% Inorganic + 50% organic) as main treatments and 5 NPK levels S₁- control, S₂- 100% RDF, S₃- 150% RDF, S₄- 200% RDF, S₅- Based on soil test value (STV) for target yield of 6t ha⁻¹

Table 3- Effect of different sources and NPK levels on RGR at different growth stages

M/S	RGR ($\text{g}^{-1} \text{m}^{-2} \text{day}^{-1}$)											
	21-45 DAS				45- 60 DAS				60- 90 DAS			
	M ₁	M ₂	M ₃	Mean	M ₁	M ₂	M ₃	Mean	M ₁	M ₂	M ₃	Mean
S ₁	0.0594	0.0581	0.0590	0.0588	0.0183	0.0180	0.0187	0.0183	0.0153	0.0151	0.0154	0.0153
S ₂	0.0652	0.0606	0.0615	0.0624	0.0231	0.0218	0.0229	0.0226	0.0173	0.0146	0.0164	0.0161
S ₃	0.0731	0.0608	0.0665	0.0668	0.0257	0.0232	0.0237	0.0242	0.0191	0.0167	0.0173	0.0177
S ₄	0.0735	0.0614	0.0670	0.0673	0.0264	0.0230	0.0246	0.0247	0.0191	0.0167	0.0186	0.0181
S ₅	0.0701	0.0605	0.0616	0.0641	0.0245	0.0209	0.0227	0.0227	0.0180	0.0163	0.0169	0.0171
Mean	0.0683	0.0603	0.0631	0.0639	0.0236	0.0214	0.0225	0.0225	0.0178	0.0159	0.0169	0.0168
SEm±		0.0014				0.0004				0.0003		
CD(p=0.05)		0.0056				0.0016				0.0010		
SEm±		0.0011				0.0002				0.0003		
CD(p=0.05)		0.0031				0.0005				0.0008		
Int I		0.0019				0.0014				0.0005		
		NS				NS				NS		
Int II		0.0033				0.0025				0.0007		
		NS				NS				NS		

M₁- Inorganic sources (NPK fertilizers), M₂- organic sources (FYM, vermicompost, biofertilizers Azotobacter and PSB) and M₃- Integrated sources (50% Inorganic + 50% organic) as main treatments and 5 NPK levels S₁- control, S₂- 100% RDF, S₃- 150% RDF, S₄- 200% RDF, S₅- Based on soil test value (STV) for target yield of 6t ha⁻¹

Table 4- Effect of different sources and NPK levels on NAR at different growth stages

M/S	NAR ($\text{g}^{-1} \text{m}^{-2} \text{day}^{-1}$)											
	21-45 DAS				45- 60 DAS				60- 90 DAS			
	M ₁	M ₂	M ₃	Mean	M ₁	M ₂	M ₃	Mean	M ₁	M ₂	M ₃	Mean
S ₁	0.0151	0.0155	0.0154	0.0153	0.0116	0.0123	0.0114	0.0118	0.0111	0.0113	0.0115	0.0113
S ₂	0.0171	0.0155	0.0168	0.0165	0.0129	0.0130	0.0129	0.0129	0.0126	0.0122	0.0126	0.0125
S ₃	0.0189	0.0158	0.0171	0.0172	0.0157	0.0130	0.0142	0.0143	0.0152	0.0125	0.0139	0.0139
S ₄	0.0193	0.0162	0.0178	0.0178	0.0160	0.0133	0.0146	0.0147	0.0158	0.0126	0.0144	0.0143
S ₅	0.0175	0.0158	0.0170	0.0168	0.0142	0.0129	0.0129	0.0133	0.0141	0.0124	0.0124	0.0130
Mean	0.0176	0.0158	0.0168	0.0167	0.0141	0.0129	0.0132	0.0134	0.0138	0.0122	0.0130	0.0130
SEm±		0.0002				0.0002				0.0002		
CD(p=0.05)		0.0010				0.0009				0.0009		
SEm±		0.0003				0.0002				0.0002		
CD(p=0.05)		0.0008				0.0007				0.0007		
Int I		0.0005				0.0004				0.0004		
		NS				0.0012				0.0012		
Int II		0.0007				0.0006				0.0006		
		NS				0.0017				0.0016		

M₁- Inorganic sources (NPK fertilizers), M₂- organic sources (FYM, vermicompost, biofertilizers Azotobacter and PSB) and M₃- Integrated sources (50% Inorganic + 50% organic) as main treatments and 5 NPK levels S₁- control, S₂- 100% RDF, S₃- 150% RDF, S₄- 200% RDF, S₅- Based on soil test value (STV) for target yield of 6t ha⁻¹

UNDER PEER REVIEW