

Original Research Article

Impact of biotic and abiotic stress on survival of lac insects (*Kerria lacca* (Kerr) on pigeonpea (*Cajanus cajan* (L.) Millsp)

Abstract

Cajanus cajan is generally grown in rainfed ~~condition~~ conditions; and is also a good annual host plant of lac ~~insect~~ insects. *C. cajan* is widely reported to have biotic stress due to insect ~~pest-pests~~ on it. Lac insect is phloem sap feeder and hence impart a biotic stress. The present field study was conducted to evaluate ~~the~~ survival of lac insects on *C. cajan* by adjusting different levels of biotic and abiotic stress on the host plant. The biotic stress due to insect pests on *C. cajan* was minimised with ~~a~~ periodic spray of contact insecticides. The varying level of biotic stress ~~i.e., No, Low, Medium, and High level was maintained on C. cajan plants with lac insects on it, i.e., No, Low, Medium, and High, was maintained on C. cajan plants with lac insects.~~ The three abiotic stresses in this experiment were soil moisture stress. It was managed the irrigation per plant through drips, ~~it-which~~ was believed ~~that-to~~ creating different levels of moisture stress in soil that will impact the host plant. The abiotic stress was of three levels i.e., Low, Medium, and High. The result reveals that ~~the~~ survival percent of Lac insect from brood lac inoculation to the harvest of lac crop was highest 37.52% on *C. cajan* with one primary branch and its secondary branches with lac insect (L₁. Low biotic stress) and 32.13% (W₃. Low soil moisture stress). The study indicates that biotic and abiotic stress play a major role in the growth of the host plants and ~~the~~ survival of *K. lacca*.

Formatted: Font: Italic

~~Keyword~~ **Keywords** – Biotic, Abiotic, Survival, Lac insect, Pigeonpea.

Introduction

~~Pigeonpea~~ **Pigeon pea** (*Cajanus cajan* (L.) Millsp) is a significant deep-rooted pulse crop that is mostly grown in rainfed regions by small and marginal farmers in South Africa and India [1]. Interestingly, this group of farmers is undernourished and has a low socioeconomic status. It ~~has been shown to be~~ is an effective lac insect host plant. [2]. The lac insect, *Kerria lacca* (Kerr), is a scale insect that is a member of the superfamilies Lacciferidae, Coccoidea, order Hemiptera, and suborder Homoptera. The world's largest producer of lac is India; other

significant producers are Indonesia, Thailand, China, Burma, and Sri Lanka. [3]. Abiotic and biotic variables, including ~~as~~ weather (rain, heat, and temperature), soil conditions (water, pH, and nutrients), insect populations, disease incidence, and management techniques (cultivar, irrigation, fertilization, and rotation), all have an impact on crop development and output [4]. The process of producing lac requires growing *K. lacca* on its host plants. As a result, in addition to environmental stress, *K. lacca*'s sap feeding causes stress to the host plant. In addition, even the lac insect faces biotic and abiotic stress in a particular ~~ee~~ system/ecosystem, which has an impact on lac crop productivity. Lac production causes stress to the host plants, as seen by the 10.9–25.3% decrease in *Z. jujube* yield [5]. Lac insect mortality is a common observation in India and is caused by soil moisture stress and pest burden on the host, particularly during the summer [6–10], which have reported ~~that~~ the effect of abiotic stress on organism/organisms. The ongoing raising of Lac insects on their host trees puts the plants under stress, which causes the trees to weaken and produce fewer coppices in the following years. The biotic stressors in lac production are predators [11]. Three primary predators of *K. lacca* include *Pseudohypatopa pulverea* Meyr (Lepidoptera: Blastobesidae), *Eublemma amabilis* Moore (Lepidoptera: Noctuidae), *Chrysopa lacciperda* Kimmins, and *C. madestes* Banks. (Chrysopidae; Neuroptera) [12–17]. In this context, the present field study was conducted to evaluate survival of lac insects on *C. cajan* by adjusting different levels of biotic and abiotic stress on the host plant.

Materials and methods


The field trial under Ph.D. thesis programme was conducted at Jawaharlal Nehru Krishi Vishwa Vidyalyaya, Jabalpur, M.P during the year *kharif* – *Rabi* season in 2020–21. The field experimentina Factorial Randomized Completely Blocked Design (RCBD) with three replications comprising of two factors viz., settlement of lac insect on a varying number of branches and varied level of irrigation on pigeonpea–pigeon pea crop. The experiment consisted of twenty-one treatment combinations with seven level of lacinsects' settlement (L_1 to L_7) and three levels of irrigation (W_1 to W_3).

Treatment details:

Factor A: Biotic stress

Settlement of lac insect on

L_1 -Any one primary branch and its secondary branches with lac



L ₂ -Any two primary branches and its - <u>their</u> secondary branches with lac insects		biotic stress
L ₃ -Any three primary branches and its - <u>their</u> secondary branches with lac insects	}	Medium biotic stress
L ₄ - Any four primary branches and its - <u>their</u> secondary branches with lac insects		
L ₅ - Any five primary branches and its - <u>their</u> secondary branches with lac insects	}	High biotic stress
L ₆ – All the primary branches and its - <u>their</u> secondary branches with lac insects		
L ₇ – Control (Plant with no lac insects)		Nobiotic stress

Factor B: Abiotic stress

W₁ - @ 2-litres h⁻¹ (high soil moisture stress)

W₂ - @ 4-litres h⁻¹ (medium soil moisture stress)

W₃ - @ 8-litres h⁻¹ (low soil moisture stress)

Schedule of operations

~~Schedule~~ The schedule of field operations during ~~the the course of~~ experiment ~~were was~~

i) Nursery raising of *C. cajan*-Nursery of *C. cajan* was raised on the substrate (Kapu + FYM in equal ratio) filled polythene bag of size 18 x 16 cm, by sowing seeds treated with *Trichoderma viridae*, Rhizobium and phosphorus solubilizing bacteria (PSB). Perforated polythene bags with seedlings were irrigated at weekly intervals. The perforated wall of polythene bags drained excess irrigation water. Seedlings were sprayed Insecticides to avoid insect pest infestation. The ~~seedlings~~ seedling's growth tips were nipped at 15 days intervals till its transplantation.

ii) Layout of the main field -The experimental layout in the main field was planned in a plot size of 62 feet x 42 feet to accommodate 63 *C. cajan* plants. Plant to plant and row to row spacing was maintained at six feet while, it was ten feet between the replications. Transplantation of *C. cajan* seedlings ~~were~~ was done in the evening hours of 16.08.2020, in polypropylene bags (PPB) filled with forty-five kilograms of substrate[18].

iii) Poly propylene bag (PPB) -The PPB weighed 125 g and had a dimension of 93 cm x 61 cm when empty. On filling the PPB with 45 kg substrate filled PPB attained[19]a dimension of 30 cm height and 125 cm circumference. The substrate was filled in the PPB on the designated spot in the layout of the experiment, such that it is not disturbed in the future. Thirty days old seedlings were transplanted in the PPB.

iv) Substrate -*C. cajan* seedlings were transplanted in polypropylene bags (PPB). The substrate was a combination of well-rotted farmyard manure (FYM) and river bed basin soil (Kapu). The substrate consisted of 30 kg of soil and 15 kg of FYM. The soil and FYM were filled in the PPB in layers i.e., soil followed by FYM. A *tasala* was used to fill the substrate in the PPB. After each filling the PPB was vigorously shaken for compactness.

v) Irrigation - Each PPB with a *C. cajan* plant was irrigated using a drip irrigation system as per the treatment schedule. There was no irrigation from July to September 2020 owing to rain. The irrigation interval was at 7 days from October 2020 to May 2021.

vi) Brood lac inoculation -*Rangeeni* brood lac purchased from Adarsh Lac Samiti, Jamankhari village, Tehsil Barghat, district Seoni, M.P. on 30.10.2020, was segregated for quality and ~~predator free~~ predator-free brood before its inoculation on *C. cajan*. Each *C. cajan* in the PPB was inoculated with a 15 g brood lac stick with the help of a twine as per the treatments.

vii) Phunki removal -Removal of left-over brood lac twigs from *C. cajan* after complete emergence of lac nymphs from female cells is *Phunki* removal operation. It was carefully removed from *C. cajan* plant 21 days after BLI without damaging the freshly settled lac insect on the branches[1],[20].

viii) Marking of slot -Usually by 30 days after brood lac inoculation (BLI), majority nymphs of *K. lacca* leaves the brood lac cells to settle on the branches of host plant. After settlement the crawlers become sedentary by inserting ~~its their~~ stylets into the phloem tissues, to become sedentary. Thirty days after BLI, branches with good lac insect settlement were selected for marking of slot. A ~~slot of 1cm width and 2.5cm length~~ 1cm width and 2.5cm length slot was marked on the branch bearing good settlement of lac insects. Three slots S₁, S₂ and S₃ were made on a single branch each of 2.5cm². These slots were tagged with the help of woolen threads of different ~~colour~~ colours for different slots. Stretching a thread between the index fingers of both ~~the hands, the insect settlement adjacent to the boundaries of the slot was carefully removed to make the slot clearly differentiated~~ hands, the insect settlement adjacent to the boundaries of the slot was carefully removed to differentiate the slot from the rest of the lac settlement on the branch[18].

ix) Digital recording -Lac insect settlement within the slot was digitally photographed with the help of a Digital Single Lens Reflex (DSLR) camera fitted with 100mm micro lens by setting it in manual mode with ISO 400 and shutter speed of 4.5-6. Several pictures of the slot were taken for clarity. Finally, the best frame was selected[18].

x) **Digital counting** - The digital images from the DSLR camera were transferred to the Laptop by inserting memory card in the port. The images were opened in the Paint 3D programme of the MS-Windows 10. After enlarging the image on the Laptop screen, the brush tool on the tool bar of the Paint 3D programme was selected. Selection of the thickness point of calligraphy pen was selected from 1x to 18x with a contrast colour of the brush tool. This was followed by placing the cursor on the individual lac insect in the image within the slot displayed on the computer screen, on a left click of the mouse, a dot of the selected thickness and colour appears on the insect. The process was repeated till all the lac insects in the slot of the image had a dot on it. At the end all the dots were counted and recorded as live lachthis was followed by saving the image in a designated folder after renaming it for retrieval in future (Patent application no. (201921007852A)[18].

xi) **Frequency of lac insect count** - Counting of live lac insects within the slots was done on 30th, 60th, 90th, 120th, 150th and 180th day after BLI during the year 2020-21.

Result and discussion

Mean number of live lac insects per 2.5 cm² of branches (MNL) on *C. cajan* (host plant)

As mentioned in xi, The MNL were recorded on 30th (28.11.2020), 60th (28.12.2020), 90th (27.01.2021), 120th (26.02. 2021), 150th (27.03.2021) and 180th (26.04.2021) day after BLI.

On 30th day after BLI

(Biotic stress)

On 28.11.2020 i.e., 30 days after BLI, the settlement of lac insects was remarkably good in all the treatments. The MNL varied from 166.26 (L₃) to 209.33 (L₆). Among the two low biotic ~~stresses~~, The MNL was 186.30 (L₁) and 177.52 (L₂). Among in the medium biotic stress, it was 166.26 (L₃) and 183.63 (L₄), while that in high biotic stress levels were 184.30 (L₅) and 209.33 (L₆). The MNL was significantly higher in (L₆) over L₃ (166.26).

Abiotic stress

The MNL on *C. cajan* at different levels of irrigation (soil moisture stress) ~~though~~ varied from 154.73 (W₂ – medium abiotic stress), 156.90 (W₁ – high abiotic stress) to 162.94 (W₃- low abiotic stress), but had no significant difference among the treatments. The irrigation was initiated on 6.10.2020, as the rain ~~were~~ ~~was~~ till 25.09.2020. During this period maximum and minimum mean temperature was 29.6°C and 11.6°C respectively. Rainfall was just 6.6mm.

The MNL due to the interaction effects of settlement of lac insect on varying number of branches (biotic) and irrigation levels (abiotic) varied from 150.78 (L₃W₁) to 235.44 (L₆W₁). The MNL was significantly higher in L₂W₁ (197.33) L₁W₂ (209), L₁W₃ (191.56), L₄W₃

(207.44), L₅W₃ (200.22), L₆W₃ (223.89) and L₆W₁ (235.44) over L₃W₁ (150.78). However, the former seven interactions were at par with each other.

On 60th day after BLI

(Biotic stress)

In comparison to 30th day BLI, on 60th day a reduction in the MNL due to biotic stress 14.99 (L₁-low biotic stress to 26.65 percent (L₆- high biotic stress) was observed. On 28.12.2020 i.e., 60 days after BLI, the MNL was varied from 128.61 (L₃- medium biotic stress) to 158.37 (L₁- low biotic stress). The MNL was significantly higher in L₆ (153.54) and L₁ (158.37) over L₃ (128.61). However, the former two were at par with each other.

Abiotic stress

The MNL on *C. cajan* at different levels of irrigation though varied from 118 (W₂- medium abiotic stress) 123.84 (W₁- high abiotic stress) to 125.24 (W₃-low abiotic stress) but had no significant difference in the MNL among all the level of irrigation. Between 30 to 60 days after BLI the percent loss in the MNL due to abiotic stress varied from 21.07 (W₁) to 23.74 (W₃). Between 30 and 60 days after BLI, the amount of water per plant was 16 litre (W₁), 32 litres (W₂) and 64 litres (W₃). During this period maximum and minimum mean temperature was 26.2°C and 8.9°C respectively. The winter rain during this period was 2.3mm. The MNL due to the interaction effects of settlement of lac insect on varying number of branches (biotic) and irrigation levels (abiotic) varied from 116.22 (L₃W₁) to 182.33 (L₁W₃). The MNL was significantly higher in L₂W₁ (153.44), L₆W₁ (174.22), L₁W₂ (158.66), and L₁W₃ (182.33) over L₃W₁ (116.22). However, the MNL in all the former four interactions ~~were~~ was at par with each other. Between 30 to 60 days after BLI the percent loss in MNL varied from 4.81 (L₁W₃) to 36.55 (L₆W₃).

On 90th day after BLI

(Biotic stress)

On 27.01.2021 i.e., 90 days after BLI, the MNL was varied from 107.64 (L₂- low biotic stress) to 126.15 (L₁- low biotic stress). The MNL was significantly higher in L₅ (122), L₆ (120.68), and L₁ (126.15) over L₂ (107.64). The MNL of all the former three treatments were at par with each other. Between 60 to 90 days after BLI the percent loss in MNL varied from 13.43 (L₃) to 21.40 (L₆).

Abiotic stress

The MNL on *C. cajan* at different levels of irrigation though varied from 99.22 (W₁- high abiotic stress) 100.63 (W₂- medium abiotic stress) to 103.26 (W₃- low abiotic stress) but had

no significant difference in the MNL among all the levels of irrigation. Between 60 to 90 days after BLI the percent loss in MNL due to abiotic varied from 14.72 (W_2) to 19.88 (W_1). The total water supply per plant varied from 36 ~~litres~~ ~~liters~~ (W_1), 72 litres (W_2) to 144 litres (W_3) between 30 to 90 days after BLI. The MNL due to the interaction effects of settlement of lac insect on varying number of branches (biotic) and irrigation levels (abiotic) varied from 99.56 (L_3W_1) to 135.11 (L_1W_2). The MNL was significantly higher in L_2W_1 (121.11), L_6W_1 (127.32), L_1W_2 (135.11), L_4W_3 (127.22), L_5W_3 (130.44) and L_6W_3 (122.58) over L_3W_1 (99.56). The MNL of all the former six interactions were at par with each other. Between 60 to 90 days after BLI, the percent loss in MNL varied from 6.97 (L_3W_2) to 27.30 (L_1W_3). During this period maximum and minimum mean temperature was 25.3°C and 10.1°C respectively, where the rainfall was just 0.9mm. In the first three ~~observation~~ ~~observations~~, the weather was ~~favourable~~ ~~favorable~~ (November to January) and the lac insects were still ~~on~~ ~~in~~ ~~its~~ ~~their~~ immature stage, therefore both biotic and abiotic factors were almost the same. During 120 to 180 days after BLI (February to April) which had extreme weather with a temperature range of 26.7°C to 37.7 °C, the lac insects were in ~~the~~ adult stage drawing more phloem sap from the host plant adding biotic stress. Thus, both type of stress was more during this period.

On 120th day after BLI

(Biotic stress)

The MNL was varied from 85.96 (L_3 - medium biotic stress) to 106.91 (L_1 - low biotic stress). The MNL was significantly higher in L_5 (96.96), L_6 (93.85) and L_1 (106.91) over L_3 (85.96). The MNL of all the former three treatments were at par with each other. Between 90 to 120 days after BLI, the percent loss in MNL varied from 15.25 (L_1) to 23.56 (L_6). There was 10.09 to 13.55 percent more MNL loss as compared to 60 to 90 BLI due to increased biotic stress.

Abiotic stress

The MNL on *C. cajan* at different levels of irrigation varied from 77.38 (W_1 - high abiotic stress) 81.52 (W_2 - medium abiotic stress) to 82.66 (W_3 - low abiotic stress). The MNL was significantly higher in (W_3) over W_2 and W_1 . Between 90 to 120 days after BLI the percent loss in MNL varied from 18.99 (W_2) to 22.01 (W_1). There was 10.71 to 29 percent more loss in MNL as compared to 60-90 BLI due to increase abiotic stress temperature and wind speed 1.4 °C and 0.03 km/h respectively. The total water supply per plant varied from 52 litres (W_1), 104 litres (W_2) to 208 litres (W_3) between 30 to 120 days after BLI. During this period the maximum and minimum mean temperatures were 26.7°C and 9.3°C respectively. There was winter rain and it was 12.6mm. ~~Total~~ ~~The total~~ rainy days was two. The MNL due to the

interaction effects of settlement of lac insects on varying number of branches (biotic) and irrigation levels (abiotic) varied from 78.22 (L₃W₁) to 124.11 (L₁W₂). The MNL was significantly higher in L₁W₁ (90.78), L₂W₁ (95.78), L₅W₁ (92.89), L₆W₁ (98.78), L₁W₂ (124.11), L₂W₂ (91.74), L₃W₂ (97.11), L₅W₂ (90.56), L₁W₃ (105.85), L₃W₃ (82.56), L₄W₃ (103.78), L₅W₃ (107.44) and L₆W₃ (100.56) over L₃W₁ (78.22). However, all the former thirteen interactions were at par with each other. Between 90 to 120 days after BLI, the percent loss in MNL varied from 8.14 (L₁W₂) to 28.05 (L₄W₁).

On 150th day after BLI

(Biotic stress)

The MNL was varied from 61.11 (L₃- medium biotic stress) to 80.03 (L₁- low biotic stress). The MNL was significantly higher in L₄ (64.33), L₅ (67.93) and L₁ (80.03) over L₃ (61.11), however, the former three were at par with each other. Between 120 to 150 days after BLI the percent loss in MNL varied from 25.14 (L₁) to 33.19 (L₆). There was 40.87 to 64.85 percent more MNL loss as compared to 90 to 120 BLI due to increased biotic stress and male emergence.

Male emergence and mating take place between 120th and 150th day of BLI while the major Lac secretion phase is after mating of female Lac insect i.e., 150th day onwards in case of Baishakhi crop of *Rangeeni* Lac. Adult female lac insects play a major role in lac production. In comparison to male lac insects, female insects have longer life i.e., emergence from egg to the harvest of lac crop at maturity. Adult male insect on the contrary has a very short life span of 3 to 5 days, when it aggressively mates with its adult females.

Abiotic stress

The MNL on *C. cajan* at different levels of irrigation varied from 53.25 (W₁- high abiotic stress) 58.41 (W₂- medium abiotic stress) to 59.38 (W₃- low abiotic stress). The MNL was significantly higher in (W₃) and (W₂) over (W₁). However, the former two were at par with each other. Between 120 to 150 days after BLI, the percent loss in MNL varied from 28.16 (W₃) to 33.18 (W₁). There was 48.28 to 50.74 percent more MNL loss as compared to 90 to 120 BLI due to abiotic stress increased temperature and wind speed 6.8°C and 0.4 km/h respectively. The total water supplied per plant varied from 68 litres (W₁), 136 litres (W₂) to 272 litres (W₃) between 30 to 150 days after BLI. The maximum and minimum mean temperature during this period was 33.5°C and 14.3°C respectively. Rainfall was 6.2mm. The MNL due to the interaction effects of settlement of lac insects on varying number of branches (biotic) and irrigation levels (abiotic) varied from 51.22 (L₆W₂) to 90.78 (L₁W₂). The MNL was significantly higher in all the interactions except L₃W₁ (7.40) and L₆W₂ (7.19). However, rest of the treatments were at par with each other. Between 120 to 150 days after BLI, the percent loss in MNL varied from 21.38

(L₁W₃) to 37.69 (L₆W₂).

On 180th day after BLI

(Biotic stress)

The MNL was varied from 52.19 (L₃. medium biotic stress) to 69.89 (L₁. low biotic stress). The MNL was significantly higher in L₅ (58.85) and L₁ (69.89) over L₃ (52.19), however, the former two were at par with each other. Between 150 to 180 days after BLI the percent loss in MNL varied from 12.68 (L₁) to 17.50 (L₄). There was 89.65 to 98.26 percent more female adult was survive as compared to 120-150 BLI.

Abiotic stress

The MNL on *C. cajan* at different levels of irrigation varied from 43.62 (W₁. high abiotic stress), 49.65 (W₂. medium abiotic stress) to 52.35 (W₃. low abiotic stress). The latter (W₃) was significantly higher than (W₁) but was at par with (W₂). Between 150 to 180 days after BLI the percent loss in MNL varied from 11.84 (W₃) to 18.09 (W₁). The total water per plant varied from 84 litres (W₁), 168 litres (W₂) to 336 litres (W₃) between 30 to 180 days after BLI. During these period maximum and minimum mean temperature was 37.7°C and 17.4°C respectively. The rain during the period was 0.6mm. The MNL due to the interaction effects of settlement of lac insects on varying number of branches (biotic) and irrigation levels (abiotic) varied from 41.22 (L₆W₂) to 79.89 (L₁W₂). The MNL was significantly higher in all the interactions except and L₆W₂ (6.46). However, rest of the interactions were at par with each other. Between 150 to 180 days after BLI, the percent loss in MNL varied from 7.61 (L₁W₃) to 20.82 (L₆W₁).

Mean survival percent

The percent survival of Lac insects is reported here is in three phases of Lac insect growth after BLI. The duration between 30th and 120th day of BLI is usually larval growth and pupal period of the Baishakhi crop of *Rangeeni* Lac insects. The overall mean percent survival of lac insects from 30 days after BLI to 180 days after BLI was highest 37.52% in (L₁. low biotic stress) followed by 31.93% (L₅. high biotic stress), 31.39% (L₃. medium biotic stress), 29.88% (L₂. low biotic stress) and 28.90% (L₄. medium biotic stress), while it was lowest 25.19% in (L₆- high biotic stress). The percent survival in (L₁. low biotic stress) was 48.94 % more as compared to (L₆- high biotic stress).

The overall mean percent survival of lac insects at different levels of irrigation (soil moisture stress) varied from 27.80 percent (W₁- high abiotic stress), to 32.13 percent (W₃- low abiotic stress). The latter was closely followed by 32.09 percent (W₂. medium abiotic stress). The percent survival in (W₃) was 15.57 % more as compared to (W₁), while in(W₂) it was 15.43 %.

The percent survival in (W₃) was 0.12 % more as compared to (W₂) and it takes 168 litres more water as compared to (W₂). The total water per plant varied from 84 litres (W₁), 168 litres (W₂) to 336 litres (W₃) between 30 to 180 days after BLI.

Discussion

Mean live lac insects (MNL) per 2.5cm² on *C. cajan*

Survival and growth of insects especially phloem feeders depend on the quantity and quality of

Table 1: Mean number of live lac insect (MNL) count per 2.5. cm² on the branches after BLI

Treatments	30 th	60 th	90 th	120 th	150 th	180 th	Mean Survival (%)
Factor A (Biotic stress)							
L ₁	186.30 (13.62)	158.37 (12.58)	126.15 (11.24)	106.91 (10.34)	80.03 (8.95)	69.89 (8.35)	37.52
L ₂	177.52 (13.29)	136.77 (11.69)	107.64 (10.39)	88.65 (9.43)	63.00 (7.96)	53.04 (7.31)	29.88
L ₃	166.26 (12.89)	128.61 (11.34)	111.33 (10.56)	85.96 (9.29)	61.11 (7.84)	52.19 (7.25)	31.39
L ₄	183.63 (13.53)	140.37 (11.87)	119.44 (10.95)	91.30 (9.57)	64.33 (8.04)	53.07 (7.31)	28.90
L ₅	184.30 (13.55)	138.86 (11.80)	122.00 (11.06)	96.96 (9.86)	67.93 (8.26)	58.85 (7.69)	31.93
L ₆	209.33 (14.44)	153.54 (12.39)	120.68 (11.00)	93.85 (9.70)	62.70 (7.93)	52.74 (7.27)	25.19
L ₇	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	-
SEm(±)	0.31	0.32	0.19	0.11	0.05	0.05	
CD(5%)	0.88	0.91	0.56	0.31	0.15	0.15	
Factor B (Abiotic stress)							
W ₁	156.90 (11.67)	123.84 (10.40)	99.22 (9.34)	77.38 (8.26)	53.25 (6.88)	43.62 (6.24)	27.80
W ₂	154.73 (11.60)	118.00 (10.16)	100.63 (9.40)	81.52 (8.46)	58.41 (7.18)	49.65 (6.62)	32.09
W ₃	162.94 (11.89)	125.24 (10.45)	103.26 (9.51)	82.66 (8.52)	59.38 (7.24)	52.36 (6.81)	32.13
SEm(±)	0.20	0.21	0.13	0.07	0.03	0.03	
CD (5%)	0.58	0.59	0.36	0.20	0.10	0.10	
Interaction							
L ₁ W ₁	158.33 (12.60)	134.11 (11.60)	110.78 (10.55)	90.78 (9.55)	66.11 (8.16)	52.89 (7.30)	33.40
L ₂ W ₁	197.33 (14.05)	153.44 (12.41)	121.11 (11.02)	95.78 (9.51)	65.56 (8.13)	53.11 (7.32)	26.91
L ₃ W ₁	150.78 (12.29)	116.22 (10.80)	99.56 (10.00)	78.22 (8.87)	54.22 (7.40)	47.56 (6.93)	31.54
L ₄ W ₁	175.67	146.11	118.44	85.22	58.89	47.56	27.07

	(13.27)	(12.11)	(10.91)	(9.26)	(7.71)	(6.93)	
L ₅ W ₁	180.78 (13.41)	142.78 (11.97)	117.33 (10.85)	92.89 (9.66)	60.22 (7.79)	50.56 (7.14)	27.97
L ₆ W ₁	235.44 (15.35)	174.22 (13.22)	127.32 (11.30)	98.78 (9.96)	67.78 (8.26)	53.66 (7.36)	22.79
L ₇ W ₁	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	-
L ₁ W ₂	209.00 (14.42)	158.66 (12.62)	135.11 (11.64)	124.11 (11.16)	90.78 (9.55)	79.89 (8.97)	38.22
L ₂ W ₂	183.00 (13.50)	120.67 (11.01)	101.70 (10.11)	91.74 (9.60)	68.34 (8.30)	57.11 (7.59)	31.21
L ₃ W ₂	182.78 (13.53)	133.89 (11.59)	124.56 (11.18)	97.11 (9.88)	70.89 (8.45)	57.33 (7.60)	31.37
L ₄ W ₂	167.78 (12.97)	137.33 (11.74)	112.67 (10.64)	84.89 (9.24)	61.22 (7.86)	52.22 (7.26)	31.13
L ₅ W ₂	171.89 (13.08)	131.15 (11.46)	118.22 (10.90)	90.56 (9.54)	66.44 (8.18)	59.78 (7.76)	34.78
L ₆ W ₂	168.67 (13.00)	144.33 (12.03)	112.16 (10.61)	82.21 (9.09)	51.22 (7.19)	41.22 (6.46)	24.44
L ₇ W ₂	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	-
L ₁ W ₃	191.56 (13.85)	182.33 (13.52)	132.56 (11.53)	105.85 (10.31)	83.22 (9.15)	76.89 (8.79)	40.14
L ₂ W ₃	152.22 (12.34)	136.20 (11.66)	100.11 (10.03)	78.44 (8.88)	55.11 (7.45)	48.89 (7.02)	32.12
L ₃ W ₃	165.22 (12.84)	135.73 (11.63)	109.89 (10.51)	82.56 (9.11)	58.22 (7.66)	51.67 (7.22)	31.27
L ₄ W ₃	207.44 (14.35)	137.67 (11.75)	127.22 (11.30)	103.78 (10.21)	72.89 (8.57)	59.44 (7.74)	28.66
L ₅ W ₃	200.22 (14.16)	142.67 (11.96)	130.44 (11.44)	107.44 (10.39)	77.11 (8.81)	66.23 (8.17)	33.08
L ₆ W ₃	223.89 (14.98)	142.06 (11.93)	122.58 (11.09)	100.56 (10.05)	69.11 (8.34)	63.33 (7.98)	28.29
L ₇ W ₃	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	-
SEm(±)	0.54	0.25	0.14	0.11	0.09	0.09	
CD(5%)	1.53	0.72	0.40	0.32	0.25	0.26	

Figures in parenthesis are $\sqrt{X+0.5}$ value. Dash represents no brood lac inoculation.

sap access to it [21-22], as well as protection from its natural enemies [23- 24]. Survival of any insect on a crop or variety indicates its compatibility [25] and preference [26-27]. There was a loss in lac insects from the first count to the last count at the lac crop maturity i.e., there was a reduction in survival of lac insects. Decline-The decline in the MNL is reported earlier by [18], [28-35] and [36]. The mean percent Lac insect loss ranged from 7.62 to 16.83 percent [28], while according to [29] it ranged from 69.14 to 74.52 percent. [31] reported it from 73.95 to 81.63 percent, while [30], [32],[33] and [37] reported it to vary from 68.10 to 73.01, 79.42 to 80.37,

39.60 to 63.41 and 22.99 to 28.85 percent respectively. The survival percent of lac insects from BLI to the harvest of the lac crop was highest (37.52) in (L₁. low biotic stress) followed by (31.93) (L₅. high biotic stress), 31.39 (L₃. medium biotic stress), 29.88 (L₂. low biotic stress), 28.90 (L₄. medium biotic stress) and 25.19 (L₆. high biotic stress). 32.13 (W₃. low abiotic stress), 32.09 (W₂. medium abiotic stress), and 27.80 (W₁. high abiotic stress). The result reveals that less lac insects on branches survival were higher. That means plants with less biotic stress helped in survival of lac insects. It may be due to less competition for food. A plant with more branches free of lac insects produced enough food for the plant itself and the phloem feeder on it. This reveals that biotic stress on the host plant impacts the survival of lac insects. The survival percent of lac on *C. cajan* reported by earlier workers varied from 24.91 to 38.13 [36]. 48.00 to 61.54 [38] at harvest. 21 to 25 percent [18] from BLI to maturity of lac crop. In the present study, it was observed that the MNL in L₁ was significantly higher throughout the observation period. In *C. cajan* plants of the treatment L₁ refers to the presence of lac insects in only one primary branch and its secondary branches in the whole plant. It means the load of lac insects in the plant was comparatively very less, while the plant was producing enough food for itself and the phloem feeders on it. This means that the host plant could tolerate the pest on it and in the abundant food availability (phloem sap) the insect on it could grow with minimum mortality. In the initial three ~~observation-observations~~ i.e., at 30th, 60th and 90th day after BLI, there was no significant difference in MNL among the different ~~level-levels~~ of irrigation (abiotic stress). The soil moisture stress during the early three observations may not be much because of the weather factor (immediate day after rains) and less demand ~~of-for~~ phloem sap by the immature lac insects [20]. However, the MNL was significantly higher in the *C. cajan* plant with less abiotic stress (soil moisture stress) i.e., in plants with high irrigation level. This means the demand of the plant for water was sufficient to meet its growth and reproductive demand as well as the phloem sap drawn by the growing lac insects. In the plant with less water supply, there must have been a soil moisture stress, which may have affected the food supply (phloem sap) to the lac insects. Constraint in food availability may have affected the survival of lac insects in W₁.

The candidate manuscript does not have a robust scientific discussion. I suggest the authors incorporate the suggested paragraphs, in this way, it would improve the scientific quality of the manuscript.

This study holds significant scientific relevance. Firstly, it contributes to our understanding of ecological interactions in agricultural ecosystems. By investigating how both biotic (predators) and abiotic (like climate conditions) stressors affect lac insect survival, the study can offer

[insights into the complex dynamics of these systems\[39\]\(Bertorelli and Olivares, 2020\). Understanding the ecological factors influencing lac insect populations can lead to better pest management practices and ultimately improve pigeonpea-pigeon pea cultivation, an important food and cash crop in many regions\[40\]\(Rodriguez et al. 2023\). Moreover, it can shed light on potential ecosystem services provided by lac insects, such as natural pest control\[41, 42\]\(Olivares et al. 2022a; Campos, 2023\).](#)

[From a social perspective, this research carries substantial importance as well. Pigeonpea-Pigeon pea is a vital source of nutrition and income for numerous communities, especially in developing countries like those in Latin America. By investigating the factors that impact lac insect survival on pigeonpea, the study may contribute to the development of more sustainable and resilient agricultural practices\[43\]\(Olivares et al. 2021\). This, in turn, can lead to increased crop yields\[44\]\(Araya-Alman et al. 2020\), economic stability\[45\]\(Pitti et al. 2021\), and improved food security\[46\]\(Montenegro et al. 2021\). Furthermore, this research can be linked to broader discussions on sustainable agriculture and the preservation of biodiversity, as it highlights the importance of considering both biotic and abiotic factors in crop management strategies\[47\]\(Olivares et al. 2022b\).](#)

[When compared to studies of environmental factors on pests and diseases in agricultural areas of Latin America, the research on lac insects and pigeonpea-pigeon pea presents a unique ecological perspective. While many studies in the region tend to focus on the direct effects of environmental conditions on crops, this research expands the scope to include intricate insect-plant interactions. This holistic approach can provide a valuable complement to existing research, as it adds depth to our knowledge of the complex relationships within agricultural ecosystems\[48, 49, 50\]\(Guevara et al. 2012; 2012; Hernandez et al. 2018e\). Consequently, this study may serve as a model for how to address similar issues in Latin American agriculture, offering innovative solutions that consider both ecological and environmental factors\[51, 52, 53\]\(Hernandez and Olivares, 2019; Hernandez et al. 2020; Hernandez et al. 2018a\).](#)

Conclusion

The study indicates that biotic and abiotic stress play a major role in the growth of the host plants and survival of *K. lacca*. The result reveals that survival percent of Lac insect from brood lac inoculation to the harvest of lac crop was highest 37.52% on *C. cajan* with one primary branch and its secondary branches with lac insect (L₁. Low biotic stress) and 32.13% (W₃. Low soil moisture stress).

References

1. Patidar R, Vajpayee S, Kakade S, Thomas M, Tripathi N, Upadhyay A, Namdev BK, Kurmi A, Sharma HL, and Kulhare PS.2021. Simultaneous Production of Both Lac and Pulse from [Pigeonpea-Pigeon pea](#) [*Cajanus cajan* (L) Millsp.] for Doubling Farmers' Income Legume Research- An International Journal, Volume Issue:()
2. Thomas M. 2003. Lac to Lakhs, Reviving self-reliance, KVK Shahdol 1-20.
3. Chamberlin JC. 1923. A systematic monograph of the Tachardiidae or lac insects (Coccidae). Bull Ento Res London, 14 (2):147 -212
4. Thomas M, Engle Y, Rathore V, Khobragade D, Kunal B and Shrivastava A. 2012. Stress management in lac production. JNKVV Res J 46(3): 277-285
5. Barman JC, Rahman MM, Pramanik MEA, Alam MA, Uddin MZ. 2007. Effect of lac cultivation on the yield of jujube of different ber (*Zizyphus jujube*) varieties. J Subtrop Agri Res and Dev Gulpurkar, Bangladesh, 5(3): 286-290
6. Nigel W A. (1998). Climate change and water resources in Britain. Climatic Change 39: 83-110
7. Gaivoronskaya NF. 2005. Production planning and cost management in crop production sectors. Ekonomika Sel'skokhozyaistvennykhi Pererabatyvayushchikh Predpriyatii 8: 18-24
8. Singh R, Singh D, Rao V U M (2007) Effect of abiotic factors on mustard aphid on Indian brassica. Indian J Agric Res 41(1): 67- 70
9. Dhiman S, Reji G, Diganta G, Das NG, Baruah I, Rabha B, Talukdar P K, Singh L. 2009. Diversity, spatio temporal distribution and biting activity of mosquitoes in Tripura State, India. Entomon 34(4): 223-232
10. Shrivastava AK, Srivastava DC, Solomon S, Srivastava MK, Singh I. (2003). Physiological characters imparting resistance to biotic and abiotic stresses in sugarcane. Sugar Tech 5: (3): 105-120
11. Varshney RK. 1976. A check list of insect parasites associated with lac. Orient Insects 10: 55-78.
12. Glower PM. 1937. Lac cultivation of India. Monograph of Indian lac Institute Ranchi pp 147
13. Narayanan ES. 1962. Pest of lac in India. Indian Lac Research Institute, Ranchi, pp 90-133
14. Mehra BP. 1965. Biology of *Chrysopa madestes* Banks (Nuropteran), Chrysopidae. Ent Soc India, 27(4): 398 407

15. Thomas M (2004) Squirrel damage to Kusumi lac in Madhya Pradesh, India. *Vaniki-Sandesh* 28(1): 14-15.
16. Malhotra CP, Katiyar RN. 1975. Control of *Eublemma amabilis* Moore, serious predators of the lac insect *Kerria lacca* (Kerr.). Screening of insecticide for their safety to lac insect. *Indian J Ent* 37: 385 -396
17. Sharma KK, Kumari K, Kumar M. 2006. Role of lac culture in biodiversity conservation: issues at stake and conservation strategy. *Curr Sci* 91(7): 894-898
18. Vajpayee S, Patidar R, Kakade S, Thomas M, Tripathi N, Bhowmick AK and Gontia AS. 2019. Effect of Population Density of *Kerria lacca* Kerr. On its Growth and Survival. *Int. J. Curr. Microbiol. App. Sci.* 2019;8(12):912-924. ISSN: 2319-7706 Volume 8
19. Patil D, Kakade S, Patel SK, Anjana G, Khichi A, Raut V, Thomas M and Tripathi N. 2022. Turmeric production in polypropylene bags for higher profitability. *Octa J. Biosci.* Vol. 10 (1):1-5
20. Khichi A, Thomas M, Patil DB, Kakade S, Raut S, Saxena AK, Upadhyay A and Tripathi N. 2023. *Rangeeni* lac production on *Cajanus cajan* (L.) MILLSP. *Journal homepage:*<http://www.plantarchives.org> DOI URL <https://doi.org/10.51470/PLANTARCHIVES.2023.v23.no1.057>
21. Cook AG and Denno RF. 1994. Planthopper/Plant Interactions: Feeding Behavior, Plant Nutrition, Plant Defense, and Host Plant Specialization. In: Denno R.F., Perfect T.J. (eds) *Planthoppers*. Springer, Boston, MA. https://doi.org/10.1007/978-1-4615-2395-6_3
22. Kehr J. 2006. Phloem sap proteins: their identities and potential roles in the interaction between plants and phloemfeeding insects. *J. Exp. Bot.* 57(4): 767-774.
23. Janghel S, Thomas M, Thakur AS, Nema S. and Sharma HL. 2014. Study on Bio Efficacy of Insecticides in the predator management of *Katki* Lac crop. *Bioeng. Biosci.* 2(2): 15-22
24. Engla Y. 2011. Study on predator management on *Baishakhi* crop of *Rangeeni* on *Zizyphus mauritiana* in Janamkhari village, Seoni district, Madhya Pradesh. M.Sc. (Ag.)
25. Horikoshi M, Beaumont R, Day F. 2016. Genome-wide associations for birth weight and correlations with adult disease. *Nature*, 538: 248-252
26. McGuinness KA. 1987. Disturbance and organisms on boulders. *Oecol.* 71: 420-430.
27. Gogi MD, Arif JM, Asif M, Abdin Z, Bashir HM. and Arshad M. 2012. Impact of nutrient management schedules on infestation of *Bemisia tabaci* on and yield of non-BT-cotton *Gossypium hirsutum* under unsprayed conditions. *Pak. Entomol.* 34(1):87-92.
28. Khobragade D. 2010. Studies on the Incidence on major predators of *Kerria lacca* Kerr.

- and their management in Baishakhi lac crop in Anuppur district, Madhya Pradesh. M.Sc. (Ag.) Thesis submitted JNKVV, Jabalpur
29. Patel B. 2013. Comparative performance of *Kusmi* and *Rangeeni* lac on Ber. M.Sc. Thesis. JNKVV, Jabalpur, M.P.
 30. Janghel S. 2013. Study on comparative efficacy of insecticides in Katki crop for predator management on *Rangeeni* lac crop on *Zizyphus mauritiana* in Malara village, Seoni District. M.Sc. Thesis. JNKVV, Jabalpur, M.P.
 31. Bhalerao KR. 2013. Study on Predator surveillance and its management on *Rangeeni* lac in Barghat block district Seoni Madhya Pradesh. M.Sc. (Ag.) Thesis submitted JNKVV, Jabalpur.
 32. Namdev BK. 2014. Study on the performance of *Aghani* crop of *Kusmi* lac on nutrient managed *Zizyphus mauritiana* under heavy rainfall condition. M. Sc. (Ag.) Thesis submitted JNKVV, Jabalpur.
 33. Ghugal SG, Thomas M and Pachori R. 2015. Performance of Katki Lac on Nutrient Managed of *Butea monosperma* (Lam.) Taub. Trends in Biosciences 8(24), Print. Pp. 6873- 6877.
 34. Gurjar R. 2016. Study on the Effect of Foliar Application of Nitrogen and PGR on *Butea monosperma* on Katki Crop Production. M.Sc. Thesis. JNKVV, Jabalpur, M.P.
 35. Sahu S. 2016. Survival and Yield of *Rangeeni* Lac insect on *Butea monosperma* (Lam) treated with different Micronutrients and Humic acid. M.Sc. (Ag) Thesis, submitted, JNKVV, Jabalpur
 36. Shah TH and Thomas M. 2018. Survival of *kusmi* lac insect (*Kerria lacca* Kerr.) on nutrient managed *Zizyphus mauritiana*. Indian Journal of Entomology, 80(1): 56-63.
 37. Kumar S, Thomas M, Lal N, Virendra and Markam VK. 2017. Effect of nutrition in Palas (*Butea monosperma* Lam.) on the survivability of lac insect. The Pharma Innovation Journal. 6(8): 320-324.
 38. Lekha BM, Goutham S, Shankar AUR. Evaluation of Lateritic Soil Stabilized with Arecanut Coir for Low Volume Pavements, 2014. doi: <http://dx.doi.org/10.1016/j.trgeo.2014.09.001>.
 39. Bertorelli, M., B.O. Olivares. Population fluctuation of *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) in sorghum cultivation in Southern Anzoategui, Venezuela. Journal of Agriculture University of Puerto Rico., 2020. 104(1):1-16. <https://doi.org/10.46429/jaupr.v104i1.18283>
 40. Rodríguez-Yzquierdo, G.; Olivares, B.O.; Silva-Escobar, O.; González-Ulloa, A.; Soto-

Formatted: Font: Italic, English (United States)

- [Suarez, M.; Betancourt-Vásquez, M. Mapping of the Susceptibility of Colombian Musaceae Lands to a Deadly Disease: *Fusarium oxysporum* f. sp. cubense Tropical Race 4. Horticulturae 2023, 9, 757. <https://doi.org/10.3390/horticulturae9070757>](#)
41. [Olivares, B.O.; Vega, A.; Rueda Calderón, M.A.; Montenegro-Gracia, E.; Araya-Almán, M.; Marys, E. Prediction of Banana Production Using Epidemiological Parameters of Black Sigatoka: An Application with Random Forest. Sustainability 2022a, 14, 14123. <https://doi.org/10.3390/su142114123>](#)
42. [Campos, B. O. Banana Production in Venezuela: Novel Solutions to Productivity and Plant Health. Springer Nature. 2023. <https://doi.org/10.1007/978-3-031-34475-6>](#)
43. [Olivares B, Rey JC, Lobo D, Navas-Cortés JA, Gómez JA, Landa BB. Fusarium Wilt of Bananas: A Review of Agro-Environmental Factors in the Venezuelan Production System Affecting Its Development. Agronomy, 2021, 11\(5\):986. <https://doi.org/10.3390/agronomy11050986>](#)
44. [Araya-Alman, M., Olivares, B., Acevedo-Opazo, C. et al. Relationship Between Soil Properties and Banana Productivity in the Two Main Cultivation Areas in Venezuela. J Soil Sci Plant Nutr. 2020; 20 \(3\): 2512-2524. <https://doi.org/10.1007/s42729-020-00317-8>](#)
45. [Pitti, J; Olivares, B; Montenegro, E. The role of agriculture in the Changuinola District: a case of applied economics in Panama. Tropical and Subtropical Agroecosystems. 2021, 25 - 1, 1 - 11. <https://www.revista.ccba.uady.mx/ojs/index.php/TSA/article/view/3815>](#)
46. [Montenegro, E., Pitti-Rodríguez, J, Olivares-Campos, B. Identification of the main subsistence crops of Teribe: a case study based on multivariate techniques. Idesia \(Arica\), 202; 39\(3\), 83-94. <https://dx.doi.org/10.4067/S0718-34292021000300083>](#)
47. [Olivares, B.O.; Rey, J.C.; Perichi, G.; Lobo, D. Relationship of Microbial Activity with Soil Properties in Banana Plantations in Venezuela. Sustainability 2022b; 14, 13531. <https://doi.org/10.3390/su142013531>](#)
48. [Guevara, E. Olivares, B., Demey, J. The Use of Climate Biomarkers in Agricultural Production Systems, Anzoategui, Venezuela. Revista Multiciencias. 2012, 12 \(2\): 136-145. <https://n9.cl/ak22r>](#)
49. [Guevara, E., Olivares, B., Oliveros, Y. & López, L. Estimation of thermal comfort index as an indicator of heat stress in livestock production in the Guanipa plateau, Anzoategui, Venezuela. Revista Zootecnia Tropical. 2013, 31 \(3\): 209-223. <https://n9.cl/ovcu9>](#)
50. [Hernández, R; Olivares, B. Arias, A; Molina, JC., Pereira, Y. Agroclimatic zoning of corn crop for sustainable agricultural production in Carabobo, Venezuela. Revista](#)

Formatted: Font: Spanish (Spain, International Sort)

Field Code Changed

Universitaria de Geografía., 2018a. 27 (2): 139-159. <https://n9.cl/12m83>

51. Hernández, R., Olivares, B., Arias, A, Molina, JC., Pereira, Y. Eco-territorial adaptability of tomato crops for sustainable agricultural production in Carabobo, Venezuela. *Idesia.* 2020. 38(2):95-102. <http://dx.doi.org/10.4067/S071834292020000200095>
52. Hernández, R; Olivares, B., Arias, A; Molina, JC., Pereira, Y. Identification of potential agroclimatic zones for the production of onion (*Allium cepa* L.) in Carabobo, Venezuela. *Journal of the Selva Andina Biosphere.*, 2018c. 6 (2): 70-82. http://www.scielo.org.bo/pdf/jsab/v6n2/v6n2_a03.pdf
- 38-53. Hernández, R. Olivares, B. Ecoterritorial sectorization for the sustainable agricultural production of potato (*Solanum tuberosum* L.) in Carabobo, Venezuela. *Agricultural Science and Technology.* 2019. 20(2): 339-354. https://doi.org/10.21930/rcta.vol20_num2_art:1462

UNDER PEER REVIEW