

## Original Research Article

Utilization of wastes lemon grass (*Cymbopogon citratus*) substrates in the cultivation of oyster mushroom; Modern view and trends

### ABSTRACT

Lemon grass wastes are abundant in both rural and urban areas and when carelessly disposed off to the environment by dumping or burning, they lead to environmental pollution, and consequently to health hazards. Growing of edible mushroom on these wastes can contribute to decrease in environment pollution. The aim of this study was conducted to compare the wastes lemon grass (*cymbopogon citratus*) substrates in the cultivation of *Pleurotus sajor-caju* and *Pleurotus florida*. Six substrate, namely, Trt1 (oil extracted lemongrass), Trt2 (oil extracted lemon grass 40% + wheat straw 30% + paddy straw 30%), Trt3 (oil extracted lemongrass 70% + wheat straw 30%), Trt4 (oil extracted lemon grass 50% + wheat straw 50%), Trt5 (oil extracted lemongrass 70% + paddy straw 30%) and Trt6 (oil extracted lemongrass 50% + paddy straw 50%) in the cultivation of PSC and PF. The results indicated that different substrate formulas gave a significant difference in complete full spawn run and days to pinhead formation were faster in Trt2 (15.5 and 13.70 days) and (18.40 and 17.40 days respectively) while Trt1 recorded the longest days (23.64 and 22.37) and (27.00 and 25.64 respectively) in PF and PSC. Number of fruiting bodies, fresh and dry weights of the mushroom was significantly higher in Trt2 than in the other substrates. The least number of fruiting bodies, fresh and dry weights were recorded in Trt1 of PF and PSC. The treatment Trt2 showed all over best result produce 884.37g and 856.33g in PSC and PF. but as far as lemon grass (Trt1) alone and its combined effect all other treatments. Combination of lemon grass with wheat straw + paddy straw also showed significant produce while, using oil extracted lemongrass 100% does not show better results.

**Keywords:** Oyster mushrooms, Utilization; Substrate, Lignocellulosic waste

### Introduction

The mushroom (*Pleurotus sp.*) is a fruiting body of fungi belonging to sub-division Ascomycotina and Basidiomycotina. They depend on organic matter for their nutrition and

**Comment [M1]:** Please be more detailed and also restructure the aim of the study. E.g. The aim of this study was to investigate the effect / compare (state the substrates you are comparing). You only mentioned one substrate in your aim but in a plural form.

**Comment [M2]:** Where are the methods used in this study? Please revisit and outline the methods (is one of the key component of Abstract) used in this work. I agree with you that the Treatments and its keys will be part of the methods.

**Comment [M3]:** Please present the Results in a detailed way, revisit. At what level of significant difference? Write full meaning of PF and PSC in the abstract.

**Comment [M4]:** Conclusion. What is your conclusion based on your results. Limitation and suggestions for further studies.



oyster mushrooms. The study was conducted to compare the effects of wastes lemongrass on the growth, yield, and nutritional composition of two oyster mushrooms *Pleurotus sajor-caju* and *Pleurotus florida*. The final aim is to find the best substrate formulas for effective cultivation of oyster mushrooms. The main objective of this study was proved that lemongrass is suitable substrate for the cultivation of oyster mushroom (Das *et al.*, 2000). Therefore present work was carried out to evaluate the lemon grass alone and in combinations with different easily available agricultural wastes like wheat straw and paddy straw.

**Comment [M7]:** There are many aims and objectives. Try and be more detailed and concise.

### MATERIALS AND METHODS

The study was to evaluate the suitability of utilization lemon grass waste combination with other substrate, wheat straw and paddy straw waste management of new model for cultivation of *Pleurotus sajor-caju* and *Pleurotus florida* was conducted at CSIR- Indian institute of Integrative Medicine Jammu (Figure1). The trial was laid out in a completely randomized design with 5 replications for each substrate treatment. The substrates used for the cultivation of the oyster mushroom were;

**Comment [M8]:** Try to be consistent, each substrate replicated 5 times, but subsequently, it was stated 10 replication s for each substrate.




Fig 1. Flow of utilization of lemongrass waste management according to a new model at the CSIR-Indian Institute of Integrative Medicine in the cultivation of *Pleurotus sajor-caju* and *Pleurotus florida*

Table 1: Base substrates and their composition as used in this experiment as substrate material

Treatment	Composition by weight
Trt <sub>1</sub>	Oil extracted lemon grass 100%
Trt <sub>2</sub>	Oil extracted lemon grass 40% + Wheat straw 30%+ Paddy straw 30%
Trt <sub>3</sub>	Oil extracted lemon grass 70% + Wheat straw 30%
Trt <sub>4</sub>	Oil extracted lemon grass 50% + Wheat straw 50%
Trt <sub>5</sub>	Oil extracted lemon grass 70% + Paddy straw 30%
Trt <sub>6</sub>	Oil extracted lemon grass 50% +Paddystraw 50%

The oil extracted lemon grass was collected from the CSIR- Indian institute of Integrative Medicine in field station chatha Jammu. The oil extracted lemon grassleaves were then chopped into 2-3 cm pieces and wheat straw + paddy straw was then chopped into 1.5-2.5 cm pieces. The wheat straw and paddy strawwas collectedfrom SKUAST- Jammu.

**Substrate Preparation:** Easily available, cheap and used agro-waste such as lemongrass, wheat straw, paddy straw containing high lignolytic, cellulotic, hemicellulotic, lignocellulotic content were used as a substrate for the cultivation of *Pleurotus* species. The oil extracted lemon grass leaves were used as an additional substrate to support the wheat straw substrate as additional nutrient source in ration of 3:1 as used previously by Hussain *et al.*, 2000. Each substrate was chopped into 2 to 3cm pieces and separately soaked in a solution of carbendazim (75 ppm) and formalin 40 per cent (500ppm) for 16-18 hours. The substrates were then separately dipped in hot water for 30 minutes for sterilization. The partially dried substrate (60-65 per cent moisture) used singly as well as in combination (1:1, 2:1 and 3:1) was mixed with wheat straw and paddy straw in following proportions in Table 1. The substrates were taken out of hot water and squeezed lightly by hands to remove excess of water. After that drained straw was mixed with spawn and spawning was performed by using polypropylene bags of 15 ×20cm and 14 × 28cm. In this stage, substrate moisture contain was 65 to 70%. Two kg (on dry weight basis) of each substrate was used to fill up in each bag. Filling and spawning of substrates was done simultaneously. Filling of substrates in the polythene bags was done in layers. Multi layered

**Comment [M9]:** Bold and shift, separate from the subsections

spawning technique was adopted for spawning of substrates and 2 per cent spawn was used and then incubated at 22-25<sup>0</sup>C for spawn run (mycelium growth) inside the cropping room. The spawned bags were kept in a dark room for the completion of spawn run. The corners and lower centre at the base of polythene bags were cut with scissors and holes were made to avoid any accumulation of water in the bags. Each treatment was replicated ten times and the bags were arranged in completely randomized design. In the fruiting house, the bags were hung on horizontal poles; water was sprayed onto the bags to keep them moist, the floors were also wetted to help increase the humidity to not less than 75-85 %, and temperature was maintained in the range of 14–18°C. Harvesting was done by cutting the bigger mushrooms at the base to allow the base itself and smaller mushrooms to continue growing.

**Comment [M10]:** Please, revisit. Write the correct times, the treatment was replicated. As noted in comment 8

**Data Collection:** The data was collected and observations were made on the following parameters: number of days from spawning to complete spawn run was recorded, and after two weeks of spawning, the spawned bags were checked daily. Days taken from complete spawn run to pinhead formation, time taken from pin formation to pin maturation and the number of days from spawning to the first harvest were recorded in days. The total number of pins (TNP) formed during the first and four flushes were recorded based on the total number of pins formed. Total number of flushes and yield of each bag and total yield (g) of the mushroom were calculated after the completion of cropping period. The yield was measured in terms of the fresh weight (kg) of the oyster mushrooms. The effectiveness of a mushroom species and substrate was measured. The biological efficiency (Yield of mushroom per kg substrate on dry wt. basis) was calculated by the following formula Chang *et al.*, (1981)

**Comment [M11]:** I suggest the subsections methods should be bold.

$$\text{Biological efficiency, BE (\%)} = \frac{\text{Fresh weight of mushroom}}{\text{Dry weight of substrate}} \times 100.$$

**Data Analysis:** The economic analysis of the different substrate treatments was determined by calculating the benefit: cost ratio of each treatment. Statistical evaluations of the above-mentioned parameters were done by one-way ANOVA using Gestate statistical software package and the mean comparisons were done using Fisher's protected CD at  $p < 0.05$ .

## RESULTS AND DISCUSSION

*Days from Spawning to Complete-Spawn Run (S-CSR):* There was a significant difference ( $p < 0.05$ ) in the number of days taken from spawning to complete spawn run (Table 2 and Fig 2.). Among the two species PSC and PF based on their fast and vigorous mycelial growth rate were selected and consequently evaluated for the cultivation parameters of *Pluratus sp.* In case of *Pleurotus sajor-caju* substrate (Trt2) took the least number of days (14.66) to complete the spawn run followed by a combination of substrate (Trt6) and Trt4 with 15.33 and 16.10 days, respectively. However, maximum spawn run period of 23.04 days was observed in Trt1 substrate alone. In *Pleurotus florida* the minimum earliest days for spawn run (12.51 days) was observed with (Trt2) substrate followed by a combination of substrate Trt6 13.70 days, and Trt4 substrate 14.06 days. Maximum spawn run period of 22.37 days in (Trt2) oil extended lemon grass alone. This is comparable with other similar studies elsewhere. Biswas and Biswas (2015) reported the completion of spawn running on wheat straw waste to be 14 days, while, Lalithadevi *et al.* (2014) recorded between 16-25 days on paddy straw.

**Comment [M12]:** I suggest you present your results before discussion, for better understanding and to be more clear.

Table 2 The oil extracted lemon grass wastes substrate treatments on mushroom growth parameters investigated of *Pleurotus sajor-caju* and *Pleurotus florida*

**Comment [M13]:** The Tables number, should be properly labeled and separated from the Table title.

Substrate	S-CSR		S-PF		F-PF	
	Species		Species		Species	
	PSC	PF	PSC	PF	PSC	PF
Trt1	23.04	22.30	27.00	25.64	37.22	35.73
Trt2	14.66	12.51	18.40	17.40	27.24	25.70
Trt3	20.70	20.41	24.44	22.66	33.17	32.03
Trt4	16.10	14.06	20.33	19.66	29.99	28.18
Trt5	18.00	17.11	22.81	22.03	32.45	30.45
Trt6	15.33	13.70	19.66	18.66	28.46	26.66
CD	2.61	2.50	3.30	3.45	1.31	1.80
S Em ( $\pm$ )	0.82	0.52	1.03	1.08	0.41	0.56

\*Mean of five replications; \*PSC- *Pleurotus sajor-caju*, \*PF- *Pleurotus florida*, \*S-CSR- Spawning to Complete-Spawn Run, \*S-PF -Spawning to First Pinhead Formation, \*F-PF fruiting body formation

*Days from Spawning to First Pinhead Formation (S-PF)*: Data patterning time taken from spawning to first pinhead formation is shown in Table 2 and Figure 2. The data reveal that there was a significant difference ( $p < 0.05$ ) in the time taken from spawning to first pinhead formation among the different substrates. In case of *Pleurotus florida* all the substrates were significant to each other. The minimum (earliest) days for pinhead formation (17.40 days) were observed with *Pleurotus sajor-caju* on wheat straw (control) Trt1 substrate followed by Trt6 (18.66 days). However, maximum pinhead formation period of 25.64 days was observed in (Trt2) lemongrass substrate alone. The time required for the formation of pinheads is comparable with other similar studies. Ahmed (1998) reported pinhead formation of oyster mushroom cultivated in different substrates to be between 23 and 27 days from spawning, while Fan *et al.* (2000) reported it to be 20-23 days. Tan (1981) recorded 23-26 days for the appearance of pinheads. Patra and Pani (1995) recorded 20-24 days on paddy straw. In *Pleurotus florida* the minimum earliest days for pinhead formation (18.40 days) was observed with wheat straw followed by Trt6 19.66 days. Maximum pinhead formation period of 27.00 days (Trt2) oil extracted lemon grass alone.




Fig2. Oyster mushroom *Pleurotus florida* and *Pleurotus sajor-caju* showing different stages on the substrate: (A) Trt1 (Oil extracted lemon grass 100%), (B) Trt2. Oil extracted lemon grass 40% + Wheat straw 30% + Paddy straw 30%, (C) Trt3 Oil extracted lemon grass 70% + wheat straw 30% (D) Trt4 Oil extracted lemon grass 50% + wheat straw 50% (E) Trt5 Oil extracted lemon grass 70% + Paddy straw 30% (F) Trt6 Oil extracted lemon grass 50% + Paddy straw 50%) in spawn run and pinhead formation.

*Days for fruiting body formation (after pinhead formation) (F-PF)*: In table 2 and figure 2 indicates that the *Pleurotus sajor-caju* and *Pleurotus florida* were non-significant in regard to days taken for fruiting body formation (after pinhead formation) in case of different substrates.

Minimum number of days recorded for fruiting body formation by *Pleurotus florida* were (25.70 days) in substrate Trt2 followed by 26.66 days in case of substrate (Trt6). However, maximum days for fruiting body formation (35.73 days) were observed with Trt1 substrate with *Pleurotus florida*. In case of *Pleurotus sajor-caju*, minimum days for fruiting body formation (27.24 days) were recorded in Trt2 substrate followed by 28.46 days for substrate (Trt6) alone. Maximum days for fruiting body formation (37.22 days) were observed in (Trt1) substrate.

Number of fruit bodies of *Pleurotus sajor-caju* and *Pleurotus florida*: Table 3 indicates mean  $\pm$  SD for each flush and the overall mean of fruit bodies of *Pleurotus sajor-caju* and *Pleurotus florida*. Maturity was not significantly different ( $p > 0.05$ ) among the flush of each treatment, while among the treatments in case of *Pleurotus sajor-caju*, maximum average of fruit bodies (41.35) was recorded in Trt1 followed by 34.60 in Trt6, whereas minimum average of fruit bodies (20.87) were recorded in Trt2. In case of *Pleurotus florida*, maximum average of fruit bodies (39.99) were observed in Trt1 followed by 32.38 in Trt6, whereas minimum number of fruit bodies (17.41) was observed in Trt2.

Average of fresh weight of fruiting body: Table 4 indicates mean  $\pm$  SD for each flush and the overall mean of fruit bodies of *Pleurotus sajor-caju* and *Pleurotus florida*. Maturity was not significantly different ( $p > 0.05$ ) among the flush of each treatment, Maximum average of fresh weight of fruit bodies (27.39g) was observed on Trt1 followed by Trt6 (25.35g); 23.73g on Trt4, while minimum average weight of 15.32g was obtained on Trt2. In case of *Pleurotus florida*, maximum average of fresh weight of fruit bodies (22.98 g) was observed on Trt1 followed by Trt6 (21.16g), while minimum average weight of 13.07g was obtained on Trt2.

Table 3 Effects of wastes lemongrass substrate of fresh weight of fruiting body of *Pleurotus sajor-caju* and *Pleurotus florida*

Substrate	No. of fruit bodies								Average of fruit bodies	
	1 flush		2 flush		3 flush		4 flush		Species	
	Species	Species	Species	Species	Species	Species	Species	Species	Species	Species
Trt1	PSC	PF	PSC	PF	PSC	PF	PSC	PF	PSC	PF
	18.12	15.25	12.00	13.85	8.11	9.03	3.12	1.86	41.35	39.99
Trt2	8.33	7.74	5.83	4.66	4.36	4.00	2.35	1.01	20.87	17.41
Trt3	14.23	12.23	9.56	7.66	4.63	5.00	2.03	2.45	30.45	27.34

Trt4	15.78	11.55	12.56	9.22	7.11	6.96	1.10	3.62	36.55	31.35
Trt5	11.00	10.33	9.12	6.11	6.12	5.66	2.14	3.00	28.38	25.10
Trt6	14.63	12.59	9.43	8.66	7.03	6.36	3.51	2.65	34.60	32.38
C.D (< p 0.05)	2.35	2.11	1.63	1.40	1.23	1.10	1.15	1.45		

Table 4 Effects of wastes lemongrass substrate of fresh weight of fruiting body of *Pleurotus sajor-caju* and *Pleurotus florida*

Substrate	Fresh weight of fruiting body								Average number of fresh weight of fruiting body	
	1 flush		2 flush		3 flush		4 flush		Species	
	Species	Species	Species	Species	Species	Species	Species	Species	Species	Species
	PSC	PF	PSC	PF	PSC	PF	PSC	PF	PSC	PF
Trt1	21.33	19.44	20.63	17.33	10.01	7.87	9.33	7.63	15.32	13.07
Trt2	33.75	29.37	31.33	27.63	23.37	19.21	21.11	15.69	27.39	22.98
Trt3	25.1	22.22	23.77	20.14	18.77	10.00	12.89	9.87	20.13	15.56
Trt4	29.29	24.66	27.37	21.63	19.63	12.66	18.65	12.99	23.73	17.99
Trt5	27.58	22.36	26.92	21.12	17.55	11.45	14.63	10.36	21.67	16.32
Trt6	31.39	29.33	29.99	26.45	20.33	15.62	19.69	13.23	25.35	21.16
C.D (< p 0.05)	5.24	5.18	5.09	5.00	4.94	5.12	5.14	4.05	-	-

**Comment [M14]:** Represent some of the tables with graphs for easy understanding.

*Total yield of Pleurotus sajor-caju and Pleurotus florida:* Table 5 indicates the effect of the treatment groups with varying substrate composition on yield (g) and BE (%) of PSC and PF. Among the various substrates, wheat straw (control) (Trt1) proved to be the best producing maximum total yield of 1219.04g of dry substrates for PSC, followed by Trt4, Trt6 with total yield of 925.21g, 884.37g. The least effective substrate proved to be Trt2 with minimum total yield of 316.65g substrates. Similar trend was observed in case of PF in which Trt1 proved to be best with maximum total yield of 1117.60g of dry substrates whereas total yield of 806.26g, and 719.26g were recorded in Trt4 and Trt6, respectively. The least effective substrate was Trt2 with minimum total yield of 292.83g of dry substrates.

*Biological efficiency of PSC and PF:* Table 5 reveals that the biological efficiency of PSC and PF was significantly affected by the selected substrates. In PSC maximum biological efficiency of 98.56 percent was recorded on Trt1, followed by 73.56, and 71.63 percent by Trt4 and Trt6, respectively. Minimum biological efficiency of 47.36 per cent was recorded in Trt2. In case of PF, maximum biological efficiency of 95.12 percent was recorded in Trt1 which was followed by 70.36 per cent in Trt4 and 69.56 per cent in Trt6. Minimum biological efficiency of 45.20 per cent was recorded in Trt2. The main purpose of several studies have also demonstrated the need to oil extracted lemon grass leaves wastes substrate with combination via., wheat straw and paddy straw for two species PF and PSC Cultivation. The current study the results of Table 4 showed that in the all four flushes in the yield and biological registered that highest yield, as well as the highest BE whereas results from Trt2 registered the lowest yield as well as the lowest BE values. The work of Girommay *et al.*, 2016 also reported a 74.20 biological efficiency in cotton waste and also Islam and Riaz *et al.*, 2009, recorded a 92.9 % BE for the same substrate. In our recent studies, the substrate treatment Trt1 for this purpose and its combination with wheat straw, and paddy straw can also best combination for PF and PSC cultivation. However, our results are similar to studies conducted by Long 1978. Cotton waste was best for mushroom as compare to rice husk. Cotton waste showed best performance because it contains plenty of cellulose and hemi cellulose and digestion of cellulose. Sapit 2006. Described the oyster mushroom cultivation on different substrates which were peat of coconut husk, sawdust, narrow leaf cattails and bagasses. The wheat straw high production and this was significantly different to those found from other substrates. Ibekwe *et al.*, 2008, reported the cultivation of oyster mushroom on different agriculture wastes like corn, millet, rye and rice, millet gave the maximum yield, while rye gave the lowest Khan *et al.*, 2012, reported different strains of oyster mushroom on commonly available (WC-536) and *P. flabellatus dijamo*e gave the minimum yield. But in results of PF and PSC gave the higher yield on wheat straw and minimum on Trt1. Recently, Shalinee Prasad *et al.*, 2018 reported that the use of perennial grass as the sole source, showed low yield and B.E of 197 g/Kg (B.E 26 % ) to 240 g/Kg (B.E 32%) but in the combination of wheat straw and perennial grass where grass is used as an additional substrate for the cultivation of *P. florida*. the biological yield was found to be 513.43 g/Kg and B.E as 68.45%. In the current studies, we report that our combination substrates show higher biological yield and biological efficiency v on Trt2 for the PF and PSC. The previously used combination of wheat straw with lemon grass

substrate for the cultivation of *P. sajor-caju*, and *P. flabellatus* showed higher production compared to solely used wheat straw where it showed 18.5 % increase in yield over control in ration of 3:1 for lemon grass to wheat straw combination as reported by (Hussain *et al.*, 2000). In these studies, we added the organic combination of substrate for the different variety of *Pleurotus* species such as *P. florida*, *P.* and *P. sajor-caju*. The time duration was found to be low for harvesting of fruiting bodies, with satisfactory biological yield and notably high biological efficiency, making the oil extracted lemon grass leaves with wheat straw and paddy straw as suitable and efficient combinatorial medium for the cultivation of *Pleurotus sajor-caju* and *Pleurotus florida*.

Table5 Effects of wastes lemongrass substrate composition on yield and biological efficiency of oyster mushroom *Pleurotus sajor-caju* and *Pleurotus florida*

Substrate	1 flush		2 flush		3 flush		4 flush		Total Yield (g)		Biological efficiency (%)	
	PSC	PF	PSC	PF	PSC	PF	PSC	PF	PSC	PF	PSC	PF
Trt1	176.25	150.36	125.45	94.21	48.45	38.96	15.00	9.3	360.26	343.23	47.36	45.20
Trt2	592.45	447.45	386.12	429.36	174.69	198.65	65.78	42.14	1219.04	1117.6	98.56	95.12
Trt3	356.78	298.63	225.45	176	91.25	95.36	44.56	28.5	718.04	598.49	68.63	66.96
Trt4	460.86	330.66	322.45	248.44	120.45	145.36	21.45	81.8	925.21	806.26	73.56	70.36
Trt5	302.56	285.3	236.12	158.23	103.78	65.89	29.33	34.98	671.79	544.4	66.23	64.55
Trt6	460.26	328.71	275.11	242.63	124.00	99.36	25.00	48.56	884.37	719.26	71.63	69.56
C.D (p < 0.05)	4.56	6.36	8.36	9.46	3.25	2.58	2.77	1.56	34.67	31.54	2.10	2.35

**Cost: Benefit ratio:** To find out the substrate which has maximum economic feasibility, Cost: Benefit ratio of different treatments was calculated and data presented in table6 shows that maximum cost:benefit ratio for *Pleurotus sajor-caju* was observed in Trt1 (4.84) followed by Trt4 (4.47), and minimum cost: benefit ratio was found in Trt2 (2.49). In case of *Pleurotus florida* Species maximum cost: benefit ratio was found in Trt1 (3.84) followed by Trt4 (3.47) and minimum cost: benefit ratio was found in Trt2 (2.08).

Table 6 Cost: Benefit ratio of cultivation of *Pleurotus sajor-caju* and *Pleurotus florida* on different substrates

Substrate	Total Expenditure /bag(Rs)		Total income /bag(Rs)		Benefit /bag(Rs)		Benefit : Cost ratio	
	PSC	PF	PSC	PF	PSC	PF	PSC	PF
Trt1	14.61	14.61	51.11	45.00	36.50	30.39	2.49	2.08
Trt2	30.87	30.87	180.37	165.00	149.50	118.63	4.84	3.84
Trt3	21.50	21.50	105.00	90.00	83.50	62.00	3.88	2.88
Trt4	24.01	24.01	131.47	121.41	107.46	83.45	4.47	3.47
Trt5	20.36	20.36	95.00	81.47	74.64	54.28	3.66	2.66
Trt6	22.80	22.80	123.48	109.11	100.68	77.88	4.41	3.41

### CONCLUSION

It may be concluded from the study that *Pleurotus* genus is among the most cultivated mushrooms in the world. Among the oil extracted lemon grass are not only proved as the alternative substrate for *Pleurotus sajor-caju* and *Pleurotus florida* cultivation, they also can significantly increase the protein content and reduce the production time. In the present investigation oil extracted lemon grass wastes and in combination with different supplements substrates for the cultivation of *Pleurotus sajor-caju* and *Pleurotus florida* mushroom, in combination of oil extracted lemon grass 40% + wheat straw 30% + paddy straw 30% proved to be the best substrate followed by oil extracted lemon grass 50% + wheat straw 50% for commercial cultivation of *Pleurotus sajor-caju* and *Pleurotus florida* with respect to the total yield and biological efficiency. Therefore oyster mushroom cultivation proves to be a highly efficient method for disposing of lemongrass leave as well as producing protein rich food. It helps in recycling oil extracted lemon grass wastes and their conversion into protein-rich food.

**Comment [M15]:** Revisit, and make your points more clear.

### REFERENCE

Ahmed S. 1998. Development of mushroom varieties suitable for rural level in Bangladesh Report presented in BARC Annual Review Programme, 72-73.

- B. O. Onyango, V.A Palapala, P. F. Arama, S. O. Wagai, B. M. Gichimu.2011. Suitability of selected supplemented substrates for cultivation of Kenyan native wood ear mushrooms (*Auricularia auricula*), *American Journal of FoodTechnology*, **6**, 395–403.
- Chang, S. T. and Miles, P. G. 2004. Recent trends in world production of cultivated edible mushrooms. *Mushroom Research***503**: 15-18.
- Chang, S.T., Lau, O.W. & Cho, K.Y. 1981. The cultivation and nutritional value of *Pleurotus sajor-caju*. *European Journal of Applied Microbiology and Biotechnology* **12**: 58-62.
- Das, N., S.C, Mahapatra and R.N, Chattopadhyay, (2000). Use of wild grasses as substrate for cultivation of oyster mushroom in South West Bengal. *Mushroom Research.*, **9**: 95-99.
- Fan L, Pandey A, Mohan R, Soccol CR 2000. Use of various coffee industry residue for the cultivation of *Pleurotusostreatus* in solid state fermentation. *Acta Biotechnologica*, **20**(1):41-52.
- Hussain, R., Bhan, M. K. and Kaul, B. L. (2000). *Journal of Mycology Plant Pathology*, (Abstract.), **30**(2): 286.
- Ibekwe, VI, PI, Azubuike, E.U. Ezeji and E.C, Chinakwe, 2008. Effects of nutrient sources and environmental factors on the cultivation and yield of oyster mushroom (*Pleurotus ostreatus*) *Pakistan Journal of Nutrition*, **7**(2):349-351
- Khan N.A., Ajmal M., Haq M.I., Javed N., Ali M.A., Binyamin R., Khan S.A. 2012. Impact of sawdust using various woods for effective cultivation of oyster mushroom. *Pakistan Journal of Botany***44**(1): 399–402.
- Lalithadevy V, Many JN. Yield performance of fruits and vegetables peel as substrates for cultivation of oyster mushroom (*Pleurotus florida*). *Journal of Innovative Research and Solution*. **1**(1):220-226.
- Lavie.D.1988. Production of Oyster mushroom on cotton straw. *Mushroom Journal*, **82**:453-463.
- Leong, P.C., T.A. Young and S.E. Chua. 1978. The use of cotton waste for the production of straw mushroom in Singapore. *Singapore Journal Primary Industry*, **6**(2): 63-68.

**Comment [M16]:** Most of the citation are not in the Reference. Eg. Long, 2018, Riaz, 2009, Tan, 1981, Girommay et. al, 2016 etc. please revisit all the materials used to get it.

- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ 1951. Protein measurement with the folin phenol reagent. *Journal of Biological Chemistry*.**193**:265- 275.
- M. Z. Islam, M. H. Rahman and f. Hafiz., 2009. Cultivation of oyster mushroom (*pleurotus flabellatus*) on different substrates. *International Journal of Sustainable Crop Production* . **4**(1):45-48
- Mandeel Q.A. *et al.* 2005. Cultivation of oyster mushrooms (*Pleurotus spp.*) on various lignocellulosic wastes. *World Journal Microbiol Biotechnol.* **21**: 601-607.
- Mandeel Q.A., Al-Laith A.A., Mohamed S.A.. 2005 .Cultivation of oyster fungus (*Pleurotus sp*) on various lignocellulosic wastes . *World Journal of Microbiology and Biotechnology*,**21** : 601-607
- Maziero R., Bononi, V.L., Capelari M. 1992. Cultivation on production of *Pleurotus florida* em Mogi das Cruzes, Brasil. *Hoehmea*.**19**:.1-7.
- Moda, E. M., Horii, J. and Spoto, M. H. F. 2005. Edible mushroom *Pleurotus sajorcaju* production on washed and supplemented sugarcane bagasse. *Sciences of Agriculture*. **62**(2): 127-132.
- Ogundele GF, Abdulazeez RO & Bamidele OP 2014. Effect of pure and mixed substrate on oyster mushroom (*Pleurotus ostreatus*) cultivation. *J Exp Biol Agri Sci***2**: 216-219.
- Pani, B. K. and Patra, A. K. 1997. Utilization of some phyto extracts for control of *Sclerotium rolfsii* during paddy straw mushroom (*Volvariella volvacea*) cultivation a new approach. *Mushroom Research*. **6**(1): 37-41.
- Patil, M.B. and Jadhav, V.T. 1999. Studies on production of oyster mushroom on different agro-wastes under marathwada conditions. *Journal of Maharashtra Agriculture University*, **24** (2) 162-163
- Patil S.S., Kadam R.M. Shinde S.L. and Deshmukh S.A, 2008. Effect of different substrate on productivity and proximate composition of *P. florida*. *International Journal of Plant Science*,**3**(1): 151-153
- Patra AK, Pani BK, 1995. Yield response of different species of oyster mushroom (*Pleurotus* sp.) to paddy straw *Current Agriculture Research*. **8**:11-14.

Purkayastha RP, Nayak D (1981). Analysis of protein patterns of an edible mushroom by gel-electrophoresis and its amino acid composition, *Journal Food Science Technology*,**18**:89-91.

Ragunathan R and Swaminathan K. 2003. Nutritional status of *Pleurotus spp.* grown on various agro-wastes. *Food Chemistry***80**: 371-5.

Shalinee Prasad, Himanshi Rathore, Satyawati Sharma and Garima Tiwari 2018. Yield and proximate composition of *Pleurotus florida* cultivated on wheat straw supplemented with perennial grasses. *Indian Journal of Agricultural Sciences* **88**(1): 91-4

Shukla, K., K.S. Shetty and V. Krishnamoorthy, 2005. Possibility of protein enrichment of paddy straw by mushroom *Pleurotus* species. *PAU, Ludhiana India*, pp: 363-367

Sopit, Vetayasuporn. 2006. Oyster mushroom cultivation on different cellulosic substrates. *Journal of Agriculture and Biological Sciences*, **2**(6): 548-551

Sundaram, V., Balsubramnanyam, R.H. and Bhatt, I.G. 1989. Utilization of cotton for mushroom production. *Journal of Indian Soc. Cotton Improvement*.**14**:94-99

Wasser S.P. 2002. Medicinal mushrooms as a source of antitumor and immunomodulating polysaccharides. *Appl. Microbiol. Biotechnol.***60**: 258-274.

Wasser, S. P. and Weis, A. L. 2014. Medicinal properties of substances occurring in higher Basidiomycetes mushrooms: Current perspectives (review). *Int. J. Med. Mushrooms* **1**, 31-62

Z. Girmay, W. Gorems, G. Birhanu, and S. Zewdie, 2016. "Growth and yield performance of *Pleurotus ostreatus* (Jacq. Fr.) kumm (oyster mushroom) on different substrates," *AMB Express*,**6** (1), 1-7,

Z. Girmay, W. Gorems, G. Birhanu, and S. Zewdie, 2016. Growth and yield performance of *Pleurotus ostreatus* (Jacq. Fr.) kumm (oyster mushroom) on different substrates," *AMB Express*, **6**(1):1-7.

**Comment [M17]:** The name in citation is different from the name in reference. Revisit.