

# Risk Factors of Hypertension Among Cardiac Patients in Bangladesh-A Single Center Study

## ABSTRACT

**Background:** Hypertension, commonly known as high blood pressure, is a pervasive health issue worldwide, and its prevalence is particularly concerning among cardiac patients in Bangladesh. Cardiovascular diseases are a leading cause of morbidity and mortality in the country, and hypertension is a well-established risk factor for these conditions. **Aim of the study:** The aim of the study was to assess the risk factors of hypertension among cardiac patients in Bangladesh. **Methods:** This cross-sectional observational study took place at the Department of Cardiology, Mugda Medical College & Hospital, Dhaka, Bangladesh, from January 2022 to December 2022. A total of 180 cases of hypertension lasting more than a year enrolled in this study as study population. A purposive sampling method was applied, and data collection utilized a semi-structured, pre-designed questionnaire. MS Office tools SPSS version 23.0 were used for data analysis. **Results:** Approximately 27% of participants had hypertension for less than 5 years, while 19% for 5 to 10 years, and 16% for 10 to 14 years. Remarkably, 28% managed hypertension for over 20 years. The most frequent risk factors were hyperlipidemia (34%), physical inactivity (24%), family history of hypertension (20%), obesity (19%), smoking (17%), and diabetes (12%). **Conclusion:** Among the cardiac patients in Bangladesh, the most potential risk factors for hypertension are hyperlipidemia, physical inactivity, family history of hypertension, obesity, smoking, and diabetes.

**Keywords:** Risk factors, Hypertension, Cardiac patients, Family history, Physical activity.

## INTRODUCTION

“Hypertension or high blood pressure is a leading modifiable risk factor for cardiovascular disease and accounts for more than 10% of the population-attributable fraction (PAF) for mortality worldwide” [1]. “Mortality and morbidity rates due to cardiovascular diseases are escalating worldwide, with disproportionately significantly worse outcomes in developing countries, due to rapid health and nutrition transitions caused by urbanization and globalization. More than 17.92 million people died in 2015 due to cardiac diseases, with the highest death rate of 9.4 million recorded in men and 8.5 million in women” [2]. “The management of HTN and HTN-related diseases, including CHD, usually varies from one country to another, as it depends on the capacity of the country to utilize the available resources for the prevention and control of the diseases” [3]. Over the last several decades, impressive gains have been made in improving hypertension detection, treatment, and control [4,5], but comparatively little work has been done in promoting primary prevention, a subject of growing interest. Encouragingly, accumulating evidence suggests that healthy lifestyle factors, such as a healthy diet and increased physical activity, contribute to lowering blood pressure [6,7], and that managing these risks can offset, at least to some extent, genetic predisposition towards hypertension and the development of subsequent cardiovascular sequelae [8]. “Further, community-based approaches can be effective in reducing blood pressure along with other cardiovascular risk factors in the population” [9,10]. The possible impact of an aging population [11], worsening levels of obesity [12], sedentary lifestyles [13], and high sodium consumption [14] on the burden of hypertension calls for a better understanding of the major risk factors associated with hypertension. It is predicted that there will be a 30% increase, with approximately 1.56 billion adults suffering from hypertension by 2025, and approximately 75% of the world's hypertensive population will reside in developing countries [15]. “Globally, complications of hypertension account for 9.4 million deaths every year out of 17 million deaths due to cardiovascular problems” [16]. “Hypertension has been identified as a significant risk factor for coronary heart disease, congestive heart failure, ischemic and hemorrhagic stroke, renal failure, and peripheral arterial diseases” [17,18]. The objective of this study was to assess the risk factors of hypertension among cardiac patients in Bangladesh.

## METHODOLOGY

This cross-sectional observational study was conducted at the Department of Cardiology, Mugda Medical College & Hospital, Dhaka, Bangladesh, from January 2022 to December 2022. It involved 180 cases of hypertension lasting more than one year. A purposive sampling technique was employed for sample selection and informed written consent was


obtained from all participants before data collection. Inclusion criteria encompassed patients who had made at least two visits to the mentioned hospital, and possessed clear and complete medical charts containing a comprehensive disease history. Exclusion criteria included patients who had only visited the hospital once, those with unavailable or unclear and incomplete medical records, pregnant women, and individuals with mental health issues. Detailed demographic and clinical data were recorded for all participants, and data analysis and presentation were carried out using MS Office tools.

## RESULTS

In this study comprising 180 participants, the age distribution revealed that a significant majority fell into the category of individuals over 45 years old, constituting 67% of the total cohort. The next largest group was those aged between 30 and 45 years, comprising 23% of the participants. In contrast, the youngest age group, consisting of individuals under 30 years old, constituted 10% of the total study population. The majority of cases (59%) were male, indicating a predominance of males in the study population. We observed that nearly two-thirds of the participants (61%) were from rural areas. The findings of this study indicate that nearly half of the participants, accounting for 47% of the total, fell into the category of individuals with a BMI of 30.0 or higher, which is considered obese. The next most prevalent group had a BMI ranging from 25.0 to 29.9, representing 34% of the study subjects. The remaining 19% of participants had a BMI within the range of 18.5 to 24.9, which is considered within the normal weight range. In analyzing the socio-economic status of our participants, we observed that nearly half of the cases (48%) were from the middle class and more than one-third (34%) were from lower-class families. Approximately 27% of our participants lived with hypertension for less than 5 years. In contrast, 19% had a duration of 5 to 10 years, while 16% had dealt with hypertension for 10 to 14 years. Notably, a significant proportion, comprising 28% of the participants, had managed hypertension for over 20 years. In distributing the risk factors among our participants, we observed that, in more than one-third (34%) of cases hyperlipidemia and in about one-fourth (24%) of cases, physical inactivity was found. Besides these, in 20%, 19%, 17% and 12% of cases family history of HTN, obesity, smoking and diabetes were observed respectively.

**Table 1:** Age distribution of Participants (N=180)

Age (Year)	n	%
<30 yrs.	18	10%
30-45 yrs.	41	23%
>45 yrs.	121	67%



*Figure 1: Column chart showed age wise participants (N=180)*




Figure 2: Pie chart showed gender wise participants distribution (N=180)




Figure 3: Ring chart showed residential status wise participants (N=180)

**Table 2:** BMI status of study subjects (N=180)

BMI (Kg/m <sup>2</sup> )	n	%
18.5–24.9	35	19%
25.0–29.9	61	34%
≥30.0	84	47%




Figure 4: Column chart showed family status wise participants (N=180)

Table 3: Duration of hypertension (N=180)

Duration (Year)	n	%
<5 Yrs.	48	27%
5-10 Yrs.	34	19%
10-14 Yrs.	29	16%
15-20 Yrs.	18	10%
>20 Yrs.	51	28%




Figure 5: Bar chart showed duration of hypertension among participants (N=180)

Table 4: The risk factors of hypertension. (N=180)

Risk factors	n	%
Hyperlipidemia	62	34%
Physical inactivity	43	24%
Family history of HTN	36	20%
Obesity	34	19%
Smoking	31	17%

Diabetes	22	12%
Chronic kidney disease	16	9%
Alcohol use	3	2%




Figure 6: Bar chart showed risk factors of hypertension among participants (N=180)

## DISCUSSION

This study aimed to assess the risk factors of hypertension among cardiac patients in Bangladesh. In this study of 180 participants, the majority (67%) were over 45 years old, with the next largest group (23%) aged between 30 and 45 years. However, some other studies have reported an increasing prevalence of coronary artery disease among the young population in Bangladesh and India. [18] In a study by Smith et al. [19], it was reported that the relationship between BP and the risk of CVD events is continuous, consistent, and independent of other risk factors. The pre-hypertensive group has a greater chance of progression to hypertension and other CVDs. The majority of our cases (59%) were male and similar findings were found in another recent study [20]. But in some studies [21,22] females were dominating. We found that nearly two-thirds of our participants (61%) were from rural areas. However, in many studies, the majority of the study subjects were found from urban areas. Furthermore, the World Health Federation reported that urban residents may face an increased risk of experiencing CVD due to poor quality city living standards. These standards include the limited availability of safe spaces for exercise or recreation, growing pressures from mass marketing, and the prevalence of unhealthy and affordable food options in urban areas. The study findings revealed that 47% of participants were classified as obese (BMI of 30.0 or higher), while 34% had a BMI between 25.0 and 29.9, and 19% fell within the normal weight range (BMI of 18.5 to 24.9). Another study indicated that overweight individuals faced a twofold risk of developing hypertension, while obese individuals had over a threefold risk compared to underweight individuals. Additionally, further research demonstrated that obesity, defined as a BMI greater than 25, carried a 2.62 times higher risk of developing hypertension when compared to those with a BMI below 25. [23] This suggests that obesity not only increases the risk of hypertension but can also predict uncontrolled hypertension. Other studies have supported these findings, showing that overweight and obese patients are less likely to have controlled blood pressure. [24] In analyzing the distribution of risk factors among our participants, it was noted that hyperlipidemia was present in more than one-third (34%) of cases, while physical inactivity was identified in about one-fourth (24%) of cases. Additionally, a family history of hypertension was present in 20% of cases, obesity in 19%, smoking in 17%, and diabetes in 12%. Some other studies [25,26] reported that the association of family history increases the risk of developing hypertension because it exposes the patient to high blood pressure, heart disease, and stroke, or their family had the same lifestyle habit. In some studies, [27,28], it was reported that physical activity in less than 10 minutes daily increased the risk of hypertension. This increase in risk was attributed to increased peripheral vascular resistance due to an increase in neuro-hormonal and

structural responses, leading to enhanced sympathetic nerve activity and a decrease in arterial lumen diameters [29]. Similarly, a study in India showed that a lack of physical exercise was the major risk factor for the development of complications of hypertension [30]. A Canadian study showed that excess body weight and living a sedentary lifestyle predispose an individual to hypertension and its complications [31].

### **LIMITATION OF THE STUDY**

This study has significant limitations. It was a single-centered study with a relatively small sample size and a short duration. Therefore, its findings may not be fully representative of the entire country's situation. Caution is necessary when applying these results to a broader national context. A larger, multi-centered study with a longer timeframe may provide a more comprehensive understanding.

### **CONCLUSION & RECOMMENDATION**

The most prevalent risk factors of hypertension among cardiac patients in Bangladesh are hyperlipidemia, physical inactivity, family history of hypertension, obesity, smoking, and diabetes. Recognizing these prevalent risk factors is crucial for developing targeted interventions and public health strategies to mitigate the burden of hypertension in this population. Addressing these factors through lifestyle modifications, improved screening, and effective management approaches is vital for improving cardiovascular health and reducing hypertension-related complications among cardiac patients in Bangladesh.

### **Consent**

As per international standards or university standards, Participants' written consent has been collected and preserved by the author(s).

### **REFERENCES**

- [1] Tanuseputro P, Manuel DG, Leung M, et al. Risk factors for cardiovascular disease in Canada. *Canadian Journal of Cardiology* 2003; 19(11):1249-59.
- [2] BeLue R, Okoror TA, Iwelunmor J, et al. An overview of cardiovascular risk factor burden in sub-Saharan African countries: a socio-cultural perspective. *Global Health* 2009; 5:10.
- [3] Whitworth JA; World Health Organization, International Society of Hypertension Writing Group. 2003 World Health Organization (WHO)/International Society of Hypertension (ISH) statement on management of hypertension. *J Hypertens* 2003; 21:1983-92.
- [4] Hemmelgarn BR, Chen G, Walker R, et al. Trends in antihypertensive drug prescriptions and physician visits in Canada between 1996 and 2006. *Canadian Journal of Cardiology* 2008; 24(6):507-12.
- [5] Campbell NR, Tu K, Brant R, et al. The impact of the Canadian Hypertension Education Program on antihypertensive prescribing trends. *Hypertension* 2006; 47(1):22-8.
- [6] Appel LJ, Champagne CM, Harsha DW, et al. Effects of comprehensive lifestyle modification on blood pressure control: main results of the PREMIER clinical trial. *JAMA* 2003; 289(16):2083-93.
- [7] Patnode CD, Evans CV, Senger CA, et al. Behavioral Counseling to Promote a Healthful Diet and Physical Activity for Cardiovascular Disease Prevention in Adults Without Known Cardiovascular Disease Risk Factors: Updated Evidence Report and Systematic Review for the US Preventive Services Task Force. *JAMA* 2017; 318(2):175-93.
- [8] Pazoki R, Dehghan A, Evangelou E, et al. Genetic Predisposition to High Blood Pressure and Lifestyle Factors: Associations with Midlife Blood Pressure Levels and Cardiovascular Events. *Circulation* 2018; 137(7):653-61.
- [9] Jousilahti P, Laatikainen T, Peltonen M, et al. Primary prevention and risk factor reduction in coronary heart disease mortality among working-aged men and women in eastern Finland over 40 years: population-based observational study. *BMJ* 2016; 352: i721.
- [10] Pennant M, Davenport C, Bayliss S, et al. Community programs for the prevention of cardiovascular disease: a systematic review. *American Journal of Epidemiology* 2010; 172(5):501-16.
- [11] Statistics Canada. Population Trends by Age and Sex, 2016 Census of Population. Available at: <https://www150.statcan.gc.ca/n1/pub/11-627-m/11-627-m2017016-eng.htm>.

- [12] Public Health Agency of Canada. Tackling Obesity in Canada: Obesity and Excess Weight Rates in Canadian Adults. Available at: <https://www.canada.ca/en/public-health/services/publications/healthy-living/obesity-excess-weight-rates-canadian-adults.html>.
- [13] Colley RC, Garriguet D, Janssen I, et al. Physical activity of Canadian adults: accelerometer results from the 2007 to 2009 Canadian Health Measures Survey. *Health Reports* 2011; 22 (1):7-14.
- [14] Garriguet D. Sodium consumption at all ages. *Health Reports* 2007; 18 (2):47-52.
- [15] Bacurau AGdM, Francisco PMSB. Reasons for non-vaccination against influenza among older adults with hypertension in Brazil: a cross-sectional study. *Sao Paulo Med J*. 2020; 138:322–325.
- [16] Zinat Motlagh SF, Chaman R, Ghafari SR, et al. Knowledge, treatment, control, and risk factors for hypertension among adults in Southern Iran. *Int J Hypertension*. 2015; 2015.
- [17] Hypertension: analysis of worldwide data. *Lancet*. 2005;365(9455):217–223.
- [18] Das SK, Sanyal K, Basu A. Study of urban community survey in India: growing trend of high prevalence of hypertension in a developing country. *International Journal of Medical Sciences*. 2005; 2(2):70 –78.
- [19] Vasan RS, Larson MG, Leip EP, Kannel WB, Levy D. Assessment of frequency of progression to hypertension in non-hypertensive participants in the Framingham Heart Study: a cohort study. *The Lancet*. 2001; 358(9294):1682 –1686.
- [20] Ayele, Habtamu, Akalu Banbeta, and Abiyot Negash. "Cardiovascular disease risk factors in hypertensive patients: a case study of Jimma University Medical Center." *Health Services Research and Managerial Epidemiology* 9 (2022): 23333928221078601.
- [21] Parikh NI, Pencina MJ, Wang TJ, et al. A risk score for predicting near-term incidence of hypertension: The Framingham Heart Study. *Annals of Internal Medicine* 2008;148(2):102-10.
- [22] Kivimaki M, Tabak AG, Batty GD, et al. Incremental predictive value of adding past blood pressure measurements to the Framingham hypertension risk equation: The Whitehall II Study. *Hypertension* 2010; 55(4):1058-62.
- [23] Jonas JB, Nangia V, Matin A, Joshi PP, Ughade SN. Prevalence, awareness, control, and associations of arterial hypertension in a rural central India population: The Central India Eye and Medical Study. *Am J Hypertens*. 2010;23(4):347–350.
- [24] Sakboonyarat B, Rangsin R, Kantiwong A, Mungthin M. Prevalence and associated factors of uncontrolled hypertension among hypertensive patients: a nationwide survey in Thailand. *BMC Res Notes*. 2019; 12 (1):1–8.
- [25] Corvol P, Jeunemaitre X, Charru A, Soubrier F. Can the genetic factors influence the treatment of systemic hypertension? The case of the renin-angiotensin-aldosterone system. *Am J Cardiol*. 1992;70(12): D14–D20.
- [26] Carmelli D, Robinette D, Fabsitz R. Concordance, discordance and prevalence of hypertension in World War II male veteran twins. *J Hypertens*. 1994;12(3):323–328.
- [27] Awoke A, Awoke T, Alemu S, Megabiaw B. Prevalence and associated factors of hypertension among adults in Gondar, Northwest Ethiopia: a community-based cross-sectional study. *BMC Cardiovasc Disord*. 2012;12(1):1–6
- [28] Giday A, Tadesse B. Prevalence and determinants of hypertension in rural and urban areas of southern Ethiopia. *Ethiop Med J*. 2011;49(2):139–147.
- [29] Hamer M. The anti-hypertensive effects of exercise. *Sports Med*. 2006;36(2):109–116.
- [30] Kumar C, Sasi Sekhar T, Sahithi B. Hypertension—the silent killer, awareness of the risk factors and complications of hypertension among hypertensives. *Int J Adv Res*. 2006;4(6):1277–1281.
- [31] Kirkland SA, MacLean DR, Langille DB, Joffres MR, MacPherson KM, Andreou P. Knowledge and awareness of risk factors for cardiovascular disease among Canadians 55 to 74 years of age: results from the Canadian Heart Health Surveys, 1986-1992. *CMAJ (Can Med Assoc J)*. 1999;161(8): S10.