

Anthelmintic Potential of *Vernonia amygdalina* on *Toxocaracanis* in Domesticated Dogs

Abstract:

Toxocariasis is a worldwide zoonotic parasitic infection and one route of preventing the disease in dogs and subsequent transmission to humans, especially children, is through treatment with anthelmintic. A study aimed at assessing the anthelmintic efficacy of *Vernonia amygdalina* leaves extract on *Toxocaracanis* in domesticated dogs was carried out where the proximate and phytochemical components of the leaf extract were determined using the standard method and Gas Chromatography-Mass Spectrometry, the anthelmintic efficacy was assessed *in vitro* and *in vivo* respectively. The study revealed that this plant contained alkaloids, saponins, tannins, glycosides, flavonoids, terpenoids, phenol and steroids. Lethal concentration (LC50) of 14.83ml/mg at 12 hours and a percentage egg reduction of 84% after 2 weeks was recorded, which revealed that the plant leaves have anthelmintic potential and this could be attributed to the presence of flavonoids which are associated with the treatment of gastric infections and also n-Hexadecanoic acid which accounts for its anthelmintic (nematicidal) effect, alongside Octadecanoic acid which is an anti-inflammatory compound.

Keywords: Anthelmintic, Toxocariasis, *Vernonia amygdalina*, Dog, Efficacy

Introduction

Zoonoses are those diseases that can be transmitted from human and animals, in this regard dogs are considered as a public health concern as they harbor various pathogens such as zoonotic helminthes including *Toxocaracanis* (Lescandoet *al.* 2010). Toxocariasis is a worldwide parasitic infection of dogs and other carnivores, where the adult worms are found in the intestine and eggs are passed with feces into the environment which becomes embryonated under optimum climate condition and infection is established upon ingestion by another host (Habluetzelet *al.*, 2003). The shedding of eggs by adult worm through the host is considered a means of infection dissemination, where human can become an accidental host by ingesting the eggs via pica and devouring on paratenic host such as chicken, cattle, pig, etc and vegetable. (Anunobiet *al.*, 2017, Meriguetiet *al.*, 2017; Healy *et al.*, 2021). Although, in humans the development to adult is hindered, but the larvae migrate through various organs resulting in syndromes such as Visceral larva migrans (VLM), Ocular Larva Migrans (OLM) and Neuron Larva Migrans(NLM) as well as covert infection and asymptomatic toxocariasis (Eslahiet *al.* 2021; Morsy, 2020).

The preventive measures required against this infection include environmental intervention strategies that limit where dogs are allowed in our houses and surroundings but unfortunately, dogs have been identified as one of man's closest companion since the ages of non-civilization where they act as emotional support, security etc., and limiting their access to man will reduce its effectiveness in fulfilling its role to man (Udoidunget *et al.*, 2018). As such, treatment with anthelmintic becomes one of the major measure of preventing the disease in dog and subsequent transmission to man especially children. (El-Menyawe and Mohammad, 2015).

In recent times, reports have emerged on the potential reduction in the effectiveness or even emergence of drug resistance through wide spread and frequent use of this anthelmintic chemotherapy (Nweze *et al.*, 2013). These have created an urgent need for newer, inexpensive and accessible drugs, and traditional herbal medicine based largely on medicinal plants which offers a major and accessible source of health care to the man and animals in most developing countries (Tandon *et al.*, 2011). Therefore, there is need for proper scientific documentation of the efficacy of some of the plants extracts used as anthelmintic as various plants contain bioactive compounds that have shown potential in combating helminth infections as reported in folklore (Egualé and Giday, 2009; Oyeyemi *et al.*, 2018; Adams *et al.* 2023). One of such plant is the bitterleaf scientifically known as *Vernonia amygdalina* which is a shrub that grows in tropical Africa and its parasitic use was first reported among chimpanzees in the wild and has been used by humans (Huffman and Seifu, 1989; Al-shaibani *et al.*, 2009), it's important to note that the scientific evidence supporting its effectiveness as an anthelmintic is limited and this work is aimed at validating its use.

Materials and Methods

The study was carried out in Akwa Ibom State, Nigeria which lies between latitude 4° 32' and 5° 53' North and longitude 7° 25' and 8° 25' East with total land area of 7.249 km². The leaves of the *Vernonia amygdalina* plant (fig.1) were collected from matured plants identified and authenticated in the herbarium unit, Department of Botany and Ecological Studies, University of Uyo. The ethanolic extraction of the leaves was carried out using maceration method (Sujon *et al.*, 2008). Dogs (puppies) aged about 7 months weighing about 7 – 8 kg were bought from vendors and used for the studies. The crude leaf extract of *Vernonia amygdalina* commonly known in the area as bitterleaf (Etidot) was screened for anthelmintic effects *in vitro* using adult

worms (*Toxocaracanis*) and also *in vivo* using experimentally infected dogs (puppies infected with *Toxocaracanis*) according to the method of Adamset *al.*, (2008).



Fig 1: Picture of *Vernonia amygdalina Delile*

Qualitative phytochemical screening of the plants leaves was carried out using standard screening tests as described by Adingra (2017).

The proximate composition of the plant leaves samples was determined using the standard methods of analysis of Association of Official Chemists (AOAC, 2010). Crude protein, crude lipid, carbohydrate, Moisture, crude fiber and ash contents in the samples were analyzed.

GC-MS analysis was carried out on an Agilent technology 7890 GC system equipped with a mass spectrometric detector (MSD). Ms model was agilent technology 5975 ms, the column used was HP-5MS agilent technology, length of the column 30 m, internal diameter 0.320 mm, thickness of 0.25 μ m. Volume of sample injected was 1 μ L. Oven temperature program with initial temperature of 80°C to hold for 2 minutes at 10°C/min to final temperature of 240°C to hold for 6 minutes with injector temperature of 250°C. The mobile phase was helium gas while the stationary phase was the column.

Detection of Components Analysis of mass spectrum GC-MS was carried out using the database of National Institute Standard and Technique (NIST) having more than 62,000 patterns. The spectrum of the unidentified components were compared with the spectrum of the identified components stored in the NIST library. The name, molecular weight, structure of the components in the test material were ascertained (Oshiobugieet *al.*, 2017)

In vitro analysis of the extract was carried out using the method of Adamset *al.* (2023). Prior to the experiment, all the puppies were examined coprologically and treated accordingly, except for few of those that had only *Toxocaracanis* eggs present in their faeces and was kept as the reservoir host, to ascertain a worm free colony. Faeces containing *Toxocaracanis* eggs was obtained from a naturally infected puppy (reservoir host). Dogs (worm free colony of puppies) were infected with *Toxocaracanis* by contaminating their food and water with the infected faecal sample that was collected from the reservoir host and kept moist for some days at room temperature for embryonation to occur, and infection was confirmed through faecal analysis after three weeks. Some of the dogs that were passing eggs in faeces were sacrificed using appropriate humane method and the small and large intestine carefully extracted from the body of the dogs. The intestines were opened with a scissor and the content washed with tap water into a clean plastic container, then the adult worms were collected carefully with the aid of the blunt head of a needle into petri dishes containing physiological saline (buffered saline). Two sets each of 5 petri dishes (groups) containing 10 adult worms each (total of 100 worms) was prepared and the plant extract was diluted at various concentrations 10%, 20%, 50% and 100% and a control with only physiological saline. 0.1ml of each diluted plant extracts (extract of *Vernonia amygdalina*) at various concentrations were pipetted into the different petri dishes (groups) leaving a control and then allowed for 12 hours at room temperature. The groupings were as follows:

Group 1: adult worms treated with 10% extract

Group 2: adult worms treated with 20% extract

Group 3: adult worms treated with 50% extract

Group 4: adult worms treated with 100% extract (crude extract)

Group 5: adult worms which received only physiological saline

The non-motile worms were counted from the various plates (groups) after 30 mins, 1 hour, 6 hours and 12 hours respectively and then percentage mortality calculated.

$$\text{Percentage death} = \frac{\text{Number of dead parasites}}{\text{Total number of parasites}} \times 100$$

Lethal concentration at 25 and 50% was calculated using probit analysis (AAT Bioquest Inc., 2022).

The *in vivo* screening was carried out according to the method of Adamset *al.* (2023), the plant ethanolic extract was tested using 4 groups of experimentally infected puppies and one group of worm free colony of puppies which were made up of two males and two females respectively i.e. in duplicate, (making a total of 40 puppies).

Three of the experimentally infected groups were treated orally with LC50 and LC25 of the extract and Mebendazole, all at the dosage of 50 mg/kg body weight divided into three doses for 3 consecutive days (as it is the recommended dosage of Mebendazole for therapeutic use in Nigerian dogs) and one group was left untreated (positive control). The groupings were:

Group 1: Infected puppies treated with LC25 obtained from the *in vitro* studies

Group 2: Infected Puppies treated with LC50 obtained from the *in vitro* studies

Group 3: Infected Puppies treated with Mebendazole

Group 4: Infected puppies without treatment

Group 5: Non-infected puppies

The efficacy of the extract on the experimental animals was determined by faecal egg count from direct smear faecal analysis using the formula:

$$\text{Faecal Egg Count Reduction (\%)} = \frac{\text{Pre-treatment egg count} - \text{post treatment egg count}}{\text{Pre-treatment egg count}} \times 100$$

(Salienet *al.*, 2019).

Haematological test which included white blood cell count (WBC), packed cell volume (PCV) and red blood cell count (RBC) was carried out on the puppies before and after the experiment. Packed cell volume (PCV) was assessed by the method of Nweze *et al.* (2013). Blood was collected from the tail veins of the dogs into heparinized capillary tubes. One end of the tube was sealed using sealer. The tubes were placed on a capillary tube rack and centrifuged with a micro-hematocrit centrifuge at 5000 revolutions/min for 5 min. The PCV was then read using a hematocrit reader. White and red blood count was carried out using the Neubauer haemocytometer counting chamber. Behavioural changes observed from the experimental animals was noted.

All data obtained were subjected to analysis using Microsoft Excel and SPSS Version 21 data packages. Chi-square test was used to compare differences based on the level of statistical significance of $P \geq 0.05$ with 95% confidence interval and results are presented in table 3.

RESULT AND DISCUSSION

The Qualitative phytochemical analysis of the plants *Vernonia amygdalina* in Table 1 revealed that the plant have some phytochemical components namely; saponins, tannins, glycosides, flavonoids, phenol and steroids. Saponins and flavonoids were the most abundant.

Table 1: Qualitative Analysis of *Vernonia amygdalina*

Phytochemical	Status
Alkaloids	+
Saponins	++
Tannins	+
Glycosides	+
Flavoniods	++
Steroids	+
Terpenoid	+
Phenols	+

++ =Abundantly Present, + = Present, - = Absent

Proximate analysis of *Vernonia amygdalina* presented in Table 2 revealed 22.8g of crude protein, 4.2g of fat, 35.9g of carbohydrate, 8.4g of moisture, 15.5g of ash and 20.2g of crude protein.

Table 2: Proximate Analysis of *Vernonia amygdalina*

Parameter	<i>Vernonia amygdalina</i>	
	Value (g/100mg)	
Crude protein	22.8	

Crude fat	4.2
Carbohydrate	35.9
Moisture	8.4
Ash	15.5
Crude fiber	20.2

The GC-MS Chromatogram of the ethanolic leave extract *Vernonia amygdalina* revealed 15 peaks of compounds as presented in Fig 2 with their percentage peak area. The major phyto-compounds identified as shown in Table 3 were Oleic acid (29.52%), Octadecanoic acid (19.3%), n-Hexadecanoic acid (8.29%), Methyl octanoic acid (5.75%) and Tetradecanoic acid (4.80%) while the minor compounds encountered were Lupeol (2.38%), E-13-Decanoic acid (1.51%), Pentafluoropropanoic acid (1.40%), 3H-Byarazol-3-one and Decanoic acid (1.28%), 15-Tetracosenoic acid (1.21%), vitamin E (1.17%), Chiloscyphone (0.81%) and 4-Octenoic acid (0.73%).




Figure 2: GC-MS Chromatogram of Ethanolic Extract of *V. amygdalina* Leaf

Table 3: Phyto- components identified in the *Vernonia amygdalina*

Peak	Retention time	% Peak Area	Compound Analyzed	Molecular Weight	Molecular formular
1.	7.628	1.28	3H-Pyrazol-3-one	126	C ₆ H ₁₀ N ₂ O
2.	10.658	1.28	Decanoic acid	186	C ₁₁ H ₂₂ O ₂
3.	11.526	8.29	n-Hexadecanoic acid	256	C ₁₆ H ₃₂ O ₂
4.	12.956	0.55	Pentanoic acid	130	C ₇ H ₁₄ O ₂
5.	13.761	5.75	Methyloctanoic acid	158	C ₉ H ₁₈ O ₂
6.	15.667	0.73	4-Octenoic acid	156	C ₉ H ₁₆ O ₂
7.	17.139	4.80	Tetradecanoic acid	228	C ₁₄ H ₂₈ O ₂
8.	18.941	1.21	15-Tetracosenoic acid	380	C ₂₅ H ₄₈ O ₂
9.	20.178	29.52	Oleic acid	282	C ₁₈ H ₃₄ O ₂
10.	20.389	19.31	Octadecanoic acid	284	C ₁₈ H ₃₆ O ₂
11.	21.658	1.40	Pentafluoropropionic acid	332	C ₁₅ H ₂₅ F ₅ O ₂
12.	23.524	1.51	(E)-13-Docosenoic acid	338	C ₂₂ H ₄₂ O ₂
13.	25.156	1.17	Vitamin E	430	C ₂₉ H ₅₀ O ₂
14.	26.499	2.38	Lupeol	426	C ₃₀ H ₅₀ O
15.	27.913	0.81	Chiloscyphone	218	C ₁₅ H ₂₂ O
Total		100.0			

Table 4: *In vitro* screening of the *Vernonia amygdalina* extracts on adult worms

Concentration (ml/mg)	Percentage (%) of Dead Adult Worms			
	N=100			
	30mins	1hr	6hrs	12hrs
10ml/mg (10%)	–	–	–	40
20ml/mg (20%)	–	–	10	60
50ml/mg (50%)	–	–	20	70
100ml/mg (100%)	–	–	20	80
Control	–	–	–	–

LC25= 3.32ml/mg, LC50=14.83ml/mg

In vitro administration of the ethanolic extract of *Vernonia amygdalina* to determine the lethal concentration shown in Table 4 revealed that the extract had no effect on the adult *Toxocaracanis* within 30mins to one hour. Observable effect was seen within 6hrs and 12hrs, where ten percent (10%) and 60% death were recorded for 20ml/mg, 20% and 70% death recorded for 50ml/mg and the crude extract(100ml/mg) recorded 20% and 80% death respectively. But with 10ml/kg, no death was observed in the first 6 hours although 40% death

was recorded at 12hours. However, the lethal concentration was calculated at 12hrs to be 14.83ml/mg.

In vivo screening of *Vernonia amygdalina* extract on puppies revealed a high percentage egg reduction of 80.6% and 84% from the Faecal Egg Count (FEC) at LC25 and LC50 respectively, while the drug (Mebendazole) showed a 90.6% reduction as shown in Figure 3




Figure 3: *In vivo* screening of *Vernonia amygdalina* extracts on puppies (Fecal Egg Count)

Table 5: Heamatological analysis of the puppies used for the *In vivo* screening

Treatments	RBCs(μ L) Mean \pm SD	PCV (%) Mean \pm SD	WBCs(μ L) Mean \pm SD
Non-infected	5.30 \pm 0.12a	42.00 \pm 1.73a	9.33 \pm 0.17cd
Infected untreated	3.70 \pm 0.12f	23.77 \pm 0.62c	14.17 \pm 0.88a
Infected treated with Mebendazole	4.40 \pm 0.06de	25.83 \pm 0.83c	11.63 \pm 0.52b
Infected treated with LC25 of <i>Vernonia amygdalina</i>	4.43 \pm 0.07cd	31.00 \pm 0.58b	9.00 \pm 0.76d
Infected treated with LC50 of <i>Vernonia amygdalina</i>	4.73 \pm 0.09bc	33.00 \pm 1.53b	10.00 \pm 0.29bcd
Total	4.51\pm0.09	25.96\pm0.89	10.82\pm0.52
p Value	<0.001*	<0.001*	<0.001*

Means along the same column with different Alphabet(s) are statistically significant at $p \leq 0.05$, ns – Not significant at $p > 0.05$, * - significant at $p \leq 0.05$.

The overall blood count of all the puppies used for the experiment as shown in Table 5 were $4.51 \times 10^6 \mu\text{L}$ for RBC, 25.96% for PCV and 10.82×10^3 for WBC. A marked reduction was observed in the pack cell volume (PCV) and Red blood count (RBC) of the infected untreated puppies (23%, $3.70 \times 10^6 \mu\text{L}$) and also those that were treated with Mebendazole (25.83%, $4.40 \times 10^6 \mu\text{L}$) while those treated with LC25 and LC50 extracts concentrations showed a slight reduction of 31, $3.443 \times 10^6 \mu\text{L}$ and 33%, $4.73 \times 10^6 \mu\text{L}$. The difference was statistically significant at $P < 0.001$.

The WBC revealed an increase in the infected untreated (positive control) ($14 \times 10^3 \mu\text{L}$) and also those treated with Mebendazole ($11.63 \times 10^3 \mu\text{L}$) while those treated with LC₂₅ and LC₅₀ extract showed a slight increase of 9.0 and $10 \times 10^3 \mu\text{L}$. The difference was statistically significant at $P < 0.001$. *Vernonia amygdalina* revealed a minimal alteration of the three hematological parameters (WBC, PCV and RBC) analyzed compared to the uninfected control group and group treated with Mebendazole.

Discussion

The study revealed that *Vernonia amygdalina* has an anthelmintic efficacy on *Toxocaracanis* in dogs which agrees with the works of Danguah *et al.* 2012; Nweze *et al.* 2013; Ejiofor *et al.*, 2017. *Vernonia amygdalina*, also known as bitter leaf, is a plant that is widely used in Africa to treat a multiplicity of diseases including schistosomiasis, amoebic dysentery, and gastrointestinal problems (Masaba, 2000). Research has reported that any part of the plant can be used, but there seems to be a preference for the leaves, which are quite bitter, the bitterness is seen as a sign of potency (Masaba, 2000; Ijeh and Ejike, 2011). It is found to contain two classes of bitter compounds namely sigmastane-type steroid glucoside and four known sesquiterpene lactones, vernodalin, vernolide, hydroxyvernolide, and vernodalol. All the pharmacological activity observed for this plant has been attributed to these bioactive compounds (Alawa *et al.*, 2003).

The reduction in fecal egg count by the leaf extract of *Vernonia amygdalina* in this study which was also reported by Adeolu *et al.* (2007) is a positive and welcomed development in our local

helminths struggle because the plants are available year-round in Nigeria. The easy access to this plant and its availability might mean reduced cost of veterinary attention which in most cases are neglected.

The study showed that the plant leaves contained alkaloids, saponins, tannin, glycosides, flavonoids, terpenoids, phenol and steroid which agrees with the report of Ekekere *et al.*, 2000; Kaur *et al.*, 2019; Ujah, *et al.*, 2021. The presence of these phytochemicals compounds especially flavonoids indicate that the plants could be useful in the treatment of helminthiasis alongside other diseases. Also, compounds like Alkaloids are implicated with antimalarial and analgesic value and can equally be used as stimulants (Arona and Tamrakar, 2017).

GC-MS analysis revealed that it contained n-Hexadecanoic acid as one of its major compounds which is a nematicidal compound that accounts for its anthelmintic effect and it also has anti-inflammatory compounds such as Octadecanoic acid and Decanoic acid. This agrees with the findings of Oshiobugie *et al.*, (2017) and Nuryanto (2018).

The leaf extract used in *in vivo* treatment revealed a minimal alteration of the three hematological parameters (WBC, PCV and RBC) examined in the study which agrees with the report of Adeolu *et al.*, (2007)

The activities shown by *Vernonia amygdalina* leaf extract are of considerable importance and justify its use as anthelmintic in folklore traditional medicine as the use of medicinal herbs in the treatment of infectious diseases has attracted the attention of scientists worldwide (Aswathi and Dhivya, 2017; Oyeyemi *et al.*, 2018).

Conclusion

Aside from the use of prescribed drugs for the treatment of toxocariasis in dogs, various plant extracts (both water and ethanolic extract) have been used by traditional/herbal medicine in the treatment of gastrointestinal helminthiasis in both humans and animals. The plant leaves of bitter leaf (*Vernonia amygdalina*) which is one of the most commonly used plants were collected, identified and anthelmintic efficacy on *Toxocaracanis* was tested *in vivo* and *in vitro*. LC50 of 14.83ml/mg at 12 hours and a percentage egg reduction of 84% after 2 weeks was recorded which revealed that the plant leaves have anthelmintic potentials with minimal alteration of

hematological parameters and this could be attributed to the presence of flavonoids which is associated with the treatment of gastric infections and also n-Hexadecanoic acid which accounts for its anthelmintic effect, alongside Octadecanoic acid which is an anti-inflammatory compound.

REFERENCES

- AAT Bioquest Inc. (2022). *Quest Graph LC50 Calculator*. <http://www.aatbio.com/tools/LC50-Calculator>
- Adams, E. G., Ekott, E. J., Usip, L. P. and Opara, K. N. (2023). Efficacy of *Azadirachta indica* in the Treatment of Gastrointestinal Helminthiasis. *Biotechnology Journal International*, 27(2): 47-55.
- Adeolu, A., Adedapo, A., Otesile, T. and Soetan, K. O. (2007) Assessment of the anthelmintic Efficacy of an Aqueous Crude Extract of *Vernonia amygdalina*. *Pharmaceutical Biology* 45(7):546-568
- Adingra, K. M. D., Konan, K. H., Eugene, J. P., and Tano, K. (2017). Phytochemical Properties and Proximate Analysis of Papaya (*Carica papaya*) Peel. *Turkish Journal of Agriculture Food Science and Technology*, 5(6): 676-680
- Alawa, C. D., Adamu, A. M., Gefu, J. O., Ajanusi, O. J., Adadu, P. A., Chiezey, N. P., Alawa, J. N. and Bowman, D. D. (2003). In vitro Screening of two Nigerian Medicinal Plants *Vernonia amygdalina* and *Anna senegalensis* for Anthelmintic Activity. *Veterinary Parasitology*, 113:73-81.
- Al-Shaibani, I.R.M., Phulan, M. S. and Shiekh, M. (2009). Anthelmintic Activity of *Fumaria parviflora* (Fumariaceae) against Gastrointestinal Nematodes of Sheep. *Int. J. Agric. Biol.*, 11(4): 431–436
- Anunobi, T. J., Okeye, I.C. and Godwin, N. (2017) Toxocariasis and public health: an epidemiology Review.
- AOAC (Association of Official Analytical Chemist) (2010). *Official Methods of Analysis* (15th Edition). Washington D.C, 12-13.
- Arona, C. and Tamakar, V. (2017). *Gmelina arborea*: Chemical Constituents, Pharmacological Activities and Applications. *Journal of Phytomedicine*, 9: 528-542
- Aswathi, P. V. and Dhivya, R. (2017). Quality Phytochemical Screening and Mosquito Replelency of *Chromolaenaodonrata* leaf extract against adult *Culex sp.* *Indo American Journal of Pharmaceutical Science*, 4(3): 698-705.

- Danquah, C. A., Koffuor, G. A., Annan, K and Ketor., E. C. (2012) The anthelmintic activity of *Vernonia amygdalina* (Asteraceae) and *Alstoniaboonei* de wild (Apocynaceae). *J Med Biomed Sci* 1(1):21–27
- Eguale, T. and Giday, M. (2009) *In vitro* Anthelmintic Activity of Three Medicinal Plants against *Haemonchus contortus*. *International Journal of Green Pharmacy* 29-34
- Ejiofor, I.I., Zaman, K. and Das, A. (2017) Effect of Extracts of *Vernonia amygdalina* in *Helminthiasis*: A Troical Negelected Disease. *Open Access Journal of Pharmaceutical Research* 1(8): 147-151
- Ekekere, G. J. and Jeyachandran, R.(2000) phytochemical and antimicrobial efficacy of *Vernonia amygdalina*. *International journal of research in applied science and engineering technology* 7(4): 91-93
- El-Menyawe, S. M. and Mohammed A. (2015).The Review of *Toxocara* in Man and Animal. *Vet Parasitol* 12(3): 231-242
- Eslahi, A. V. Badri, M., Nahavandi, K. H., Houshmand, E., Dalvand, S. and Riahi, S. M. (2021) Occurrence of *Dictyophymerenela* in Road Killed Canids of Iran and its Public Health Implication. *Vet Parasitol* 24(1): 568-587
- Habluetzel, A. Traidi, C., Ruggieri, S., Attili, A. R., Scuppa, P., Marchetti, G., Esposito, F. (2003) An Estimation of *Toxocaracanis* Prevalence in Dogs, Environmental Egg Contamination and Risk of Human Infection in the Marche Region of Italy. *Vet Parasitol* 113: 243-252
- Healy, S. R., Morgan, E. R., Prada, J. M. and Betson, M. (2021) Brain food: rethinking food-borne toxocariasis. *Parasitology*. 16: 20-26.
- Huffman, M.A. and Seifu, M (1989) Observation on the Illness and Consumption of a Possibly Medicinal Plant *Vernonia amygdalina* (Del) by Wild Chimpanzee in the Mahale Mountains National park, Tanzania. *Primates* 30:51-63
- Ijeh, I. I. and Ejike, C. E. C. C. (2011) Current perspectives on the medicinal potentials of *Vernonia amygdalina* Del. *Journal of Medicinal Plant Resource* 5:1051-1061
- Kaur, D., Kaur, N., and Chopra, A. (2019). A comprehensive review on phytochemistry and pharmacological activities of *Vernonia amygdalina*. *Journal of Pharmacognosy and Phytochemistry*, 8, 2629-2636.
- Lescano S A., Nakhle., M. C., Ribeiro, M. C., Chieffi, P. P. (2012) IgG antibody responses in mice coinfectd with *Toxocaracanis* and other helminths or protozoan parasites. *Rev Inst Med Trop Sao Paulo*. 54:145–152

- Masaba, S. (2000). The Anthelmintic Activity of *Vernonica amygdalina Del* (Compositae). *Transaction of the Royal Society of Tropical Medicine and Hygiene*, 94(6): 694-670
- Merigueti, Y. F. F. B., Santarem, V. A., Ramires, L.M., Da silveira, B. A., DaCosta B. Eserra, L. V., Nuci, A.L., De Paula Esposte, T. M. (2017) protective and risk factors associated with the presence of Toxocarasp eggs in dog fur. *Vet parasitol* 244:39-43
- Morsy T. A. (2020)Toxocariasis: Visceral and Ocular Larva Migrans. *Journal of the Egyptian Society of Parasitology* 50(10): 41-18
- Nuryanto, M. K., Paramita, S and Iskandar. A.(2018). Membrane Stabilization Activity As Anti-Inflammatory Mechanisms Of *Vernonia Amygdalina* Leaves Extracts. *IOP Conf. Ser.: Earth Environ. Sci.* **144**
- Nweze, N. E., Arine, O. and Ngongeh, L.A. (2013). Anthelmintic Potential of Three Plants Used in Nigeria Ethnoveterinary Medicine. *Pharmaceutical Biology* 5(3): 311-315
- Oshiobugie, M. J., Olaniyi, A. M. and Raphael, A. O. (2017) AAS and GC-MS Analysis of Phytocomponents in the Leaf, Stem and Root of *Azadirachta indica A. Juss* (Dongoyaro). *British Journal of Pharmaceutical Research* 15(4): 1-12
- Oyeyemi, I. T., Akinlabi, A. A. Adewumi, A., Aleshinloye, A. O. (2018) *Vernonia amygdalina*: A Folkloric herb with anthelmintic properties. *International journal of advance res.* 9(4): 477-483
- Salien, A. M., Kassuku, A. A., Magwisha, H., Edward, C., Marisa, J. and Ngovi, C. (2019). Determination of Anthelmintic Resistance in Goats and Sheep Using Faecal Egg Count Reduction Test At Luguruni Farm, Dares Salaam, Tanzania. *Tanzania Veterinary Association Proceedings*, 34:87-90
- Sujon, M. A., Mostofa, M., Jahan, M. S., Das, A. R. and Rob, S. (2008). Studies on Medicinal Plants against Gastrointestinal Nematodes of Goats. *Bangl. J. Vet. Med.*, 6 (2): 179–183
- Tandon, V., Yadav, A. K., Roy, B. and Das, B. (2011) *Emerging Trends in Zoology: Phytochemicals as Cure for Worm Infections in Traditional Medicine System*. Narendra Publication House, New Delhi. 351-378
- Udoigung, N. I., Adams, E. G., Ekpo, E. J., and Opara, K. N. (2018) A Survey on Intestinal Nematodes of Dogs in Uyo, Akwa Ibom State, Nigeria. *Journal of Parasitology and Vector Biology* 10(3): 39
- Ujah, I.I. Nsude, C.A.Ani, O.N. Alozieuwa, U.B., Okpako, I.O. and Okuwur, A. E. (2021) Phytochemical of Neem plant (*Azadirachta indica*) explains its use in traditional medicine and pest control. *GCS Biological and Pharmaceutical science* 14(2): 163-171

UNDER PEER REVIEW