

# HEALTH EDUCATION IMPACT ON STIGMA AND DISCRIMINATION AGAINST PEOPLE LIVING WITH HUMAN IMMUNODEFICIENT VIRUS INFECTION AMONG ADOLESCENTS IN SOUTH WESTERN NIGERIA- A COMMUNITY BASED EXPERIMENTAL STUDY

## ABSTRACT

**Introduction:** Young people in Nigeria constitute an important group due to their vulnerability and sexual inexperience and there are still relatively high levels of misconceptions and stigma. This study was therefore carried out to investigate effect of health education on stigma and discrimination of PLWHAs among adolescents in Ogun State, Nigeria.

**Methodology:** This was a quasi-experimental community-based study. The study was carried out in three (3) phases. A multistage random sampling technique was used in choosing the required samples for this study. Two Local Governments which was selected in the State formed the experimental and control groups. A semi-structured pretested interviewer-administered questionnaire was used for data collection. Data analysis was done using SPSS statistical software version 19.

**Results:** A total of 215 participants were recruited into the study, 31.2% were males, 52.6% were adolescents aged 10-14 years and 47.4% were aged 15-19 years, while 67.3% were Christians, both the control and experimental groups were similar [ $p=0.128$ ]. Stigma scores at baseline were also correlated with knowledge scores. There was a weak negative but significant correlation between knowledge and stigma ( $r = -0.27, p < 0.001$ ). There was a significant reduction in stigma scores for both the intervention ( $t = 4.92, p < 0.001$ ) and control groups at 4 months post intervention ( $t = 4.33, p < 0.001$ ). There was only a significant difference for family type, where the mean stigma scores were higher among respondents from polygamous homes and those with widowed parents compared to those from monogamous homes ( $F = 3.22, p = 0.008$ ).

**Conclusion:** The study shows that adolescents from dysfunctional families such as polygamous homes and widowed parents had a significantly higher **score**. Establishment of Youth Friendly

Clinics that would focus more on adolescents from dysfunctional homes is needed among adolescents in Nigeria and other low income population.

Key words: Stigma, Health Education intervention, HIV/AIDS, Adolescents, Nigeria

## INTRODUCTION

Young people in Nigeria constitute an important group due to their vulnerability and sexual inexperience. Even though there is a high level of awareness about HIV among young people in Nigeria, there are still relatively high levels of misconceptions and stigma [1-2]. In a nationwide survey, only about a third of respondents had comprehensive knowledge about HIV while less than a quarter had accepting attitudes towards people living with HIV AIDS. Improvement of knowledge about HIV transmission and prevention and correction of stigma and discriminatory attitudes will help in the fight against HIV among young persons in Nigeria [3-4].

Stigma and discrimination are two major problems often faced by people living with HIV and AIDS in many developing countries, including Nigeria [3]. Stigma and discrimination shown to persons living with and affected by HIV and AIDS can worsen the spread and the impact of the HIV and AIDS epidemic. A major contributor to this stigma and discrimination is the criminalisation of high risk groups like commercial sex workers, drug addicts/abusers and men who have sex with men (MSMs) in our society which thus drives them underground and further increasing their vulnerability to HIV infection [4-6]. As a result of fear of discrimination and stigma, many individuals are afraid of seeking HIV testing to know their HIV status while persons living with HIV and AIDS (PLWHAs) may be less inclined to declare and openly acknowledge their HIV sero-status. This can lead to continued under-reporting of the epidemic, increased transmission, and limited access to treatment, care and support programmes. On the other hand, stigma and discrimination violate the human rights and dignity of people living with HIV and AIDS and those affected by the epidemic [4,7].

HIV/AIDS-related stigma and discrimination remains a major hindrance to knowing one's HIV status as well as accessing care if positive and remains one of the key drivers of the HIV epidemic in Nigeria which also include low personal risk perception (which is very common amongst adolescents) and multiple concurrent sexual partnerships[7-10].This notwithstanding,

there are few studies that quantifies the burden of effect of stigma and discrimination on access to care of HIV in Nigeria. Hence there is need for intervention studies on the adolescents that would focus on their development of accepting attitudes to people living with HIV/AIDS (PLWHAs). There is generally an increased awareness about HIV/AIDS among adolescents in Nigeria as evidenced by findings from recent national surveys [9-11].

Programmes aimed at reducing stigma and discrimination against people living with HIV or people at risk of HIV infection should address the actionable causes of stigma and discrimination and empower people living with and vulnerable to HIV. Use of media, including advertising campaigns, entertainment designed to educate as well as to amuse (“edutainment”), and integration of non-stigmatizing messages into TV and radio shows; this was reinforced by a Nigerian-based study to assess the relationship between media saturation, communication exposure and HIV stigma in Nigeria and found that accepting attitudes towards people living with HIV were more prevalent among men than among women [10-13]. Exposure to HIV-related communication on the media was associated with increased knowledge about HIV, which is in turn a strong predictor of accepting attitudes. Communication exposure also had a significant and positive association with accepting attitudes towards people living with HIV. In contrast, community media saturation was not strongly linked with accepting attitudes for either sex. The findings strongly suggest that media-based HIV programs constitute an effective strategy to combat HIV/AIDS-related stigma and should therefore be intensified in Nigeria.

Other effective communication strategies involve engagement with religious and community leaders, and celebrities; Inclusion of non-discrimination as part of institutional and workplace policies in employment and educational settings [10, 14-15]. Measurement of HIV-related stigma including in health care settings and communities; and Peer mobilization and support developed for and by people living with HIV aimed at promoting health, well-being and human right [11, 12]. This study was therefore carried out to investigate the effect of health education on stigma and discrimination among adolescents in Ogun state Nigeria.

## **METHODOLOGY**

### **Study Area**

The State is populated mainly by the Yoruba ethnic group with several sub-groups mainly The Sub-groups are mainly the Egba, Yewa, Awori, Egun, Ijebu, Remo, Ikale and Ilaje, Anago, Ketu and Ohori. There are also other non-Yoruba ethnic groups that settled in the State owing largely to its proximity to Lagos (the economic capital of Nigeria) and the presence of several industries. The occupations of the inhabitants are majorly farmers, civil servants, factory workers and self-employed businesses (e.g. traders, commercial transporters including motorcycle riders).

Each Local Government has at least a secondary healthcare facility as well as several primary health care facilities spread throughout the political wards in the LGA. There are three (3) tertiary health facilities in Ogun State: Olabisi Onabanjo University Teaching Hospital, Sagamu; Federal Medical Centre, Abeokuta and National Neuropsychiatric Hospital, Aro (Abeokuta). HIV Counselling and Testing (HCT) services are rendered in most of these health facilities as well as other reproductive health services. Ogun State presently has a total of ninety-five (95) HCT centres.

### **Study design**

The study was a quasi-experimental community-based study to determine the effect of health education on HIV/AIDS risk behaviours and stigmatisation practices among adolescents in the selected Local Governments in the State. Two Local Governments (Ado-Odo/Ota & Ifo), randomly selected from two of the senatorial zones in the State formed the experimental and control groups.

The study was carried out in three (3) phases – Pre-intervention, Intervention and Post-Intervention phases. Phase one (pre-intervention) involved cross-sectional comparative descriptive study, while phase two involved comprehensive health education intervention in the experimental/intervention group only. Phase three (post-intervention) involved comparative study between the experimental and control group immediate post-intervention as well as four (4) months post-intervention.

### **Pre-Intervention activities**

1. Constituting research team: the lead investigator was assisted by trained assistants which included a coordinator of a local community-based organisation (CBO) involved in

HIV/AIDS youth programmes; a trained peer educator; a secondary school teacher trained in family life and HIV education (FLHE); two (2) volunteer persons living positively with HIV (PLPs); a health worker in each of the selected Local Government Areas involved with HIV/AIDS programme.

2. House enumeration in the selected political wards for the study (was conducted in conjunction with town-planning officials of the LGA) as part of the process of selection of participants for the study. These were done between 2<sup>nd</sup> and 4<sup>th</sup> weeks in Dec. 2014.
3. Informed consent of the parents/guardians of the adolescents to allow them fully participate at all stages of the study were obtained.
4. Informed consent of the participants themselves were obtained (which emphasised their voluntary participation and permission to withdraw further participation from the study at any time if they so desired).
5. There was pre-testing of 20 questionnaires before the main study was conducted using some adolescents resident in Ipokia LGA which was different from the two Local Governments already selected for the main study. Amendments were made to the research instrument to minimise ambiguity and improve clarity. This was also to ensure validity and reliability of the research instrument.
6. There was a baseline survey to determine the knowledge of the adolescents in the two groups (i.e. control and intervention) about modes of transmission and prevention of HIV as well as their perception of risk of contracting the virus and their level of stigma towards persons living with HIV; this represented the pre-training assessment for the intervention group and the initial assessment for the control group.

### **Intervention**

Training sessions were for two days and were in four (4) modular units which were: (1) knowledge about HIV/AIDS and other STIs; (2) Personal Risk Perception; (3) HIV Status, Stigma and Discrimination and (4) Care and Support and Rights of People Living with HIV/AIDS (PLWHAs). The first two modules i.e. (1) and (2) above constituted the training for Day 1; while the latter two modules i.e. (3) and (4) above constituted the training for the second day. The baseline survey for the control group (in Ifo LGA) was held on Sat. 10<sup>th</sup> Jan. 2015 at Ilepa community hall at 3 – 5pm.

## Post-Intervention Activities

- These were conducted in two phases: immediate post-intervention and four (4) months post-intervention.
- The post-intervention evaluation was carried out using same questionnaire that had been used during the pre-intervention evaluation to determine immediate gain (immediate post-intervention) and the residual gain (four months post-intervention) in HIV/AIDS-related Knowledge Attitude and Practice (KAP) after the initial assessment in the intervention and control groups respectively.
- Evaluation of the effects of the training was done using calculated scores for the various variables during analysis.
- There was Focus Group Discussions (FGD) conducted in all the political wards used for the study. This was done soon after the immediate post-intervention activities were concluded.

## Sample size determination

The minimum sample size (n) was determined by the statistical formula for comparing proportions between two groups thus:

$$n = D(Z_{\alpha/2} + Z_{\beta})^2 \times \{P_1(1 - P_1) + P_2(1 - P_2)\} / (P_1 - P_2)^2$$

Where,  $Z_{\alpha/2}$  = Critical value of the standard normal deviate, ( $\alpha$ ) of 5% = 1.96.

$Z_{\beta}$  = Critical value of the standard normal deviate, type II error ( $\beta$ ) of 10% = 1.28.

$D$  = Design effect for the sampling design used = 1.5

- $P_1 - P_2$  = difference in stigma perception between the experimental and control groups to be detected = 20%

Hence,  $n = 1.5 (1.96 + 1.28)^2 \times \{0.638(1 - 0.638) + 0.438(1 - 0.438)\} / (0.638 - 0.438)^2$ .

$n = 1.5 \times 10.4976 \times 0.474232 / 0.04$ .

$n = 154.29 \equiv 155$ .

Hence, to cater for attrition rate of 10%, the minimum sample size is  $155 / (1 - 10\%) = 172$  for each group (i.e. intervention and control); Further adjustment for 10% non-response gives a

minimum of  $172/0.9 = 192$ . However in this study 250 participants were recruited in each group.

### **Subject Selection and Sampling Methods**

A multistage random sampling technique was used in choosing the required samples for this study. **Stage I (Senatorial zone selection):** Ogun West and Ogun Central senatorial zones were selected from 3 senatorial zones of the state by balloting

**Stage II (LGA selection):** From each of the two (2) selected senatorial zones, one LGA was selected by simple random sampling technique by balloting Hence Ado-Odo/Ota Local Government was selected to be the intervention (experimental) group whilst Ifo LGA the control.

**Stage III (Ward selection):** From the list of political wards in each of the selected LGAs, two (2) wards with contiguous borders were selected by simple random sampling technique Hence for the experimental LGA (Ado-Odo/Ota), of the sixteen (16) wards, Iju ward was first selected; A similar procedure was deployed in the control LGA (Ifo) in which of the eleven (11) wards, Olose ward was selected by balloting

**Stage IV (Houses selection):** At the level of the selected Wards, house enumeration was carried out by the research team and some officials from the town-planning unit of the Local Governments. From the combined total number of houses counted in the experimental and control Wards clusters (i.e. the two (2) wards selected in each LGA), a systematic random sampling technique (using a sample interval) was deployed to choose 250 houses in each of the experimental and control wards respectively. This sample interval was determined by dividing the total number of houses enumerated in both the experimental (4,016) and the control (2,857) wards by the sample size in the experimental and control wards respectively i.e. 250 {hence  $4,016/250$  and  $2,857/250$  for the experimental and control wards respectively}. Hence the sample interval was 17 and 12 for the experimental and control wards respectively. The first house was determined by using the table of random numbers to pick a house from the house enumeration list.

**Stage V (selection of individual adolescent participant):** Only one adolescent (age 10-19yrs) was studied per house and this was randomly selected by a simple random sampling technique carried out by balloting. When there was no adolescent in a selected house, then the next house

to it was considered but subsequent house selection was made without prejudice to predetermined sampling interval.

**Exclusion Criteria:** Only adolescents whose parents/guardians were permanent residents and who had been residing with them (parents/guardian) in the area for at least 6 months prior to the time of study were included in the study.

Consent was obtained to participate in the study from both the parents/guardian of the adolescents as well as the adolescents themselves. Others were excluded from the study.

### **Research Instrument and Data collection**

A semi-structured interviewer-administered questionnaire (adapted from the National HIV/AIDS and Reproductive Health Survey (NARHS Plus) 2007) was used for data collection (Appendix 1). This was pre-tested before the main study was conducted using some adolescents resident in Ipokia Local Government, Ogun State. Amendments were made to some aspects of the instrument that were found ambiguous or lacked clarity.

The sections of the questionnaire included: Section A consisting of socio-demographic characteristics. Section B which consists of questions on knowledge of the adolescents about modes of transmission and prevention of HIV; and Section C which consists of questions to assess their level of stigma and discrimination towards persons living with HIV.

### **Data management and analysis**

The questionnaires were checked for proper completion upon collection from participants. The data were entered into SPSS statistical software version 19. The data were cleaned for errors and data edited. Composite variables (aggregate scores) for knowledge and stigmatising attitudes were computed from items on the questionnaire. For knowledge about HIV, responses from 29 items were combined. These items included knowledge and misconceptions about HIV transmission, HIV prevention, and treatment for HIV positive individuals. A correct response was given a score of 1 and an incorrect one scored 0. Thus the 'yes' responses for the questions

on correct modes of transmission and prevention were scored 1 while this was reversed for the misconceptions. The total obtainable knowledge score was 29.

Concerning stigma and discrimination, stigmatizing attitudes got a score of 1 while accepting attitudes were scored 0. There were 16 items related to attitudes to family members, non-family members, classmates and teacher with HIV, thus the total obtainable stigma score was 16. Data were summarized using means and standard deviation for normally distributed quantitative data and median and range for skewed data. Qualitative data was summarized using frequencies and proportions. Baseline comparisons in the distributions of categorical socio-demographic variables and HIV knowledge, testing and stigma were tested using the Chi square test. The mean scores (knowledge and stigma scores) were compared between baseline and immediate post intervention for the intervention group and between baseline and 4 months post intervention for the two groups using the Paired t test.

In order to determine the association between stigma and socio-demographic characteristics, mean stigma scores were compared across levels of socio-demographic characteristics. For variables with two categories, independent samples t test was used to compare the groups while One-way ANOVA was used to compare variables with more than two categories. Multiple linear regression of stigma scores was also done on selected variables. The association between risky sexual behaviour and socio-demographic characteristics was tested using Chi square tests. Level of significance for all tests was 5%.

## RESULT

### **Socio-demographic characteristics**

Table 1 shows the distribution of socio-demographic characteristics between the intervention and control groups. This shows that 31.2% were males, 52.6% were adolescents aged 10-14 years and 47.4% were aged 15-19 years, while 67.3% were Christians. The two groups were similar concerning gender ( $X^2 = 2.31$ ,  $p = 0.128$ ), age ( $X^2 = 2.69$ ,  $p = 0.101$ ), parents' marital status ( $X^2 = 0.12$ ,  $p = 0.729$ ), family type ( $X^2 = 5.57$ ,  $p = 0.062$ ), mother's children ( $X^2 = 2.39$ ,  $p = 0.303$ ), father's children ( $X^2 = 1.50$ ,  $p = 0.473$ ), father's education ( $X^2 = 2.88$ ,  $p = 0.237$ ), mother's education ( $X^2 = 4.75$ ,  $p = 0.093$ ) and religion ( $X^2 = 1.16$ ,  $p = 0.282$ ).

### **Comparison of change in responses to knowledge items at 4 months between intervention and controls**

Knowledge about HIV was compared at baseline and 4 months post intervention for the intervention and control groups and the results are shown in Tables 2. Generally there were more significant improvements in knowledge at 4 months for the intervention compared to control groups. For example the knowledge about HIV transmission improved more for the intervention while misconceptions reduced among the intervention group compared to controls. In the intervention group, the proportion of respondents that thought an individual could contract HIV by sharing eating utensils significantly reduced from 24.4% to 11.1% ( $p = 0.001$ ). Similarly a lower proportion post intervention (4.9%) compared to baseline (12%) thought witchcraft was a mode of HIV transmission ( $p = 0.024$ ).

### **Comparison of change in stigma and discrimination at 4 months between intervention and controls**

Table 3, the changes in stigmatizing attitudes at 4 months are shown for the intervention and control groups. Compared to controls, the intervention group had significant improvements for those items related to attitudes to family member with HIV. For most other items there were no clear differences between the intervention and control groups.

Among the intervention group, the proportion of respondents willing to care for HIV positive relative significantly increased from 88.8% to 97.6% ( $p = 0.001$ ). Also the proportion willing to share meals with PLWHA increased from 52% to 69.7% ( $p = 0.002$ ). There were similar large improvements for willingness to sit beside PLWHA ( $p < 0.001$ ), and to buy food from food seller with HIV ( $p < 0.001$ ) among the intervention group.

Stigma scores significantly reduced immediate post intervention for the experimental group ( $t = 9.53$ ,  $p < 0.001$ ). There was also a significant reduction in stigma scores for both the intervention ( $t = 4.92$ ,  $p < 0.001$ ) and control groups at 4 months post intervention ( $t = 4.33$ ,  $p < 0.001$ ). The reduction in stigma scores were computed and compared between the intervention and control groups. There was no significant difference in the reduction in stigma scores between the intervention and control groups ( $p = 0.952$ ) (Table 2).

### **Association between stigma and socio-demographic characteristics (bivariate and multivariate analysis)**

The differences in stigma scores across levels of categorical variables are shown in Table 4. There was only a significant difference for family type, where the mean stigma scores were higher among respondents from polygamous homes and those with widowed parents compared to those from monogamous homes ( $F = 3.22$ ,  $p = 0.008$ ). Post hoc tests for pairwise comparisons only found significant differences between widowed parents and monogamous homes. Stigma scores at baseline were also correlated with knowledge scores. There was a weak negative but significant correlation between knowledge and stigma ( $r = -0.27$ ,  $p < 0.001$ ). The variables significant at 20% on t tests and F tests (gender, family type and father's education) were entered into a multiple linear regression. The regression coefficients and 95% confidence intervals are shown in Table 5. Respondents from polygamous homes or widowed parents had significantly higher mean stigma scores ( $p = 0.049$ ) compared to those from monogamous homes. Gender and father's education were not significant.

## DISCUSSION

Misconceptions about modes of HIV transmission certainly have a role to play in accepting attitudes towards those with the disease, thus efforts at reducing misconceptions are necessary in reducing stigma. In this study the intervention educated the students and tried to correct the misconceptions. Less than 80% knew about oral sex and anal sex as modes of HIV transmission. Though these modes of transmission become risky when there is no protection, campaign messages need to emphasize the high HIV risk associated with these sexual behaviours. The stigma associated with oral and anal sex especially in the traditional African society suggests that innovative approaches to discussion of oral and anal sex are needed. Compared to the modes of HIV transmission, the knowledge about HIV prevention was lower. In fact only avoidance of sharp objects was known by over 80% of the respondents. This study is unique because it is one of the very few studies in Nigeria with a main focus on the effect of an intervention on stigma. Several Nigerian studies have investigated the effect of different interventions on HIV AIDS knowledge, sexual behaviour, and knowledge of Reproductive Health issues such as premarital sex, pregnancy prevention, abortion and STIs. [16-22]

Regarding accepting attitudes towards people living with HIV (PLHIV) vis-a-vis stigma and discrimination, the respondents in this study had fair levels of accepting attitudes, as the mean

scores were 6 in the intervention and 8 among controls at baseline out of a total stigma score of 16. While there were very positive attitudes concerning caring for family member with HIV and reminding HIV positive friend to take drugs, the responses to most other items on attitude were less positive. In fact less than a quarter of respondents indicated willingness to buy food from HIV positive food seller. These are higher prevalence than those found in other studies [ 23-24] Generally less than two thirds would share meals with or sit beside HIV positive non-family member, share materials or writing materials with classmate. These results indicate that more campaigns are needed to correct these stigmatizing behaviours among adolescents.

Another related finding is the high proportion (about 70%) that would keep the status of a family member with HIV secret. Keeping the HIV status of a family member secret portends negative consequences for the HIV positive individual as early access to treatment and other benefits of HIV care are missed once HIV status is kept secret. A related finding is that less than two thirds will encourage family member to openly acknowledge HIV status. Interventions such as the one in this study, especially with new and innovative methodologies are urgently needed to correct several stigmatizing and discriminatory attitudes found among the adolescents in this study. It appears that the relatively high level of knowledge does not translate to accepting attitudes towards persons with HIV.

The study also revealed significant associations between stigma, family type and HIV knowledge. The lowest stigma scores were found among adolescents from monogamous homes compared to those from other homes. This finding could be an indication that those from monogamous homes are more tolerant, as they reside in homes with smaller families, with fewer siblings and perhaps better bonded with family members. Thus they may be more likely to care for family members and other people needing help. The inverse association between HIV knowledge and stigmatizing attitudes to PLWHA found in this study has been reported by previous studies among antenatal attendees in South-south region of Nigeria and among adults in Ghana [20-21]. It is apparent that lower levels of misconceptions about HIV transmission and prevention would translate to more tolerant behaviour towards PLWHA. The implication of this association is that efforts at improving HIV knowledge must be intensified, as stigma levels are also likely to reduce.

The association between stigma and father's education was not significant but lower stigma scores were found among those whose fathers had tertiary education. The educational level of

the parent is likely to positively influence the child's level of information about social and health issues generally and this could reduce their levels of stigma. Studies have also shown significantly lower levels of stigma among those with higher education [25-27]. Concerning gender differences in stigma, males had lower stigma scores than females, however the difference was not statistically significant. Previous studies in Nigeria have reported higher accepting attitudes among males [28-30].

The intervention instituted appeared beneficial among the intervention group especially when the responses immediate post-intervention are compared to baseline. The mean stigma score reduced from 6 at baseline to 2.8 immediate post, but rose to 3.7 at 4 months post intervention. Looking at some of the individual stigma items, the proportions willing to sit beside person with HIV, to buy food from HIV infected shopkeeper, to share writing materials, or share meals with classmate reduced between immediate post intervention and 4 months post. This indicates that some of the stigmatizing attitudes are ingrained in the minds of the adolescents and will require regular exposure to interventions to correct some of the stigmatizing attitudes. A previous systematic review has noted that the stigma reduction interventions usually work on a small scale and their impacts are short term [30-33]. One way to ensure regular exposure to education about HIV is the inclusion of HIV AIDS as compulsory subjects to be taken by all students for those in school, while regular community interventions should also focus on out of school adolescents. The intervention in this study didn't seem to markedly reduce stigma and more innovative interventions need to be tried in future studies. Some studies have highlighted the importance of community factors on accepting attitudes towards PLWHA [34-36]. This suggests that interventions targeting whole communities as well as individuals such as adolescents could be more successful in reducing stigma. Such large-scale interventions need to be tried by HIV AIDS programmes and governmental organizations. Furthermore, this study shows a significant loss to follow up, significant number of the participants did not report for the 4 month post intervention assessment. This is one of the limitations of this study which is expected in community based follow up studies.

## CONCLUSION

This study is one of the few interventions focusing on HIV stigma among adolescents. Stigma has an important role to play in the transmission of HIV and in the utilization of health care

services such as HIV testing. Innovative interventions and campaigns at the community level focusing on reducing stigmatization are needed that will consider the peculiar socio-cultural factors in the design of such interventions as dynamics of change in the community may be more important in shaping attitudes and behaviour rather than individual factors. Stigma reduction interventions need to focus more on adolescents from polygamous homes and those with poor HIV knowledge as this study has shown higher stigmatizing attitudes among them.

Establishment of Youth Friendly Clinics in the communities that would focus more on meeting the peculiar health needs of adolescents especially against the backdrop that in this study, it was a general consensus of the respondents that these facilities though very beneficial to them, were none existent in their communities. Government at all levels should ensure stricter enforcement and compliance with the various laws related to protecting people living with HIV/AIDS (PLWHAs) against stigma and discrimination as this can serve as a disincentive to those (persons, groups or corporate establishments) in the habit of meting out such treatment to these individuals. Government at all levels should begin to responsibly take ownership of all HIV/AIDS related programmes in their domain and channel resources at their disposal towards meeting the peculiar HIV/AIDS related health needs of their community members not least the adolescents

#### **ETHICAL APPROVAL AND CONSENT:**

Permission was sought from the community leaders in the study area for easy access to the community. Ethical approval was obtained from the Ethics Committee of the Federal Medical Centre (FMC), Abeokuta with No: FMC/470/HREC/07/2014. Written informed consent was obtained from the participants and their parents.

#### **REFERENCES.**

1. Nigeria National HIV AIDS and Reproductive Health Survey (NARHS PLUS) 2007:63-139.

2. Federal Ministry of Health [Nigeria] (2013). National HIV & AIDS and Reproductive Health Survey, 2012 (NARHS Plus II). Federal Ministry of Health Abuja, Nigeria.
3. National Agency for the Control of AIDS (NACA) Nigeria. Global AIDS Response Country Progress Report Nigeria GARPR 2015. 2015:15.
4. Babalola S. Readiness for HIV testing among young people in northern Nigeria: the roles of social norm and perceived stigma. *AIDS Behav.* 2007; 11(5):759-69
5. UNAIDS. Key programmes to reduce stigma and discrimination and increase access to justice in national HIV responses. 2012
6. Bassey EA, Abasiubong F, Ekanem U, Abasiatai AM. Attitude of antenatal attendees to people living with HIV/AIDS in Uyo, South-south Nigeria. *Afr Health Sci.* 2007; 7(4):239-43.
7. Visser MJ. HIV/AIDS prevention through peer education and support in secondary schools in South Africa. *SAHARA J.* 2007; 4(3):678-94.
8. Babalola S, Fatusi A, Anyanti J. Media saturation, communication exposure and HIV stigma in Nigeria. *Soc Sci Med.* 2009; 68(8):1513-20
9. Ogun State Specific AIDS Reproductive and Child Health Survey (SPARCS) 2011:3-12.
10. Marcu EA, Dinescu SN, Pădureanu V, Dumitrescu F, Diaconu R. Perinatal Exposure to HIV Infection: The Experience of Craiova Regional Centre, Romania. *Healthcare (Basel).* 2022 Feb 6;10(2):308. doi: 10.3390/healthcare10020308.
11. Moher D, Dulberg CS, Wells GA. Statistical Power, Sample Size and their reporting in randomised controlled trials. *JAMA.* 1994; 272:122-124.
12. Esiet AO, Esiet U, Philliber S, Philliber WW. Changes in knowledge and attitudes among junior secondary students exposed to the family life and HIV education curriculum in Lagos State, Nigeria. *Afr J Reprod Health.* 2009 Sep; 13(3):37-46.
13. Akpabio II, Asuzu MC, Fajemilehin BR, Ofi B. Effects of parental involvement in HIV/AIDS preventive education on secondary student knowledge about transmission and

prevention in Akwa Ibom State, Nigeria. *Int Q Community Health Educ.* 2008-2009. 29(1):71-87.

14. Cai Y, Hong H, Shi R, Ye X, Xu G, Li S, Shen L. Long-term follow-up study on peer-led school-based HIV/AIDS prevention among youths in Shanghai. *Int J STD AIDS.* 2008; 19(12):848-50.
15. Fawole IO, Asuzu MC, Oduntan SO, Brieger WR. A school-based AIDS education programme for secondary school students in Nigeria: a review of effectiveness. *Health Educ Res.* 1999; 14(5):675-83.
16. Ye XX, Huang H, Li SH, Xu G, Cai Y, Chen T, Shen LX, Shi R. HIV/AIDS education effects on behaviour among senior high school students in a medium-sized city in China. *Int J STD AIDS.* 2009 Aug; 20(8):549-52. doi: 10.1258/ijsa.2008.008471
17. Huang H, Ye X, Cai Y, Shen L, Xu G, Shi R, Jin X. Study on peer-led school-based HIV/AIDS prevention among youths in a medium-sized city in China. *Int J STD AIDS.* 2008 May; 19(5):342-6. doi: 10.1258/ijsa.2007.007208
18. Cheruiyot C, Magu D, Mburugu P, Sagwe D. Uptake and utilization of institutional voluntary HIV testing and counseling services among students aged 18-24 in Kenya's public Universities. *Afr Health Sci.* 2019 Dec;19(4):3190-3199. doi: 10.4314/ahs.v19i4.42
19. Bamise OF, Bamise CT, Adedigba MA. Knowledge of HIV/AIDS among secondary school adolescents in Osun state, Nigeria. *Niger J Clin Pract.* 2011; 14(3):338-44.
20. National Population Commission (NPC) [Nigeria] and ICF International. 2014. Nigeria Demographic and Health Survey 2013. Abuja, Nigeria, and Rockville, Maryland, USA: NPC and ICF International.
21. Tenkorang EY1, Owusu AY. Examining HIV-related stigma and discrimination in Ghana: what are the major contributors? *Sex Health.* 2013 Jul; 10(3):253-62. doi: 10.1071/SH12153.

22. Chovwen CO, Ita M. Influence of gender, self-consciousness and stigmatisation on perceived acceptance among people living with HIV in Nigeria. *Afr J AIDS Res.* 2007 Apr; 6(1):87-90. doi: 10.2989/16085900709490402.
23. Brown L1, Macintyre K, Trujillo L. Interventions to reduce HIV/AIDS stigma: what have we learned? *AIDS Educ Prev.* 2003 Feb; 15(1):49-69.
24. Apinundecha C1, Laohasiriwong W, Cameron MP, Lim S. A community participation intervention to reduce HIV/AIDS stigma, Nakhon Ratchasima province, northeast Thailand. *AIDS Care.* 2007 Oct; 19(9):1157-65.
25. French H, Greeff M, Watson MJ, Doak CM. A comprehensive HIV stigma-reduction and wellness-enhancement community intervention: a case study. *J Assoc Nurses AIDS Care.* 2015 Jan-Feb; 26(1):81-96. doi: 10.1016/j.jana.2014.03.007. Epub 2014 Apr 4.
26. Catalani C, Castaneda D, Spielberg F. Development and Assessment of Traditional and Innovative Media to Reduce Individual HIV/AIDS-Related Stigma Attitudes and Beliefs in India. *Front Public Health.* 2013 Jul 22; 1:21. doi: 10.3389/fpubh.2013.00021. eCollection 2013.
27. Ishikawa N1, Pridmore P, Carr-Hill R, Chaimuangdee K. The attitudes of primary schoolchildren in Northern Thailand towards their peers who are affected by HIV and AIDS. *AIDS Care.* 2011 Feb; 23(2):237-44. doi: 10.1080/09540121.2010.507737.
28. Ye XX, Huang H, Li SH, Xu G, Cai Y, Chen T, Shen LX, Shi R. HIV/AIDS education effects on behaviour among senior high school students in a medium-sized city in China. *Int J STD AIDS.* 2009 Aug; 20(8):549-52. doi: 10.1258/ijsa.2008.008471
29. Babalola S. Readiness for HIV testing among young people in northern Nigeria: the roles of social norm and perceived stigma. *AIDS Behav.* 2007; 11(5):759-69
30. Chiao C1, Mishra V, Sambisa W Individual- and community-level determinants of social acceptance of people living with HIV in Kenya: results from a national population-based survey. *Health Place.* 2009 Sep; 15(3):712-20. doi: 10.1016/j.healthplace.2008.12.001.
31. Ehiri JE, Anyanwu EC, Donath E, Kanu I, Jolly PE. AIDS-related stigma in sub-Saharan Africa: its contexts and potential intervention strategies. *AIDS Public Policy J.* 2005 Spring-Summer; 20(1-2):25-39.

32. Odimegwu CI, Adedini SA, Ononokpono DN. HIV/AIDS stigma and utilization of voluntary counselling and testing in Nigeria. *BMC Public Health*. 2013 May 13; 13:465. doi: 10.1186/1471-2458-13-465
33. Arnold R, Maticka-Tyndale E, Tenkorang E, Holland D, Gaspard A, Luginaah I, HP4RY Team. Evaluation of school- and community-based HIV prevention interventions with junior secondary school students in Edo State, Nigeria. *Afr J Reprod Health*. 2012 Jun; 16(2):103-25
34. Ajuwon AJ, Brieger WR. Evaluation of a school-based reproductive health education program in rural South Western, Nigeria. *Afr J Reprod Health*. 2007; 11(2):47-59.
35. Mba CI, Obi SN, Ozumba BC. The impact of health education on reproductive health knowledge among adolescents in a rural Nigerian community. *J ObstetGynaecol*. 2007; 27(5):513-7.
36. Okanlawon FA, Asuzu MC. Effect of peer education intervention on secondary school adolescents' reproductive health knowledge in Saki, Nigeria. *Afr J Med Sci*. 2011; 40(4):353-60.

Table 1: Comparison of socio-demographic characteristics of adolescents

Variable	Intervention (n=215)	Control (n=215)	Chi square	P value
Gender				
Male	67(31.2)	82(38.1)	2.31	0.128
Female	148(68.8)	133(61.9)		
Age (years)				
10-14	113(52.6)	96(42.4)	2.69	0.101
15-19	102(47.4)	119(55.3)		
Parents' marital status				
Currently married			0.12	0.729
Others	175(81.8) 39(18.2)	173(80.5) 42(19.5)		
Family type				
Monogamous	161(75.2)	143(66.8)	5.57	0.062
Polygamous	48(22.4)	58(27.1)		
Widowed parent	5(2.3)	13(6.1)		
Mother's children				
1-3	70(32.6)	68(31.8)	2.39	0.303
4-5	119(55.3)	109(50.9)		
6+	26(12.1)	37(17.3)		
Father's children				
1-3	63(29.6)	53(25.1)	1.50	0.473
4-5	91(42.7)	90(42.7)		
6+	59(27.7)	68(32.2)		
Father's education				
Primary and below	33(15.5)	38(17.8)	2.88	0.237

Secondary	71(33.3)	84(39.3)		
Tertiary	109(51.2)	92(43.0)		
Mother's education				
Primary and below	49(22.9)	54(25.1)	4.75	0.093
Secondary	75(35.0)	92(42.8)		
Tertiary	90(42.1)	69(32.1)		
Religion				
Islam	70(32.7)	81(37.7)	1.16	0.282
Christianity	144(67.3)	134(62.3)		

Table 2: Comparison of knowledge and misconceptions about HIV transmission at baseline and 4 months post intervention for the intervention and control groups

Variable	Intervention			Control		
	Baseline	4 months post*	P value	Baseline	4 months post*	P value
Sexual intercourse						
Yes	183(100)	179(97.8)	0.989	186(96.9)	186(96.9)	0.999
No	0	4(2.2)		6(3.1)	6(3.1)	
Blood transfusion						
Yes	179(97.8)	177(96.7)	0.754	184(95.8)	182(94.8)	0.815
No	4(2.2)	6(3.3)		8(4.2)	10(5.2)	
Mother to unborn child						
Yes	147(81.2)	142(78.5)	0.597	153(79.7)	138(71.9)	0.091
No	34(18.8)	39(21.5)		39(20.3)	54(28.1)	
Oral sex						
Yes	116(64.1)	131(72.4)	0.159	155(80.7)	135(70.3)	0.024
No	65(35.9)	50(27.6)		37(19.3)	57(29.7)	
Sharing sharp objects like razor						
Yes	173(95.6)	175(96.7)	0.791	187(97.4)	182(95.3)	0.424
No	8(4.4)	6(3.3)		5(2.6)	10(4.7)	
Sharing needles						
Yes	172(95.0)	180(99.4)	0.021	188(97.9)	186(96.9)	0.754
No	9(5.0)	1(0.6)		4(2.1)	6(3.1)	
Anal sex						
Yes	151(85.3)	150(84.7)	0.999	154(80.2)	164(85.4)	0.237
No	26(14.7)	27(15.3)		38(19.8)	28(14.6)	

<b>Misconceptions</b>						
<b>Sharing toilets</b>						
Yes	44(24.0)	30(16.4)	0.087	77(40.1)	49(25.5)	0.003
No	139(76.0)	153(83.6)		115(59.9)	143(74.5)	
<b>Sharing eating utensils</b>						
Yes	44(24.4)	20(11.1)	0.001	102(53.1)	29(15.1)	<0.001
No	136(75.6)	160(88.9)		90(46.9)	163(84.9)	
<b>Mosquito bites</b>						
Yes	90(49.2)	40(21.9)	<0.001	109(56.8)	96(50.0)	0.218
No	93(50.8)	143(78.1)		83(43.2)	96(50.0)	
<b>Witchcraft</b>						
Yes	22(12.0)	9(4.9)	0.024	7(3.6)	20(10.4)	0.015
No	161(88.0)	174(95.1)		185(96.4)	172(89.6)	
<b>Kissing</b>						
Yes	58(31.7)	68(37.2)	0.302	90(46.9)	54(28.1)	<0.001
No	125(68.3)	115(62.8)		102(53.1)	138(71.9)	
<b>Hugging</b>						
Yes	18(9.8)	8(4.4)	0.076	26(13.5)	11(5.7)	0.017
No	165(90.2)	175(95.6)		166(86.5)	181(94.3)	

\*At 4 months post intervention there were 183 participants in the intervention group and 192 in the control group that remained in the study

\*\*For the question on anal sex, 177 respondents answered

Table 3: Comparison of responses to stigma and discrimination items at baseline and 4 months post intervention among the intervention and control groups

<b>Variable</b>	<b>Intervention</b>			<b>Control</b>		
	<b>Baseline</b>	<b>4 months post</b>	<b>P value</b>	<b>Baseline</b>	<b>4 months post</b>	<b>P value</b>
<b>Desirable attitudes</b>						
<b>Willing to care for family member with HIV</b>						
Yes	150(88.8)	165(97.6)	0.001	174(91.6)	181(95.3)	0.210
No	19(11.2)	4(2.4)		16(8.4)	9(4.7)	
<b>Willing to share meals with PLWHA</b>						
Yes	91(52.0)	122(69.7)	0.002	52(27.8)	62(33.2)	0.343

No	84(48.0)	53(30.3)		135(72.2)	125(66.8)	
Willing to sit beside PLWHA						
Yes	110(65.5)	142(84.5)	<0.001	82(43.2)	124(65.3)	<0.001
No	58(34.5)	26(15.5)		108(56.8)	66(34.7)	
Willing to buy food from food seller with HIV						
Yes	39(23.1)	75(44.4)	<0.001	19(10.6)	23(12.8)	0.627
No	130(76.9)	94(55.6)		161(89.4)	157(87.2)	
Agree that HIV positive female teacher be allowed to teach						
Yes	126(69.6)	147(81.2)	0.015	89(47.1)	149(78.8)	<0.001
No	55(30.4)	34(18.8)		100(52.9)	40(21.2)	
Willing to sit with HIV positive classmate						
Yes	120(67.4)	147(82.6)	0.001	90(47.1)	122(63.9)	0.003
No	58(32.6)	31(17.4)		101(52.9)	69(36.1)	
Willing to share writing materials with HIV positive classmate						
Yes	117(66.1)	129(72.9)	0.188	80(42.6)	104(55.3)	0.021
No	60(33.9)	48(27.1)		108(57.4)	84(44.7)	
Willing to play with HIV positive classmate						
Yes	135(78.5)	146(84.9)	0.152	85(44.5)	125(65.4)	<0.001
No	37(21.5)	26(15.1)		106(55.5)	66(34.6)	
Willing to share meals with HIV positive classmate						
Yes	92(53.8)	122(71.3)	0.001	45(23.6)	79(41.4)	0.001
No	78(46.2)	49(28.7)		146(76.4)	112(58.6)	
Willing to share toilet with HIV positive classmate						
Yes	113(66.1)	124(72.5)	0.222	44(23.0)	78(40.8)	0.001
No	58(33.9)	47(27.5)		147(77.0)	113(59.2)	
Willing to help a support group of HIV positive people						
Yes	147(81.2)	156(86.2)	0.253	144(75.4)	168(88.0)	0.002
No	34(18.8)	25(13.8)		47(24.6)	23(12.0)	

Willing to participate in a support group meeting						
Yes	112(61.9)	150(82.9)	<0.001	72(37.9)	109(57.4)	0.001
No	69(38.1)	31(17.1)		118(62.1)	81(42.6)	
Willing to remind HIV positive friend/family member to take drugs						
Yes	171(94.5)	177(97.8)	0.180	178(92.7)	178(92.7)	0.999
No	10(5.5)	4(2.2)		14(7.3)	14(7.3)	
<b>Undesirable attitudes</b>						
Will stop being friends if found out close friend has HIV						
Yes	36(19.7)	21(11.5)	0.036	73(39.5)	26(14.1)	<0.001
No	147(80.3)	162(88.5)		112(60.5)	159(85.9)	

Table 4: Comparison of stigma scores across categories of socio-demographic variables for all participants (intervention and controls combined) at baseline

Variable	Mean stigma score	SD	N	Independent samples t OR F test**	P value
Gender					
Male	6.74	4.17	135	1.57	0.118
Female	7.40	3.84	255		
Age (years)					
10-14	7.10	3.83	185	0.33	0.744
15-19	7.23	4.07	205		
Parents' marital status					
Currently married	7.13	4.00	312	0.52	0.605
Others	7.39	3.84	77		
Family type					
Monogamous	6.93	4.02	274	3.22	0.041*
Polygamous/Widowed parent	8.85	3.76	114		
Mother's children					

1-3	7.39	3.89	127	0.27	0.761
4-5	7.05	4.06	204		
6+	7.16	3.87	58		
<b>Father's children</b>					
1-3	7.01	3.92	107	0.18	0.832
4-5	7.16	4.08	161		
6+	7.33	3.88	118		
<b>Father's education</b>					
Secondary and below	7.49	4.10	208	1.64	0.173
Tertiary	6.79	3.92	179		
<b>Mother's education</b>					
Primary and below	7.25	3.99	100	0.07	0.934
Secondary	7.08	4.00	146		
Tertiary	7.21	3.95	143		

\*Post hoc tests showed significant differences between participants from monogamous homes and those with widowed parents ( $p = 0.017$ )

\*\*Independent samples t test was used to compare mean stigma scores of dichotomous variables while the F test was used to compare the means of variables with more than two categories (multichotomous variables).

**Table 5: Multiple linear regression of stigma scores at baseline on variables**

<b>Variable</b>	<b>Regression coefficient</b>	<b>95% CI for regression coefficient</b>	<b>P value</b>
<b>Gender</b>			
Female	0.52	-0.21 to 1.27	0.161
Male			
<b>Family type</b>			
Polygamous/Widowed parent	0.83	0.01 to 1.67	0.049
Monogamous			
<b>Father's education</b>			
Secondary and below	0.71	-1.46 to 0.04	0.065
Tertiary			