

## **Population dynamics of mustard aphid, *Lipaphis erysimi* (Kaltenbach) on cauliflower in relation to biotic and abiotic factors**

### **Abstract**

The field experiment were conducted at Horticulture farm, S.K.N. College of Agriculture, Jobner during *Rabi* 2021-22 to know the population dynamics of mustard aphid, *Lipaphis erysimi* (Kaltenbach). The incidence of aphid started in the first week of December and second week of December remained active throughout the crop season during both the years . The population of aphid gradually increased and reached to its peak in the second week of February and first week of February in *Rabi*, 2021-22 respectively. The correlation studies indicated that the aphid population had non significant correlation with temperature, relative humidity and sunshine hours in both the years. Both the predators, ladybird beetle and syrphid fly had significant positive relationship with aphid population during both the years.

**Keywords:** Aphid, Correlation, Cauliflower.

### **Introduction**

Cauliflower (*Brassica oleracea* var. *botrytis* Linn) is one of the most significant vegetable crops in India and the rest of the world. It is a member of the *Brassicaceae* family. In India, cauliflower ranks third in term of vegetable production. The centre of origin is Mediterranean coastal area (Choudhary, 1996). The area under cauliflower in India is 473 thousand hectares with a production of 9283 metric tonnes (Anonymous, 2021-22). Cauliflower is mostly grown in Uttar Pradesh, Bihar, Assam, West Bengal, Maharashtra and Haryana states. India loses about 30% of its crops because of pests and diseases every year (Deeplata Sharma and Rao, 2012). cauliflower aphid, *Lipaphis erysimi* Kalt. (Aphididae: Hemiptera) are the most important pests causing severe yield loss to cauliflower every 12 months. Aphids are most communal and destructive pests of brassicaceous crops internationally and often cause heavy losses in yield (Shylesha *et al.*,

2006). Due to varying climatic conditions and shifting pest status, the population dynamics of significant insect pests are crucial. The peak period for bug pest interest provides ideas for creating defences against them. Determining how weather variables affect the frequency of insect pests on cauliflower will help to create a forecasting system that will enable the use of scheduled preventative measures. Natural enemies such as predators and parasitoids provide the important surroundings carrier of conservation via biological manipulate (Gardiner *et al.*, 2011).

### **Material and Methods**

The experimental study conducted at Horticulture farm, S.K.N. College of Agriculture, Jobner (Rajasthan) during *Rabi*, 2021-22 to know the population dynamics of aphid on cauliflower. The cauliflower crop, variety F<sub>1</sub> hybrid cauliflower 456 was transplanting on 16<sup>th</sup> November 2021-22 and 17<sup>th</sup> November 2022 in five plots. The plot size was 3.0 m x 2.25 m with row to row and plant to plant distance of 60 cm x 45 cm, respectively.

### **Method of observations**

The estimation of aphid population was based on the numerical count method as described by Lal (1998). The ten plants of cauliflower were randomly selected from each plot and tagged. Total number of aphids on ten plants were counted visually with the help of magnifying lens at weekly interval and converted into aphids per plant. For recording the aphid population at early plant stage, leaves were grasped at the petiole by thumb and four fingers and twisted until entire underside of the leave clearly visible.

### **Interpretation of data**

To interpret the results of population dynamics of aphid on cauliflower and its natural enemies the simple correlation was computed between aphid population, natural enemies and abiotic factors, *i.e.* minimum & maximum temperatures, relative humidity and sunshine hours was worked out using following formula. The following formula was used for calculating the correlation coefficient (Gupta, 1996).

$$r = \frac{N \sum xy - (\sum x)(\sum y)}{\sqrt{N \sum x^2 - (\sum x)^2 \cdot N \sum y^2 - (\sum y)^2}}$$

Where,

r = Simple correlation coefficient

x = Independent variables *i.e.* abiotic components

y = Dependent variables *i.e.* pests

N = Number of observations

## Results and Discussion

The mean population of aphid and the standard meteorological week wise weather parameters are presented in Table 1 (*Rabi, 2021-22*) and Table 2 (*Rabi, 2022*). The findings of the present study and the related discussion are explained hereunder. The infestation of aphid, *L. erysimi* commenced in 49<sup>th</sup> and 50<sup>th</sup> SMW (Standard Meteorological Week) *i.e.* first week of December and second week of December remained active throughout the crop season during the years 2021-22 respectively. Initially, the population of aphid was 7.40 and 9.40 per plants which gradually increased and reached to its peak with 161.20 and 269.80 aphids per plant in the second week of February (7<sup>th</sup> SMW) and first week of February (6<sup>th</sup> SMW) in *Rabi, 2021-22* respectively. After reaching the peak, the population of aphid started to decline during both the years. The current findings are in accordance with those of Bhavani and Punnaiah (2006), Varmora *et al.* (2009), Patra *et al.* (2013), Saranya *et al.* (2013), Sahu *et al.* (2017), Dwivedi *et al.* (2018), Mishra *et al.* (2018), Yadav *et al.* (2018), Lal *et al.* (2020), Pradhan *et al.* (2020), Arvind (2021) and Amarchand *et al.* (2022), who were reported the incidence of aphid started from the different weeks of December and the peak population of aphid from the February month which support the present results. The somewhat variation in commencement of

incidence and peak period as reported by above researchers might be due to the difference in agro climatic conditions of the locality and time of sowing.

The aphid population had non significant negative correlation with maximum temperature ( $r = -0.169$  and  $r = -0.308$ ), minimum temperature ( $r = -0.226$  and  $r = -0.020$ ) in both the year. The relative humidity showed non significant negative correlation ( $r = -0.075$ ) in the first year (2021-22) while, non significant positive correlation ( $r = 0.423$ ) in the second year (2022-23). The sunshine hours showed non significant positive correlation ( $r = 0.379$ ) in the first year (2021-22) while, non significant negative correlation ( $r = -0.297$ ) in the second year (2022-23). The aphid population had positive significant correlation with ladybird beetle ( $r = 0.929$  and  $r = 0.967$ ) and syrphid fly ( $r = 0.917$  and  $r = 0.962$ ) in both the years.

The present investigation on association of aphid population with the biotic and abiotic factors are in conformity with the findings of Varmora *et al.* (2009) was reported non significant negative correlation with maximum temperature and bright sunshine hours and relative humidity showed non significant positive correlation with pest population. Arvind (2021) reported the aphid population showed non- significant negative correlation with minimum and maximum temperature and non significant positive correlation with relative humidity.

Bana *et al.* (2012) reported that relative humidity and sunshine hours showed non significant correlation. Pradhan *et al.* (2020) reported relative humidity had non significant negative correlation with aphid population. Amarchand *et al.* (2022) reported that minimum temperature had non significant negative correlation, whereas, relative humidity had non significant positive with aphid population. Jandial and Kumar (2007) reported that non significant correlation existed between maximum and minimum temperature, relative humidity and aphid population. Triphati *et al.* (2005), Varmora *et al.* (2009) Pardhan *et al.* (2020) Amarchand *et al.* (2022), Dotsara *et al.* (2022) were found that the population of ladybird beetle and

syrphid fly had a significant positive correlation with aphid population and these findings are full support with present findings.

### **Natural enemies**

The incidence of ladybird beetle, *C. septempunctata* and syrphid fly, *Xanthogramma spp.* commenced in first week of December and second week of December in both the years. The peak population of ladybird beetle and syrphid fly in the second week of February and first week of February during both the years. The correlation studies indicated that the ladybird beetle and syrphid fly population had non significant correlation with all the weather parameters during both the years.

The current findings are in accordance with those of Vakaria and Patel (2005) and Sharma *et al.* (2022) who were reported the incidence of ladybird beetle started from the first week of December and the peak population of ladybird beetle from the third week of February and supports the present results. Patra *et al.* (2013), Dwivedi *et al.* (2018) and Amarchand *et al.* (2022) reported that the *C. septempunctata* and syrphid fly population was maximum in the third and second week of December. Bana *et al.* (2012), Mishra *et al.* (2018) and Sharma *et al.* (2022) reported that the *C. septempunctata* population was maximum in the third and fourth week of January. The work of studies on correlation of predators population with the weather parameters are supported by the findings of Arvind (2021) Amarchand *et al.* (2022) and Dotasara *et al.* (2022) reported that the non significant correlation with all weather parameters.

### **Conclusion**

The important conclusions drawn from present investigation made on population dynamics of aphid and its natural enemies. The infestation of aphid started in the first week of December and second week of December remained active throughout the crop season during both the years . The population of aphid gradually increased and reached to its peak in the second week of February and first week of February in *Rabi*, 2021-22 respectively. The aphid, ladybird beetle and syrphid fly had non significant

correlation with maximum temperature, minimum temperature relative humidity and sunshine hours during both the years. Both the predators ladybird beetle and syrphid fly had significant positive relationship with aphid population during both the years.

## References

- Amar Chand, Khinchi S.K., Kumawat K.C., Hussain A. and Sharma S.L. 2022. Quantitative and qualitative status of insect pests of mustard, *Brassica juncea* (L.) Czern and Coss and their natural enemies. *The Pharma Innovation Journal*, 11 (1): 412-415.
- Anonymous, 2021-22. Indian Horticulture Database. National Horticulture Board. Gurgaon, pp. 1& 2.
- Arvind 2021. Seasonal abundance and eco-friendly management of major insect pests of mustard, *Brassica juncea* (Linn.) Czern and Coss. M.Sc. (Ag.) thesis, submitted to SKNAU, Jobner.
- Bana J.K., Jat B.L. and Bajya D.R. 2012. Seasonal incidence of major pests of cabbage and their natural enemies. *Indian Journal of Entomology*, 74 (3): 236-240.
- Bhavani B. and Punnaiah K.C. 2006. Population dynamics of cabbage aphid, *Lipaphis erysimi* (Kalt.) and its control with selected insecticides in cabbage. *Pestology*, 30 (11): 24-29.
- Choudhary B. 1996. In "vegetables". Published by the director, National book trust, India: 72-74.
- Dotasara S.K., Kumawat K.C. and Swami D. 2022. Population dynamics of mustard aphid *Lipaphis erysimi* (kaltenbach) and its natural enemies. *Indian Journal of Entomology*, 84 (4): 979-980.
- Dwivedi S.A., Singh R.S. and Gharde S.K. 2018. Populations build up of mustard aphid and their natural enemies in relation to biotic and abiotic factors. *Plant Archives*, 18 (2): 2495-2500.

- Gardiner M.M., Fiedler A.K., Costamagna A.C. and Landis D.A. 2011. Integrating conservation biological control into IPM systems. In: Radcliffe E.B., Hutchison W.D. (Eds.), Integrated Pest Management. Cambridge University Press, Cambridge, pp. 151–162.
- Gupta S.C. 1996. Correlation, Fundamentals of Statistics. Himalaya Publishing House, Mumbai, 510-587.
- Jandial V.K. and Kumar A. 2007. Seasonal incidence and population fluctuation of mustard aphid, *Lipaphis erysimi* (Kalt.) in relation to ecological parameters. *Indian Journal of Entomology*, 69: 162-167.
- Lal J., Swaminathan R., Meena A.K. and Nagar R. 2020. Seasonal incidence of major insect pests of cabbage, *Brassica oleracea* var. *capitata* L, *Journal of Entomology and Zoology Studies*, 8 (3): 387-391.
- Lal O.P. 1998. Notes summer school on “Advance Technologies in Important Vegetable Crops, including Cole Crops”. May 4-24, I.A.R.I. New Delhi, pp. 63-66.
- Mishra S.K. and Kanwat P.M. 2018. Seasonal incidence of mustard aphid, *Lipaphis erysimi* (Kalt) and its major predator on mustard and their correlation with abiotic factors. *Journal of Entomology and Zoology Studies*, 6 (3): 831-836.
- Patra S., Dhote V.W., Alam K.F., Das B.C., Chatterjee M.L. and Samanta A. 2013. Population dynamics of major insect pests and their natural enemies on cabbage under new alluvial zone of West Bengal. *The Journal of Plant Protection Sciences*, 5 (1): 42-49.
- Pradhan P.P., Borkakati R.N. and Saikia D.K. 2020. Seasonal incidence of insect pests and natural enemies of mustard in relation to meteorological parameters. *Journal of Entomology and Zoology Studies*, 8 (1): 1538-1542.
- Saranya V.S.L., Rana B.S. and Murdia A. 2017. Seasonal incidence of major insect pests of cauliflower. *Indian Journal of Applied Entomology*, 31 (2): 85–89.

- Sharma D. and Rao D.V. 2012. A field study of pest of cauliflower cabbage and okra in some areas of Jaipur. *International Journal of Life Sciences Biotechnology and Pharma Research*, 1 (2): 2250-3137.
- Shylesha A.N., Thakur N.S.A., Pathak, K.A., Rao K.R., Saikia K., Surose S., Kodandaram N.H. and Kalaishekar A. 2006. Integrated management of insect pest of crops in north eastern hill region. *Technical Bulletin No. 19. ICAR RC for NEH Region*, Umiam, 50p.
- Sharma P., Sharma A., Jat S.M., Kumawat K.C. and Badriprasad 2022. Seasonal abundance of diamondback moth and natural enemies on cabbage. *The Pharma Innovation Journal*, 11 (1): 1376-1379.
- Sahu V., Yadav R.R., Singh R.P. and Bareliya P.K. 2017. seasonal incidence of mustard aphid, *Lipaphis erysimi* in Jhansi during *Rabi* crop season. *Plant Archives*, 17 (2) 1505-1507.
- Tripathi A., Mishra D.S. and Sharma R.C. 2005. Role of biotic and abiotic factors on mustard aphid, *Lipaphis erysimi* (Kalt.). *Farm Science Journal*, 14: 84-85.
- Varmora J.M., Raghvani K.L., Joshi M.D., Makadia R.R., Dalwadi N.G. and Boricha H.V. 2009. Population dynamics of aphid, *Lipaphis erysimi* (Kalt.) and predators on cabbage. *Asian Sciences*, 4 (1 &2): 56-58.
- Vekaria M.V. and Patel G.M. 2005 Seasonal abundance of *Lipaphis erysimi* (Kalt.) and their natural enemies on important cultivars of mustard in North Gujarat. *Indian Journal of Entomology*, 67: 369- 377.
- Yadav N. and Agrawal N. 2018. Seasonal abundance of *Brevicoryne brassicae* L. and their predators under different field of cabbage in relation to weather parameters. *Journal of Entomology and Zoology Studies*, 6 (5): 1098-1101.

**Table 1: Population dynamics of mustard aphid, *Lipaphis erysimi* (Kaltenbach) on cauliflower in relation to biotic and abiotic factors in *Rabi*, 2021-22**

SMW	Date of observations	Temperature (°C)		Mean relative humidity (%)	Sunshine (hrs)	Aphid/plant	Mean population/ 10 plant	
		Max.	Min.				<i>C. septempunctata</i>	Maggots of <i>Xanthogramma spp.</i>
48	27/11/2021	27.20	8.30	59	6.40	0	0	0
49	4/12/2021	24.30	7.00	65	5.10	7.4	1.8	1
50	11/12/2021	22.90	3.60	60	6.70	23.2	3.6	2.2
51	18/12/2021	22.40	-0.30	55	7.50	46.8	6.2	4.8
52	25/12/2021	22.70	6.00	65	3.80	78.6	8.8	5.6
1	2/01/2022	20.30	6.10	73	4.10	98.8	9.2	6
2	9/01/2022	17.30	4.50	69	5.00	63.8	9.6	6.4
3	16/01/2022	18.90	3.90	65	6.00	99.6	10.2	7.8
4	23/01/2022	18.90	3.40	66	6.30	113.4	10.6	8.4
5	30/01/2022	23.50	4.10	61	8.60	126.6	11	8.8
6	6/02/2022	23.50	5.10	58	8.90	143.8	11.6	9.2
7	13/02/2022	26.30	4.70	50	9.70	161.2	12	9.8
8	20/02/2022	29.00	7.40	51	9.00	91.2	6.4	3.2
9	27/02/2022	30.1	8.0	49	9.3	47.2	3.8	1.2
Correlation coefficient with max. temp.						-0.169	-0.482	-0.497
Correlation coefficient with min. temp.						-0.226	-0.415	-0.489
Correlation coefficient with relative humidity						-0.075	0.180	0.167
Correlation coefficient with Sunshine (hrs.)						0.379	0.125	0.161
Correlation coefficient with <i>C. septempunctata</i>						0.929**		
Correlation coefficient with Maggots of <i>Xanthogramma spp.</i>						0.917**		

SMW= Standard meteorological weeks

\* Significant at 5 per cent level

\*\* Significant at 1 per cent level

**Table 2: Population dynamics of mustard aphid, *Lipaphis erysimi* (Kaltenbach) on cauliflower in relation to biotic and abiotic factors in Rabi, 2022-23**

SMW	Date of observations	Temperature (°C)		Mean relative humidity (%)	Sunshine (hrs)	Aphid/plant	Mean population/ 10 plant	
		Max.	Min.				<i>C. septempunctata</i>	Maggots of <i>Xanthogramma spp.</i>
48	29/11/2022	27.90	4.80	49.00	8.90	0	0	0
49	6/12/2022	25.50	3.70	55.00	8.10	0	0	0
50	13/12/2022	26.60	4.60	54.00	8.90	9.4	2	1.6
51	20/12/2022	25.50	4.60	58.00	8.10	27.8	3.8	2.8
52	27/12/2022	23.30	2.20	56.00	7.90	77.8	6.8	5.8
1	3/01/2023	20.70	-0.10	60.00	7.50	126.6	10.2	6.6
2	10/01/2023	24.50	4.90	55.00	7.30	175.2	10.8	7
3	17/01/2023	20.10	-0.50	55.00	8.30	194.6	11	7.4
4	24/01/2023	21.30	3.60	61.00	4.10	215.4	11.8	8.8
5	31/01/2023	22.00	5.20	69.00	7.30	237.2	12.6	9.6
6	7/02/2023	26.90	6.90	56.00	9.00	269.8	14.8	13.2
7	14/02/2023	28.40	5.80	51.00	9.40	199.2	9.6	7.8
8	21/02/2023	31.90	9.80	49.00	8.60	112.8	7.6	4.8
9.	28/02/2023	32.10	12.00	49.00	8.80	88.2	4.6	3.2
Correlation coefficient with max. temp.						-0.308	-0.401	-0.345
Correlation coefficient with min. temp.						-0.020	-0.124	-0.077
Correlation coefficient with relative humidity						0.423	0.484	0.471
Correlation coefficient with Sunshine (hrs.)						-0.297	-0.341	-0.270
Correlation coefficient with <i>C. septempunctata</i>						0.967**		
Correlation coefficient with Maggots of <i>Xanthogramma spp.</i>						0.962**		

SMW= Standard meteorological weeks

\* Significant at 5 per cent level

\*\* Significant at 1 per cent level




Fig.1 Population dynamics of major insect pests of cauliflower in relation to biotic and abiotic factors,rabi 2021-22




Fig .2 Population dynamics of major insect pests of cauliflower in relation to biotic and abiotic factors, Rabi 2022-23

UNDER PEER REVIEW