

Short Research Article

EFFECT OF *Foeniculumvulgare*MINGLED DIET UPON GROWTH, REPRODUCTION AND SPAWNING PERFORMANCE OF GUPPY FISH

Abstract

Breeding and rearing of ornamental fish is an important sector of aquaculture, which has increased income and employment in the country in recent years. Introducing new methods and developing new food supplements can impact efficiency and sustainable development in this new technology.

In the present study, phytoestrogenic compounds of ethanolic extract of *F.vulgar*were evaluated on growth performance and fertility rate and gonadal tissue changes of guppy for its introduction as a nutritional supplement in aquaculture.

120 juvenile guppies were distributed three times in 12 glass aquaria at 10 fish/tank. Four treatment doses were prepared with different *Foeniculumvulgare* seed extract concentrations i.e. 50, 100, 150µl/g while the fourth dose was control with 0µl *Foeniculumvulgare* extract. These diets were called D0, D1, D2 and D3.

This diet was fed to fish, twice daily, at 3% of body weight, for 90 days. All groups were tested for desired parameters after fifteen days.

Analysis of results showed significant increase in weight gain and fertility rate in group T2 (100µl/g).

There was an increase in gonad cells in the test groups as compared to the control group and an increase in phytoestrogens in the test groups simultaneously.

In follicles, increasing granulose layer with increasing number of ovules at different developmental stage (O1-O7) was observed in test groups.

The increase in fertility rate in the test groups may be due to the increase in estrogenic compound levels which may be due to the action of the extract in the diet.

The observation made in the present investigation shows that *F.vulgar* can act as a phytoestrogenic compound at low concentration and therefore *F. vulgar* is considered as a potential and natural fish feed additive in commercial aquaculture. Can be used to increase fertility rate.

1.Introduction

The fisheries division is playing a pivotal role in nourishment, dietary security, profit earning, and to enhance the national economy. Among fishes, some are edible and some are ornamental and might be perceived as decorative fishes.

They are likewise called “living jewels” due to their friendly behavior, beautiful color, and shape (Nasser and Rajkumar, 2001). All over the world, the trade of aquatic animals and their products has been increased in the last few years.

Ornamental fish keeping is considered an interesting, easy, and relaxant hobby throughout the world. The worldwide exchange of aquatic organisms has been

proposed as the wellspring business. Ornamental fishes are very attractive aquatic species in the world.

Ornamental fish is a key element of the aquatic environment and considered an ecosystem maintainer in the aquatic environment (Rodríguez, 2006). *Poeciliareticulata* is native species from North Brazil and South America (Ojanguren et al., 2005).

Comment [MF1]: Not found among references

Poeciliareticulata is known as “Guppy fish” belongs to the family Poeciliidae, it is a small ornamental fish also called livebearer, inhibited in a freshwater environment. *Poeciliareticulata* is a commercially important fish due to its attractive colors and variation in the tail that makes it the most trading ornamental aquarium fish in the world (Egset et al., 2011).

Poeciliareticulata start to breed from February till while its peak in July. Female *Poeciliareticulata* are livebearers and give birth to live young. The rearing period of *Poeciliareticulata* in the Southeastern U. S. starts in May and finishes in September and October (Hidebrand, 1921).

Comment [MF2]: Not found among references

At the age of 10-12 weeks, females become mature while males mature at the age of 7 weeks. Female *Poeciliareticulata* show polyandry, female can mate with multiple males. Females can give birth 20-200 young in a single spawn. The gestation period of *Poeciliareticulata* is about 4-6 weeks (Coad, 2017).

Seeds availability is only possible by the reproduction. Natural reproduction is not enough to meet the seed availability according to demand. Induce breeding is the best way to enhance the reproduction, to produce the seeds.

Availability of seeds can be possible by reproduction as well as artificially induced breeding in hatchery (Sudha and Gokula, 2015). Due to the lack of proper environmental condition, reproduction and growth of fishes is disturbed, and it is difficult to meet the demand of fish seed by natural resources. So, to fulfill this demand synthetic stimulators are used for fish breeding (Ojima and Iwata, 2009).

For egg development and spawning, female need appropriate amount of lipid and protein. Diet play a significant job for development and reproduction of ornamental fish, and various feeds have been utilized for its raising.

Supplement feed, hormones, anti-infection agents, and numerous herbal products are used for enhancements for fecundity and gametes maturation of ornamental fish (Ghosh et al., 2007).

There are many ways to stimulate the reproduction of fish. Reproduction of fish can be provoked by using one of these products in ornamental fish, Ovaprim a liquid form injection which is a mixture of GnRH and domperidone in the liquid of propylene glycol.

Ovaprim used as a spawning abet to induce evolution (Yanong et al., 2010). Gonadosomatic index (GSI) is the ratio between body weight and gonads weight of fish.

Gonadosomatic index involves to identify the spawning season of fish. While fecundity is the relation between numbers of eggs produced in each spawn and body weight of fish. It is most important parameters used in the study of reproduction of fish (Hunter and Macewicz, 2001).

Comment [MF3]: Not found among references

Gonadosomatic index and fecundity measures are helpful in measuring population dynamics, population carrying capacity and productivity (Shankar and Kulkarni, 2005).

Comment [MF4]: Not found among references

It evaluate the reproductive system of a fish by calculating the total body size. It is considered as the most important parameter of fish reproductive biology, provides us a comprehensive view about the fish breeding and the reproductive status of related species, especially in finding the fish breeding seasons. Fertility and effectiveness are estimated by Gonadosomatic Index and fecundity (Chavan and Muley, 2014).

Comment [MF5]: Not found among references

To check the effect of traditional herb *Foeniculumvulgareon* growth rate, reproductive performance of *Poeciliareticulata*.

To evaluate the influence of *Foeniculumvulgareon* spawning of female *Poeciliareticulata*.

2.Materials and Method

Complete Random Design (CRD) was used for sampling purposes to evaluate the variable of growth (weight gain, length gain, FCR, and survival rate) and Reproduction (Fecundity, GSI, Hatching rate, and spawning).

The present study was conducted in the Laboratory, Department of Zoology, The University of Lahore, Sargodha Campus. *Poeciliareticulata* stock of 120 was kept in the 12-glass aquarium of size of 12×12×18. Juvenile *Poeciliareticulata* having size range of 0.19 ± 0.02 were divided into four experimental groups. Each glass aquarium for experimental treatment contains 10 juvenile guppies (30 in triplicate). Fish feed used during the experiment was commercial.

Comment [MF6]: cm or grams

Sample Size

Present experiment was performed for 3 months (90 days) after being acclimatized in laboratory conditions for two weeks. Initial stocks of 120 juvenile *Poeciliareticulata* having an average weight of 0.19 ± 0.007 were purchased from a commercial fish shop in Sargodha.

Comment [MF7]: g

Fish were identify by following the Coad, (2017). Guppies were kept in the four aquariums to make them acclimatized the laboratory condition for two weeks before the research. After two weeks, fingerlings were divided into four Aquarium manually @ ten fish per aquarium.

During this period juveniles were observed and diseased juveniles were separated from the healthy fishes, then remaining stocks were divided into four treatments having an average weight of 0.18 ± 0.01 g (T₁), 0.19 ± 0.01 g (T₂), 0.19 ± 0.01 g (T₃), and 0.19 ± 0.01 g (T₄).

Poeciliareticulata were provided with a feed named "INCH GOLD" a commercial feed of 3% of body weight twice a day i.e. 8:00 am and 6:00 pm. Aquarium water was changed weekly to maintain a healthy and clean water environment.

Foeniculumvulgare SEEDS EXTRACT PREPARATION

Foeniculumvulgare seeds were purchased from the herbal shop in Sargodha, Pakistan. *Foeniculumvulgare* seeds were identified by the Department of Botany, Lahore University, Sargodha Campus. *Foeniculumvulgare* seeds were grinded with a blender to get a fine powder.

To get *Foeniculumvulgare* seed extract, *Foeniculumvulgare* seed powder (50 g) was taken and 96% ethanol mixed with a ratio of 1: 1. The extract was kept at room temperature for 24 hours with continuous stirring.

What-man filter paper No.1 was used to filter the extract. The obtained extract was diluted with distilled water to obtain the final solution (Nazari and Roozbehani, 2015).

Treatment Feed Preparation

Feed was prepared by followings (Haghighi et al, 2014). Commercial feed which is “Inch gold” used in feed and mixed with *Foeniculumvulgare*seeds extract in Petri dish. Four treatments diets with different *Foeniculumvulgare*seeds extract concentration were prepared i.e. 50, 100, 150µl/g while fourth diet was control having 0µl *Foeniculumvulgare*extract.

Comment [MF8]: Not found among references

These diets were termed as D₀, D₁, D₂, and D₃. Feed pellets were dry at room temperature. The pellets were air dried and stored in air tight container at room temperature until fed. This feed was fed to fishes, twice a day, at the 3 % of body weight, for 90 days. From all groups required parameter evaluation was checked after fifteen days.

Seed Analysis

*Foeniculumvulgare*seed extract was analyzed through gas chromatography-mass spectroscopy system (Hammouda et al., 2014).

Comment [MF9]: Not found among references

GC-MS system contains an auto-sampler unit, injector use, and a gas chromatograph with a mass spectrometer. With a ratio of 1: 100, approximately 2µl samples were inserted into split mode. Helium gas was used as the carrier with a pressure of (65.2 kPa).

The ionization energy used to identify the compounds by GC-MS was 70 eV. The temperature in the Column oven was 80⁰C to 220⁰C. The flow rate of helium gas was used as 1.5 l ml/min as a carrier gas.

The average temperature range for the injector and MS transfer line was 220⁰C to 290⁰C. Total time required for separation was 65 minutes. The MS capillary column was extracted the *Foeniculumvulgare*seed extract compounds (Alam, 2019).

List 1 :*Foeniculumvulgare*seed extract composition (%) by GC-MS

Chemical compound	%
4-Hexen-2-one	0.20

3-Hydroxytetrahydropyran	0.65
Acetic acid	0.10
Fenchone	11.68
Delta-3-carene	0.52
Camphor	Trace
Tetradecane	0.90
Benzaldehyde-4-methoxy	10.01
L-Histidine	Trace
Benzene -1- methoxy-4- (2-propene)	2.52
Fenchyl acetate	0.11
Cyclohexane	0.41
1-Methyl-2-methylene-4-isopropyl	Trace
Trans-anethole	64.49
1-Tetradecene	0.74
1-Methoxy-3-(ethenylcarbonyl)B	0.81
Hexatriacontane	1.54
10-Nonadecanone	3.17

shows the chemical component of fennel seed extract. With the aid of GC–MS, 18 compounds were described. Trans-anethole (64.49 percent), Fenchone (11.68 percent), and Benzaldehyde-4-methoxy (10.01) were the main constituents of the ethanolic extract of fennel seed. Furthermore, the fennel seed contained significant amounts of different minor constituents, with a contribution of less than 10%.

3.Results

GROWTH PARAMETERS

4.4.1 Weight gain and length gain

Table 1: Growth data of *Poeciliareticulata* record after 14 days Trail.

Difference between Initial reading and after 14 days reading of samples						
	Initial	Final	weight gain	Initial	Final	length

Treatments	Wt(g)	Wt(g)	(W _f - W _i)	Length (cm)	Length (cm)	gain L _f -L _i
T ₀	0.17	0.21	0.04	1.93	2.2	0.27
T ₁	0.19	0.24	0.05	1.95	2.4	0.45
T ₂	0.19	0.27	0.08	1.97	2.7	0.73
T ₃	0.18	0.24	0.06	1.92	2.5	0.58

Table 2:Growth data of *Poeciliareticulata* record during Trail.

Total Weight gain and length gain durig experiment (90 days)							
Sr. No	Treatment	Initial Wt(g)	Final Wt (g)	weight gain (W _f - W _i)	Initial Length (cm)	Final Length (cm)	Net length gain L _f -L _i
1	T ₀	0.17	0.31	0.12	1.93	3.07	1.41
2	T ₁	0.19	0.38	0.19	1.95	3.73	1.78
3	T ₂	0.19	0.44	0.25	1.97	4.09	2.12
4	T ₃	0.18	0.39	0.21	1.92	3.66	1.74

Calculated results for weight gain and length gain were presented in Table 2. Statistical analysis shows that maximum weight gain was observed in T₂ while minimum growth rate was recorded in T₀ (Control group) respectively.

The increase in length was also observed during the experiment to check the effect of *Foeniculumvulgare* extract on total length gain. Length increase is presented in Table 2. Analysis of increase in length presented in graph (Fig 1) shows that the maximum length was attaining by Treatment T₂ and T₀ achieved minimum length. Growth performance results showed that with a concentration of 100µl of *Foeniculumvulgare* seed extract containing has a positive effect (P<.0.05) on the growth parameters of *Poeciliareticulata*.




Figure 1: Increase growth rate observation fed with *Foeniculumvulgare* concentrations.

Weight gain, height increase, survival rate, and feed conversion ratio were recorded in (Table 3). Growth parameters were recorded after fifteen days. The statistically analyzed data showed that as the concentration of *Foeniculumvulgare* increases the weight and length were also increases significantly ($p < 0.05$).

Table 3: Growth parameters of *Poeciliareticulata* fed diets containing different concentrations of *Foeniculumvulgare* for 90 days.

Growth parameters	T ₀ (0 μ l)	T ₁ (50 μ l)	T ₂ (100 μ l)	T ₃ (150 μ l)
Initial weight (g)	0.17 \pm 0.008 ^a	0.18 \pm 0.01 ^a	0.19 \pm 0.008 ^a	0.19 \pm 0.007 ^a
Final weight (g)	0.31 \pm 0.03 ^c	0.38 \pm 0.01 ^b	0.44 \pm 0.03 ^a	0.39 \pm 0.03 ^b
Weight gain (g)	0.12 ^c	0.21 ^b	0.25 ^a	0.19 ^b
Initial Length(cm)	1.93 \pm 0.13 ^a	1.95 \pm 0.16 ^a	1.97 \pm 0.1 ^a	1.92 \pm 0.14 ^a
Final Length (cm)	3.07 \pm 0.1 ^c	3.66 \pm 0.2 ^b	4.10 \pm 0.3 ^a	3.73 \pm 0.2 ^b
Net length gain	1.14 ^c	1.71 ^b	2.12 ^a	1.81 ^b
Feed conversion	3.11 \pm 0.8 ^a	2.52 \pm 0.4 ^b	2.24 \pm 0.1 ^c	2.93 \pm 0.2 ^{ab}

ratio				
Survival rate (%)	100 ± 0.00 ^a	100 ± 0.00 ^a	100 ± 0.00 ^a	100 ± 0.00 ^a

Note: Values given are mean ± standard error. Means with the same letter in the same column are not statistically significantly ($P < .05$) different

4.4.2 Feed conversion ratio

According to the results, data is presented in Table 3, different concentrations of fennel extract has a positive effect on the FCR of feed fed by *Poeciliareticulata* in experimental treatments. FCR values were recorded as in T₀ (3.11±0.8), T₁ (2.52±0.4) T₂ (2.24±0.1), and T₃ (2.93±0.2), respectively. A significant decrease in the FCR value of T₂ is observed when compared with other groups in the experiment. Fennel seed extract contains anethole which is a digestive stimulator, a reason for the decrease in FCR. Graphical presentation (Fig 2) of Feed Conversion Ratio shows that highest in T₀ while it was lowest in T₂. The results indicate that T₂ has significantly differed from T₁, T₃, and T₀. The results about FCR showed that *Foeniculumvulgare* reduces the Feed conversion ratio.




Figure 2. Mean values of feed conversion ratio under varying stocking concentration of fennel extract on *Poeciliareticulata*

4.4.3 Survival rate

Graphical presentation of survival rate (Fig 3) showed that fennel extract did not negatively affect the Survival rate of *Poeciliareticulata* in every experimental

group. The survival rate in all experimental and control were consistently high ($P>0.05$). The survival rate of all treatments is recorded as 100 %. It is inferred that *F. vulgare* seed extract has no adverse effect on the Survival rate of *Poeciliareticulata*.




Figure 3: Mean values of survival rate under varying stocking concentrations of fennel extract on *Poeciliareticulata*

4.5 REPRODUCTION PARAMETERS

4.5.1 Gonadosomatic index

Table 4 : Age, fish, and gonad weight, Gonadosomatic index (GSI) of *P.reticulata*

Sr.no	Treatments	T ₀	T ₁	T ₂	T ₃
1	Age in days	180-192	180-198	180-210	180-195
2	Fish weight	0.79±0.008 ^c	0.93± 0.008 ^b	1.04± 0.01 ^a	0.93 ± 0.008 ^b
3	Gonads weight	0.023±0.002 ^d	0.041±0.004 ^c	0.065±0.001 ^a	0.053±0.003 ^b
4	GSI	6.07 ± 0.11 ^c	7.03 ± 0.04 ^b	8.69 ± 0.15 ^a	7.54 ± 0.05 ^{ab}

Note: Values (mean \pm SEM) of GSI are superscripted by alphabets within the same line are different significantly ($P < 0.05$).

Final data of reproduction parameters are described in Tables 4. The gonadosomatic index was measured by the weight of the fish and the gonad's weight of the fish. Gonad's weight gain at maturity is shown in Fig 4. The highest Gonadosomatic index was observed in T₂ followed by T₁, T₃, and T₀ separately. The groups that contain *F. vulgare* seed extract, the Gonadosomatic index (GSI) is significantly high ($P < 0.05$). The highest GSI was observed in T₂ (8.69 ± 0.2) and lowest GSI was found in T₀ (6.07 ± 0.2) respectively.




Figure 4: Effect of *Foeniculum vulgare* in diet containing different concentrations on Gonadosomatic index (Mean \pm SD) of *Poeciliareticulata*

4.5.2 Fecundity

Table 5: The impact of different concentrations of *Foeniculum vulgare* extract on reproductive parameters of *Poeciliareticulata* (♀) for 90 days

Sr. no.	Parameter	Treatments			
		T ₀	T ₁	T ₂	T ₃
1	Fish weight	0.79 \pm 0.008 ^c	0.93 \pm 0.008 ^b	1.04 \pm 0.01 ^a	0.93 \pm 0.008 ^b
2	Fish length	3.9 \pm 0.2 ^d	4.4 \pm 0.12 ^c	5.6 \pm 0.08 ^a	5.1 \pm 0.04 ^b
3	Absolute	5.7 \pm 0.12 ^d	7.9 \pm 0.12 ^c	9.4 \pm 0.19 ^a	8.3 \pm 0.15 ^b

	fecundity				
4	Relative fecundity	46.8 ± 1.20 ^c	53.5 ± 0.98 ^b	62.3 ± 1.87 ^a	57.1 ± 1.71 ^{ab}
5	Gonadosomatic index	6.07 ± 0.11 ^d	7.03 ± 0.04 ^c	8.69 ± 0.15 ^a	7.54 ± 0.05 ^b
6	Hatching (%)	62.3 ± 1.69 ^c	81.2 ± 0.08 ^b	91.3 ± 4.1 ^a	85.5 ± 2.3 ^b

Note: Values (mean ± SEM) superscripted by various letter sets inside a similar line are differ essentially (P < 0.05).

Findings of reproductive parameters are presented in Table 5 and Fig 5. Significant differences were found among treatments in Gonadosomatic index (P<0.05); although absolute fecundity and relative fecundity of guppy fish significantly increased (P<0.05) in T₂ fed with 100 µl of *Foeniculumvulgare*extract. .

As the gonads weight increase, relative fecundity and absolute fecundity also increase. The numbers of eggs were highest count in T₂. This showed that fennel seed extract with 100µl concentration in diet has significant results (p<0.05) on the Gonads initial maturity and fecundity.

The present study reported that, the gestation period of *Poeciliareticulata* was 25-35 days with an average of 28 days.

In *Poeciliareticulata*, treatment is interior and happens through mating of couples demonstrating explicit mating conduct. Male play out S-formed stance known as 'sigmoid display' and orientates himself before the females toward the start of courtship (Krumholz, 1948).




Figure 5: Comparison of absolute fecundity and relative fecundity changes in test groups and control during experiment.

4.5.3 Hatching

Results for hatching rate (%) were presented in Table 5. According to graph (Fig 6) highest value was observed in T₂ ($p < 0.05$) which show that increased significantly followed by T₁, T₃ and T₀ respectively. Results were best in T₂ which contains 100 μ l extract of *Foeniculumvulgare*. Highest and lowest hatchling rate were respectively in T₂ and control group. According to result *Foeniculumvulgare* has a positive effect towards increase in hatching percentage.




Figure 6: Number of young ones released by control and experimental groups with varying concentrations of *Foeniculumvulgare*.

4.5.4 Spawning performance

Table 6: Evaluation of spawning response of *Poeciliareticulata*

Parameters	Treatments			
	T ₀	T ₁	T ₂	T ₃
Survival rate	100 ±	100 ±	100 ±	100 ±
percentage	0.00 ^a	0.00 ^a	0.00 ^a	0.00 ^a
Number of larva	45±3.5 ^d	74± 1.4 ^c	91± 0.81 ^a	82± 1.6 ^b
Number of ova	34±2.4 ^d	68±2.9 ^c	81±2.1 ^a	75±2.4 ^b
Spawning time(hours)	88±0.81 ^a	65±0.81 ^c	39±2.94 ^d	76±1.63 ^b

Note: Values (mean SEM) superscripted by alphabets are significantly different ($P < 0.05$).

To evaluate the influence of *Foeniculumvulgare* on the spawning performance of Guppy fish results were presented in the Table 5. Graphical presentation (Fig 7) presented that guppy fish laid more egg when treated with Ethanolic extract of *Foeniculumvulgare* with 100µl concentration. Treatment (T₂) when treated with 100µl *Foeniculumvulgare* resulted a significantly positive effect ($p < 0.05$) 91±2.1 on no of young ones produced by *Poeciliareticulata*. Present findings indicated that herbal extract of *Foeniculumvulgare* is a positively effective stimulator for spawning of *Poeciliareticulata*.




Figure 7: graphical comparisons of different doses of *Foeniculumvulgare* mingle diet on spawning performance in *Reticulata*.

4. DISCUSSION

The success of aquaculture depends on a number of parameters, the most important is a suitable diet that contains the complete balanced nutritional diet for the average growth of fish (Salehi, 2008).

The aim of this research was to investigate the effect of *F. vulgare* extract mingles diet on the growth and reproductive performance of *Poeciliareticulata*. According to results of the present study, FCR decreased in T₂ as compared to other treatments. *Foeniculumvulgare* seed extract contains the Anethole and Estragole perform the function as a digestive stimulant may be a reason for the decrease in FCR (Frankic et al., 2009).

More than this growth of *Poeciliareticulata* significantly ($p < 0.05$) positive increase in T₂. The growth parameter, the survival rate was uniformly the same in all treatments which are about 100% in all treatments.

This justified that *Foeniculumvulgare* has no ($p > 0.05$) effect on the survival rate. There are similarities between the current study and those investigated by Sotoudeh & Yeganeh (2017). The finding of the research is also co-related with the

Comment [MF10]: Why was Treatment 2 the best in terms of growth, while in Treatment 3 a greater percentage of the tested material was used?

Comment [MF11]: Not found among references

Comment [MF12]: Not found among references

work of Yilmaz et al. (2012). His research investigated that dietary thyme improved the growth performance of *Dicentrarchus labrax*.

Water parameters recorded during the experiment were temperature, alkalinity, dissolved oxygen, TDS, TSS, and EC. Water temperature recorded during the experiment was range from 24⁰C to 30⁰C.

Environmental parameter, the temperature was accordingly to the work of Dawes, (1991). The results of this study are congruous with the finding of Yilmaz et al, (2012). They evaluated that a dietary cumin level of 1% provides the best survival rate for tilapia, *Oreochromis mossambicus*, with no inauspicious effect on growth performance simultaneously.

Tribulus Terrestris showed the same results with a significant difference ($P < 0.05$) in weight gain (Yeganehet al., 2017). The present investigation showed similar findings as *Thymus vulgaris*, *Rosmarinus officinalis*, and *Trigonella foenumgraecum* had a positive response towards the growth performance of *Dicentrarchus labrax* (Yilmaz et al., 2010).

After the chemical analysis of *Foeniculum vulgare* seed extract resulted that *Foeniculum vulgare* extract contains the trans-anethole (64.49 %) which is similar in structure to 17-beta- estradiol.

The presence of trans-anethole may cause a significant increase in estrogenic hormones; lead to an increase in reproductive performance and fecundity (Albert-Puleo, 1980). An increase in the reproduction activities of *Poeciliareticulata* fed feed containing *F.vulgare* seed extract, a reason for increasing the level of estrogenic hormones lead to increase the reproductive performance. A similar result was reported by Nazari and Roozbehani, (2015). Their finding showed that the fertility rate of *Poeciliareticulata* enhanced when used *Foeniculum vulgare* extract in the diet.

Among dietary treatments used in this research, T₂ show positive increase in GSI (8.69 ± 0.2) Why was Treatment 2 the best in terms , while in Treatment 3 a greater percentage of the tested material was used?, Fecundity (9.4 ± 0.19), and hatching rate. Dada and Adeparusi (2012) intimate that diets with *Sesamum indicum* supplement and seed powder of *Croton zambesicus* were improved female *C. gariepinus* GSI.

Comment [MF13]: Not found among references

Comment [MF14]: Not found among references

Comment [MF15]: Not found among references

Comment [MF16]: Not fo Why was Treatment 2 the best in terms , while in Treatment 3 a greater percentage of the tested material was used? and among references

Dada and Ajilore (2009) distinguished an increase in egg diameter and fecundity of *C. gariepinus* when treated with *Garcinia* seeds extract. Sadeghpour et al. (2015) commented that increase in serum level estrogen in mice female when injected with extracts of *Foeniculumvulgare*.

Comment [MF17]: Not found among references

Comment [MF18]: Not found among references

The findings of another research are also consistent with this current research, fecundity of guppy was directly proportional to the bodyweight of the fish, which means that the fecundity increased with the increase in body weight (Shahjahan et al., 2013). The use of injection of ovaprim shown the highest fecundity of *Pterophyllumscalare* suggested by Chatterjee, Patra, and Talwar, (2013).

Comment [MF19]: Not found among references

Comment [MF20]: Not found among references

Ghosh et al. (2007) reported that incorporation of probiotics in feed influenced the reproductive performance of livebearers in terms of high fecundity, high Gonadosomatic index, high fry survival rate, reduction in fry mortality and deformity, and higher average weight and length gain of fish fry.

Nielsen and Baatrup, (2006) treated *Poeciliareticulata* with estrogens that enter into the aquatic system, results of this research showed that no significant difference was seen in the Gonadosomatic index (GSI).

Comment [MF21]: Not found among references

A similar observation was found with *Cyprinuscarpio* when treated with *Ferula coskunii* (Yilmaz et al., 2006). Current research finding indicated that *Poeciliareticulata* treated with *Foeniculumvulgare* seed extract, increased number of young's ones produced by guppy fish in T₂ were 91±2.1 while the control group produced 45±2.5. Dada (2012) investigated that feed supplement with *G. kola* seed powder improved hatchability and fecundity of *C. gariepinus*.

Successful spawning of *C.punctatus* at 0.3 and 0.5ml/kg and 3000IU/kg weight of HCG was noted by Kather and Sridhar (2002). For *H.fossilis* the utilization of ovaprim came about greatest Successful body spawning (Chatterjee et al., 2013). Effective spawning of *C.punctatus* is when treated with HCG. *Natrummuriaticum* showed a positive effect on the spawning performance of *Poeciliareticulata* (Sudha and Gokula, 2018).

Comment [MF22]: Not found among references

Comment [MF23]: Not found among references

The findings of previous research are similar to the results of the current investigation. There are also similarities between the work of (Sudha and Gokula, 2015) when *Puntiusconchonus* treated with similar herbal medicine *Natrummuriaticum* on spawning response.

5. CONCLUSIONS

The present study investigated that *F.vulgare* seeds extract has a significant effect on the growth performance of *Poeciliareticulata*.

Under the report of the current investigation, *F.vulgare* seed extract has a significant positive effect on reproductive parameters.

Increasing the level of estrogen hormones in fish fed with *F.vulgare* ethanol extract, due to the presence of trans-anethole, reproduction activities and fecundity of the fish was increased.

Considering, fennel seed extract within the diet has a significant increase in the Gonad's initial maturity Gonadosomatic index increase is directly proportional to a female's body size.

As the gonad's weight increases, relative fecundity, and absolute fecundity also increase the number of eggs.

It is suggested that herbal extracts contain estrogenic compounds as in *Foeniculumvulgare* can be used as a supplement in fish diets.

In the future, the use of plant extract in aquaculture will be an effective tool to attain durable, economically safe fish production.

References

- Coad, B. W. (2017). Review of the livebearer fishes of Iran (Family: Poeciliidae). Iranian Journal of Ichthyology, 4(4), 305-330.
- Dada, A. A. (2012). Effect of Garcinia kola seed meal on egg quality of the North African catfish (*Clariasgariepinus*) (Burchell, 1822) broodstock. African Journal of Food, Agriculture, Nutrition and Development, 12(5), 6447-6459.
- Egset, C. K., Bolstad, G. H., Rosenqvist, G., Endler, J. A., & Pelabon, Cz/(2011). Geographical variation in allometry in the guppy (*Poeciliareticulata*). Journal of Evolutionary Biology, 24(12), 2631-2638.
- Frankic, T., Voljc, M., Salobir, J., & Rezar, V. (2009). Use of herbs and spices and their extracts in animal nutrition. ActaAgricSlov, 94(2), 95-102.
- Ghosh, S., Sinha, A., & Sahu, C. (2007). Effect of probiotic on reproductive 18-526.methods by GC/MS. African Journal of Traditional, Complementary and Alternative Medicines, 11(2), 277-279.

Comment [MF24]: Rearrange references according to alphabet

Nasser, A. K. V., & Rajkumar, U. (2001). Ornamental fish-prospects for culture. Souvenir issued on the Occasion of the inauguration of Visakhapatnam RC of CMFR, 24-26.

Nazari, A., & Roozbehani, S. (2015). Influence of fennel *Foeniculum vulgare* extract on fertility, growth rate and histology of gonads on guppy *Poecilia reticulata* Turkish Journal of Fisheries and Aquatic Sciences, 15(3), 457-463.

Ojima, D., & Iwata, M. (2009). Central administration of growth hormone-releasing hormone triggers downstream movement and schooling behavior of chum salmon (*Oncorhynchus keta*) fry in an artificial stream. Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology, 152(3), 293-298.

Rodriguez, R. D. R. (2006). Ornamental fish: economic, ecologic and health implications of the trade. JAINA Boletín Informativo, 16(1), 6-27.

Salehi, H. (2008). Benefit-cost analysis for fingerling production of kutum *Rutilus rutilus* (Kamensky, 1901) in 2005 in Iran. Aquac. Asia Mag, 13(3), 35.

Sudha, C., & Gokula, V. (2015). Induced spawning of rosy barb *Puntius conchonus* by *Natrum muriaticum*. IJAPBC, 4(4), 801-804

Yanong, R. P., Martinez, C., & Watson, C. A. (2010). Use of Ovaprim in ornamental fish aquaculture. EDIS, 2010(2).

Yilmaz, E., Genc, M. A., Cek, S., Mazlum, Y., & Genc, E. (2006). Effects of orally administered *Ferula coskunii* (Apiaceae) on growth, body composition and histology of common carp, *Cyprinus carpio*. Journal of Animal and Veterinary Advances, 5(12), 1236-1238.

Yilmaz, S., Ergün, S., & Turk, N. (2012). Effects of cumin-supplemented diets on growth and disease (*Streptococcus iniae*) resistance of tilapia (*Oreochromis mossambicus*). The Israeli Journal of Aquaculture – Bamidgah, 64. p5

Yilmaz, S., Sebahattin, E., & Celik, E. (2010). Effects of herbal supplements on growth performance of sea bass (*Dicentrarchus labrax*): Change in body composition and some blood parameters. Energy (kJ/g), 1(3), 217-222

Comment [MF25]: This is a reference...it should be classified as a reference

Comment [MF26]: It is written in italics

UNDER PEER REVIEW