

Effect of Potassic fertilizer, FYM and Vermicompost on Growth Parameters, Yield Attributes and Yield of Pigeon Pea (*Cajanus cajan*) Crop in Chitrakoot Area

Abstract

A field experiment was conducted during *Kharif* season of 2022-2023 at Rajoula Agriculture farm, of Mahatma Gandhi Chitrakoot Gramodaya Vishwavidyalaya Chitrakoot, Satna (M.P.) to evaluate the effect of potassic fertilizer, FYM and vermicompost on growth parameters, yield attributes and yield of pigeon pea. The present experiment having 11 treatment combinations replicated thrice in randomized block design. Pigeon pea variety Rajeshwari Phule-12 was grown with recommended agronomic practices. On the basis of the results emanated from present investigation, it could be concluded that application of T₁₁ [100 % N & P, K- 0 % + 100 % K through Vermicompost] significantly recorded maximum growth parameters such as plant population plot⁻¹ (138.42), plant height (211.38 cm), number of leaves (208.6) and number of branches plant⁻¹ (32.8) and maximum yield attributing characters such as number of pod plant⁻¹ (103.6), number of seed pod⁻¹ (4.4) and seed index (9.68 g). The result showed highest grain yield (17.69 q ha⁻¹) and straw yield (52.64 q ha⁻¹), with 100% RDF + soil application of treatment T₁₁ [100 % N & P, K- 0 % + 100 % K through Vermicompost] in comparison to all the treatments.

Keywords: FYM, Pigeon pea, Yield and Vermicompost.

1. Introduction

“Pulses are rich source of protein, vitamins, fibres and minerals (iron, zinc, magnesium) and some essential amino acids which is play a vital role in human’s health. Pulses the second most importance food crops after cereals in food security. Pulses are a good source of dietary protein. Significant part of protein in Indian diet come from pulses. Pulses improve human health as well as soil health by improves its nitrogen content through symbiotic nitrogen fixation from atmosphere and mushrooming the soil microorganism population in soil”. (Yadav *et al.*, 2017).

“Madhya Pradesh is the largest producer of pulses accounting for 16.95 % of total pulses production. In Madhya Pradesh, pulses grown in an area about 48.67 lakh hectare with a

production of 5.30 mt and the average productivity is 1084 kilogram per hectare”. (**Agricultural Statistics at a Glance, 2021**). “In Madhya Pradesh, pigeon pea grown in an area about 4.04 lakh hectare with a production of 4.72 lakh ton and the average productivity is 1170 kilogram per hectare”. (**Annual Progress Report 2021-22, DPD Bhopal**)

Pulses are blessed with an unique natural process of biological nitrogen fixation through which inert nitrogen present in atmosphere is converted into plant usable form. It is assumed that available soil nitrogen and fixed nitrogen by pulse crop is sufficient to meet out its major nitrogen requirement and minimum is required to be supplemented through chemical fertilizer. (**Reay and Reay, 2015**)

“Nitrogen is most important plant growth promoting element. It increases the leaf area production that results in higher photosynthetic activity. The carbohydrates produced through photosynthesis provide energy to rhizobia bacteria. Poor nitrogen supply will affect the plant growth and there by photosynthesis which affect symbiosis”. (**Singh et al., 2017**)

“Phosphorus is a component of adenosine triphosphate (ATP), which is the primary energy currency in cells. ATP stores and transfers energy within the plant, supporting essential processes like photosynthesis, respiration, and nutrient uptake. Phosphorus is essential for cell replication and the formation of new tissues, supporting overall plant growth. Phosphorus is crucial for root development in pigeon peas. Adequate phosphorus levels promote the growth of strong and healthy roots, enabling better absorption of water and essential nutrients from the soil” (**Hawkesford et al., 2012**).

Potassium helps regulate the osmotic balance in plant cells. It plays a crucial role in maintaining cell turgor pressure, which influences cell expansion, water uptake, and overall plant hydration. Adequate potassium levels ensure proper water uptake and help plants withstand drought conditions. Potassium is required for the activation of numerous enzymes involved in various metabolic pathways within the plant. These enzymes are essential for photosynthesis, respiration, protein synthesis, and other vital cellular processes. **Garg & Manchanda, (2009)**.

“FYM contains many species of living organisms which release phytohormones as GA₃, IAA and CYT which stimulates plant growth, absorption of nutrients. There is a growing interest among the farmers to cultivate crops under organic farming because of the escalating cost of inorganic fertilizers, decreased soil fertility, environmental and health concerns due to pesticide usage and expected premium prices for organically grown crops” (**Ramesh et al., 2005**). “One of

the important aspects of organic farming is the soil fertility or nutrient prepared from farm wastes is an important way of recycling nutrients to the management to optimize the crop productivity". (Ramesh et al., 2005).

"Vermicomposting is a type of composting in which certain species of earthworm used to enhance the process of organic waste conversion and produce a better end-product. Vermicomposting is a mesophilic process utilizing microorganisms and earthworms. Earthworms feeds the organic waste materials and passes it through their digestive system and gives out in a granular form which is known as vermicompost. The use of manures from livestock and the composts soil" (Chanda *et al.* 2011). Vermicompost enhances the availability of oxygen, soil porosity and water infiltration, maintains soil temperature, influences soil microbial activity, and improves soil water holding capacity. It also promotes plant growth, yield and quality (Arora *et al.* 2011).

Keeping in view the significance of potassic fertilizers in combination with organic manures on growth parameter, yield components and yield of pigeon pea present investigation was undertaken at Rajoula Agriculture farm, of Mahatma Gandhi Chitrakoot Gramodaya Vishwavidyalaya Chitrakoot, Satna (M.P.)

2. Materials and Methods

2.1 Experimental Site

The experiment was carried out at Rajoula Agriculture farm, Mahatma Gandhi Chitrakoot Gramodaya Vishwavidyalaya Chitrakoot, Satna (M.P.) which lies in the semi- arid and sub-tropical region of Madhya Pradesh between 25.148° North latitude and 80.855° East longitude. The altitude of town is about 190-210 meter above mean sea level.

2.2 Edaphic Condition

The soil was moist, well drained with uniform plane topography. The soil of the experimental field was alluvial in origin, sandy loam in texture and slightly alkaline in reaction having pH 7.25 (1:2.5 soil: water suspension method given by Jackson, 1973), low in organic carbon percentage in soil is 0.22 per cent (Walkley and Black's rapid titration method given by Walkley and Black, 1934), low in available nitrogen 98.95 kg ha⁻¹ (Alkaline permanganate method given by Subbiah and Asija, 1956), medium in available phosphorus as sodium bicarbonate-extractable P was 16.32 kg ha⁻¹ (Olsen's calorimetrically method, Olsen *et al.*, 1954), high in available potassium was 262.99 kg ha⁻¹ (Flame photometer method given by Hanwey and Heidel, 1952),

2.3 Experimental Details

The experiment was laid out in randomized block design and replicated thrice comprising with 11 treatment combinations.

Table no.-1: Treatment details

Treatment	Treatment combination
T₁	100 % RDF
T₂	100 % N & P or 50 % K through MOP + 50 % K through FYM (Toxicity of K)
T₃	100 % RDF + 100 % K through FYM
T₄	100 % N & P + 50 % K MOP + 50 % K through Vermicompost (Toxicity of K)
T₅	100 % RDF + 100 % K through Vermicompost
T₆	100 % N & P, K- 50 % + 50 % K through FYM
T₇	100 % N & P, K- 50 % + 50 % K through Vermicompost
T₈	100 % N & P, K- 0 % + 50 % K through FYM
T₉	100 % N & P, K- 0 % + 100 % K through FYM
T₁₀	100 % N & P, K- 0 % + 50 % K through Vermicompost
T₁₁	100 % N & P, K- 0 % + 100 % K through Vermicompost

2.4 Fertilizer application

FYM and vermicompost were applied as basal dose as per treatment. After the layout of experimental plot, the fertilizers were weighed and applied in the plots and thoroughly mixed with soil. As per the experimental recommended doses of Nitrogen, Phosphorus, Potassium and Sulphur were applied to all the plots. Recommended dose of Nitrogen, Phosphorus and Potassium were applied through Urea, DAP and MOP (20:60:40 kg ha⁻¹)

2.4.1 Seed and sowing: A seed rate of 12 kg per hectare of pigeon pea variety Prakhar (K 1055) was used and sowed on 28th July 2022. Row to row distance was 60 cm.

2.4.2 Harvesting: When the plants turned yellow and dried up, on 24th January, 2023, the crop was harvested. “After measuring the weight of the air-dried bundles with a spring balance, the harvested crop was tied in labelled bundles and threshed using a tractor-drawn thresher. Plot-by-plot records of the grain and straw yields were made and translated to q/ha”. [24]

2.5 Observations recorded

2.5.1 Grain yield ($q\ ha^{-1}$): “The total weight of clean and dried grains from each plot was weighed with the help of electronic balance in kg/ha and converted into q/ha”. [24]

2.5.2 Straw yield ($q\ ha^{-1}$): “Straw yield of each plot can be obtained by deducting the grain yield from the respective biological yield and expressed in q/ha”. [24]

2.6 Statistical Analysis: “The data on various characters studied during the course of investigation were statistically analyzed for randomized block design. Wherever treatment differences were significant (“F” test), critical differences were worked out at five per cent probability level. The data obtained during the study were analyzed statistically using the methods” advocated by **Gomez and Gomez (1984)**.

3. Result and Discussion

3.1. Growth Parameters

Data pertaining to growth parameters mainly no. of plants plot⁻¹, plant height (cm), number of leaves and number of branches plant⁻¹ are presented in table no. 2 clearly revealed that application of inorganic fertilizers and organic manures in different modes increased these attributes significantly over control. The results revealed that the plant population plot⁻¹ varied in between 128.33 to 132.96 and all the treatments were non-significantly superior to treatment T₁ [100 % RDF]. The treatment combination T₁₁ [100 % N & P, K- 0 % + 100 % K through Vermicompost] gave the maximum plant population plot⁻¹ (132.96). Number of leaves of pigeon pea varied in between 165.2 to 208.6 and all the treatments were significantly superior to T₁ [100 % RDF]. The treatment combination T₁₁ [100 % N & P, K- 0 % + 100 % K through Vermicompost] gave the maximum no. of leaves (208.6). Number of branches plant⁻¹ of pigeon

pea varied in between 22.4 to 32.8 and all the treatments were significantly superior to T₁ [100 % RDF]. The treatment combination T₁₁ [100 % N & P, K- 0 % + 100 % K through Vermicompost] gave the maximum no. of branches plant⁻¹ (32.8). These findings are further supported by **Brar et al. (2004), Verma et al. (2018), and Gupta et al. (2022),**

Table no.-2: Effect of different treatment combinations on growth parameters of pigeon pea

Treatment	Plant population plot⁻¹	Plant height (cm)	No. of leaves	No. of branches plant⁻¹
T₁	128.33	191.62	165.2	22.4
T₂	129.35	193.48	168.9	23.1
T₃	130.89	196.85	176.5	24.2
T₄	130.25	195.49	173.6	23.6
T₅	131.41	198.43	180.3	24.8
T₆	133.71	204.79	193.2	28.0
T₇	134.90	206.34	197.3	29.1
T₈	132.42	200.31	184.6	25.3
T₉	136.53	209.94	202.5	30.7
T₁₀	132.96	202.46	189.4	26.5
T₁₁	138.42	211.38	208.6	32.8
S.E.m±	1.34	5.29	3.58	1.75
C.D. (P= 0.05)	NS	15.87	10.74	5.26

3.2. Yield Components

Data pertaining to yield attributing parameters mainly no. of pod plant⁻¹, no. of seed pod⁻¹ and seed index (g) are presented in table no. 3 clearly revealed that application of inorganic fertilizers and organic manures increased these attributes significantly over control except seed index (g). Maximum no. of pod plant⁻¹ (103.6), no. of seed pod⁻¹ (4.4) and seed index (9.68 g) were recorded under the treatment T₁₁ [100 % N & P, K- 0 % + 100 % K through

Vermicompost] followed by treatment T₉ [100 % N & P, K- 0 % + 100 % K through FYM] with the value 100.5, 4.3 and 9.42 g respectively and the minimum no. of pod plant⁻¹ (85.6) no. of seed pod⁻¹ (3.5) and seed index (8.60 g) was recorded under the treatment T₁ (100 % RDF). These findings are further supported by **Beena *et al.* (2011), Achakzai *et al.* (2012), and Mishra *et al.* (2015),**

Table no.-3: Effect of different treatment combinations on yield attributes of pigeon pea

Treatment	No. of pods plant ⁻¹	No. of Seed pod ⁻¹	Seed index (g)
T ₁	85.6	3.5	8.60
T ₂	87.9	3.5	8.71
T ₃	90.6	3.7	8.85
T ₄	88.5	3.6	8.79
T ₅	92.4	3.8	8.91
T ₆	96.7	4.1	9.15
T ₇	98.9	4.2	9.26
T ₈	93.5	3.9	8.97
T ₉	100.5	4.3	9.42
T ₁₀	95.4	4.0	9.05
T ₁₁	103.6	4.4	9.68
S.E.m±	3.58	0.09	0.08
C.D. (P= 0.05)	10.75	0.27	NS

3.3 Productivity Parameters

It was observed that application of different inorganic fertilizers and organic manures both enhanced the grain yield and straw yield of pigeon pea significantly over its present in table no. 4. Maximum grain yield (17.69 q ha⁻¹) was recorded under the treatment T₁₁ [100 % N & P, K- 0 % + 100 % K through Vermicompost] followed by treatment T₉ [100 % N & P, K- 0 % + 100 % K through FYM] with the value 17.06 q ha⁻¹ and the minimum grain yield (11.25 q ha⁻¹) was

recorded under the treatment T₁ (100 % RDF). Maximum stover yield (52.64 q ha⁻¹) was recorded under the treatment T₁₁ [100 % N & P, K- 0 % + 100 % K through Vermicompost] followed by treatment T₉ [100 % N & P, K- 0 % + 100 % K through FYM] with the value 51.87 q ha⁻¹ and the minimum stover yield (43.65 q ha⁻¹) was recorded under the treatment T₁ (100 % RDF). These findings are further supported by the findings of **Gaikwad *et al.* (2011)**, **Chaudhary *et al.* (2016)** and **Pradeep *et al.* (2017)**

Table no.-4: Effect of different treatment combinations on yields of pigeon pea

Treatment	Grain yield (q ha⁻¹)	Stover yield (q ha⁻¹)
T₁	11.25	43.65
T₂	11.89	43.99
T₃	13.48	45.92
T₄	12.26	44.58
T₅	14.95	46.35
T₆	16.25	49.63
T₇	16.89	50.21
T₈	15.28	47.52
T₉	17.06	51.87
T₁₀	15.97	48.90
T₁₁	17.69	52.64
S.E.m±	0.81	2.05
C.D. (P= 0.05)	2.38	6.16

5. Conclusion

The experimental results indicated that superiority in regard to growth parameters, yield components and productivity parameters *viz.* grain yield (q ha⁻¹), straw yield (q ha⁻¹), with the

use of treatment combination 100 % N & P, K- 0 % + 100 % K through Vermicompost gave in soil ensure highest growth parameters, yield components and productivity, of barley crop as comparison to all the treatments.

6. References

1. **Achakzai. K.M., and Mane. L.J. (2012)** Studies on Growth, Yield and Quality of mung bean as Influenced by Different Levels. *Int.J.Curr.Microbiol.App.Sci* **7(9)**: 2303-2307.
2. **Arora, V. K., Singh, C. B., Sidhu, A. S. and Thind, S. S. (2011).** Irrigation, tillage and mulching effects on soybean yield and water productivity in relation to soil texture. *Agricultural Water Management*. **98(4)**, 563-568.
3. **Beena Saeed, H., Gul, S. Wahab, Y., Durrani, B. Haleema, M., Ayub, A., Muhammad, A. Said and I. Ahma, (2011).** Effect of phosphorus and potassium on seed production of Berseem. *African J. Biotech.* **10**: 63-65.
4. **Brar, M. S., N. Kaur, Racina and Sharma, A. (2004).** Effect of graded doses of potassium on Pea. *J. Pot. Res.* **20**: 100-108.
5. **Chanda, G.C., Bhunia, G. and Chakraborty, S.K. (2011).** The effect of vermicompost and other fertilizers on cultivation of Tomato plants. *Journal of Horticulture and Forestry*. **3**, 42-45.
6. **Chaudhary, M., Singh, S., Babu, S. and Prasad, M. (2011)** Effect of Integrated Nutrient Management on yield and yield attributes of blackgram (*Phaseolus mungo L.*) in an Inceptisol of Varanasi. *International Journal of Applied and pure Science and Agriculture*, Volume 02, Issue 6, e-ISSN: 2394-5532, p-ISSN: 2394-823X.
7. **Gaikwad, R. D., P. R. Patil, N. K. Medhe and V. A. Mane, (2011).** Verification of yield targeting equations with and without organic matter for optimization of yield Pigeon pea. *Green Farming*. **2 (5)**: 530-532.
8. **Garg, N., & Manchanda, G. (2009).** Role of arbuscular mycorrhizae in the alleviation of ionic, osmotic and oxidative stresses induced by salinity in *Cajanus Cajan (L.) Millsp.* (Pigeon pea). *Journal of Agronomy and Crop Science*, **195(2)**, 110-123.
9. **Gomez, K. A., & Gomez, A. A. (1984).** *Statistical procedures for agricultural research.* John wiley & sons.

10. **Gupta, G., Dhar, S., Kumar, A., Choudhary, A. K., Dass, A., Sharma, V. K., & Rajawat, M. V. S. (2022).** Microbes-mediated integrated nutrient management for improved rhizo-modulation, pigeon pea productivity, and soil bio-fertility in a semi-arid agro-ecology. *Frontiers in Microbiology*, *13*, 924407.
11. **Hanway, J.J; and Heidel, H. (1952).** Soil analysis methods as used in Iowa State College, Soil Testing Laboratory. *Iowa Agriculture* **54**: 1-31.
12. **Hawkesford, M., Horst, W., Kichey, T., Lambers, H., Schjoerring, J., Møller, I. S., & White, P. (2012).** Functions of macronutrients. In *Marschner's mineral nutrition of higher plants* (pp. 135-189). Academic press.
13. **Jackson, M.L. (1973).** Soil chemical analysis. Prentice Hall of India Pvt. Ltd, New Delhi.
14. **Mishra, K. S., Parihar, S. S., and Namdeo, K. M. (2015).** Effect of organic and inorganic sources of nutrients on growth, yield and economics of Pigeon pea. *Annals of Plant and Soil Research* *17*(4): 374-376.
15. **Olsen, S.R, Cole, C.V., Watanable, F. S. and Dean, L. A. (1954).** Estimation of available phosphorous in soil by extraction with sodium bicarbonate. *USDA, Cric.* **930**:19- 23
16. **Pradeep, S., Ullasa, M. Y. and Naik A. H. K. (2017).** Effect of different organic nutrient management practices on growth and yield of pigeon pea (*Cajanus cajan* L. Millsp.). 10th *International Conference on Agriculture & Horticulture*, 2017 London, UK.
17. **Ramesh, P, Mohan Singh, A. and Subba Rao. (2005).** Organic Farming: it is relevance to the Indian context. *Current Sci.* *88*(4).
18. **Reay, D., & Reay, D. (2015).** A brief history of nitrogen. *Nitrogen and Climate Change: An Explosive Story*, 9-25.
19. **Singh, K., Rani, A., Padder, S. A., & Gera, R. (2017).** Plant Growth Promoting (PGP) Attributes of stress tolerant Rhizobial isolates from root nodules of Pigeon pea [*Cajanus Cajan* (L.) Millspaugh] growing in Haryana, India. *Int. J. Curr. Microbiol. Appl. Sci.* *12*, 461-473.
20. **Subbiah, B.V. and Asija, C.L. (1956).** A rapid procedure for the estimation of available N in Soil. *Curr. Sci.* **25**:259-260.

21. **Verma, S., Singh, A., Pradhan, S. S., Singh, J. P., & Verma, S. K. (2018).** Effects of organic formulations and synthetic fertilizer on the performance of pigeon pea in eastern region of Uttar Pradesh. *Bangladesh Journal of Botany*, 47(3), 467-471.
22. **Walkley, A. and Black, C. S.A. (1934).** Old piper, S.S. soil and plant analysis. *Soil Sci.* 37:29- 38.
23. **Yadav, K.V., Singh, D.P., Sharma, S.K. and Kishor, K. (2017).** Use of phosphorus for maximization of summer mungbean [*Vigna radiata* (L.) Wilszeck] productivity under sub humid condition of Rajasthan, India. *Journal of Pharmacognosy and Phytochemistry*, 6(4): 01-03.
24. **Tiwari T, Tiwari US, Sharma S, Pandey HP, Sachan R, Kumar D, Kumar S, Singh KK.** Effect of Biofortification with Zinc and Iron on Growth Parameters, Yield Attributes and Yield of Barley (*Hordeum vulgare* L.). *International Journal of Plant & Soil Science*. 2023 Jul 29;35(18):1148-57.