

# The Potential Role of Sleep in Post-Stroke Motor Learning

---

## Abstract

Research in the scientific community has been focusing on identifying the essential aspects that play a role in the movement rehabilitation process in an effort to facilitate motor recovery that is not only efficient but also risk-free. Survivors of strokes may suffer from a variety of movement abnormalities, including weakness, spasticity, dystonia, tremor, chorea, and parkinsonism. The severity of these impairments may differ depending on the size and location of the neuronal lesion(s). It is essential to realize that the use of hypnotics resulted in a momentary improvement in the quality of sleep experienced by those participating in the trial. This improvement may have been a factor that contributed to the fact that patients with insomnia were able to obtain functional and cognitive outcomes that were statistically comparable to those of individuals who did not suffer from insomnia.

*Keywords: parkinsonism, neuronal lesion, dystonia, motor recovery*

## 1. Introduction

Most humans complete surprisingly complex motor behaviours every day (e.g. walking, running, eating). We execute them efficiently and the perceptual and behavioural aspects of the tasks blend together seamlessly. For the 4% of individuals suffering from a functional disability in the UK, (equates to approximately 2.65 million individuals; GOVUK., 2014) however, performing even the most simple motor behaviours can be an arduous task. Stroke is most common cause of adult onset disability. There are an estimated 1.2 million stroke survivors in the UK with approximately 110,000 new strokes each year (Stroke Association, 2018). Of these survivors, two-thirds are expected suffer from long term movement disability (NICE, 2019). Currently, NHS costs for stroke care is £26 billion a year (NICE, 2019) with £5.2 billion spent on social care (Stroke Association, 2017) and £3.4 billion spent on early-supported discharge and rehabilitation [1-4].

Depending on size and location of consequent neuronal lesion(s), stroke survivors may suffer from a variety of movement problems including: weakness, spasticity, dystonia, tremor, chorea and parkinsonism. Falls (average of 6.55 falls suffered per person, per year; equates to circa 720,500 annual falls; Weerdesteyn et al., 2008) and fall related injuries (e.g. disabilities,


fractures) are common complications among stroke survivors with 5% (36025) of overall falls expected to result in a fracture. Thus, patients may undertake motor rehabilitation therapy to assist with recovery from the initial deficit, or, to assist with recovery from a secondary disability suffered as a consequence of stroke [5-8].

In the pursuit of safe and efficacious motor recovery, scientific research has attempted to identify key variables that influence movement rehabilitation. In recent years, promising (albeit limited) evidence suggests that sleep is one key modulator of motor learning and therefore, post-stroke rehabilitation outcomes[9]. Interestingly, it is estimated that 50% of stroke patients suffer from insomnia (with N3 sleep particularly curtailed; Jirakittayakorn and Wongsawat., 2018) and that 5.66% (40,780) of stroke-related falls can be attributed to daytime sleepiness and inattention [10]. Therefore, it is not unreasonable to suggest that sleep-enhancement therapy could be used to assist stroke patients suffering from a primary or secondary post-stroke movement disorder [11].

### **Empirical review**

The rationale underpinning this review, therefore, was to examine and synthesize most recent and relevant literature concerning sleep and post-stroke rehabilitation outcomes. Before reviewing pertinent literature, however, it is first important to consider mechanism(s) by which sleep deprivation (SD) and sleep enhancement may influence post-stroke motor learning [12-14].

Many researchers hypothesize that post-stroke SD originates from hypothalamic-pituitary-adrenal (HPA) axis hyperactivity. Hyperactivity can be caused by either deficit itself or by secondary consequences such as stress or anxiety. In any case, parvocellular neurosecretory neurons within the paraventricular nucleus (PVN) of the hypothalamus are signalled to accelerate production and secretion of corticotrophin-releasing factor (CRF; a neurohormone). This triggers a chemical cascade resulting in uncontrolled glucocorticoid binding activity (see figure 1).



*Figure 1: Schematic Depiction of the biomolecular processes underpinning HPA-axis hyperactivity. Parvocellular neurosecretory neurons within the paraventricular nucleus (PVN) of the hypothalamus are signalled to accelerate the production and secretion of corticotrophin-releasing factor (CRF; a neurohormone) into the hypophyseal portal system (Smith, 2006). CRF is then transported into the anterior pituitary gland, where it binds to its type-1 receptor (CRFR-1) to activate adenylate cyclase. This stimulates adrenocorticotrophic hormone (ACTH) release from the pituitary corticotropes. ACTH then binds melanocortin type 2-receptor in parenchymal cells of the adrenocortical zona fasciculata. This initiates the cAMP pathway which facilitates glucocorticoid (e.g. noradrenaline and dopamine) secretion from the adrenal cortex (see figure 1; Smith, 2006). Substantially elevated levels of noradrenaline and dopamine inhibit NREM sleep and trigger uncontrolled stimulation of lower-affinity  $\alpha$ 1-receptors and D1-receptors respectively. –*

High glucocorticoid binding activity then induces excessive cAMP signalling (Arnsten, 2009) which enables extensive, coordinated shunting of cAMP network inputs (Delmas and Brown, 2005). This promotes amygdala functions (brain area which stimulates fear conditioning and consolidation of emotionally relevant information i.e. ‘bottom-up’ reflexive regulation) but impairs PFC functions (brain area which protects representational knowledge from interference of external or internal distractions, inhibits inappropriate actions and promotes task-relevant operations i.e. ‘top-down’ thoughtful regulation; Arnsten, 2009) [15, 16]. The dorsolateral PFC (DLPFC) interacts with sensory and motor cortices to regulate attention, working memory and action. Thus, a reduction in both persistent firing and tuning of DLPFC neurons causes an individual to attend to irrelevant stimuli in place of relevant stimuli which (further) impairs sleep-onset (cycle of decline), working memory, concentration, alertness

and consequently performance [17]. Ergo, motor learning is not as efficacious whilst deprived of sleep [18].

### **Sleep-induced motor learning**

Sleep-induced motor learning can be divided into 2 key phases [19]: sleep before learning (SBL) and sleep after learning (SAL). Sleep (primarily stage-2 SWS) before learning primes the cerebral cortex (short-term memory store) and hippocampus (consolidates novel memories) for receiving new information by transferring and consolidating newly acquired, relevant memories into neocortical structures (long-term memory store) to restore hippocampal encoding capacity for next day of practice [20, 21]. SAL then enables newly encoded memories to be consolidated and stored as engrams within the CNS (akin to that of a computer 'file transfer') to facilitate future performance [22].

According to the classical model of sleep dependent ML, structures contained within medial temporal lobe (MTL; most notably the hippocampal complex) are central to both SBL and SAL [23]. More specifically, these structures link together patterns of cortical activation that were present during initial encoding to facilitate formation and retrieval of novel motor memories. Alongside binding patterns of cortical activation, hippocampus has been suggested to play a key role in offline reactivation of such networks. It is hypothesized that repetition of these reactivation processes through multiple bouts of nightly sleep cycles strengthen and thereby, reinforce initially weak neocortical connections [24, 25]. Thus, initial dependence on MTL binding subsides as newly acquired information is progressively integrated into neocortical circuits (LTM; see figure 2). Over time, therefore, offline reinforcement enables newly acquired information to be activated within the cortex, independently of the hippocampus.

An advanced version of this classical hypothesis has since been proposed by Buzsaki [26]. Buzsaki [27] suggests a model of consolidation that is dependent upon two key states of hippocampal activity. The first pertains to a state of wakeful 'recording' which shifts to a second state of offline 'playback'. Playback lasts for up to 3-hours and is characterised by bursts of neural activity that occur during SWS and are termed 'sharp-waves' or 'sleep spindles' (conducted within hippocampal place cells and cortical structures respectively). Interestingly, these bursts of activity are replayed at a speed that is approximately 20 times faster than prior online experience and may represent a specific spatial location that occurred

during ‘recording’[28, 29]. Thus, neuronal activity experienced whilst awake may be replayed nocturnally during SWS, potentially representing motor memory processing.

Together, therefore, both versions of classical hypothesis suggest two key predictions regarding offline motor learning [30]. First, is that offline-consolidation promotes and strengthens formation of cortico-cortical connections to produce motor memories that are more resistant to interference (SAL). Second, is that offline strengthening of these cortico-cortical connections reduces dependence on MTL binding by increasing hippocampal encoding capacity (SBL).





Figure 2: Schematic Representation of Offline Learning (Walker, 2009) –



In support of sleep-induced learning, Siengsukon and Boyd [31] observed that sleep may enhance post-stroke movement learning. Researchers split stroke patients and healthy controls into either a sleep (baseline test in evening and post-test in morning following sleep) or no-sleep group (baseline test in morning and post-test that evening). Each group were

instructed to complete a continuous tracking task (CTT; use of a hand-driven joystick to track a target moving horizontally across a computer screen). Interestingly, sleep-induced improvements in spatial tracking accuracy (more negative score denotes less error; -1.4 vs -0.3 = 21.4% improvement;  $p=0.014$  vs  $p=0.556$  respectively) and temporal tracking accuracy (positive scores indicate improved time lag of tracking at retention; 60ms vs -2ms = 31% improvement;  $p=0.036$  vs  $p=0.962$ ) were only evident in the sleep-stroke group. This indicates that sleep enhances ML in stroke patients but not healthy individuals.

In agreement with Siengsukon and Boyd [32], Joa et al., 2017 conducted a multi-centre observational study on mild-moderate stroke patients ( $n=280$ ) and reported that Berg Balance Scale (Korean version) score improvements were significantly lower in a disturbed-sleep group (assessed by a health professional using Diagnostic Statistical Manual of Mental Disorders criteria to define patients with any sleep disturbance) compared to a normal sleep group. This effect was even maintained after adjusting for confounders such as age, sex and hypnotics usage. Interestingly, when results were analysed according to stroke-severity, significant improvements in balance were only maintained by moderate-stroke sufferers and not by mild-stroke sufferers. This suggests that sleep disturbance has a negative influence on functional recovery following moderate, but not mild stroke.

Similarly, Iddagoda et al., [33] reported an inverse association between poor sleep quality (assessed through PSQI; completed at baseline (pre-stroke) and prior to discharge (post-stroke)) and rehabilitation outcomes as assessed by Functional Independence Measure (assesses degree of disability depending on patients score in 18 motor and cognitive function categories;  $R_s. -0.317$ ,  $P = 0.005$ ) in a prospective cohort study of 104 Adult, Australian stroke patients from two major stroke units in Western Australia. This suggests that poor sleep quality impedes post-stroke motor recovery.

Interestingly, it was also reported by Iddagoda et al., [34] that sedatives were used by 18.2% of patients with no impact on either sleep quality or rehabilitation being identified. Similar findings have also been reported by Kim et al., [35], who identified no significant effect of hypnotics-use on sleep-patterns in a sample of subacute stroke patients ( $n=30$ ) with insomnia (experimental group; hypnotics consumed) vs those without insomnia (control; placebo) at 3-weeks follow-up. It should be noted, however, that hypnotics administration did temporarily enhance sleep among the experimental condition, which may have contributed toward insomnia patients obtaining functional and cognitive outcomes that were (statistically)

comparable to those achieved by non-insomnia patients. Given that improvements in sleep were not maintained at 3-weeks follow-up, however, it is doubtful that optimal sleep-induced motor benefits were attained by the insomnia group.

This is concerning given that hypnotics are the mainstay prescription for post-stroke insomnia. More worrying still is that hypnotics use has been linked with a 2-fold increase in depression, over 3-times as many in-hospital falls, a 4-fold increase in overall mortality [36] and a consequential overall cost (of hypnotics) to the NHS of approximately £72 million per annum [37]. Moreover, recent research has also identified sleep induced by some hypnotics (e.g. zolpidem) may actually damage neuronal connections and impair memory rather than enhance learning [38]. This suggests that stroke patients will actually experience detriments and not benefits to their motor performance after using hypnotics to enhance sleep.

There are noteworthy limitations within discussed research that may limit reliability of results. First, is recurrent utilisation of subjective self-report tools when assessing sleep (e.g. PSQI). Falck et al., [39] identified that stroke survivors sleep for 0.25-hours less than they subjectively report. Therefore, true sleep-post stroke movement learning effects may not have been accurately depicted within research conducted until present. In future, experimenters should utilise objective tools (e.g. polysomnography) to minimise measurement errors whilst assessing sleep.

Second, is cross-sectional design of multiple studies. Cross-sectional research utilises a 'snapshot' measurement of both exposure (sleep) and outcome (post-stroke movement learning). Therefore, it is not possible to infer causal relationships from such research because reverse-causality cannot be denied (i.e. cannot refute that alterations in movement learning influenced sleep), and because other confounding variables (e.g. diet) may have influenced result outcomes. Forthcoming research should utilise longitudinal designs to ease issues related to reverse-causality, and also, so that effects of sleep on post-stroke movement learning effects can be more accurately assessed.

A final issue is that none of the studies were crossover or single subject design. Because there are multiple variables (e.g. age, diet) that may confound the sleep-movement learning relationship, it cannot be ascertained that individuals from distinct groups would have reacted to different conditions (e.g. sleep vs no-sleep) in same way. Future research studies should utilise randomized cross-over designs so that confounding variables have less influence on result outcomes.

## **Conclusion**

In conclusion, discussed research suggests a promising role of sleep for post-stroke motor learning. However, current evidence is derived primarily from cross-sectional studies which utilised self-reported sleep assessments. Due to such methodological limitations, a beneficial role of sleep for movement learning after stroke cannot be established. In future, research studies should utilise either prospective, longitudinal or randomized crossover designs and objective sleep assessments so that true effects of sleep on post-stroke motor learning can be more accurately portrayed.

References:

UNDER PEER REVIEW

1. Hasan F, Gordon C, Wu D, Huang HC, Yuliana LT, Susatia B et al. Dynamic prevalence of sleep disorders following stroke or transient ischemic attack: systematic review and meta-analysis. *Stroke*. 2021;52(2):655–63. A systematic review and meta-analysis that evaluated the prevalence and evolution of main sleep disorders (sleep-disordered breathing, insomnia, restless legs syndrome, and periodic limb movements of sleep) from acute to chronic phase of stroke.
2. Lo EH, Albers GW, Dichgans M, Donnan G, Esposito E, Foster R et al. Circadian biology and stroke. *Stroke*. 2021;52(6):2180–90. A review that assessed molecular, cellular, and physiological pathways in circadian biology in relation to the clinical consequences in stroke.
3. Wang H, Sun J, Sun M, Liu N, Wang M. Relationship of sleep duration with the risk of stroke incidence and stroke mortality: an updated systematic review and dose-response meta-analysis of prospective cohort studies. *Sleep Med*. 2022;90:267–78. A meta-analysis on studies that evaluated the correlation between the risk of stroke and sleep duration.
4. Bassetti CLA, Randerath W, Vignatelli L, Ferini-Strambi L, Brill AK, Bonsignore MR et al. EAN/ERS/ESO/ESRS statement on the impact of sleep disorders on risk and outcome of stroke. *Eur J Neurol*. 2020;27(7):1117–36. A statement of four major European medical organizations (European Academy of Neurology, European Respiratory Society, European Stroke Organization, European Sleep Research Society) on current knowledge and management of sleep disorders in terms of stroke risk and outcome.
5. Lisabeth LD, Sanchez BN, Lim D, Chervin RD, Case E, Morgenstern LB et al. Sleep-disordered breathing and poststroke outcomes. *Ann Neurol*. 2019;86(2):241–50. A large cohort study evaluating the impact of sleep-disordered breathing on functional and neurological outcome in patients with stroke.
6. Tanayapong P, Kuna ST. Sleep disordered breathing as a cause and consequence of stroke: a review of pathophysiological and clinical relationships. *Sleep Med Rev*. 2021;59:10149. A review with an educational value evaluating the pathophysiological pathway of sleep-disordered breathing in patients with stroke.
7. Baillieux S, Bailly S, Detante O, Alexandre S, Destors M, Clin R et al. Sleep-disordered breathing and ventilatory chemosensitivity in first ischaemic stroke patients: a prospective cohort study. *Thorax*. 2022;77:1006–14. A large cohort of study describing the peculiar ventilatory chemosensitivity characteristics of patients with acute stroke.
8. Lapow JM, Dicipinigaitis AJ, Pammal RS, Coghill GA, Rechester O, Feldstein E, et al. Obstructive sleep apnea confers lower mortality risk in acute ischemic stroke patients treated with endovascular thrombectomy: National Inpatient Sample analysis 2010–2018. *J Neurointerv Surg*. 2021;neurintsurg-2021-018161. <https://doi.org/10.1136/neurintsurg-2021-018161>. The first study that evaluated the effect of sleep-disordered breathing on the outcome in terms of mortality in patients with acute ischemic stroke treated with mechanical thrombectomy.
9. Brown DL, Jiang X, Li C, Case E, Sozener CB, Chervin RD, et al. Sleep apnea screening is uncommon after stroke. *Sleep Med*. 2019;59:90–3. A study that showed as just a minority of patients with ischemic stroke are screened for sleep apnea.
10. Boulos MI, Kamra M, Colelli DR, Kirolos N, Gladstone DJ, Boyle K, et al. SLEAP SMART (sleep apnea screening using mobile ambulatory recorders after TIA/stroke): a randomized controlled trial. *Stroke*. 2022;53(3):710–8. A randomized control study evaluating the efficacy, feasibility and cost of home sleep apnea test compared to the polysomnography.
11. Bernardini A, Brunello A, Gigli GL, Montanari A, Saccomanno N. AIOSA: An approach to the automatic identification of obstructive sleep apnea events based on deep learning. *Artif Intell Med*. 2021;118:102133. A study that applied artificial intelligence techniques in order to

- automatically identify obstructive sleep apnea relying on routinely recorded vital signs in stroke units.
12. Lin HJ, Yeh JH, Hsieh MT, Hsu CY. Continuous positive airway pressure with good adherence can reduce risk of stroke in patients with moderate to severe obstructive sleep apnea: an updated systematic review and meta-analysis. *Sleep Med Rev.* 2020;54:101354. A meta-analysis demonstrating the efficacy of continuous positive airway pressure for stroke prevention in patients with moderate to severe obstructive sleep apnea and good adherence.
  13. Kotzian ST, Saletu MT, Schwarzsinger A, Haider S, Spatt J, Kranz G, et al. Proactive telemedicine monitoring of sleep apnea treatment improves adherence in people with stroke—a randomized controlled trial (HOPES study). *Sleep Med.* 2019;64:48–55. A randomized controlled trial that demonstrated that telemedicine ameliorates the adherence of apnea treatment in post-stroke patients.
  14. Yeghiazarians Y, Jneid H, Tietjens JR, Redline S, Brown DL, El-Sherif N, et al. Obstructive sleep apnea and cardiovascular disease: a scientific statement from the American Heart Association. *Circulation.* 2021;144(3):e56–e67. A statement of American Heart Association on management of obstructive sleep apnea in patients with cardiovascular diseases.
  15. Wang YH, Wang J, Chen SH, Li JQ, Lu QD, Vitiello MV, et al. Association of longitudinal patterns of habitual sleep duration with risk of cardiovascular events and all-cause mortality. *JAMA Netw Open.* 2020;3(5):e205246. A cohort study on a large population demonstrating that short sleep as well as unstable sleep pattern are associated with higher risk of cardiovascular events.
  16. Tobaldini E, Fiorelli EM, Solbiati M, Costantino G, Nobili L, Montano N. Short sleep duration and cardiometabolic risk: from pathophysiology to clinical evidence. *Nat Rev Cardiol.* 2019;16(4):213–24. A review describing the effects of short sleep duration on cardiovascular health and diseases.
  17. Rollo E, Brunetti V, Scala I, Callea A, Marotta J, Vollono C, et al. Impact of delirium on the outcome of stroke: a prospective, observational, cohort study. *J Neurol.* 2022. <https://doi.org/10.1007/s00415-022-11309-2>.
  18. Baylan S, Griffiths S, Grant N, Broomfield NM, Evans JJ, Gardani M. Incidence and prevalence of post-stroke insomnia: a systematic review and meta-analysis. *Sleep Med Rev.* 2020;49:101222. A systematic review and meta-analysis on insomnia characteristics and changes in insomnia prevalence over time in post-stroke patients.
  19. Jaramillo V, Jendoubi J, Maric A, Mensen A, Heyse NC, Eberhard-Moscicka AK, et al. Thalamic influence on slow wave slope renormalization during sleep. *Ann Neurol.* 2021;90(5):821–33. A study evaluating by means of high density electroencephalography and polysomnography patients with thalamic stroke showing a role of thalamus in the synaptic homeostasis.
  20. Wang XX, Feng Y, Tan EK, Ondo WG, Wu YC. Stroke-related restless legs syndrome: epidemiology, clinical characteristics, and pathophysiology. *Sleep Med.* 2022;90:238–48. A review on epidemiology, clinical characteristics, pathophysiology, and impact on functional outcome of stroke-related RLS.
  21. Tellenbach N, Schmidt MH, Alexiev F, Blondiaux E, Cavalloni F, Bassetti CL, et al. REM sleep and muscle atonia in brainstem stroke: a quantitative polysomnographic and lesion analysis study. *J Sleep Res.* 2022:e13640. <https://doi.org/10.1111/jsr.13640>.
  22. Alexiev, F., et al., *Sleep-disordered breathing and stroke: chicken or egg?* *J Thorac Dis*, 2018. **10**.
  23. Baillieux, S., et al., *Sleep apnoea and ischaemic stroke: current knowledge and future directions.* *Lancet Neurol*, 2022. **21**.
  24. Barlinn, K., et al., *Autotitrating bilevel positive airway pressure in large vessel steno-occlusive stroke patients with suspected sleep apnea: a multicenter randomized controlled study.* *Front Neurol*, 2021. **12**.

25. Bernardini, A., et al., *OSASUD: A dataset of stroke unit recordings for the detection of obstructive sleep apnea syndrome*. *Sci Data*, 2022. **9**.
26. Bernasconi, C., et al., *SAS CARE 2 - a randomized study of CPAP in patients with obstructive sleep disordered breathing following ischemic stroke or transient ischemic attack*. *Sleep Med X*, 2020. **2**.
27. Bivard, A., et al., *MIDAS (modafinil in debilitating fatigue after stroke): a randomized, double-blind, placebo-controlled, cross-over trial*. *Stroke*, 2017. **48**.
28. Blanchard, M., et al., *Hypoxic burden and heart rate variability predict stroke incidence in sleep apnoea*. *Eur Respir J*, 2021. **57**.
29. Boulos, M.I., et al., *Trials in sleep apnea and stroke: learning from the past to direct future approaches*. *Stroke*, 2021. **52**.
30. Brown, D.L., et al., *Sleep for stroke management and recovery trial (Sleep SMART): rationale and methods*. *Int J Stroke*, 2020. **15**.
31. Brown, D.L., et al., *Prediction of sleep-disordered breathing after stroke*. *Sleep Med*, 2020. **75**.
32. Brown, D.L., et al., *Sleep-disordered breathing is associated with recurrent ischemic stroke*. *Stroke*, 2019. **50**.
33. Brown, D.L., et al., *Overnight rostral fluid shifts exacerbate obstructive sleep apnea after stroke*. *Stroke*, 2021. **52**.
34. Brunetti, V., et al., *Autonomic nervous system modifications during wakefulness and sleep in a cohort of patients with acute ischemic stroke*. *J Stroke Cerebrovasc Dis*, 2019. **28**.
35. Camilo, M.R., et al., *Diagnostic accuracy of positive airway pressure device for sleep apnea detection in acute stroke patients*. *Stroke*, 2020. **51**.
36. Cananzi, S.G., et al., *Obstructive sleep apnea intensifies stroke severity following middle cerebral artery occlusion*. *Sleep Med*, 2020. **67**.
37. Cao, W., et al., *The association between sleep breathing impairment index and cardiovascular risk in male patients with obstructive sleep apnea*. *Nat Sci Sleep*, 2022. **14**.
38. Castello-Branco, R.C., et al., *Association between risk of obstructive sleep apnea and cerebrovascular reactivity in stroke patients*. *J Am Heart Assoc*, 2020. **9**.
39. Chen, C.Y. and C.L. Chen, *Recognizable clinical subtypes of obstructive sleep apnea after ischemic stroke: a cluster analysis*. *Nat Sci Sleep*, 2021. **13**.
40. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4263505/#!po=5.68182>
41. <https://nyaspubs-onlinelibrary-wiley-com.libproxy.ncl.ac.uk/doi/full/10.1111/j.1749-6632.2009.04416.x> = key
42. [https://www.stroke.org.uk/sites/default/files/costs\\_of\\_stroke\\_in\\_the\\_uk\\_report\\_-\\_executive\\_summary\\_part\\_2.pdf](https://www.stroke.org.uk/sites/default/files/costs_of_stroke_in_the_uk_report_-_executive_summary_part_2.pdf) = stroke association