

Original Research Article

**Indoor air quality and hazards associated with laundry operations in
Ibadan metropolis, Nigeria**

Abstract

Background and Objective: Launderers are exposed to potentially harmful chemicals during washing ironing and dyeing; exposures potentially result has negative health effects, ranging from skin and throat irritation to carcinogenicity, and negative effects on the environment.

Comment [HI1]: Delete

Aims: Limited information existed on indoor air quality (IAQ) in laundry environment in developing countries especially Nigeria. This study assessed indoor air quality hazards associated with laundry operations in Ibadan metropolis, Nigeria

Comment [HI2]: Delete

Materials and Methods: A descriptive cross-sectional survey was used in the study that involving a simple random selection of 50 out of 100 consenting laundry operators in five local governments, Ibadan North (IBN); Ibadan North-East (IBNE); Ibadan South-West (IBSW), Ibadan North-West (IBNW) and Ibadan South-East (IBSE).

Comment [HI3]: Delete

Results: Temperature and Relative Humidity (RH) were monitored using Multi Testers N21FR. The IAQ parameters: Particulate Matters (PM₁₀) concentration, carbon dioxide (CO₂), airborne Total Bacteria Count (TBC) and Total Fungi Count (TFC) were determined using a Met One GT 321, CO₂ meter and an air sampler respectively. Measurements were taken in the morning and evening for 8 weeks. Values obtained for temperature, RH, PM₁₀, TBC and CO₂ were respectively compared with WHO limits of 26°C, 60%, 50µg/m³ and ≤5x10²CFU/m³ and 1,000ppm (ASHRAE)respectively. Data

Comment [HI4]: Delete

were analysed using descriptive statistics with SPSS (version 20), Pearson Correlation and Chi-square at $p < 0.05$. The respective mean indoor temperature ($^{\circ}\text{C}$), RH (%), PM_{10} ($\mu\text{g}/\text{m}^3$) and CO_2 (ppm) levels for morning across the local governments were IBN (27.4 ± 2.5 , 67.5 ± 6.8 , 7.2 ± 1.1 , 337.9 ± 57.8); IBNE (27.5 ± 2.0 , 65.5 ± 3.5 , 7.6 ± 9.6 , 377.4 ± 83.3); IBSW (27.6 ± 2.3 , 68.7 ± 4.0 , 7.2 ± 0.8 , 326.5 ± 44.7); IBNW (26.8 ± 2.6 , 64.9 ± 6.0 , 7.2 ± 9.6 , 363.5 ± 75.3) and IBSE (27.6 ± 1.5 , 64.9 ± 3.2 , 7.5 ± 1.2 , 371.6 ± 79.6) while evening readings were IBN (28.2 ± 2.4 , 66.7 ± 6.7 , 7.2 ± 9.6 , 409.2 ± 68.3); IBNE (28.4 ± 2.1 , 68.3 ± 4.2 , 8.0 ± 0.9 , 352.0 ± 54.7); IBSW (28.4 ± 1.8 , 69.2 ± 4.5 , 8.0 ± 0.8 , 326.2 ± 41.2); IBNW (28.4 ± 2.1 , 68.8 ± 4.4 , 7.8 ± 0.9 , 356.2 ± 62.5) and IBSE (27.8 ± 1.4 , 66.3 ± 4.2 , 7.9 ± 0.9 , 352.0 ± 54.7). The mean indoor morning TBC were 82.9 ± 9.0 , 141.4 ± 10.2 , 88.5 ± 7.0 , 163.9 ± 12.6 and 88.5 ± 7.0 in Ibadan North, Ibadan North West, Ibadan North East, Ibadan South West and Ibadan South East respectively and mean indoor afternoon TBC were 82.8 ± 6.1 , 155.7 ± 6.3 , 87.9 ± 4.4 , 183.7 ± 8.9 and 87.9 ± 4.4 in Ibadan North, Ibadan North West, Ibadan North East, Ibadan South West and Ibadan South East respectively. Both indoor and outdoor mean total bacterial counts (TBC) CFU/m^3 values were much lower than the $\leq 5 \times 10^2 \text{CFU}/\text{m}^3$ recommended guideline limit by WHO. Indoor RH and temperature were above WHO guideline limit, while PM_{10} and CO_2 are below the WHO and ASHRAE guideline limit respectively and the TBC and TFC were within the WHO guideline limit in all the study locations. Therefore, improved cross ventilations are advocated to enhance indoor air quality in the laundries.

Conclusion: In laundry environment in Ibadan metropolis, Nigeria, IAQ appeared compromised. Mechanical ventilation systems in laundry environment need to be improved with respect to operation and maintenance to better maintain thermal comfort

Comment [HI5]: Find a way of summarizing this information

Comment [HI6]: Delete

parameters and reduce levels of chemical pollutants emitted by launderers and product use during services.

Comment [HI7]: (1)The abstract is lengthy
(2) Don't paragraph the abstract

Keyword: Indoor air quality, Health hazard, Laundry

1. INTRODUCTION

Clean air is a basic requirement of life [1]. Air is essential to life. It is important that human breath in good and quality air to avoid been faced with the health problems of polluted air intake. The indoor air quality can be degraded and polluted by human activities within the building and also by the nature of surrounding outdoor air. Air pollutants have several health impacts which range from mild to severe symptoms. It is important to ensure the air within buildings that are less polluted to prevent risk of exposing occupants to pollutants.

Comment [HI8]: Correct the sentence

Comment [HI9]: Correct this sentence

In Nigeria, there was a time a families dying overnight as a result of generator fumes.

Comment [HI10]: Correct this sentence

Indoor Air quantity is very significant because degradation of this quality leads to acute exposure to air pollutants within a confined area.

Air is an essential requirement for live without which living organisms would die. Quality air is essential for health. This has become a thing of concern across the globe because most human activities are carried out within buildings such as offices, schools, restaurants homes and so on. The quality of air within all buildings (laundry buildings are inclusive) should be maintained to ensure healthy living.

Comment [HI11]: Separate the two words with comma

According to Godson et al., indoor air pollution poses health risks to all, and the severity depends on age, sex, physiological conditions and individual predisposition. Indoor air pollution usually occurs due to pollution from surrounding outdoor air, building

Comment [HI12]: Incomplete citation

materials and activities within the **building**. Ventilation is a major factor within the building upon which the severity of air pollution is dependent. Health issues associated with indoor air pollution include dizziness, sneezing, coughing, nasal congestion, wheezing, and diseases of the **lungs**. **Evidences** indicated that developing countries are at higher risk of poor indoor air quality due to poor ventilation system in most homes, buildings and inefficient cooking materials [2].

Comment [HI13]: Insert reference

Comment [HI14]: Insert reference

Comment [HI15]: Use suitable word

In the last several years, a growing body of scientific evidence has indicated that the air within homes and other buildings can be more seriously polluted than the outdoor air in even the largest and most industrialized **cities**. Other research indicates that people spend approximately 90 percent of their time **indoors**. Thus, for many people, the risks to health may be greater due to exposure to air pollution indoors than outdoors.

Comment [HI16]: Insert reference

Comment [HI17]: Insert reference

In addition, people who may be exposed to indoor air pollutants for the longest periods of time are often those most susceptible to the effects of indoor air pollution. Such groups include the young, the elderly and the chronically ill, especially those suffering from respiratory or cardiovascular **disease**.

Comment [HI18]: Insert reference

Indoor Air Quality (IAQ) is an increasing concern in the world today. In fact, the mere presence of people in a building can significantly alter indoor air quality. The quality of air inside homes, offices, schools, laundry centers, public buildings, health care facilities or other private and public buildings where people spend a large part of their life is an essential determinant of healthy life and people's well-being [3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, and 15]. Problems in the indoor air of workplaces are issues which occupational health care often needs to consider when evaluating the health risks of a

work environment. On the other hand, good indoor air quality has a beneficial effect on the health of employees, the social atmosphere at work, and productivity in offices. [16, 17, 18]. It was reported that up to 30% of employees in new or renovated buildings expressed an unusually high number of complaints concerning the work environment, enabling classification of the buildings as “sick” [19]. This appeared to be a problem especially in countries with a colder climate. Working in these problem buildings may cause respiratory symptoms (stuffy and irritated nose, rhinitis, cough, sore throat, and shortness of breath), skin symptoms, as well as general symptoms (fatigue, headache, fever), all of which are typical to the sick building syndrome (SBS) [20, 21, and 22].

Comment [HI19]: Inappropriate way of beginning a sentence

Indoor pollution sources that release gases or particles into the air are the primary cause of indoor air quality problems in several building. Inadequate ventilation in laundry environment can increase indoor pollutant levels by not bringing in enough outdoor air to dilute emissions from indoor sources and by not carrying indoor air pollutants out of the home. High temperature and humidity levels can also increase concentrations of some pollutants. The impact of indoor air pollution on man may consist of undesired health effects of different types, ranging from sensory annoyance or discomfort to severe health injuries.

Comment [HI20]: Use space instead of home

Comment [HI21]: This is not the way to end an introduction section of your paper. What do you intend to do?

2.0 MATERIALS AND METHODS

2.1 STUDY AREA

The study area was Ibadan. Ibadan is the largest indigenous city in West Africa and is located in the South Western part of Oyo State of Nigeria. The population of central Ibadan, including five local governments; Ibadan North, Ibadan North-East, Ibadan

North-West, Ibadan South-East and Ibadan South West. LGAs, is 1 338 659 according to census results for 2006, covering an area of 128 km².

Comment [HI22]: Inadequate description of a study area. You need to provide information regarding the Latitude, Longitude, and the climate of the study area.

2.2 DESCRIPTION OF SAMPLING LOCATIONS

Twenty (20) laundry houses were purposefully selected from each local government area. This gave a total of 100 laundry houses from 5 Local Government areas. In each of the laundry houses, 5 respondents were randomly selected for completion of the questionnaire.

Comment [HI23]: The number of people in each of the laundry house was not specified. Hence, selecting 5 respondent from each of the laundry house may not give a good representation. And if so, you may have problem of external validity.

Comment [HI24]: I suggest your MATERIALS AND METHODS SHOULD START HERE AS SECTION 3.0

2.3 DATA COLLECTION

The study involved four methods of data collection namely air quality monitoring (Particulate matter, temperature, relative humidity, CO₂, microbial load), survey (questionnaire administration and observational checklist), laboratory analysis and statistical methods (Data management and statistical analysis). Temperature and Relative Humidity (RH) were monitored using Multi Testers N21FR. The IAQ parameters: Particulate Matters (PM₁₀) concentration, carbon dioxide (CO₂), airborne Total Bacteria Count (TBC) and Total Fungi Count (TFC) were determined using a Met One GT 321, CO₂ meter and an air sampler respectively. Values obtained for temperature, RH, PM₁₀, TBC and CO₂ were respectively compared with WHO limits of 26°C, 60%, 50µg/m³ and ≤5x10² CFU/m³ and 1,000ppm (ASHRAE) respectively.

2.4 DATA MANAGEMENT AND STATISTICAL ANALYSIS

Data was entered and analyzed using descriptive statistics with SPSS (version 20), Pearson Correlation and Chi-square at p<0.05. Descriptive statistics was used to

summarize data. Mean \pm Standard Deviation (SD) and range was calculated for Temperature, Relative Humidity (RH), Particulate Matters (PM₁₀), microbial burden and carbon dioxide (CO₂) for morning and afternoon and compared with WHO guidelines of 26°C, 60%, 50µg/m³, $\leq 5 \times 10^2$ CFU/m³ and ASHRAE guideline of 1,000ppm respectively.

3.0 RESULTS & discussion

3.1 METEOROLOGICAL PARAMETERS OF INDOOR AND OUTDOOR AIR QUALITY OF THE SELECTED

LAUNDRIES

3.1.1 RELATIVE HUMIDITY (%).

The results of indoor and outdoor relative humidity of the selected laundry houses are presented in Table 1. Both the indoor and outdoor mean relative humidity (%) values were higher than the WHO recommended guideline limit of 60%.

3.1.2 TEMPERATURE READINGS (°C).

The results of the indoor and outdoor temperature of the selected laundry houses are presented in Table.2.Both the indoor and outdoor mean temperature (°C) values were higher than the WHO recommended guideline limit of 26°C.

3.1.3 AIR CO₂ CONCENTRATION (PPM).

The results of the indoor and outdoor CO₂ concentrations of the selected laundry houses are presented in Table.3. Both the indoor and outdoor mean CO₂ concentrations (ppm) values were much lower than the ASHRAE recommendation limits of 1000ppm.

3.1.4 INDOOR CONCENTRATION OF PARTICULATE MATTER

Comment [HI25]: Use 'and and i suggest this section to be 4.0

Comment [HI26]: Use upper case letters

Comment [HI27]: Never use an orphan section. Write something for this section before you start 3.1.

Comment [HI28]: Disjointed

Comment [HI29]: Same as HI26

The mean concentration for indoor PM₁₀ readings ($\mu\text{g}/\text{m}^3$) for both morning and afternoon periods in the selected local governments is represented in table.4. Both the indoor and outdoor mean PM₁₀ ($\mu\text{g}/\text{m}^3$) values were much lower than the WHO recommendation guideline limit of 50 $\mu\text{g}/\text{m}^3$.

UNDER PEER REVIEW

Table.1: Mean Indoor and Outdoor Relative Humidity (%) across the five local governments (%)

Location		Air Relative Humidity (%)						WHO Limit
		Morning			Afternoon			
		Min	Max	Mean±SD	Min	Max	Mean±SD	
Indoor	Ibadan North	52.6	78.0	67.5±6.8	60.1	78.8	66.7±6.7	60
	Ibadan Northwest	56.1	75.1	64.9±6.0	63.3	77.9	68.8±4.4	
	Ibadan Northeast	60.8	70.8	65.5±3.5	63.2	75.6	68.3±4.2	
	Ibadan Southwest	62.7	75.6	68.7±4.0	62.5	77.0	69.2±4.5	
	Ibadan Southeast	59.0	69.0	64.9±3.2	60.9	75.1	66.3±4.2	
Outdoor	Ibadan North	58.2	77.2	66.5±5.9	55.2	63.0	60.7±2.4	
	Ibadan Northwest	59.2	73.3	66.4±3.6	58.2	69.1	65.6±3.9	
	Ibadan Northeast	55.7	77.8	66.8±5.4	57.4	68.6	64.5±3.9	
	Ibadan Southwest	58.0	78.0	67.5±6.5	57.1	68.6	62.8±3.7	
	Ibadan Southeast	57.9	75.8	66.1±4.5	55.6	66.9	62.4±4.0	

Table.2: Mean Indoor and Outdoor Temperature Readings across the five local governments (⁰C)

	Location	Indoor air temperature(⁰ C)						WHO Limit
		Morning			Afternoon			
		Min	Max	Mean±SD	Min	Max	Mean±SD	
Indoor	Ibadan North	24.8	33.7	27.4±2.5	25.5	34.2	28.2±2.4	26
	Ibadan Northwest	23.4	32.9	26.8±2.6	26.8	33.8	28.4±2.1	
	Ibadan Northeast	24.6	31.2	27.5±2.0	24.0	30.0	27.1±2.3	
	Ibadan Southwest	24.8	33.0	27.6±2.3	26.7	33.0	28.4±1.8	
	Ibadan Southeast	24.5	29.5	27.6±1.5	25.1	29.9	27.8±1.4	
Outdoor	Ibadan North	23.8	33.9	26.1±2.8	25.9	33.6	27.6±2.3	
	Ibadan Northwest	25.5	33.7	26.8±2.4	27.4	34.2	28.7±2.0	
	Ibadan Northeast	24.6	31.2	27.5±2.0	24.0	30.0	27.1±2.3	
	Ibadan Southwest	24.8	33.2	27.1±2.5	26.2	36.3	29.1±2.8	
	Ibadan Southeast	23.0	30.0	26.0±2.4	23.9	28.1	25.6±1.6	

Table.3: Mean Indoor and Outdoor Air CO₂ Concentration across the five local governments (ppm).

	Location	Air CO ₂ (ppm)						ASHRAE Limit
		Morning			Afternoon			
		Min	Max	Mean±SD	Min	Max	Mean±SD	
Indoor	Ibadan North	244.0	419.0	337.9±57.8	316.0	446.	355.0±45.7	1000
	Ibadan Northwest	251.5	476.0	363.5±75.3	288.0	473.	356.2±62.5	
	Ibadan Northeast	268.0	511.5	377.4±83.3	284.5	439.	352.0±54.7	
	Ibadan Southwest	257.5	378.0	326.5±44.7	260.5	393.	326.2±41.2	
	Ibadan Southeast	262.0	472.5	371.6±79.6	299.0	356.	321.7±19.8	
Outdoor	Ibadan North	240.	476.0	358.5±82.1	323.	502.	409.2±2.3	
	Ibadan Northwest	239.	394.0	327.9±56.0	330.	490.	395.5±56.3	
	Ibadan Northeast	246.	384.5	326.8±51.9	347.	380.	370.0±10.8	
	Ibadan Southwest	250.	470.5	363.6±80.8	336.	514.	424.5±66.5	
	Ibadan Southeast	240.	384.	311.3±45.5	339.	371.	355.0±10.7	

Table 4: Mean Indoor and outdoor Concentration of PM₁₀ across the five local governments (µg/m³).

	Location	Air PM ₁₀ (µg/m ³)						WHO limits
		Morning			Afternoon			
		Min	Max	Mean±SD	Min	Max	Mean±SD	
Indoor	Ibadan North	6.4	10.3	7.2±1.1	7.2	9.6	7.2±9.6	50
	Ibadan Northwest	7.2	9.6	7.2±9.6	6.8	9.8	7.8±0.9	
	Ibadan Northeast	6.9	9.9	7.6±9.6	7.1	9.9	8.0±0.9	
	Ibadan Southwest	6.8	9.3	7.2±0.8	7.2	9.8	8.0±0.8	
	Ibadan Southeast	6.8	9.7	7.5±1.2	7.0	9.8	7.9±0.9	
Outdoor	Ibadan North	5.8	10.2	7.4±1.6	6.1	10.3	8.0±1.9	
	Ibadan Northwest	5.9	10.2	7.7±1.6	5.9	10.5	7.7±1.9	
	Ibadan Northeast	5.9	10.3	7.7±1.5	6.1	10.6	7.9±1.7	
	Ibadan Southwest	5.8	10.2	7.6±1.5	6.3	10.4	8.0±1.7	
	Ibadan Southeast	5.8	10.2	7.7±1.5	6.1	10.6	7.8±1.7	

3.2 Microbial burden

3.2.1 Indoor and outdoor airborne Bacterial burden among five local governments.

The mean indoor and outdoor values of total bacterial counts (TBC) CFU/m³ in the selected laundry houses are presented in table 5. Both the indoor and outdoor mean total bacterial counts (TBC) CFU/m³ values were much lower than the $\leq 5 \times 10^2$ CFU/m³ recommended guideline limit of the WHO standards.

4.0 CONCLUSION

In laundry environment in Ibadan metropolis, Nigeria, IAQ appeared compromised. Mechanical ventilation systems in laundry environment need to be improved with respect to operation and maintenance to maintain thermal comfort parameters and reduce levels of chemical pollutants emitted by launderers and product use during services.

Relative humidity for both indoor and outdoor in the morning and afternoon in all the laundries was above the WHO standards of 60% and this may be attributed to the fact the study was carried in Nigeria. Although the sanitary practices were good, the temperature and relative humidity were all higher than the permissible limits and these made the wastes and water to impact indoor air quality. PM₁₀ of all laundries visited were significantly lower than WHO guideline limit of 50µg/m³ and CO₂ was significantly lower than the comfort level of 1000ppm in all laundries visited as stipulated by ASHRAE. This is an indication that there is adequate ventilation in the laundries and shows that the CO₂ outdoor has a positive effect on the indoor CO₂. Indoor and outdoor meteorological factors both in the morning and afternoon exceeded the comfort level of 19°C–26°C and even the WHO guideline limit of 18°C–26°C but as a result of the

weather condition in this part of the world, Nigeria precisely, the high level of temperature observed might not have any significant effect on the people in the laundry.

5.0 RECOMMENDATIONS

The following recommendations are given based on the research findings and these include programs that change the behavioural patterns of the launderers;

1. Educating launderers regarding indoor air quality is a must because launderers need to be provided with the information about sources and effects of indoor air pollution under their control and about the proper operation of ventilating system that can alert their employers and/or take action to reduce their personnel expression.
2. Sources of microbial contaminants and areas where water collection and leakage had occurred (including roofs, HVAC cooling coils and fan coil unit if any) should be cleaned up appropriately. Also, prompt removal of porous organic materials (damp insulation in ventilation system, mouldy ceiling tiles, mildewed carpets) should be encouraged.
3. Routine monitoring of the indoor air quality should be carried out bi-annually in the laundry locations so as to ensure optimal work efficiency by the Environmental Health Officers Registration Council of Nigeria (EHORECON), Oyo State Chapter and National Environmental Standards and Regulations Enforcement Agency (NESREA).
4. Biocides and bacillocids should be used periodically to mop the floor of the laundry facilities so as to control bio-aerosol concentrations.

Comment [HI30]: Reframe this recommendation

Comment [HI31]: Same as HI30

Comment [HI32]: Same as HI30

5. The association of launderers in Oyo State should enforce weekly and monthly environmental sanitation to their members in order to promote hygienic and sanitary environment.

UNDER PEER REVIEW

Table 5: Mean Total Bacterial Counts (TBC) of the laundries among the five local governments(CFU/m³).**Comment [HI33]:** Insert this table after Table 4

	Location	Air microbial quality (CFU/m ³)						WHO Limit
		Morning			Afternoon			
		Min	Max	Mean±SD	Min	Max	Mean±SD	
Indoor	Ibadan North	66.5	91.0	82.9±9.0	72.5	91.0	82.8±6.1	≤5x10 ²
	Ibadan Northwest	129.5	158.0	141.4±10.2	142.5	161.	155.7±6.3	
	Ibadan Northeast	79.5	98.5	88.5±7.0	79.5	95.5	87.9±4.4	
	Ibadan Southwest	147.0	188.5	163.9±12.6	169.5	196.	183.7±8.9	
	Ibadan Southeast	75.5	98.5	88.5±7.0	79.5	95.5	87.9±4.4	
Outdoor	Ibadan North	77.5	144.0	104.0±23.1	68.5	170.	120.4±4.6	
	Ibadan Northwest	132.0	180.5	153.4±16.2	144.5	181.	161.2±12.9	
	Ibadan Northeast	68.5	97.0	84.8±9.0	81.5	149.	91.9±20.0	
	Ibadan Southwest	147.0	188.5	163.9±12.6	169.5	196.	183.7±8.9	

Ibadan Southeast

79.5

98.5


88.5±7.0

79.5

95.5

87.9±4.4

UNDER PEER REVIEW



References

- [1] WHO, (2010): WHO guidelines for indoor air quality: selected pollutants. The WHO European Centre for Environment and Health, Bonn Office, WHO Regional Office for Europe coordinated the development of these WHO guidelines.
- [2] GODSON, R. A., Oyewale, M. M. and Gregory, A. F. (2015). Indoor Air Quality and Risk Factors Associated with Respiratory Conditions in Nigeria. <http://do.doi.org/10.5772/59864>
- [3] FRANCK, U., Herbarth, O., Roder, S., Schlink, U., Borte, M., Diez, U., Krämer, U. & Lehmann, I. (2011): Respiratory effects of indoor particles in young children are size dependent. *Science of the Total Environment*, Vol.409, No.9, pp. 1621-1631, ISSN 0048-9697.
- [4] CANHA, N.; Freitas, M.C.; Almeida, S.M.; Almeida, M.; Ribeiro, M.; Galinha, C. & Wolterbeek, H.T. (2010): Indoor school environment: easy and low cost to assess inorganic pollutants. *Journal of Radioanalytical and Nuclear Chemistry*, Vol.286, No.2, pp. 495-500, ISSN 1588-2780
- [5] WOLKOFF, P., Nojgaard, J., Franch, C., Skov, P. (2006): The modern office desiccates the eyes? *Indoor Air*, 16.4:258-265.
- [6] SALIBA, N. A., Atallah, M. & Al-Kadamany, G. (2009): Levels and indoor-outdoor relationship of PM₁₀, and soluble inorganic ions in Beirut, Lebanon. *Atmospheric Research*, Vol.92, No.1, 131-137, ISSN 0169-8095

- [7] FRAGA, S., Ramos, E., Martins, A., Samúdio, M. J., Silva, G., Guedes, J., Fernandes, E. O. & Barros, H. (2008): Indoor air quality and respiratory symptoms in Porto schools. *Revista Portuguesa de Pneumologia*, Vol.14, No.4, pp. 487-507, ISSN 0873-2159.
- [8] FROMME, H., Twardella, D., Dietrich, S., Heitmann, D., Schierl, R., Liebl, B. & Rüdén, H. (2007) Particulate matter in the indoor air of classrooms-exploratory results from Munich and surrounding area. *Atmospheric Environment*, Vol.41, No.4, pp. 854-866, ISSN 1352-2310
- [9] GUO, H., Lee, S.C., Chan, L.Y. & Li, W.M. (2004): Risk assessment of exposure to volatile organic compounds in different indoor environments. *Environmental Research*, Vol.94, No.1, pp. 57-66, ISSN 0013-9351
- [10] KOSONEN, R. & Tan, F. (2004): The Effect of Perceived Indoor Air Quality on Productivity Loss. *Energy and Buildings*, Vol.36, No.10, pp. 981-986, ISSN 0378-7788
- [11] LEE, S.C., Li, W.M. & Ao, C.H. (2002): Investigation of indoor air quality at residential homes in Hong Kong—case study. *Atmospheric Environment*, Vol.36, No.2, pp. 225–237, ISSN 1352-2310
- [12] LEE, S.C., Guo, H., Li, W.M. & Chan, L.Y. (2002): Inter comparison of air pollutant concentrations in different indoor environment in Hong Kong. *Atmospheric Environment*, Vol.36, No.12, pp. 1929–1940, ISSN 1352-2310.
- [13] LEE, S.C., Ho, K.F., Chan, L.Y., Zielinska, B. & Chow, J.C. (2001): Polycyclic aromatic hydrocarbons (PAHs) and carbonyl compound in urban atmosphere of Hong Kong. *Atmospheric Environment*, Vol.35, No.34, pp. 5949–5960, ISSN 1352-2310

- [14] LI, W.M., Lee, S.C. & Chan, L.Y. (2001): Indoor air quality at nine shopping malls in Hong Kong. *Science of the Total Environment*, Vol.273, No.1-3, pp. 27–40, ISSN 0048-9697.
- [15] ALLEN, A.G. & Miguel, A.H. (1995): Indoor organic and inorganic pollutants: in-situ formation and dry deposition in south eastern Brazil. *Atmospheric Environment*, Vol.29, No.23, pp. 3519-3526, ISSN 1352-2310
- [16] HAAHTELA T., Reijula K. (1998): Diseases caused by indoor air problems and their impact [in Finnish]. *Finnish Medical Journal*; 53:1899–1914.
- [17] FISH SC, Maynard RD, Costa LF, Cardozo A, Schierholt R. (1997): Estimates for improved productivity and health from better indoor environments. *Indoor Air*; 7:158–160
- [18] LAHTINEN S.H., Sosted, H., Andersen, K.E. and Johansen, J.D. (2002): Psychological dimensions of solving an indoor air problem. *Indoor Air*; 12:33–46.
- [19] WHO (1986): Indoor air quality research. EURO reports and studies, no. 103. Copenhagen: WHO Regional Office for Europe, 1986.
- [20] WHO (1983): Indoor air pollutants: exposure and health effects. Report on a WHO meeting. EURO reports and studies, No. 78. Copenhagen: WHO Regional Office for Europe.
- [21] FINNEGAN M.J., Pickering CAC., and Burge PS. (1984): The sick building syndrome: prevalence studies. *BMJ*; 289:1573–1574.

[22] JAAKKOLA J.J.K., Miettinen P. (1995): Type of ventilation system in office buildings and sick building syndrome. *Association of Medical Epidemiology Journal*. 141:755–65.

UNDER PEER REVIEW