

# **A Comprehensive Review on Climate Change Adaptation Strategies and Challenges in Agriculture**

## **Abstract**

This review examines the impact of climate change on Indian agriculture and adaptation strategies. Climate change, driven by human activities, poses challenges like rising temperatures and extreme weather events. Rising temperatures can induce heat stress in crops, leading to reduced yields and poor quality produce. Farmers use adaptation measures like altering planting dates and developing climate-resilient crops. Changing rainfall patterns, erratic or insufficient rainfall can trigger drought conditions, parching farmlands and causing crop failures. On the other hand, excessive rainfall can result in flooding, which damages crops and soil, disrupting agricultural operations especially during the monsoon. Soil conservation, precision agriculture, and urban food production initiatives promote food security and resource recycling. Addressing water scarcity requires improved irrigation and efficient water management. Climate change affects agricultural pests population and threatening global food security. Various adaptation strategies, including traditional practices, resource-conservation technologies, and socio-economic interventions, are being implemented. Climate-smart agriculture technologies like precision agriculture increase yields and resilience. Success depends on regional suitability, economic viability, and collective implementation. Agriculture operates within a complex socio-ecological system with uncertainties in policy, economics, and climate. Site-specific climate-smart agriculture practices are crucial for smallholders' resilience and food security. Publicly provided agricultural extension services can help adopt these technologies, but barriers like financial constraints and cultural factors must be considered. This review emphasizes the need for comprehensive, context-specific approaches to address vulnerabilities in Indian agriculture to climate change and ensure a sustainable future for food production and smallholder livelihoods.

## **Introduction**

Climate change is the biggest environmental issues facing the modern world. Global climate change is caused by the emission of greenhouse gases (GHG), which include rising levels of gases like carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), and nitrous oxide (N<sub>2</sub>O). Changes in rainfall

patterns, a rise in sea level, and the migration of climatic areas as a result of rising temperatures are just a few of the swings that climate change will bring about. Due to shifting climatic trends, it is anticipated that the intensity of droughts, storms, and floods would increase.

Agriculture, a sector heavily dependent on weather conditions, is particularly susceptible to the impacts of climate change (Hasan *et al.*, 2016). Climate change poses a significant threat to the productivity of this sector, leading to economic and physical vulnerabilities. Climate-influenced factors such as shifting rainfall patterns, rising temperatures, altered sowing and harvesting times, fluctuating water availability, and changing site suitability all exert a substantial influence on agricultural output (Shakoor *et al.*, 2011). These changes are expected to have far-reaching consequences, affecting food production, water resources, biodiversity, and livelihoods. In India, where a large portion of the population resides in rural areas and relies heavily on natural resources for their sustenance, sustainable resource management becomes imperative for long-term economic well-being. The timing of the monsoon plays a crucial role in Indian agriculture, influencing the agricultural market and essential commodities. Any deviations in rainfall patterns impact agriculture, thereby affecting the nation's economy and food security (Ahmad *et al.*, 2011).

The potential for worldwide climate change looms as a significant consequence of the anticipated doubling of atmospheric CO<sub>2</sub> levels and increased trace gas concentrations (Tangle, 1988). The direct influence of weather and climate on agricultural production cannot be understated. Potential shifts in temperature, precipitation, and CO<sub>2</sub> concentration are poised to significantly impact crop development. Given that climatic factors such as rainfall and temperature serve as critical inputs for the crop sector, any alterations or fluctuations in these variables are bound to exert substantial effects on crop yields (Barnwal and Kotani, 2013).

Climate change poses a grave threat to agriculture and the security of India's food supply, with particular emphasis on the critical role of water resources. In a country where 55% of cultivated lands lack irrigation facilities, water stands as the most vital agricultural resource (Kumar and Gautam, 2014). The demand for water has surged over time, propelled by urbanization, population growth, rapid industrialization, and other developmental pursuits. Moreover, changes in agricultural and land-use practices, excessive groundwater extraction, and alterations in irrigation and drainage have disrupted the hydrological cycle across various

climate zones and river basins in India. As a result, water availability emerges as the linchpin of agricultural productivity. Throughout much of India, the quality and availability of water remain significant impediments to successful agriculture, underscoring the pressing need for comprehensive water resource management in the face of climate change.

### **Effects of Climate Change on Agricultural Productivity**

The effect of Climate Change on Agriculture can be seen everywhere in the World (Yadav *et al.*, 2023). In India, the scientific community has employed a variety of techniques to assess the potential effects of climatic variability and change on agriculture. Historically, methods such as analogue analysis and statistical tools have been used to analyze historical data and understand the impacts of climatic variability. More recently, controlled environmental facilities like open-top chambers, free air carbon-dioxide enrichment facilities, phytotron setups, and greenhouses are being increasingly utilized to investigate how factors like temperature, humidity, and CO<sub>2</sub> levels affect crop growth and yield (Aggarwal, 2008). These controlled environments offer precise conditions for research, providing valuable insights into the potential impacts of changing climatic conditions on crop production and aiding in the development of adaptive strategies for agriculture in the face of climate change.

A defining issue of our time is climate change, primarily driven by human activities such as the combustion of fossil fuels and deforestation. This global challenge has far-reaching implications, and one of the sectors significantly impacted is agriculture, which plays a crucial role in ensuring food security, livelihoods, and economies worldwide. Climate change exerts a profound influence on agricultural productivity and sustainability, manifesting through shifts in temperature, altered precipitation patterns, extreme weather events, and rising sea levels. Temperature, in particular, emerges as a critical environmental factor influencing crop growth, development, and yields, affecting the pace at which crops progress through their phenophases. On one hand, crops require a minimum temperature threshold to complete specific life cycle stages, while exceptionally high or low temperatures, especially during crucial phases like anthesis, can detrimentally impact crop growth and production (Luo, 2011).

As global temperatures continue to rise, crop growth and productivity face direct consequences. The optimal temperature ranges for many crops are shifting, leading to adjustments in planting and harvesting times. Elevated temperatures can accelerate plant development, disrupt pollination, and enhance evaporation rates, placing plants under

increased water stress. Moreover, any potential benefits arising from higher atmospheric CO<sub>2</sub> concentrations may be counteracted by rising air temperatures, further complicating the picture. Crop growing seasons may also shorten in response to these changes. Consequently, the interplay between temperature and CO<sub>2</sub> concentration presents a complex challenge in predicting their combined effects on plant growth and agricultural outcomes (Streck, 2005).

Drought, emblematic of the mounting trend in extreme weather events, is progressively on the rise (Mazdiyasi and AghaKouchak, 2015), resulting in diminishing agricultural yields (Lesk *et al.*, 2016). To combat the adverse impacts of drought, comprehensive monitoring and prediction systems have been established, providing essential data for early warning and mitigation strategies aimed at bolstering crop resilience and productivity (Hao and AghaKouchak, 2014). Concurrently, adaptation measures have been devised to mitigate these impacts and shape the extent to which climate change affects crop production, including adjustments in planting schedules, transitioning to more suitable crop varieties, developing novel strains, and altering cultivation practices (Leng and Huang, 2017). These adaptive strategies are pivotal in fortifying agricultural systems against the increasing challenges posed by a changing climate.

Climate change-related changes in precipitation patterns are having an increasing impact on crop growth, agricultural productivity, and food security. Rainfall and snowfall, both of which constitute precipitation, supply the vital water source required for plant growth and maintenance. By delivering water for the two main crop-growing seasons, Kharif (summer) and Rabi (winter), rainfall that occurs over India during the summer monsoon season (the major rainy season typically starts in June and ends in September) has a huge impact on the agricultural production of the nation. The entire foodgrain output in India and the nation's economy, which is heavily dependent on agriculture, are both impacted by variations in monsoon rainfall (Krishna Kumar *et al.* 2004). Precipitation patterns are changing as a result of climate change, including changes in rainfall frequency, intensity, and distribution. There are greater hazards of flooding and soil erosion in some areas due to more frequent and strong rainfall events. Others have protracted droughts that diminish agricultural productivity and cause a shortage of water.

### **Crop Adaptation Strategies**

Farmers are adapting to these challenges, often through autonomous adaptation strategies, such as altering cropping patterns to include short-duration crops like pulses, vegetables,

flowers, and fruits that can withstand warming and drying climates. Planned adaptation measures, like providing farmers with new drought-resistant seed varieties, are also crucial. Farmers are modifying crop types, planting times, and irrigation practices to cope. Information dissemination about innovative management techniques, like revised crop calendars, is vital, as are enhancements to water storage infrastructure like desilting farm ponds (Dhanya and Ramachandran, 2016).

Farmers are increasingly cultivating short-duration, heat-tolerant crops such as pulses, maize, millets, and floriculture crops in response to climate variability. Phenological shifts in crops, including early blooming and fruiting, have been observed due to high temperatures and moisture stress, impacting both crop production and farmers' livelihoods (Brown, 2006).

Plant breeders face the challenge of developing crops that can thrive in changing conditions. They are using advanced breeding techniques like Rapid Generation Advance (RGA), Doubled Haploid (DH), Shuttle Breeding, Marker Assisted Back Crossing (MABC), Marker Assisted Recurrent Selection (MARS), Genomic Selection (GS), and Speed Breeding to accelerate genetic gains and create climate-resilient crop varieties (Beghey *et al.*, 2016). Shuttle breeding, for instance, enables an additional generation by growing crops in different locations each year. Genomic Assisted Breeding Approaches are promising for developing climate-resilient crops, reducing generation times, and deciphering how different crop species respond to stress. However, plant breeders must now consider a broader range of environmental factors, including increased temperature, water supply variability, UV radiation, and salinity, when developing new crop varieties (Brettell, 2008).

### **Soil Management in a Changing Climate**

Soil, a critical component of land, is an indispensable factor for crop production, and its health is intricately linked to the challenge of climate change. The importance of soil in sustaining global food production cannot be overstated, and its role becomes even more crucial in the context of climate change mitigation and adaptation.

Climate change impacts soil in several ways. Rising temperatures, altered precipitation patterns, and increased occurrences of extreme weather events can lead to soil degradation, erosion, and reduced fertility. These changes can disrupt agricultural practices and threaten food security, as highlighted by Blanco-Canqui (2008) and Lashgarara *et al.* (2008).

Therefore, addressing climate change is not only crucial for reducing greenhouse gas emissions but also for preserving soil quality.

Sustainable intensification, as proposed by Godfray et al. (2010), is a key strategy to address both food security and climate change. It involves optimizing agricultural practices to enhance production efficiency while minimizing environmental impacts. Precision agriculture (PA) is a vital component of sustainable intensification. PA relies on data-driven approaches to maximize crop yields while minimizing resource use, helping to reduce greenhouse gas emissions associated with agriculture.

Furthermore, climate change and agriculture are intricately linked through the issue of non-point source pollution, as mentioned by Cordell et al. (2011). Climate-related events like heavy rainfall can exacerbate the runoff of pollutants from agricultural fields into water bodies, contributing to water pollution. Sustainable farming practices, including those employed in precision agriculture, can help mitigate this pollution risk.

In the context of climate change, it's essential to note that the disruption of the phosphorus (P) cycle, as mentioned by Abelson (1999), is not only a concern for nutrient management but also has implications for water quality and ecosystem health. PA can play a role in optimizing nutrient use, thereby reducing the environmental impact and contributing to sustainable land management.

Finally, urbanization, as discussed by Drescher (2002), is a trend that has its own climate implications. Urban areas, with their growing populations, contribute significantly to greenhouse gas emissions. However, urban agriculture practices, including rooftop gardens and the recycling of biosolids and wastewater, as highlighted by Whittinghill and Rowe (2012), can provide sustainable alternatives for food production. These practices can help reduce the carbon footprint of urban areas and increase their resilience to climate change by promoting local food production.

### **Water Management and Irrigation**

Water is an essential element for all living organisms, including humans, animals, and plants, making it fundamental to life itself. Its importance extends to various critical aspects, such as food security, livestock husbandry, industrial processes, biodiversity preservation, and environmental conservation. Freshwater is the only source of water that can be used across a spectrum of essential needs, including drinking, industrial applications, agriculture, and more.

However, the accessibility and availability of clean drinking water remain challenges for millions of people in India and around the world, and this issue is exacerbated by various factors.

In many regions, particularly in areas with limited water resources, water demand has reached a critical juncture. Irrigation should aim to restore soil water in the root zone to a level at which crop can fully meet its evapotranspiration requirement (Kumar *et al.*, 2023). Despite the Earth's abundance of water, factors like overexploitation of water resources, inadequate water supply infrastructure, and the impact of climate change contribute to water scarcity (Mancosu, 2015). The 21st century confronts us with a profound global problem: the lack of access to water, driven by causes including overpopulation, agricultural demands, water pollution, and ineffective governance (Chakkaravarthy and Balakrishnan, 2019).

Climate change compounds the challenges to freshwater resources, affecting both their quantity and quality. India, in particular, faces heightened demand for water due to urbanization, industrialization, the agricultural sector's need for increased water, and a rapidly growing population (Amarasinghe *et al.*, 2007). In response, efforts have been made to promote water efficiency, guided by the principle that "more crop per drop" through improved management practices (Allan, 1999).

Enhancing allocative and irrigation water efficiency constitutes better water management. The approach varies based on irrigation technology, environmental conditions, land characteristics, and the timing of water application, often tied to appropriate water pricing for agricultural use (Omezzine and Zaibet, 1998). Irrigation scheduling, a critical component, influences water use efficiency (WUE) by determining the timing and quantity of irrigation, with plant water status assessments, soil moisture monitoring, and crop evapotranspiration estimations serving as methods for achieving this. However, soil-based assessments have limitations, as they provide point-based data subject to spatial and temporal variability (Gillies, 2005).

Addressing water scarcity is a paramount global challenge (Jury and Vaux, 2005). To alleviate water shortages, accurate monitoring of water use, including improved estimation of return flows and evapotranspiration, is essential.

### **Pest and Disease Dynamics in a Warming Climate**

Global climate change exerts a substantial impact on agriculture, directly and indirectly affecting agricultural insects and pests. Indirect effects encompass changes in the interactions between pests, their environment, and other insect species, including natural enemies, competitors, vectors, and mutualists (Prakash *et al.*, 2014). Direct impacts on pests encompass alterations in reproduction, development, survival, and dispersal. As noted by Yamamura and Kiritani (1998), even a modest 2°C temperature increase can lead to insects experiencing one to five additional life cycles in a single season. Elevated carbon dioxide (CO<sub>2</sub>) concentrations can affect insect feeding rates and host chemical defenses, as suggested by Arora and Dhawan (2013), thereby influencing population densities, growth rates, fecundity, and consumption rates of insect pests (Fuhrer, 2003).

Climate change-induced factors like flooding, extreme weather events, and the creation of new ecological niches can lead to shifts in the population dynamics of soil-dwelling insects. These changes provide opportunities for insect pests to establish, multiply, and migrate across different geographic regions, potentially resulting in new and severe pest challenges for farmers in the years ahead. Cross-border movement of crop pests poses a threat to food security, with global implications (FAO, 2020).

Temperature plays a pivotal role in influencing pest population dynamics, impacting metabolism, metamorphosis, mobility, and host availability (Shrestha, 2019). Soil-dwelling species are less affected by temperature fluctuations than aboveground insects due to the thermal buffering capacity of soil (Bale *et al.*, 2002). For instance, aphids may emit fewer alarm pheromones at higher temperatures, potentially increasing their susceptibility to predation by parasitoids and insect predators (Awmack *et al.*, 1997).

Precipitation patterns, including changes in quantity, intensity, and frequency, are significant indicators of climate change. Altered rainfall patterns, such as droughts and floods, can directly impact insects that overwinter in the soil, affecting their survival and diapause patterns. Flooding and heavy rains can also wash away insect eggs and larvae (Pathania *et al.*, 2020). Staley *et al.* (2007) investigated the effects of drought and increased summer rainfall on wireworms, soil-dwelling pests that pose a substantial threat to crops like potatoes, corn, and sugar beets. Their findings suggest that increasing summer rainfall events can lead to a rapid rise in wireworm populations in the upper soil layers (Gregory *et al.*, 2009). These observations underscore the complex and multifaceted relationship between climate change and agricultural pest dynamics.

## Mitigation and Adaptation Strategies

Adaptation is a critical policy option in response to the effects of climate change, particularly in sectors highly sensitive to climatic conditions like agriculture (Fankhauser, 1996). Agriculture's intrinsic sensitivity to climate makes it one of the most vulnerable industries to the risks and impacts of climate change (Reilly, 1995).

A wide range of agricultural adaptation strategies have been proposed to mitigate the anticipated negative effects of climate change. In arid and semi-arid regions, farmers have developed traditional management strategies over generations to enhance soil water retention, reduce sensitivity to drought, and combat soil erosion (Hulme, 1999). One such strategy involves raised beds with tied ridges, which helps retain rainwater, reduce runoff, and enhance water infiltration (Altieri and Nicholls, 2017).

Adaptive agricultural practices encompass a combination of conventional and agroecological management systems, including practices such as biodiversification, improved soil management, and water harvesting (Altieri and Nicholls, 2017). These practices lead to resilient soils and cropping systems, ensuring food security in the face of climate change while also promoting soil health, quality, and carbon sequestration. Key adaptation strategies can be categorized as resource-conservation technologies, cropping-system technologies, and socioeconomic or policy interventions (Venkateswarlu and Shanker, 2009).

In sub-Saharan Africa, adapting to climate change involves implementing sequential cropping systems and adjusting sowing dates according to climate conditions, minimizing crop yield losses (Verhotet *et al.*, 2007). Agroforestry practices in Kenya offer opportunities to reduce greenhouse gas accumulation in the atmosphere, benefiting both adaptation and mitigation efforts (Malhi *et al.*, 2021). Additional straightforward adaptation measures include changing planting dates and crop cultivars (Aggarwal, 2008).

Conservation agriculture, characterized by minimal soil disturbance, crop rotation, and soil cover, holds promise for reversing soil degradation caused by conventional tillage and sequestering carbon. However, it should be noted that the carbon sequestration potential of no-till cultivation is limited (Powlson *et al.*, 2014).

Aerobic rice cultivation, aided by micro-irrigation technologies, offers a sustainable approach to rice production, conserving irrigation water and reducing methane emissions (Parthasarathi *et al.*, 2012). Precision farming, which involves more efficient fertilizer management, has the

potential to improve nitrogen use efficiency and reduce greenhouse gas emissions (Ladha *et al.*, 2007; Bhatia *et al.*, 2012).

Implementing climate-smart agriculture techniques can boost crop yields and farmer incomes, with laser land leveling cited as an example that reduced expenses and limited losses from climatic fluctuations (Pal *et al.*, 2020). However, the success of these adaptation and mitigation techniques depends on factors like their applicability to specific regions, public acceptance, economic viability, and technical complexity. These strategies are most effective when multiple interventions are employed in synergy to support one another.

### **Challenges**

According to Darnhofer (2010), agriculture is a complex, dynamic socio-ecological system that is subject to policy, economic, and climatic uncertainty. Site-specific climate-smart agriculture (CSA) practices will play a critical role in boosting the income of the most vulnerable populations, minimizing the impact of climate change on food security, and improving smallholders' resilience (FAO 2018, 2017).

The adoption of technologies like CSA is likely to be supported by publicly offered agricultural extension services (Aker 2011). These services distribute vital information to farmers (Lamontagne-Godwin *et al.*, 2018). However, several obstacles hinder the adoption of climate-smart practices, including financial limitations, labor shortages, land and water scarcity, insufficient transportation resources, and low farmer organization membership (Deressa *et al.*, 2008).

Barriers to adoption also encompass institutional, socioeconomic, and cultural factors. These include the need for financial capital (Burbi *et al.* 2016), lack of knowledge, unsatisfactory land tenure, market failures, and inadequate infrastructure (Altieri and Nicholls 2012). Additional considerations involve the suitability of innovations for end-users, labor requirements, access to external inputs, and the use of crop residues for animal feeding (Giller *et al.*, 2009).

Another significant hurdle is the adoption process itself, particularly with regard to its social dimensions, such as age, gender, and diversity. For example, women and men farmers may not have equal access to, use of, or benefits from certain practices (Archer 2003). Similar disparities may occur among farmers with varying income levels, education, family size, land tenure, religious affiliations, places of origin, or connections to institutions and influential

individuals. When assessing the suitability of CSA practices and the obstacles to their implementation, these aspects must be considered.

Furthermore, the level of institutional support within a region can influence the speed at which CSA practices are adopted, especially those requiring higher initial startup costs or technical expertise. Investments in infrastructure, extension services, and healthcare within agricultural communities can significantly impact farmers' willingness to take risks and, consequently, their adoption of new practices (Below *et al.*, 2010).

## **Conclusion**

The Indian scientific community employs various methods, including historical data analysis and controlled environmental facilities, to assess climate variability's impact on agriculture. Human-driven climate change significantly affects agriculture through temperature shifts, extreme weather events, and changing precipitation patterns. Rising temperatures can accelerate plant growth but may also cause water stress, while extreme weather events like drought reduce crop production. Adaptation measures, such as altering planting dates and developing new crop varieties, help mitigate these impacts. Changing precipitation patterns during the Indian monsoon season significantly influence foodgrain yield and the country's economy. Climate change leads to shifts in rainfall patterns, affecting food productivity and increasing flooding and drought risks. It directly and indirectly impacts crop production with increased extreme events. Farmers adapt by shifting to short-duration crops, while Genomic Assisted Breeding aims to develop climate-resilient varieties. Soil is crucial for global food production, especially in developing countries. Sustainable intensification, using precision agriculture and nutrient management, enhances productivity while reducing environmental impact. Urban food production, including rooftop gardens, addresses food insecurity and promotes resource recycling. Water scarcity is a growing global challenge due to mismanagement, pollution, climate change, and population growth. Improved irrigation techniques, better management, and efficient water use are essential for addressing this issue. Accurate measurement of water use, including return flows and evapotranspiration, is crucial to mitigate water scarcity challenges and their impacts. Climate change significantly impacts agricultural pests by altering their life cycles, behavior, and interactions with the environment. Rising temperatures accelerate pest reproduction, while increased CO<sub>2</sub> affects feeding rates and host plant dynamics. Different plant responses to elevated CO<sub>2</sub> influence herbivory patterns. Climate-induced ecological shifts create new pest challenges, posing

threats to global food security. Adaptation is crucial, with strategies including traditional and agroecological practices, resource-conservation technologies, cropping-system technologies, and socio-economic interventions. These approaches enhance soil health, increase carbon sequestration, improve crop yields, and reduce greenhouse gas emissions. Water-saving techniques like drip irrigation and precision agriculture are valuable. The adoption of climate-smart agriculture technologies, such as laser land leveling, can increase resilience, depending on regional suitability, economic viability, and collective implementation. Agriculture operates within a complex socio-ecological system filled with uncertainties. Site-specific climate-smart agriculture (CSA) practices are essential for mitigating climate change's impact, but barriers include financial constraints, labor shortages, and infrastructural deficits. Institutional, socio-economic, and cultural factors play crucial roles, and adoption challenges depend on factors like age, gender, income, and institutional support. Publicly provided agricultural extension services can facilitate CSA technology adoption.

## REFERENCES

Abelson, P. H. (1999). A potential phosphate crisis. *Science*, 283(5410), 2015-2015.

- Aggarwal, P. K. (2008). Global climate change and Indian agriculture: impacts, adaptation and mitigation. *Indian Journal of Agricultural Sciences*, 78(11), 911.
- Ahmad, J., Alam, D., & Haseen, M. S. (2011). Impact of climate change on agriculture and food security in India. *International Journal of Agriculture, Environment and Biotechnology*, 4(2), 129-137.
- Aker, Jenny C. 2011. "Dial "A" for agriculture: a review of information and communication technologies for agricultural extension in developing countries." *Agricultural Economics* 42 (6):631-647.
- Altieri, M. A., & Nicholls, C. I. (2017). The adaptation and mitigation potential of traditional agriculture in a changing climate. *Climatic change*, 140, 33-45.
- Amarasinghe, U.A., Shah, T., Turrall, H. and Anand, B.K. 2007. India's water future to 2025-2050: Business as usual scenario and deviations. Research Report 123, IWMI. pp. 52.
- Archer ERM. 2003. Identifying Underserved End-User Groups in the Provision of Climate Information. *Bulletin of the American Meteorological Society* 84(11): 1525–1532.
- Arora, R. and Dhawan, A.K. 2013. "Climate change and insect pest management". In *Integrated pest management*. Ed(s): A.K. Dhawan, Balwinder Singh, Manmeet B. Bhullar, R. Arora, Scientific Publisher, Jodhpur, India: 44-60.
- Awmack, C., Woodcock, C., & Harrington, R. (1997). Climate change may increase vulnerability of aphids to natural enemies. *Ecological Entomology*, 22(3), 366-368.
- Bale, J. S., Masters, G. J., Hodkinson, I. D., Awmack, C., Bezemer, T. M., Brown, V. K., ... & Whittaker, J. B. (2002). Herbivory in global climate change research: direct effects of rising temperature on insect herbivores. *Global change biology*, 8(1), 1-16.
- Barnwal, P., & Kotani, K. (2010). Impact of variation in climatic factors on crop yield: A case of rice crop in Andhra Pradesh, India (Working Paper No. 17). Economics and Management Series.
- Begheyn, R.F., Lubberstedt, T. and Studer, B. 2016. "Haploid and doubled haploid techniques in perennial ryegrass (*Lolium perenne*L.) to advance research and breeding." *Agronomy* 6(4):60.
- Below T, Artner A, Siebert R, Sieber S. 2010. Micro-level Practices to Adapt to Climate Change for African Smallscale Farmers. Washington, D.C.: International Food Policy Research Institute (IFPRI).
- Bhatia, A., Pathak, H., Jain, N., Singh, P. K., & Tomer, R. (2012). Greenhouse gas mitigation in rice–wheat system with leaf color chart-based urea application. *Environmental Monitoring and Assessment*, 184, 3095-3107.
- Birkenholtz, T. (2017). Assessing India's drip-irrigation boom: efficiency, climate change and groundwater policy. *Water International*, 42(6), 663-677.

- Blanco-Canqui, H., & Lal, R. (2008). Corn stover removal impacts on micro-scale soil physical properties. *Geoderma*, 145(3-4), 335-346.
- Bongiovanni, R., & Lowenberg-DeBoer, J. (2004). Precision agriculture and sustainability. *Precision agriculture*, 5, 359-387.
- Brettell, R. 2008. "Breeding crops of a changing climate." ICARDA Caravan; Special Issue on Climate Change 25:9–12.
- Brown ME. 2006. Assessing natural resource management challenges in Senegal using data from participatory rural appraisals and remote sensing. *World Dev.* 34(4):751– 767.
- Burbi, Sara, R. N. Baines, and J. S. Conway. 2016. "Achieving successful farmer engagement on greenhouse gas emission mitigation." *International Journal of Agricultural Sustainability*:1-18.
- Chakkaravarthy, D. N., & Balakrishnan, T. (2019). Water scarcity-challenging the future. *International Journal of Agriculture, Environment and Biotechnology*, 12(3), 187-193.
- Childers, D. L., Corman, J., Edwards, M., & Elser, J. J. (2011). Sustainability challenges of phosphorus and food: solutions from closing the human phosphorus cycle. *Bioscience*, 61(2), 117-124.
- Cordell, D., Rosemarin, A., Schröder, J. J., & Smit, A. L. (2011). Towards global phosphorus security: A systems framework for phosphorus recovery and reuse options. *Chemosphere*, 84(6), 747-758.
- Darnhofer, Ika. 2010. "Strategies of family farms to strengthen their resilience." *Environmental Policy and Governance* 20 (4):212-222.
- Darnhofer, Ika. 2014. "Resilience and why it matters for farm management." *European Review of Agricultural Economics* 41 (3):461-484.
- De Costa, W. A. J. M., & Sangakkara, U. R. (2006). Agronomic regeneration of soil fertility in tropical Asian smallholder uplands for sustainable food production. *The Journal of Agricultural Science*, 144(2), 111-133.
- Deressa T, Hassan RM, Alemu T, Yesuf M, Ringler C. 2008. Analyzing the Determinants of Farmers' Choice of Adaptation Methods and Perceptions of Climate Change in the Nile Basin of Ethiopia. Washington, D.C.: International Food Policy Research Institute (IFPRI).
- Dhaka, B. L., Chayal, K., Poonia, M. K., & Kendra, K. V. (2010). Analysis of farmers' perception and adaptation strategies to climate change. *Libyan Agriculture Research Center Journal International*, 1(6), 388-390.

- Dhanya, P., & Ramachandran, A. (2016). Farmers' perceptions of climate change and the proposed agriculture adaptation strategies in a semi arid region of south India. *Journal of Integrative Environmental Sciences*, 13(1), 1-18.
- Drescher, A. W. (2002, September). Food for the cities: urban agriculture in developing countries. In *International Conference on Urban Horticulture 643* (pp. 227-231).
- Fankhauser, S. (1996). The potential costs of climate change adaptation. In *Adapting to climate change: An international perspective* (pp. 80-96). New York, NY: Springer New York.
- FAO. 2017. Climate-Smart Agriculture Sourcebook Summary. Rome. FAO. 2018. Climate-Smart Case studies 2018: Successful approaches from different regions. Rome: Food and Agriculture Organisation of the United Nations.
- FAO. Climate Related Transboundary Pests and Diseases. Available online: <http://www.fao.org/3/a-ai785e.pdf> (accessed on 19 December 2020).
- Fuhrer, J. (2003). Agroecosystem responses to combinations of elevated CO<sub>2</sub>, ozone, and global climate change. *Agriculture, Ecosystems & Environment*, 97(1-3), 1-20.
- Gebbers, R., & Adamchuk, V. I. (2010). Precision agriculture and food security. *Science*, 327(5967), 828-831.
- Giller, Ken E., Ernst Witter, Marc Corbeels, and Pablo Tittonell. 2009. "Conservation agriculture and smallholder farming in Africa: The heretics' view." *Field Crops Research* 114 (1):23-34.
- Gillies, M. H., & Smith, R. J. (2005). Infiltration parameters from surface irrigation advance and runoff data. *Irrigation Science*, 24, 25-35.
- Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F., ... & Toulmin, C. (2010). Food security: the challenge of feeding 9 billion people. *science*, 327(5967), 812-818.
- Graham, R. D., Welch, R. M., & Bouis, H. E. (2001). Addressing micronutrient malnutrition through enhancing the nutritional quality of staple foods: principles, perspectives and knowledge gaps.
- Gregory, P. J., Johnson, S. N., Newton, A. C., & Ingram, J. S. (2009). Integrating pests and pathogens into the climate change/food security debate. *Journal of experimental botany*, 60(10), 2827-2838.
- Hao, Z., & AghaKouchak, A. (2014). A nonparametric multivariate multi-index drought monitoring framework. *Journal of Hydrometeorology*, 15(1), 89-101.
- Hasan, M. M., Sarker, M. A. R., & Gow, J. (2016). Assessment of climate change impacts on Aman and Boro rice yields in Bangladesh. *Climate Change Economics*, 7(03), 1650008.

- He, J., McHugh, A. D., Li, H. W., Wang, Q. J., Li, W. Y., Rasaily, R. G., & Li, H. (2012). Permanent raised beds improved soil structure and yield of spring wheat in arid north-western China. *Soil use and management*, 28(4), 536-543.
- Hulme, M., Barrow, E. M., Arnell, N. W., Harrison, P. A., Johns, T. C., & Downing, T. E. (1999). Relative impacts of human-induced climate change and natural climate variability. *Nature*, 397(6721), 688-691.
- Ibrahim M, Florkowski WJ, Kolavalli S. 2012. The Determinants of Farmer Adoption of Improved Peanut Varieties and Their Impact on Farm Income: Evidence from Northern Ghana. In: Selected Paper prepared for presentation at the Agricultural and Applied Economics Association Annual Meeting, Seattle, WA, August 12-14, 2012. Seattle, WA, p 1–16.
- IPCC. 2007. Climate Change 2007: The physical science basis. Contribution of Work Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate, Cambridge University Press, United Kingdom.
- Ismail, A.M., Singh U.S., Singh S., Dar M.H., and Mackill, D.J. 2013. “The contribution of submergence-tolerant (Sub1) rice varieties to food security in flood-prone rainfed lowland areas in Asia”. *Field Crops Research* 152(1):83- 93.
- Johnson, S. N., Anderson, E. A., Dawson, G., & Griffiths, D. W. (2008). Varietal susceptibility of potatoes to wireworm herbivory. *Agricultural and Forest Entomology*, 10(2), 167-174.
- Jury, W. A., & Vaux Jr, H. (2005). The role of science in solving the world's emerging water problems. *Proceedings of the National Academy of Sciences*, 102(44), 15715-15720.
- Krishna Kumar K, Rupa Kumar K, Ashrit R, Deshpande N R and Hansen J W 2004 Climate impacts on Indian agriculture; *Int. J. Climatol.* 24 1375–1393.
- Kumar, R., & Gautam, H. R. (2014). Climate change and its impact on agricultural productivity in India. *Journal of Climatology & Weather Forecasting*, 2(1), 1-3.
- Kumar, S., Kumar, S., Kumar, R., Pathak, D., Kumar, D., Kumar, A., ... & Tiwari, H. (2023). Assessment of Physico-chemical Properties of Soil as Influenced by Different Moisture Regimes and Nitrogen Sources in Wheat Crop. *Int. J. Plant Soil Sci*, 35(19), 765-772.
- Ladha, J. K., Bains, J. S., Gupta, R. K., & Balasubramanian, V. (2007). On-farm evaluation of leaf color chart for need-based nitrogen management in irrigated transplanted rice in northwestern India. *Nutrient cycling in agroecosystems*, 78, 167-176.
- Lal, R., Delgado, J. A., Groffman, P. M., Millar, N., Dell, C., & Rotz, A. (2011). Management to mitigate and adapt to climate change. *Journal of Soil and Water Conservation*, 66(4), 276-285.
- Lamontagne-Godwin, J., F. E. Williams, N. Aslam, S. Cardey, P. Dorward, and M. Almas. 2018. "Gender differences in use and preferences of agricultural information sources in Pakistan." *The Journal of Agricultural Education and Extension* 24 (5):419-434.

- Lashgarara, F., Mirdamadi, S. M., Hosseini, S. J. F., & Chizari, M. (2008). The Role of Food- Security Solutions in the Protection of Natural Resources and Environment of Developing Countries. *Annals of the New York Academy of Sciences*, 1140(1), 68-72.
- Leng, G., & Huang, M. (2017). Crop yield response to climate change varies with crop spatial distribution pattern. *Scientific Reports*, 7(1), 1463.
- Lesk, C., Rowhani, P., & Ramankutty, N. (2016). Influence of extreme weather disasters on global crop production. *Nature*, 529(7584), 84-87.
- Luo, Q. (2011). Temperature thresholds and crop production: a review. *Climatic change*, 109(3-4), 583-598.
- Malhi, G. S., Kaur, M., & Kaushik, P. (2021). Impact of climate change on agriculture and its mitigation strategies: A review. *Sustainability*, 13(3), 1318.
- Mancosu, N., Snyder, R. L., Kyriakakis, G., & Spano, D. (2015). Water scarcity and future challenges for food production. *Water*, 7(3), 975-992.
- Mazdiyasn, O., & AghaKouchak, A. (2015). Substantial increase in concurrent droughts and heatwaves in the United States. *Proceedings of the National Academy of Sciences*, 112(37), 11484-11489.
- Mizina, S. V., Smith, J. B., Gossen, E., Spiecker, K. F., & Witkowski, S. L. (1999). An evaluation of adaptation options for climate change impacts on agriculture in Kazakhstan. *Mitigation and Adaptation Strategies for Global Change*, 4, 25-41.
- Ortiz, R., Trethowan, R., Ferrara, G.O., Iwanaga, M., Dodds, J.H., Crouch, J.H., Crossa, J. and Braun, H.J. 2007. "High yield potential, shuttle breeding, genetic diversity, and a new international wheat improvement strategy". *Euphytica* 157(3):365-384.
- Pal, B. D., Kapoor, S., Saroj, S., Jat, M. L., Kumar, Y., & Anantha, K. H. (2020). *Impact of laser land levelling on food production and farmers' income: evidence from drought prone semi-arid tropics in India* (Vol. 1960). Intl Food Policy Res Inst.
- Pandey, P. R., Zaidi, S. F. A., Kumar, S., Pathak, D., Shukla, G., & Pal, R. (2023). Depth Wise Studies of Physico-Chemical Properties of Soil under Different Land Use System at Eastern UP, India. *International Journal of Plant & Soil Science*, 35(18), 762-772.
- Parthasarathi, T., Vanitha, K., Lakshamanakumar, P., & Kalaiyarasi, D. (2012). Aerobic rice-mitigating water stress for the future climate change. *Int J Agron Plant Prod*, 3(7), 241-254.
- Pathania, M., Verma, A., Singh, M., Arora, P. K., & Kaur, N. (2020). Influence of abiotic factors on the infestation dynamics of whitefly, Bemisia tabaci (Gennadius 1889) in cotton and its management strategies in North-Western India. *International journal of tropical insect science*, 40, 969-981.

- Powlson, D. S., Stirling, C. M., Jat, M. L., Gerard, B. G., Palm, C. A., Sanchez, P. A., & Cassman, K. G. (2014). Limited potential of no-till agriculture for climate change mitigation. *Nature climate change*, 4(8), 678-683.
- Prakash, A., Rao, J., Mukherjee, A. K., Berliner, J., Pokhare, S. S., Adak, T., ... & Shashank, P. R. (2014). *Climate change: impact on crop pests*. Odisha: Applied Zoologists Research Association (AZRA), Central Rice Research Institute.
- Shakoor, U., Saboor, A., Ali, I., & Mohsin, A. Q. (2011). Impact of climate change on agriculture: empirical evidence from arid region. *Pak. J. Agri. Sci*, 48(4), 327-333.
- Shrestha, S. (2019). Effects of climate change in agricultural insect pest. *Acta Scientific Agriculture*, 3(12), 74-80.
- Staley, J. T., Hodgson, C. J., Mortimer, S. R., Morecroft, M. D., Masters, G. J., Brown, V. K., & Taylor, M. E. (2007). Effects of summer rainfall manipulations on the abundance and vertical distribution of herbivorous soil macro-invertebrates. *European Journal of Soil Biology*, 43(3), 189-198.
- Streck, N. A. (2005). Climate change and agroecosystems: the effect of elevated atmospheric CO<sub>2</sub> and temperature on crop growth, development, and yield. *Ciência Rural*, 35, 730-740.
- Tangley, L. 1988. Preparing for climate change. *BioScience* 38: 14-18.
- Venkateswarlu, B., & Shanker, A. K. (2009). Climate change and agriculture: adaptation and mitigation strategies. *Indian Journal of Agronomy*, 54(2), 226-230.
- Verchot, L. V., Van Noordwijk, M., Kandji, S., Tomich, T., Ong, C., Albrecht, A., ... & Palm, C. (2007). Climate change: linking adaptation and mitigation through agroforestry. *Mitigation and adaptation strategies for global change*, 12, 901-918.
- Whittinghill, L. J., & Rowe, D. B. (2012). The role of green roof technology in urban agriculture. *Renewable Agriculture and Food Systems*, 27(4), 314-322.
- Yadav V., Kumar S., Pal R. & Pathak D., (2023). Impact of Climate Change on Soil Properties and Agricultural production. *Agrospheres:e- Newsletter*, 4 (8), 1- 4.
- Yamamura, K. and Kiritani, K. 1998. "A simple method to estimate the potential increase in the number of generations under global warming in temperate zones. *Applied Entomology and Zoology* 33(2):289-298.