

## **Influence of seed cake as organic fertilizer on growth, dry matter production and root development of two cultivars of C4 plant, Maize (*Zea mays*) under lab conditions**

Formatted: Font: Italic

### **Abstract:**

A pot culture experiment was conducted with the C4 plant (*Zea mays*) in the lab during the period of vegetative growth to evaluate the influence of organic fertilizer in the form of mustard oil seed cake on growth, biomass yield and root development of two maize cultivars, hybrid yellow and hybrid purple. The seed cake fertilizer treatments imposed in the experiment were control (T0), others at 2.6%, 3.5% and 4.4% of soil weight respectively against T1, T2 and T3 treatment. The results indicated moderately elevated seed cake application showed considerable improvement in vegetative growth and the plant biomass components including height and other morphological parameters as leaf ~~number-no~~, length of ~~leaf~~leaves, ~~no-number~~ of node and internodal distance per plant, root yields, fresh weight ~~&-and~~ dry matter production including the root to shoot ratio of the plants, in both the cultivars and in most of the cases found highest in hybrid yellow compared to hybrid purple. Among the plant components, shoot, root dry ~~wt~~ ~~weight~~ and root- ~~to~~ shoot ratio had the greatest decrease under fertilizer deficiency at control (T0) treatment. Among the different doses of seed cake applications, the highest values recorded at moderate treatments for most of the parameters, both at vegetative growth and post harvesting after ~~3-three~~ months of experiment.

Formatted: Font: Italic

The result shows that morphological parameters, biomass yield and its components of the plant were maximum when fertilized with seed cake at 2.6% and 3.5% of soil weight in treatments ~~T1 and &T2~~, found to be the perfect doses of fertilizer for the maize vegetative growth and biomass yield compared to T0 and other treatments. So, based on the result it can be concluded that T1 and T2 treatment is the best fertilizer treatment for the hybrid yellow maize yield.

**Key words:** Maize, organic fertilizer, mustard oil ~~Seed-seed~~ cake, growth, dry matter, ~~Root~~ ~~root~~ development

### **Introduction:**

Bangladesh is an agrarian country, where crops especially maize (*Zea mays* L.), one of the most important, demandable and widely grown cereal crops in recent years. Globally, maize is among the three leading staple food grains-(Pandita et al 2023, Anon 2006) and contains the highest genetic potential among all other grains (Choudhary et al 2020). As the leading producer, the United States, contributes to over 40% of the world's maize production (Moniruzzaman, 2009). It is a C4 plant, called "Queen of Cereals" ranks next to wheat and rice in context of grain production grown in irrigated and rainfed areas (Anon 2008, Irshad et al., 2002). C4 plants are well suited for hot and dry climates and produce higher amount of energy than C3 plant. As a C4 plant, hybrid maize is an emerging high value cereal crop in Bangladesh. It can be harvested as fodder within 50 days, green cobs within 60-80 days and grain within 100-130 days of planting. It has one of the highest numbers of consumers not only in Bangladesh but also worldwide (Dewan et al. 1998, Hasan et al., 2018, Adnan, et al., 2021). It has a huge commercial potential

Formatted: Font: Italic

**Comment [SP1]:** Correct as per journal guideline.

**Comment [SP2]:** Correct as per journal guideline

**Comment [SP3]:** Correct as per journal guideline

**Comment [SP4]:** Correct as per journal guideline

**Comment [SP5]:** Correct as per journal guideline

**Comment [SP6]:** Correct as per journal guideline

**Comment [SP7]:** Correct as per journal guideline

**Comment [SP8]:** Correct as per journal guideline

due to subtropical monsoonal climate of Bangladesh suitable for maize cultivation (Ali et al. 2008). Maize production in Bangladesh increased significantly in between 2011- 2020, which peaked in 2019 at 17.14% before falling to 14.63% in 2020 (Hasan, 2008; Ahmed and Islam, 2018; Islam and Hoshnain 2022). In 2020 was 9 tons per ~~heeter~~-hectare with an average yearly growth rate of 2.98% (Rahman and Zhang, 2018). Demand and production in Bangladesh would be more than doubled between 2050. Maize production is mainly concentrated in the ~~agroecological-Agro-Ecological zones-Zones~~ (AEZs) of ~~northern~~Northern, ~~western~~-Western and Eastern districts of Bangladesh (Ali et al., 2009).

Comment [SP9]: Incomplete sentence

Maize as a potential and multipurpose cereal grain crop for human consumption with a variety of prepared food items as well as fodder and feed for animals, livestock, poultry and especially uses of baby corn by humans including processed products as -loaf, flat bread, bun, popcorn, corn flex, cornmeal, corn flour are increasing day by day across the country and all over the world with great nutritional value of the grain having 66.70% starch, 10% protein, 4.8% oil, 8.5% fiber, 3% sugar and 7% ash (Mehta & Dias, 1999, Chaudhry, 1983). It grows at diverse seasons and having different types as normal yellow/white, hybrid yellow, hybrid purple, sweet corn etc.

Comment [SP10]: Correct as per journal guideline

Comment [SP11]: Pl make 3-4 simple sentences from this.

Morphological features including growth and development are generally hampered by disruption in the cell cycle machinery including cultivation under rainfed areas with lower rainfall than the critical level to obtain optimum yield (Sah et al., 2007-). Generally loamy soil texture is suitable for producing maize. However, adding fertilizer might boost up plant growth, where organic fertilizer is more suitable and beneficial for plant and soil health (Ayeni et al., 2012 and 2015).

In a developing country like Bangladesh, the cultivated soils are generally deficient in Organic matter and major essential nutrients as nitrogen (N), phosphorus (P) and ~~Potassium-potassium~~ (K). So low crop productivity in the mentioned country is common feature because of very low ~~Organic manure~~ and nutrient contents, poor soil physical condition including unbalanced use of synthetic fertilizers and low nutrient-use-efficiency (Rashid, 1994; Anon, 2006).

Farmers using both organic and inorganic/synthetic fertilizers in their crop fields however organic fertilizer boosts the plant growth in case of maize more than inorganic fertilizer (Adediran et al. 2005, Ayeni et al., 2012). The excessive and long-term use of chemical fertilizers deteriorate the quality and contaminate the soil including degradation of -air, surface as well as groundwater through leaching of nutrients affects human body as the chemicals moves through the food chain, responsible for developing chronic diseases in humans. -On the other hand, organic fertilizers are slow releasing and beneficial for soil, plant growth but also for human health. (Sugiyanto, 2011; Sharma and Chetani, 2017; Lokanadhan et al., 2012).

Therefore, the objective of this present research work ~~is-was~~ to assess the benefits of adding seed cake as organic fertilizer with application of different doses to investigate the growth, dry matter and root development performances and find out a suitable maize variety out of two in terms of growth, plant height, leaf ~~number~~, biomass yield, root development and root ~~to~~ shoot ratio performances under controlled conditions.

## Materials and methods:

A pot culture experiment was conducted from June -September, 2022 to investigate the influence of mustard oil seed cake application on maize plant growth and development, performed- under controlled environment -at the Laboratory, Department of ~~environmental~~ Environmental science ~~Science~~ and ~~management~~ Management, North South University, Bangladesh. The pots, were kept at a temperature of 25 °C under 14/10 hour light and dark photoperiod. The experiment was laid out in a completely randomized block design with four fertilizer levels as treatments and ~~3~~ three replications.

Formatted: Font color: Black

Seeds ~~was~~ were sown on 170 g of soil ~~was~~ filled in each plastic container with 5.5 inches height and 2.5 inches width. To conduct the lab trials, the treatments imposed on the soil were T0 (control), T1, T2 and T3. The fertilizer was added in 2-time intervals, one during the sowing and another after 32 days of germination.

The first doses of organic fertilizer was added in the soil during the sowing of seeds at the rate of no fertilizer (T0), 1.5g/pot (T1), 2 g/pot (T2) and 2.5g/pot (T3) gm respectively and the 2<sup>nd</sup> doses on day 32 at the rate of control (T0), T1 (3 g/pot), T2 (4 g/pot) and T3 (5 g/pot). As a result, the total seed cake fertilizer added during the experiment were control and others 4.5g/pot (2.6%), 6g/pot (3.5%) and 7.5 g/pot (4.4%) by soil weight respectively against T0, T1, T2 and T3 treatments.

The experiment was done with sandy loam soil for 81 days, where the hybrid yellow and hybrid purple maize cultivars were the test crops, and mustard oil seed cake as organic fertilizer, used in this experiment to evaluate the influence of fertilizer on vegetative growth, dry matter production, and root development of the plants

#### Identification of the plant:

~~Scientific name: *Zea mays* L., Kingdom: Plantae, Division: Magnoliophyta, Class: Liliopsida, Order: Poales, Family: Poaceae, Genus: *Zea*, Species: *mays*~~

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

#### Analysis of Mustard oil seed cake:

Total Nitrogen (N)-3.01%, Phosphorus (P)- 1.15% and Potassium (K)- 1.22%

Comment [SP12]: Methods for estimation of these need to be mentioned along with references.

#### Physiochemical properties of soil:

The soil ~~has~~had the following characteristics. Texture – Sandy loam. Sand- 61.26%, Silt - 32.62% and Clay – 6.12%. pH -6.9, EC - 793  $\mu$ S/cm, ~~NaCl~~NaCl - 1.0%, moisture content - 2%, Organic carbon (OC) -0.81%, Total Nitrogen- 0.05%, Available Phosphorus – 33.84 ppm, Exchangeable potassium – 0.09 meq/100 g soil

**Comment [SP13]:** Methods for estimation of these need to be mentioned along with references

Mustard oil seed cake as a potential soil amendment and organic fertilizer could enhance the maize growth and biomass yield. Therefore, experiment with ~~2~~two varieties of maize plants were carried at lab condition to study the effects of organic fertilizer at four levels on the hybrid purple and hybrid yellow cultivars on yield components.

**Formatted:** Justified



**Fig 1.** Seeds were soaked in water (ESM lab) Department of Environmental Science and ~~management~~Management, North South University



**Figure 2. 11 days old seedlings at the lab**

In these 81\_days' experimental periods, different parameters of plants, such as plant height, number of leaves, leaf length, number of nodes and internodal distance etc were measured from time to time and finally fresh and dry weight of shoot, root including root shoot ratio after harvesting of the plants. The results represent the average of 3-three replications.

**Comment [SP14]:** There should not be any etc. in research articles. All the parameters should be mentioned here long with methods used for estimation and units in which it was measured.


This study was conducted to find out the best dose of organic fertilizers for the maize yield so that a specific amount of dose can be applied to find the best result.

**Comment [SP15]:** Repetition. May be deleted.


## **Results and Discussion:**

### **Morphological features of hybrid yellow and hybrid purple cultivars at 32 and 72 days with different fertilizer treatments at vegetative growth:**

After 32 days of germination, result shows that highest plant height was found in yellow hybrid at treatment T2 (Fig 3) followed by T3, T1 and lowest at T0, also same in case of T2 treatment at purple hybrid, which was highest followed by T1, T3 and T0 at the lowest. In both the cultivars control treatment means no added fertilizer showed lowest plant height (Fig 3).




**Fig. 3 Effect of fertilizer treatments on plant height**




**Fig 4. Effect of fertilizer treatments on leaf length**

In case of leaf length, highest value was obtained at-in T1 in hybrid purple followed by hybrid yellow at T3 and T2 & and T3 at-in hybrid purple (Fig 4)



**Fig 5. Effect of fertilizer treatments on no-number of leaves per plant**

Different fertilizer treatments showed that the no of leaves was found higher at T2 and T3 -and lower at T1 and T0 in both the cultivars (Fig 5)




**Fig 6. Effect of fertilizer treatments on Plant node and internodal distance**


Plant node Number of nodes per plant was found highest at T3 both in yellow and purple hybrid followed by T2, T0 but -less in no at T1 in case of purple, however, in hybrid yellow, it was found same at T0 and T2. Also, internodal distance was -highest in T3 in both yellow and purple cultivars (Fig 6).

**Morphological parameters after 72 days of germination:**

Results after 72 days were found same as was after 32 days where plant height was highest at treatment T2 in hybrid yellow followed by T2 and T1 in hybrid purple and lowest at T0 in both the cultivars. In case of purple hybrid, among the treatments, found highest at T2 but was lower in comparison to hybrid purple, followed by T1 (Fig 7)




**Fig 7. Effect of fertilizer treatments on plant height**




**Fig 8. Effect of fertilizer treatments on leaf length**

In case of leaf length, it was found highest in yellow hybrid at T2 treatment followed by T2 in purple and T3, T1 & T0 in yellow hybrid (Fig 8)



**Fig 9. Effect of fertilizer treatments on leaf number**

Fertilizer treatments showed highest number of leaves leaf no- in both the cultivars at T3 followed by T2 and lowest at T0 and the node number of nodes per plant was found highest at T1 in yellow hybrid followed by purple at T3 and same in all others among the treatments and between the cultivars (Fig 9).




**Fig 10. Effect of fertilizer treatments on plant node and internodal distance**

In case of internodal distance, like node no also found highest in yellow hybrid at treatment T1 followed by T3, T2 and T0, whereas found lowest in case of purple in all the treatments compared to yellow one. Purple ~~variety shows~~ showed less internodal distance compared to yellow one (Fig 10)


**Parameters after 81 days of germination at harvesting :**

After harvesting at 81 days, shoot fresh and dry weight were found highest at T3 followed by T2 in hybrid yellow compared to hybrid purple whereas at T0 both the cultivars showing the lower performances compared to other treatments (Fig 11)




**Fig 11. Effect of fertilizer treatments on shoot fresh and dry weight**

In ~~Hybrid-hybrid~~ yellow, shoot fresh wt (8.99\_g) and dry weight (2.1\_g) was higher at T3 treatment followed by T2 and lowest in T0 in both the entities.



**Fig 12. Effect of fertilizer treatments on root fresh and dry weight**

In case of root performance parameters, highest values were obtained in root fresh and dry weight at T3 in hybrid purple followed by T2 in hybrid yellow and lowest at control in both the cultivars (-Fig 12)



**Fig 13. Effect of fertilizer treatments on root shoot ratio**

Root to shoot ratio was found highest (1.78 g) in hybrid yellow at T1 treatment, followed by T2 (1.76 g), and T3 (1.75 g) in hybrid purple whereas no differences was found at-in control treatment which was lowest at T0 (1.08 g) in both the cultivars (Fig 13).

Organic fertilizers improve soil health through moderating the physiochemical properties with improving the water retention capacity of soil and improves soil fertility by increasing soil organic carbon which affects positively on production of crops (He et al., 2022)

The results of this trial indicated organic fertilizer increased organic matter content of the soil. However moderate rates of fertilizer application is cost ~~saving~~ effective and favored vegetative growth as well as biomass including formation of strong root structure. This result also confirms the finding of ~~(Lewis (2019), who found organic fertilizer as seed cake improves the nutrient availability in soil and enhances plant growth and development.~~ The incorporation of mustard oil seed cake has the ability of increasing the essential nutrient contents to improve the physiochemical properties of soil and thus aided growth as well as development of the plants. In the research investigation the fertility status of the soil justifies adding of seed case as organic low-cost soil amendment to improve fertility of soil and thus growth and biomass yield in plants.

The results of this experiment are in agreement with other researchers (Adediran et al.,2005; Cahill et al., 2007; Rafiq et al., 2010; Soro et al.,2015; Jjagwe et al., 2020) ~~those~~ who found that slow releasing organic fertilizer has a positive impact more than the inorganic fertilizer to improve nitrogen use efficiency (NUE) and boost the growth, development and yield of maize crops

In the experiment, fertilizers were applied at 2.6% and 3.5% of soil weight in Treatment 1 and 2, found to be the perfect doses of fertilizer for the maize yield compared to controlled and other treatments.

The results indicated that the application of organic fertilizer significantly affected the plant morphology as plant height, leaf length including number of node per plant and number of internodes per plant, fresh and dry ~~weight~~ of straw and root yields, in both the cultivars, and was found more in hybrid yellow compared to ~~hybrid purple.~~ The interaction between seed cake (at different doses) recorded the highest values for most of the parameter at T1 and T2 during the ~~3~~ three months of experiment at lab scale.

The result ~~shows~~ showed that under controlled environment morphological parameters and biomass yield and its components of maize were maximum when fertilized with moderately higher amount of seed cake, mostly at T1 and T2 treatments, indicated considerable ~~increases~~ increase at ~~in~~ vegetative growth and the plant biomass components and ~~had~~ the greatest decrease under deficiency, ~~however~~ However, number of leaves per plant did not show much changes even ~~leaf no showed not much decreased~~ under fertilizer deficiency as T0 shows some

tendency of the crop to adjust with lower fertilizer levels in production of leaves for photosynthesis activities in plants.

It is evident from the results that the moderate fertilizer treatments in yellow hybrid performed better in boosting the maize growth including parameters of shoot and root development. These results also confirm the finding of Islam et al (2007), who postulates that application of moderate amount of seed cake provides better performance in cereal crop compared to higher dose from both yield and economic viewpoints.

So, based on the results it can be concluded that T1 and T2 treatment is the best fertilizer treatment for the hybrid yellow maize yield.

**Comment [SP16]:** How can you recommend this treatment best for yield? Did you study any of the yield parameters?



**Fig 14. Harvested hybrid yellow and hybrid purple cultivar after 3 months**

Plants of the Hybrid yellow variety were found healthier compared to that of the hybrid purple in terms of above ground and belowground parts of the plants (Fig 14)





Fig 15. Root structures of Hybrid-hybrid yellow with different treatments (after harvesting)

Root structures of hybrid yellow found more stronger compared to hybrid purple (-Fig 15 and 16)



**Fig 16. Root structures of Hybrid purple with different treatments (after harvesting)**

### **Conclusion:**

At fertilizer treatments T1 and T2, the cost will be ~~deducted~~ reduced and production will be increased, implying the maximum benefits at a minimum cost. For this reason, it will be very helpful for the small farmers as well as the general people who want to grow maize in the pot or on rooftop gardens in an effective way. The findings will serve as a stepping stone for more studies on maize cultivation in the future that will consider our limitations and find variable solutions.

From the results it is found that organic fertilizer application can provide significant positive enhancement to maize growth and development as compared to duration especially in vegetative growth and biomass yield. It is recommended that moderate application of mustard oil seed cake as organic fertilizer should be encouraged to the farmers for vegetative growth of maize at the expanse of dry matter production. So, based on the result it could be concluded that in a developing country like Bangladesh, effective fertilizer treatment could be cost-effective with resourceful output in large agricultural areas to maintain their sustainability.

#### **Declaration of competing interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

**Acknowledgements:** This research was supported by the Environmental Laboratory, Department of Environmental Science and Management, North South University, Dhaka, Bangladesh. Author thanks to Soil Research Development Institute (SRDI), Bangladesh, for analyzing the samples of soil and seed cake.

#### **References:**

- Adediran, J. A., Taiwo, L. B., Akande, M. O., Sobulo, R. A., & Idowu, O. J. (2005). Application of organic and inorganic fertilizer for sustainable maize and cowpea yields in Nigeria. *Journal of plant nutrition*, 27(7), 1163-1181.
- Adnan, K. M., Sarker, S. A., Tama, R. A. Z., & Pooja, P. (2021). Profit efficiency and influencing factors for the inefficiency of maize production in Bangladesh. *Journal of Agriculture and Food Research*, 5, 100161.
- Ahmed, M. W., & Islam, M. N. (2018). Moisture sorption characteristics of selected commercial flours (wheat, rice and corn) of Bangladesh. *American Journal of Food Science and Technology*, 6(6), 274-279.
- Ali, M. Y., Waddington, S. R., Hodson, D. P., Timsina, J., & Dixon, J. (2008). *Maize-rice cropping systems in Bangladesh: Status and research opportunities* (No. 559-2016-38850).
- Ali, M. Y., Waddington, S. R., Hodson, D. P., Timsina, J., & Dixon, J. (2009). *Maize-rice cropping systems in Bangladesh: Status and research needs*. *Journal of Agricultural Science and Technology*, 3(6), ISSN 1939-1250, USA
- Anonymous. 2006. The impact of deregulation on the fertilizer sector and crop productivity in Pakistan. pp. 46-47.
- Anonymous. 2008. Agricultural Statistics of Pakistan, Ministry of Food, Agriculture and Livestock (MINFAL), Govt. of Pakistan.

- Ayeni, L. S., Adeleye, E. O., & Adejumo, J. O. (2012). Comparative effect of organic, organomineral and mineral fertilizers on soil properties, nutrient uptake, growth and yield of maize (*Zea mays*). *International Research Journal of Agricultural Science and Soil Science*, 2(11), 493-497.
- Cahill, S., D. Osmond, C. Crozier, D. Israel and R. Weisz. 2007. Winter wheat and maize response to urea ammonium nitrate and a new urea formaldehyde polymer fertilizer. *Agron. J.*, 99(6): 1645-1653
- Chaudhary, A.H. 1983. Effect of population and control of weeds with herbicides in maize. *Field Crop Abst.*, 35(5): 403.
- Choudhary, M., Grover, K., and Singh. M (2020). Maize significance in Indian food situation to mitigate malnutrition. 98(2), *Cereal chemistry*. Wiley online library
- Dewan, S. F. K., Mondal, A. A., & Salam, M. A. (1998). Prospects and challenges of expanding maize production in Bangladesh.
- Hasan, M. F. (2008). Economic efficiency and constraints of maize production in the northern region of Bangladesh. *J. innov. dev. strategy*, 2(1), 18-32.
- Hasan, M. R., Rahman, M. R., Hasan, A. K., Paul, S. K., & Alam, A. H. M. J. (2018). Effect of variety and spacing on the yield performance of maize (*Zea mays* L.) in old Brahmaputra floodplain area of Bangladesh. *Arch. Agric. Environ. Sci*, 3(3), 270-274.
- He H, Peng M, Ru S, Hou Z and Li J (2022). A suitable organic fertilizer substitution ratio could improve maize yield and soil fertility with low pollution risk. *Front. Plant Sci.* 13:988663
- Islam M.M., Anwar M.P., Rahman, M.M & Islam, A.K.M.M (2007). Influence of Mustard oil cake on performance of fine rice CV. Chinigura. *Intl. J. BioRes* 3(6):50-54
- Irshad, M., S. Yamamoto, A.E. Eneji, T. Endo and T. Honna. 2002. Urea and manure effect on growth and mineral contents of maize under saline conditions. *J. Plant Nutr.*, 25(1): 189-200.
- Islam, M. R., & Hoshain, S. (2022). A Brief Review on the Present Status, Problems And Prospects of Maize Production in Bangladesh. *Research in Agriculture Livestock and Fisheries*, 9(2), 89-96.
- Jjagwe J, Chelimo K, Karungi J, Komakech AJ, Lederer J. (2020). Comparative Performance of Organic Fertilizers in Maize (*Zea mays* L.) Growth, Yield, and Economic Results. *Agronomy*. 10(1):69.
- Lokanadhan, S., Muthukrishnan, P., & Jeyaraman, S. (2012). Neem products and their agricultural applications. *Journal of Biopesticides*, 5, 72.
- Lewis, K.L., Hons, F.M. and Gentry, T.J. (2019), Beneficial Use of White and Indian Mustard Seed Meals to Enhance Plant Growth and Nutrient Uptake. *Agrosystems, Geosciences & Environment*, 2: 1-10 180042.
- Moniruzzaman, M. S. R. S., Rahman, M. S., Karim, M. K., & Alam, Q. M. (2009). Agro-economic analysis of maize production in Bangladesh: a farm level study. *Bangladesh Journal of Agricultural Research*, 34(1), 15-24
- Mehta C.D., & F.F. Dias. (1999). Maize perspective in India. Wiley. 51( 2-3): 52-57
- Pandita, D., Parthasarathy, S., Dhivyapriya, D., Premkumar, R., Pandita, A., Wani, S.H. (2023). Genome Diversity in Maize. In: Wani, S.H., Dar, Z.A., Singh, G.P. (eds) *Maize Improvement*. Springer, Cham.

- Rafiq, M. A., Ali, A., Malik, M. A., & Hussain, M. (2010). Effect of fertilizer levels and plant densities on yield and protein contents of autumn planted maize. *Pak. J. Agri. Sci*, 47(3), 201-208.
- Rahman, K. A., & Zhang, D. (2018). Effects of fertilizer broadcasting on the excessive use of inorganic fertilizers and environmental sustainability. *Sustainability*, 10(3), 759.
- Rashid, A. 1994. Annual report of micronutrient project. Land Resources Research Institute (LRRRI), NARC, Islamabad, Pakistan.
- Sharma, A., & Chetani, R. (2017). A review on the effect of organic and chemical fertilizers on plants. *Int. J. Res. Appl. Sci. Eng. Technol*, 5, 677-680.
- Soro, D., Ayolié, K., Zro, F. G. B., Yéboua, F. Y., Kouadio, H. K. K., Bakayoko, S., & Yatty, J. (2015). Impact of organic fertilization on maize (*Zea mays*) production in a ferralitic soil of centre–West Cote D'ivoire. *Journal of Experimental Biology*, 3(6), 556-65
- Shah, Z., Z. Shah, M. Tariq and M. Afzal. 2007. Response of maize to integrated use of compost and urea fertilizers. *Sarhad J. Agric.*, 23(3): 667-673.
- Sugiyanto, C. (2011). The benefits of organic fertilizer application: Case Indonesia: Garut and Sragen regencies. Indonesia Regional Science Association, IRSA 12 Conference in Universitas Andalas, Padang, West Sumatra, Indonesia.