

**EVALUATION OF THE BIOACTIVITY AND TOXICITY PROPERTIES OF *Funtumia elastic* (Preuss) Stapf. (TSN 30176), USED IN NIGERIA ETHNOMEDICINE.**

**Proposition 1:** Evaluation of the antibacterial activity and toxicity properties of *Funtumia elastica*, used traditional medicine in Nigeria

**Proposition 2:** Evaluation of the antibacterial activity and toxicity properties of *Funtumia elastica* fractions, used traditional medicine in Nigeria

**ABSTRACT**

Background: The urgent need for new and novel antibacterial chemotherapy to combat the worrisome emergence of antimicrobial resistance necessitated the exploration of forest resources, especially the plants, for possible breakthrough. *Funtumia elastic* is a promising plant species in this regards, with a lot of ethnopharmacological claims.

**Aims:** This study was designed to evaluate the antibacterial activity, acute and sub-acute toxicity studies of *Funtumia elastica* which is used in the ethnomedical treatment of infections.

**Methods:** The leaves and stem bark of *F. elastica* were collected, pulverized, and Soxhlet extracted with distilled ethanol. Subsequently, the ethanol extracts were partitioned with n-hexane, dichloromethane, and ethyl acetate according to polarity. The dichloromethane fractions were subjected to column chromatography coupled with thin layer chromatography. Antimicrobial susceptibility testing and crude extracts were studied on clinical isolates using agar well diffusion. Minimum inhibitory concentrations (MIC) and minimum bactericidal concentrations (MBC) were also determined. Acute and subacute toxicity studies were evaluated on the dichloromethane extract (DCM) using female Swiss albino mice.

**Results:** The DCM and ethylacetate extracts of *F. elastica* were found to have some degree of activities on clinical isolates. Bioactivity guided fractionation of the DCM extract yielded only fraction 2 been active on the clinical isolates at 3.5625mg/ml. out of nine (9) fractions There were no signs of acute toxicity at the maximum dose of 5000mg/kg body weight. Biochemical parameters showed no significant changes in the liver enzymes, this infers that the extract might not have any negative impact on the liver with regards to metabolism.

**Conclusion:** The moderate antibacterial activities of *Funtumia elastica* extracts justified its ethnomedicinal uses and potential to furnish new antimicrobials.

**KEY WORDS:** *Funtumia elastic*, ethnomedicine, Phytochemistry, Bioactivity, Toxicity.

## INTRODUCTION

The use of medicinal plant in treatment of sickness and disease is dated back to ancient times, and it is still being used in the 21<sup>st</sup> century (Apkuaka and Ezem, 2011). Medicinal plants have been tagged a reservoir for phytochemicals in the production of drugs. Although they are mostly consumed wholly, research has proven that this act could have some deleterious effects on humans (Ahn, 2017). There has been a rapid emergence of resistant bacteria occurring worldwide, endangering the efficacy of antimicrobial agents (Okeke, 2005). Hence there is urgent need to look into medicinal plants for new and effective antimicrobials.

*Funtumia elastica* (Preuss) commonly called Lagos silk rubber and “Ire” in Yoruba land Southwest Nigeria, belongs to the family Apocynaceae. Members of this family are native to the Europe, Asia, Africa, Australia and America. Many species of the Apocynaceae family are found in the tropical forest, but some grow on temperate regions as perennial herbs. Preliminary phytochemical screening has revealed that *F. elastica* bark contains hydrolysable tannins,

sapogenetic glycosides, steroids, and saponins, while the leaves contain hydrolysable tannins, flavonoids, starch, and alkaloids with the tannin content of the leaves and stem bark being 2.4% and 1.3% w/w (relative to the dried material), respectively (Agyare *et al.*, 2012).

Decoction of the leaf of *Funtumia elastica* has been found to specifically inhibit the growth of many moulds, including *Aspergillus* sp, *Penicillium* sp, *Candida* sp, as well as the fungi that cause ring worm. It is also reportedly used in the treatment of wounds (Adekunle and Ikumapayi 2006). Burkill, (1995) has documented that ethanolic extract of *F. elastica* (Preuss) is traditionally used to treat whooping cough, inflammatory diseases such as asthma, blennorhea and painful menstruation. The scientific scrutiny of *Funtumia elastica* is as a result of the need to look for other sources of novel antimicrobial agents particularly from medicinal plants based on ethno-pharmacological information. The present study, was, therefore, designed to evaluate antibacterial activity of fractions from the crude extract of *Funtumia elastica* as well as to assess its acute and subacute toxicity.



**Figure 1** *Funtumia elastica* trees (Source: Author`s work)

## **MATERIALS AND METHODS**

### **Collection, identification and Extraction of Plant Materials.**

*Funtumia elastica* was collected from University of Ibadan Botanical garden. The plant was earlier identified and authenticated at the herbarium of Federal Research Institute of Nigeria (FRIN), Ibadan with voucher number FHI 112304. The plant parts (leaves and stem bark) were air-dried under shade at ambient temperature, pulverized into fine powder. The pulverized plant materials were weighed and subjected to exhaustive Soxhlet extraction with distilled ethanol. Extraction solvents were collected, concentrated and dried under reduced pressure using rotary evaporator (Xian Ltd, China) at 45°C, weighed and stored at room temperature before use. A 200g of each part of the crude extract was dissolved in a mixture 450ml of ethanol and 150ml of distilled water and was partitioned with n-Hexane, Dichloromethane and Ethyl acetate according to the polarity.

#### **Animal stock**

Female Swiss albino mice aged 6-7 weeks, weighing 25-30 g were used. The animals were fed on grower feed with free access to water under standard conditions of light (12 h light, 12 h dark), humidity and temperature.

#### **Phytochemical screening**

Phytochemical screening was carried out to detect the presence of secondary metabolites such as anthraquinones, tanins, saponins, alkaloid, phlobatanins, steroids, cardiac glycosides, reducing sugar, phenol, glycosides and resins using methods described by Harborne (1973).

#### **Test Microorganisms**

Strains of clinical isolates of *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Escherichia coli*, *Acinetobacter baumannii* and *Klebsiella spp* were gotten from Clinical Microbiology Laboratory University College Ibadan.

UNDER PEER REVIEW

## **Antimicrobial Susceptibility Test**

The agar cup diffusion technique was adopted using the method of (Adeniyi *et al.*, 1996). Each organism was sub-cultured in TSB (define) overnight, to have them in their log phase of growth. A volume of 0.1ml of each organism from the overnight culture was added to 9.9ml of sterile distilled water to give a 1 in 100 dilution. This was properly mixed to measure homogeneity.

Using a sterile 1ml pipette, 0.2ml of this dilution was withdrawn, introduced into 20ml molten Mueller Hinton agar at 45°C and poured into sterile Petri dish (diameter of Petri dish). The Petri dish was allowed to set and surface aseptically dried of all moisture. A sterile cork-borer of 6 mm diameter was used to bore equidistant wells on the agar and a 0.2ml volume of the extracts of various concentrations introduced into their respective wells. The positive (Gentamicin 10µg, add the name of all other antibiotics used with their concentration or mass, see table 2) and negative controls (20% DMSO) were also introduced in their respective wells. The Petri dishes were then allowed to stand on the bench for an hour to allow for pre-diffusion of extracts and controls. The Petri dishes were incubated at 37°C for 24 hours, after which zones of inhibitions were measured. All tests were performed in duplicates.

## **Determination of Minimum Inhibitory Concentration (MIC)**

Minimum inhibitory concentrations (MICs) were performed with modifications of agar dilution method as previously explained (Adeniyi *et al.*, 2004). Two milliliter (2ml) of each of the reconstituted extracts fractions, prepared to give concentrations of 200 mg/ml, 100 mg/ml, 50 mg/ml and 25 mg/ml, were added to 18 ml of molten Mueller-Hinton agar and shaken by rolling it between palms. The properly mixed agar and extract was poured into sterile Petri dishes, allowed to solidify and the surface dried in sterile oven to remove moisture. A loopful of

overnight culture of each test bacterial isolate was streaked on the surface of the agar. The Petri dishes were appropriately incubated for 24 hours at 37°C and examined for the presence or absence of growth. The lowest concentration that prevented growth of a bacterial isolate was taken as the minimum inhibitory concentration (MIC). All tests were performed in duplicates.

**Determination of Minimum Bactericidal Concentration (MBC)???? See result Table 5: Minimum Bactericidal Concentration (MBC) (mg/ml) of *Funtumia elastica* fractions**

**Acute toxicity studies**

You will need to justify the choice of Dichloromethane leaf extract of *Funtumia elastica* for Acute toxicity and Sub-acute toxicity studies (example: Acute toxicity studies of most active crude extract). The method of Lorke (1983), was adopted in determining the effect of acute oral administration of Dichloromethane leaf extract of *Funtumia elastica*. The study was carried out in three phases. In the first phase, three mice were administered 10, 100 and 1000mg/kg body weight of the extract orally (via a cannula), respectively. The mice were observed for signs of adverse effects and death for 24 hours. In the second phase of the study, the procedure was repeated using another set of three mice given 1500, 2000 and 2500mg/kg body weight of the extract respectively. The mice were also observed for signs of toxicity and mortality for 14 days. Based on the results gotten from the second phase, 3000mg/kg body weight and 5000mg/kg body weight of the extract were administered to another set of two mice in the third phase. The mice were observed for signs of toxicity and mortality for the first 4 hours and thereafter daily for 14 days.

**Sub-acute toxicity studies (example: Sub-acute toxicity studies of most active crude extract)**

The method of Aniagu *et al.*, (2005), was employed in the sub-acute toxicity study of crude Dichloromethane leaf extract of *Funtumia elastica*. Thirty (30) mice were selected by stratified randomization and divided into six groups of five mice each. Group one, two, three, four and five were administered 25, 50, 100, 200 and 400mg/kg body weight of the crude Dichloromethane leaf extract of *Funtumia elastica* respectively for 28 days. The sixth group was administered 5% DMSO in water used for dissolving the extract as control group. The first day of dosing was taken as D0 whereas the day of sacrifice was designated as D28. The feed intake of the mice in each group was measured once daily over the 28 days period. After sacrifice, heparinized blood sample were taken for hematological examination. The serum from non-heparinized blood was carefully collected for blood chemistry and analysis. All organs were preserved in 10% formal saline solution for histological examination. The effects of the sub-chronic administration of the crude extract for 28 days on the mice were investigated.

## **Chromatographic Methods**

### **Column chromatography of most active crude extract**

The most active extract was further purified by subjecting it to column chromatography of height, length and internal diameter 95cm by 54cm by 2cm respectively. The one hundred and fifty grams (150g) silica gel was packed wet with *n*-Hexane on the column clamped in a vertical position. As solid phase settles down, the column is gently tapped with a spatula until required height was attained. The sample (10 g of crude Dichloromethane leaf extract) was dissolved in a volatile *n*-Hexane and mixed with a little silica gel to form a slurry. Gradient elution was carried out starting from least polar solvent (that is 100% *n*- Hexane, *n*-Hexane/Ethylacetate, 100% Ethylacetate, Ethylacetate/Dichloromethane, 100% Dichloromethane,

Dichloromethane/methanol and 100% Methanol). Sub-fractions were obtained and pooled based on their thin layer chromatography (TLC) profiling.

### **Thin Layer Chromatography.**

The dissolved fractions of the column chromatography were spotted on TLC pre-coated silica gel adsorbent plate activated in an oven at a temperature of 110°C for about an hour, with the aid of capillary tubes at a distance of 1cm from the base. The spots numbered (F1, F2 .....??) were equidistant from each other and were allowed to dry on the plates and were subsequently placed in a tank saturated with different solvent system such as *n*-Hexane: Ethylacetate, *n*-Hexane: Dichloromethane etc, at different ratios. The plates were developed using ascending techniques and the developed plates were brought out of the tanks, solvent front marked, before drying the plates. Then using ultraviolet lamp ( $\lambda_{max}$  of 254 and 365 nm), the compound was clearly observed and marked. Their retardation factors ( $R_f$ ) were obtained using the formula:

$$R_f = \frac{\text{Distance moved by the sample}}{\text{Distance of the solvent front}}$$

### **Contact bioautography**

The fractions gotten from column chromatography were spotted on TLC plate and using appropriate solvent system were developed to separate the phyto-chemicals (Repetition, can be detected). The developed TLC plate were placed in a Petri dish and cooled molten agar that has been seeded with the organism of interest were poured on them and were allowed to solidify.

The plates were incubated at 25°C for 24hrs. Zones of inhibition were observed. For a clearer observation, the plates were sprayed with tetrazolium salt and re-incubated for 24hour at 25°C or 37°C??. Clear white zones, against a purple background indicated that the plant extract has antimicrobial activity against the test organism.

Statistical analysis????

## RESULTS

Dichloromethane fractions consistently gave the highest yield both in the leaves and stem-bark extracts, while the least percentage yield was recorded in the aqueous fraction for the leaves, and n-Hexane for the stem-bark respectively. (To be reviewed because in the stem-bark, the aqueous fraction (37.33%) has a high yield compared to that of the Dichloromethane fraction (20.97%)).

Table 1: Yield and characteristics of crude extract of *Funtumia elastica*

Sample	Weight (g)	%Yield	Characteristic features
Leaves			
n-Hexane	29.588	17.93	Coffee brown oily extract
Dichloromethane	88.263	53.49	Dark brown powdery extract
Ethylacetate	25.699	15.57	Deep coffee gummy extract
Aqueous	21.357	12.94	Dark greenish extract
Bark			
n-Hexane	26.759	12.68	Greasy brown extract
Dichloromethane	44.257	20.97	Dark brown gummy extract

Ethylacetate	61.2	29	Dark brown extract
Aqueous	78.78	37.33	Light brown crispy extract

The majority of the tests organisms were susceptible to Ofloxacin, Gentamicin, ciprofloxacin and Nitrofurantoin. All the bacteria studied were resistant to Augmentin and Ceftazidime. Only *P. aeruginosa* (P2) showed a slight sensitivity in the presence of and Cefotaxime (CTX).

**Table 2:** Anti-biogram of pathogenic organisms

DRUGS	Diameter zones of inhibition (in mm)									
	EC 3528	PA 27853	9A	4B	12B	P2	P4	EC1	EC23922	ACENI 1
<b>AUG</b>	-	-	-	-	-	-	-	-	-	-
<b>OFL</b>	24	-	18	20	24	24	25	-	18	-
<b>CXM</b>	11	-	-	-	-	-	-	-	-	-
<b>GEN</b>	17	15	17	15	-	24	12	-	-	-
<b>CTX??</b>	-	-	-	-	-	9	-	-	-	-
<b>CAZ</b>	-	-	-	-	-	-	-	-	-	-
<b>CPR</b>	27	31	-	16	-	26	35	-	19	24
<b>NIT</b>	26	-	25	30	20	26	-	-	28	21

**Key:**

EC 3528 - *E. coli* or

EC35218 ???

PA 27853 – *P. aeruginosa*

9A *E. coli*

4B - *E. coli*

12B – *K. pneumoniae*

AUG= Augmentin

OFL= Ofloxacin

CXM= Cefuroxim

GEN= Gentamicin

CTX= Cefotaxime

CAZ= Ceftazidime

P2 – *P. aeruginosa*  
P4 – *P. aeruginosa*  
EC1- *E. coli*  
EC 23922- *E. coli*

CPR= Ciprofloxacin  
NIT= Nitrofuratoin

ACENI 1- *A. baumannii*

-: No diameter or not applicated ???

No diameter zones of inhibition on isolates of *Staphylococcus aureus* ??? (see table 3).

Concentrations or mass for antibiotics

The dichloromethane and ethylacetate fractions of ethanolic crude extract of *Funtumia elastica* exhibited different degrees of activities while the aqueous and *n*-hexane fractions showed no activity on the pathogenic organisms.

**Table 3: Antibacterial susceptibility of different fractions of *F. elastic* (All fractions???) see table**

**4 title**

	DIAMETER ZONES OF INHIBIT																										
	m)																										
	DCM LEAVES					DCM BARK					EHTYLACETATE BARK					EHTYACETATE LEAVES											
	1	2	6.	3.12	5	D	MS	2	12	2	12	Gen	1	2.	6.2	3.1	25	5	5	25	Gent	5	.5.	5	5	10	G
	M	M	M	5	10μ	20	M	M	M	M	10μ	M	M	M	M	10μ	M	M	M	M	μ	20					
	g	g	g	mg	g	%	g	g	g	g	g	g	g	g	g	g	g	g	g	g	g	g	g	g	g	g	%
	1											1															
PA27853	0	8	7	-	-	-	8	7	-	-	-	13	1	9	-	-	5	12	7	-	-	-					
	2	2	1				2		1			1															
SA29213	3	1	9	17	15	-	0	17	0	8	15	15	3	8	7	19	5	13	7	-	18	-					
	2	2	1				2		1			1															
SA6571	5	2	8	14	19	-	2	20	6	9	20	14	2	7	-	19	6	11	-	-	19	-					
	1						1																				
EC35218	5	8	-	-	19	-	3	7	-	-	18	12	-	-	-	18	2	-	-	-	18	-					
	1	1					1																				
EC23922	5	1	7	7	10	-	5	11	7	-	10	11	-	-	-	22	4	12	-	-	18	-					
P4	-	-	-	-	15	-	7	-	-	-	18	10	-	-	-	14	-	-	-	-	15	-					
	1																										
ACINE1	6	9	7	-	-	-	8	7	-	-	-	10	-	-	-	-	4	12	-	-	-	-					
	1						1																				
4B	6	9	7	-	15	-	1	9	7	7	16	10	-	-	-	15	1	9	-	-	17	-					
	1	1	1																								
12B	3	1	0	8	14	-	9	8	-	-	13	10	8	-	-	21	-	-	-	-	17	-					
9A	-	-	-	-	19	-	-	-	-	-	18	15	2	7	-	21	0	8	8	-	21	-					

	1	1					1															
P2	2	0	9	8	19	-	0	8	-	-	18	-	-	-	-	18	-	-	-	16	-	
	1	1					1															
EC1	6	3	8	7	17	-	4	11	8	-	18	-	-	-	-	17	-	-	-	-	17	-

**Key:**

PA27853 - *Pseudomonas aeruginos*  
SA29213 – *Staphylococcus aureus*

SA6571- *Staphylococcus aureus*

EC35218 - *Escherichia coli*

EC23922 – *Escherichia coli*

P4 - *Pseudomonas aeruginosa*

ACINE1 – *Acinetobacter baumannii*

EC35218 - *Escherichia coli* (Repetition)

4B – *Escherichia coli*

12B – *Klebsiella pneumoniae*

9A- *Escherichia coli*

P2-*Pseudomonas aeruginosa*

EC1- *Escherichia coli*

DCM- Dichloromethane

DMSO- Dimethylsulfoxide

Table 3 does not present the results for the hexane and aqueous fractions. This may be related to the download of the manuscript. However, the author should better present this table for a better comparison.

Check the extract concentrations unit for the inhibition diameter texts (mg/ml or mg???)

The minimum inhibitory concentrations (MIC) of the four (4) partitioned crude extracts (dichloromethane leaves, dichloromethane bark, ethylacetate leaves and ethylacetate bark) against the test organisms ranges between <3.125 mg/ml to 25 mg/ml.

**Table 4:** Minimum Inhibitory Concentration of the four fractions of the plant extracts against the test bacterial isolates

Organisms	Dichloromet	Ethyl	Dichloromet	Ethyl	Gentamicin
	hane	acetate	hane	acetate	
	leaf (mg/ml)	leaf (mg/ml)	bark (mg/ml)	bark (mg/ml)	(µg/ml)
<i>Staphylococcus aureus</i> ATCC					
29213	3.125	6.25	3.125	6.25	5
<i>Staphylococcus aureus</i> ATCC					
6571	3.125	12.5	3.125	6.25	5
<i>Escherichia coli</i> ATCC 35218	6.25	25	12.5	25	5
<i>Escherichia coli</i> ATCC 23922	3.125	12.5	6.25	25	5
<i>Escherichia coli</i> ???	>25	6.25	>25	12.5	<1.25
<i>Escherichia coli</i> ???	6.25	12.5	3.125	25	5
<i>Escherichia coli</i> ???	3.125	>25	6.25	>25	<1.25
<i>Klebsiella pneumonia</i>	3.125	>25	12.5	12.5	5
<i>Pseudomonas aeruginosa</i>					
ATCC 27853	6.25	6.25	12.5	6.25	>10
<i>Pseudomonas aeruginosa</i> ???	>25	>25	25	25	5
<i>Pseudomonas aeruginosa</i> ???	12.5	>25	12.5	>25	<1.25
<i>Acinetobacter baumannii</i>	>25	12.5	12.5	25	>10

Note: Table 2 (10 strains), Table 3 (12 strains), Table 4 (12 strains), Table 5 (11 strains??)

UNDER PEER REVIEW

The minimum bactericidal concentrations of the various fractions of the leaves and stem-bark extracts of *Funtumia elastic*, shows that values ranged between the MIC and 2 X MIC values for each organism

**Table 5:** Minimum Bactericidal Concentration (MBC) (mg/ml) of *Funtumia elastica* fractions

Organisms	PA 27853	SA 29213	SA 6571	EC 35218	EC 23922	P4	ACINE 1	4B	12B
<b>Samples</b>									
DCM LEAVES	12.5	3.125	3.125	25	6.25	50	12.5	12.5	3.125
DCM BARK	25	3.125	3.125	25	12.5	50	25	3.125	25
E.A BARK	12.5	3.125	12.5	50	50	50	50	50	25
E.A LEAVES	12.5	12.5	25	50	25	50	25	25	50

**Key:**

PA 27853- *Pseudomonas aeruginosa*

SA 29213- *Staphylococcus aureus*

EC 35218 – *Escherichia coli*

EC 23922 – *Escherichia coli*

P4- *Pseudomonas aeruginosa*

ACINE 1- *Acinetobacter baumannii*

4B – *Escherichia coli*

12B – *Klebsiella pneumoniae*

9a – *Escherichia coli*

P2 – *Pseudomonas aeruginosa*

EC1 – *Escherichia coli*

Among the fractions spotted on the TLC plate, the F2 fraction from the dichloromethane extract of the leaves of *F. elastic* was the most active !!!?. Results of the antibacterial susceptibility of this fraction 2 (F2), showed *Staphylococcus aureus* ATCC 29213 was most susceptible to the fraction.

**Table 6:** Antibacterial susceptibility of Fraction 2 of Dichloromethane fraction leaves extract of *F. elastica* against some bacterial isolates

Organisms	Diameter zone of inhibition (mm)							Gent 10µg	DMSO 20%
	6.25 mg	3.125 mg	1.625 mg	0.781 mg	0.39 mg	0.195 mg			
<i>Staphylococcus aureus</i>									
ATCC 29213	20	15	10	-	-	-	12	-	
<i>Escherichia coli</i> ATCC									
23922	14	12	-	-	-	-	10	-	
<i>Pseudomonas aeruginosa</i>									
ATCC 27853	14	12	-	-	-	-	15	-	



Key: DMSO- Dimethylsulfoxide. Gent.- Gentamycin.

why, this table 6 does not show the results of the other bacteria tested???

It would also be interesting to add few comments to the results of the tables


Check the extract concentrations unit for the inhibition diameter texts (mg/ml or mg???)

**Make good texts that announce the Plates**



**Plate 1:** Histological evaluation of effects of Dichloromethane extract of *Funtumia elastica* using hematoxylin and eosin staining on liver tissues of control in the acute toxicity study in mice. Photomicrograph of stained sections of liver tissues using hematoxylin and eosin shows mild diffuse vacuolar degeneration of hepatocytes. A = X100 magnification; B = X400 magnification. (This part can be well commented, interpreted and put before the figure)

**Make good texts that announce the Plates**



(A)

(B)

**Plate 2:** Histological evaluation of effects of Dichloromethane extract of *Funtumia elastica* using hematoxylin and eosin staining on kidney tissues of control in the acute toxicity study in mice. Photomicrograph of stained sections of kidney tissues using hematoxylin and eosin shows no visible lesions seen. A = X100 magnification; B = X400 magnification. (This part can be well commented, interpreted and put before the figure)

UNDER PEER REVIEW

**Make good texts that announce table 7 and 8**

A significant increase was seen in the packed cell volume (PCV) and hemoglobin concentrations as well as red blood cell and white blood cell count.

**Table 7:** Effect of sub-acute toxicity of Dichloromethane leaves extract of *F. elastic*, on hematological indices in mice.

Parameters	Control	25mg/kg	50mg/kg	100mg/kg	200mg/kg	400mg/kg
PCV (%)	45.67 ± 0.33	51.33 ± 0.67*	55.33 ± 2.67*	47.33±0.67	49±1	53±1*
Hemoglobin (g/dl)	15.07 ± 0.13	17.40 ± 0.21*	18.40 ± 0.3*	15.83±0.03*	16.63±0.07*	16.87±0.07*
RBC(mil/mm <sup>3</sup> )	7.36 ± 0.30	8.57 ±0.01*	8.42 ± 0.07*	8.07 ± 0.1*	8.39±0.02*	8.46±0.05*
WBC(mm <sup>3</sup> )	4697±46.7	4250±50	5027 ±367	3507±256.7*	4400 ± 100	2683±183.3*
Platelet(mm <sup>3</sup> )	132350±1650	123268±1268	195680±6053*	95000±5000*	115000±5000	103030±8970*
Lymphocytes (%)	73±0.67	78.0 ± 1.0	79.0 ±3.53	76.0 ± 1.0	74.67 ± 0.33	79.67 ± 1.67
Neutrophils (%)	23.67 ± 0.67	21.67 ± 0.33	17.67 ± 1.33	24.67 ± 1.67*	19.67 ± 1.33	20.33 ± 0.33
Monocytes (%)	2.17 ± 0.17	1.17 ± 0.17*	0.93 ± 0.07*	1.13 ± 0.13*	1.83 ± 0.17	2.27 ± 0.27
Eosinophil (%)	1.17 ± 0.17	0.83 ± 0.33	2.83 ± 0.17*	0.83 ±0.17*	1.83 ±0.17	2.13±0.13*

Data are expressed as Mean ± SEM (n=3). Comparisons were made using one-way ANOVA followed by post-hoc

Newman-Keuls test. \*P< 0.05 when compared with control.

**Key:**

PCV= Packed Cell Volume; WBC= White blood cell ; RBC= Red blood cell

There was no significant alterations in the serum ALT, AST, and ALP and other biochemical parameters analyzed at various doses of the extract after 28 days administration on the animal models.

**Table 8:** Effect of sub-acute toxicity of Dichloromethane leaves extract of *F. elastica* on serum biochemical parameters

Parameters	Control	25mg/kg	50mg/kg	100mg/kg	200mg/kg	400mg/kg
Total Protein(g/dl)	6.4± 0.21	5.93 ± 0.56	6.33 ± 0.17	5.6 ± 0.70	5.6± 0.70	6.1±0.10
Albumin(g/dl)	2.5 ± 0.1	2.47 ± 0.13	2.37 ± 0.13	2.43 ± 0.13	2.37±0.13	2.33±0.17
Globulin(g/dl)	4.0 ± 0.1	3.7 ± 1.2	3.7 ± 0.3	3.47 ± 0.07	3.9±0.2	3.6±0.1
A-G Ratio	0.53±0.03	0.67± 0.03	0.57 ± 0.03	0.47 ± 0.03	0.53± 0.07	0.3±0.1
AST(μl)	41.67±1.67	37.33±0.67	34.67±2.33	35.33±0.67	35.0±2.0	33.67±2.33
ALT(μl)	30.33±1.33	27.67±0.67	26.33±0.67	26.0 ±2.0	26.33±2.67	24.0±1.0
ALP(μl)	89.0 ± 1.0	94.3 ± 3.67	99.0 ± 3.0	81.67±0.67	91.0±3.0	91.33±4.67
BUN (mg/dl)	16.07±0.27	15.23±0.37	15.03±0.47	15.03±0.57	15.93±0.07	16.27±0.03
Creatinine(mg/dl)	0.53 ± 0.07	0.43 ± 0.07	0.47 ± 0.03	0.4 ± 0.1	0.5±0.1	0.47±0.03

Data are expressed as Mean ± SEM (n=3). Comparisons were made using one-way ANOVA followed by post-hoc Newman-Keuls test. \*P< 0.05 when compared with control.

**Key:**

A-G Ratio = Albumin-Globulin Ratio

BUN = Bilirubin

AST = Aspartate transaminase

ALT = Alanine transferase, ALP = Alkaline phosphatase

No \*P<0.05 in table???

UNDER PEER REVIEW

## DISCUSSIONS

The test organisms are susceptible to Ofloxacin, Gentamicin, ciprofloxacin and Nitrofurantoin but are resistant to Augmentin, Cefotaxime and Ceftazidime. If these organisms are resistant to second and third generation Cephalosporin, there is a need for alternative antibiotics for treatment (Gashe *et al.*, 2018). The dichloromethane and ethylacetate fractions of ethanolic crude extract of *Funtumia elastica* exhibited different degrees of activities while the aqueous and *n*-hexane fractions showed no activity on the pathogenic organisms. This may be due to the fact that these mid-polar solvents have greater capacities to pull the bioactive compounds from the crude extract during partitioning. This finding is in agreement with that reported by Adekunle and Ikumapayi (2006), who found the extremely polar aqueous fractions of the same plant non-active against some fungal species. The minimum inhibitory concentrations (MIC) of the four (4) partitioned crude extracts (dichloromethane leaves, dichloromethane bark, ethylacetate leaves and ethylacetate bark) against the test organisms ranges between <3.125 mg/ml to 25 mg/ml. Though Osei-Djarbeng *et al.*, (2010) reported MIC ranges of 0.5 mg/mL to 2 mg/mL for the Ethyl-acetate fraction of extract of the same plant against related bacterial strains, the small differences observed could be attributed to strain susceptibility variation amongst the pathogens. The extracts were found to be most effective against *S. aureus* ATCC 29218, *S. aureus* ATCC 6571 and *P. aeruginosa* ATCC 27853 with the MIC range of <3.125 to 6.25 mg/ml but were found to be least effective against *E. coli* ATCC 35218, *P. aeruginosa* (strain P2) and *P. aeruginosa* with the MIC range of 12.5 mg/ml to 25mg/ml. It is interesting to note that *S. aureus* ATCC 29213 was the most susceptible to all the fractions with the MIC value of <3.125 mg/ml except for ethylacetate leaf extract where it had MIC value of 6.25 mg/ml. Again this pattern of susceptibility which was replicated in the sub-fraction of the Dichloromethane fraction, agrees

with Osei-Djarbeng *et al.* (2010), who reported that *S. aureus* NCTC 7447, showed the highest susceptibility in all the fractions of the extracts tested in their study. This could be as a result of the mechanism of action of the bioactive compound(s) in this fraction or the ability of the pathogen to take up sufficient bioactive concentration of the compound(s) into its system. *Pseudomonas aeruginosa* and *Escherichia coli* did not show any significant susceptibility to the ethylacetate leave and bark even at the highest concentration tested. This might not be unconnected to the absence of the bioactive compound(s) present in other fractions of the extracts; or the organism ability to resist the mechanisms of action of these extracts. This is however not surprising since *P. aeruginosa* is notorious for resisting antimicrobial agents (Agyare *et al.*, 2013). The antibacterial susceptibility testing of different fractions of the dichloromethane leaf revealed activity at fraction 2, with *S. aureus* being the most susceptible as pointed out earlier.

A significant increase was seen in the packed cell volume (PCV) and hemoglobin concentrations as well as red blood cell and white blood cell count. This indicates a potential hematopoietic potential of the dichloromethane extract with varying increases in the doses. This goes along to support a study (Akah *et al.*, 2009) which found out those medical and / or medicinal compounds and drugs have been shown to alter physiological range of hematological parameters. Platelet count was seen to reduce as the dose of the extract increases, reflecting a possible dose-dependent effect on the bleeding clotting system (Agyare *et al.*, 2013).

From this study, various biochemical parameters were assessed in validating the toxicity of *Funtumia elastica* in various systems of the body, using animal models: Alkaline phosphate (ALP), Alanine amino transferase (ALT), Aspartate aminotransferase (AAT), Bilirubin, Albumin and Globulin activities were measured in the serum to evaluate potential pathological effects of

*Funtumia elastica* on the liver. However, no alterations were seen in the serum ALT, AST, and ALP of various doses of the extract after 28 days administration. These enzymes are used to access possible damage to the liver (Fahmy *et al.*, 2014). Findings from this study indicates that the extract may not have any negative impact on the functionality of the liver with regards to metabolism. Serum albumin, bilirubin and globulin showed no significant changes in all doses. This indicates that the extract may not have any effect on the state of the liver (Fahmy *et al.*, 2014).

## **CONCLUSION**

The results obtained from the antimicrobial activities of *Funtumia elastica* leaves and bark show that the plant extracts have antibacterial properties and that can yield effective compound for the formation of a new and novel antibiotics, since they inhibited the growth of the tested pathogenic organisms. Further work is ongoing to isolate and characterize the actual bioactive compounds.

## REFERENCCEES

Adekunle, A. A and Ikumapayi, A. M. 2006. Antifungal property and phytochemical screening of the crude extracts of *Funtumia elastica* and *Mallotus oppositifolius*. *West Indian Med J*, 55, 219–223. should be written as follows:

Adekunle AA, Ikumapayi AM. Antifungal property and phytochemical screening of the crude extracts of *Funtumia elastica* and *Mallotus oppositifolius*. *West Indian Med J*. 2006; 55: 219–223.

Author must retake this whole part according to the authors guide

Adekunle, A. A., and Ikumapayi, A. M. (2006). Antifungal property and phytochemical screening of the crude extracts of *Funtumia elastica* and *Mallotus oppositifolius*. *West Indian Medical Journal*, 55(4), 219.

Adeniyi, B. A., Fong, H. H., Pezzuto, J. M., Luyengi, L and Odelola, H. A. 2004. Antibacterial activity of diospyrin, isodiospyrin, and bisisodiospyrin from *Diospyros piscatorial* (Gurke) [Ebenaceae]. *Phytotherapy Research*, 14:112-117.

Adeniyi, B. A., Odelola, H. A and Oso, B. A. 1996. Antimicrobial potentials of *Diospyros mespiliformis* (Ebenaceae). *African Journal of medicine and Medical Sciences*. 25: 2221-224

Agyare, A. F., Asase, M. I., Lechtenberg, M. S., Niehues, A. Z., Deters, I. G and Hensel, A. 2009 “An ethnopharmacological survey and in vitro confirmation of ethnopharmacological use of medicinal plants used for wound healing in Bosomtwi-Atwima-Kwanwoma area, Ghana,” *Journal of Ethnopharmacology*, vol. 125, no. 3, pp. 393–403

- Agyare, C., Koffuor, G. A., Boakye, Y. D., & Mensah, K. B. (2013). Antimicrobial and anti-inflammatory properties of *Funtumia elastica*. *Pharmaceutical biology*, 51(4), 418-425.
- Ahn, K. B. 2017. "The worldwide trend of using botanical drugs and strategies for developing global drugs". *BMB Report* 50 (3): 111-116.
- Ajani, A. A and Ayoola, S. P. 2014. Acute toxicity of Piscicidal plant extracts (*Adeniaciss ampeloides*) on tilapia (*Sarothero dongalilaeus*) juveniles. *Iranian journal of Energy and Environment*, 1(3): 246-254.
- Akah, P. A., Almeji, J. A., Salawu, O. A., Okoye, T. C and Offiah, N. V. 2009. Effects of *Vernonia amygdalina* on biochemical and hematological parameters in diabetic rat. *Asian Journal of Medical Sciences*, 1: 108-113
- Akpuaka, M. U. and Ezem, S. N. 2011. Preliminary Phytochemical Screening of some Nigerian Dermatological plants. *Journal of Basic Physical Research* 17:165-171
- Aniagu, S. O., Nwinyi, F. C., Akumka, D. D and Ajoku, G. A. 2005. Toxicity studies in rats fed nature cure bitters. *Afr. J. Biotechnol.* 4(1): 72-78.
- Burkill, H. M. 1995. The Useful Plants of West Tropical Africa, 2nd edition, Vol 3:11. London: *Royal Kew Botanical Gardens*, 522-527.
- Fahmy, S. A., Amel, M. S and Enas, M. A. 2014. Antifungal and antihepatotoxic effects of Sepia ink extract against oxidative stress as a risk factor of invasive pulmonary Aspergillois in neutopenic mice. *Afr J Tradit Complement AlternMed.* 11(3):148-159

Gashe, F., Mulisa, E., Mekonnen, M., & Zeleke, G. (2018). Antimicrobial resistance profile of different clinical isolates against third-generation cephalosporins. *Journal of pharmaceuticals*, 2018.

Harbone, J. B .1973. *Phytochemical Methods* 3<sup>rd</sup> ed. Chapman and Hall Ltd.135-203

Lorke, D. 1983. A new approach to practical acute toxicity testing. *Archives of Toxicology*. 53:275-287.

Okeke, I. N., Laxminarayan, R., Bhutta, Z, A 2005. Antimicrobial resistance in developing countries. Part 1: recent trends and current status. *The Lancet Inf. Dis.* 5:8; 481493.

Osei-Djarbeng, S. N., Cutler, S. J., & Cutler, R. R. (2010). of *Funtumia elastica* (Preuss) Stapf (Apocynaceae) Leaf Extracts. *Methodology*.

UNDER PEER REVIEW