

Original Research Article

Determinants of institutional Delivery in Bangladesh: A Multilevel Analysis of a Nationwide Population-Based Survey

ABSTRACT

Background: A substantial number of mothers and infants lose their lives due to complications during childbirth. Institutional delivery can lower the number of fatalities. This study aims to explore the prevalence of institutional delivery and its multifaceted associated factors in Bangladesh.

Methods: The data were extracted from Bangladesh Demographic and Health Survey (BDHS), conducted during the period of 2017-18. Following data preprocessing, a total of 4974 women were included in this study. This study attempts to identify and assess the variables (at the individual and community levels) having the significant impact on the institutional delivery by using a multilevel binary logistic regression model.

Results: Approximately half of the Bangladeshi women's deliveries were done at the medical center. The study found that women aged 20 or older had a 1.53 times greater likelihood of using healthcare facilities during childbirth (OR 1.53, [1.31-1.78]) than those under 20. Women with higher levels of education were more than twice as likely to choose institutional delivery as those with no education, and the women whose husbands had higher education had the same likelihood. The likelihood of using an institutional delivery service was also boosted by women's higher socioeconomic status. Women with exposure to any form of media had a 36% (OR 1.36, [1.16-1.60]) higher likelihood of accessing healthcare facilities during childbirth. Rural women had a 20% lower likelihood of receiving institutional delivery. The utilization of institutional delivery was positively correlated with religion, geographical region, child ever born, and antenatal care visits (ANC). This study discovered significant variations at the community level.

Conclusion: In order to increase the prevalence of institutional delivery in Bangladesh, the study findings suggest that policymakers should focus on addressing socioeconomic and demographic factors, particularly on women's education, husbands' education, age at first delivery and frequency of ANC visits.

KEYWORDS: Institutional delivery, BDHS, ANC, Socioeconomic and Demographic factors, Community level.

1. INTRODUCTION

According to the World Health Organization (WHO), maternal mortality is the death of a pregnant woman as a result of pregnancy-related issues, preexisting conditions that are aggravated by pregnancy, or the medical management of these conditions [1]. The global maternal mortality ratio (MMR) decreased by 34% between 2000 and 2020, from 342

maternal deaths per 100,000 live births to 223 deaths per 100,000 live births [2]. In 2020, a woman died every two minutes from pregnancy- or childbirth-related complications. Nearly 95 percent of all maternal deaths took place in low and lower-middle-income countries. The majority of these fatalities were avoidable. Sub-Saharan Africa and Southern Asia accounted for over 87% (253 000) of the expected global maternal deaths in 2020 [2].

Maternal death was most frequently caused by high blood pressure during pregnancy, unsafe abortions, severe bleeding, and infections following delivery [3]. A third of all antepartum causes, such as intrapartum and postpartum hemorrhage, were linked to risky home delivery methods [4-5]. Institutional delivery is an essential strategy for lowering maternal mortality. Deliveries that are aided in a hospital or other setting and carried out under the supervision of qualified healthcare professionals can reduce delivery problems and maternal fatalities [6–9].

Although there has been a noticeable improvement in South Asia over the past 20 years' in relation to delivery in hospitals, the rate is still relatively low when compared to other regions. The main barriers that prevent women from seeking healthcare during childbirth are inadequate healthcare services, financial limitations, physical distance, cultural norms, and a lack of information [10–11]. Home births were more likely to occur in unsafe and unclean conditions for South Asian women. It posed a major risk to the health of both the mothers and their newborns [12].

Institutional delivery is not a common practice in Bangladesh, similar to many other low- and middle-income countries (LMICs) in the South Asian region. In Bangladesh, deliveries typically occurred at home, and around half of live births occurred outside of hospitals [13]. One-third of the women in a research conducted in rural Bangladesh claimed to have experienced difficulties at the birth of their most recent child [14].

A number of studies used fixed-effect models, such as binary logistic regression, conditional logistic regression, and multivariate statistical analyses, to determine the variables affecting institutional delivery [15–20]. The majority of the studies focused mostly on individual-level factors. But community variables can have a significant impact on healthcare. A few studies in South Asia and Africa considered both individual and community-level factors. However, this study used a multilevel binary logistic regression model with a random intercept to examine the factors associated with institutional delivery in Bangladesh, taking into account individual and community-level variables. It also determined the prevalence of institutional delivery among women aged 15 to 49 in Bangladesh as well as any potential variations at the community level related to delivery outcomes.

2. MATERIAL AND METHODS

2.1 DATA SOURCE

The data is obtained from the eighth iteration of the Bangladesh Demographic and Health Survey (BDHS), which was gathered through the DHS program [21]. The BDHS survey is a nationwide, cross-sectional study that took place between October 2017 and March 2018, aiming to provide a representative snapshot of the population's characteristics during that period. The BDHS was carried out by a local research institute, the National Institute of Population Research and Training (NIPORT), in collaboration with the Medical Education and Family Welfare Division of the Ministry of Health and Family Welfare.

2.2. SAMPLE DESIGN

The population living in non-institutional dwelling units in Bangladesh was sampled using a two-stage cluster sampling technique. Bangladesh is divided into eight administrative divisions, and each of these divisions is subsequently subdivided into zilas (districts) and upazilas (townships). At the urban level, an upazila is segmented into union parishads and mouzas. Conversely, at the rural level, it is divided into wards and mohallas, which are subdivisions of the wards. In the first stage, 675 enumeration areas (250 in urban areas and 425 in rural areas) were selected with probability proportional to EA (enumeration area) size. In the second stage of sampling, a systematic sample of an average of 30 households per EA was selected to provide statistically reliable estimates of key demographic and health variables for the country as a whole, for urban and rural areas separately, and for each of the eight divisions. A total of 20,160 households were selected for the survey. The information on the child's prenatal and postpartum treatment was extracted from the Kids Record (KR) data file. The cohort's initial size was 8759. All the missing values were removed. Finally, a weighted sample of 4974 respondents was considered for data analysis.

2.3. DEPENDENT VARIABLE

The study's binary outcome variable was "institutional delivery," which was divided into two categories: "Yes" and "No." If a delivery occurred in a hospital, whether it was a public or private facility, it was labeled as a "facility-based" delivery; otherwise, it fell into the "No" category.

2.4. INDEPENDENT VARIABLES

This study took into account both individual level and community level factors as explanatory variables. The individual level factors were: age at first birth, women's education, husband's education, 4+ ANC visits, health care decision making, child ever born and exposure to mass media. On the other hand, the community level factors were: wealth index, religion, place of residence and administrative division.

2.5. STATISTICAL ANALYSIS

The data from the BDHS 2017-18 for this study were cleaned, coded, and analyzed using the statistical tools SPSS version 25 (IBM Corporation, Armonk, NY, USA). Descriptive analysis was employed in this study to obtain a broad comprehension of the sample's traits and attributes. The chi-square test of independence was used to examine the association between the selected covariates and institutional delivery. The multilevel binary logistic regression model contained components from both the individual and community levels. Individual-level determinants (respondents) were nested within the communities (clusters). We calculated the odds ratios, *P* values, and 95% confidence intervals for each factor's impact on institutional delivery. Intra cluster Correlation (ICC) was used to express random effects to assess the fitness of the model. The model was constructed by considering the binary response variable Y_{ij} , which represents "Institutional Delivery" (1 if women i in community j receive institutional delivery services and 0 otherwise) [22, 23]. The two-level random intercept binary logistic regression model, which accounts for women at level 1 and communities (clusters) at level 2, can be expressed as follows:

$$\text{logit}(\pi_{ij}) = \log\left(\frac{\pi_{ij}}{1 - \pi_{ij}}\right) = \beta_{0j} + \sum_{k=1}^m \beta_k X_{ijk};$$

$$i = 1, 2, \dots, n_j, j = 1, 2, \dots, d$$

$$\text{With } \beta_{0j} = \beta_0 + \mu_{0j}; \mu_{0j} \sim \text{iid } N(0, \sigma_{\mu_0}^2)$$

Where, $\pi_{ij} = \Pr(Y_{ij}=1)$ is the probability that the woman i in community j takes institutional delivery services, X_{ijk} is the values of m explanatory variables for women i in community j , β_k is a vector of regression coefficients to be estimated, and β_0 is a fixed component. μ_{0j} is the random error at the community level.

3. RESULTS AND DISCUSSION

Table 1 provides insights into the sociodemographic characteristics of sampled women in Bangladesh. It indicates that the majority of the mothers delivered their children during their teenage years (under the age of 20). In terms of education, the largest proportion of women had completed secondary level of education (49%), followed by those with primary level of education (27.5%), and those with higher level of education (17.2%). Only a small proportion of women lacked any formal education (6.2%). Additionally, the majority of women's husbands had received primary or secondary levels of education, with each category accounting for one third of the respondents. **Table 1** also demonstrates that about two out five women were poor (41%) and almost the same proportion were rich (39.9%). Nearly two third of the women were exposed to mass media (65.8%) and nearly half of them received 4+ ANC visits (47.2%). Furthermore, most of the women were Muslim (91.8%), and a majority of them used to make health care decision self or jointly (72.8%). A majority of the women were residing in rural region (73.2%), and had two or more children (49.8%).

Table 1: Distribution of Bangladeshi women's sociodemographic characteristics

Variables	N (%)
Age at first birth	
<20 years	3501 (70.4)
≥ 20 years	1473 (29.6)
Women's education	
No education	310 (6.2)
Primary	1370 (27.5)
Secondary	2437 (49.0)
Higher	858 (17.2)
Husband's education	
No education	680 (13.7)
Primary	1678 (33.7)
Secondary	1696 (34.1)
Higher	921(18.5)
Wealth index	
Poor	2040 (41.0)
Middle	948 (19.1)
Rich	1985 (39.9)
Exposure to mass media	

Not exposed	1703 (34.2)
Exposed	3271 (65.8)
4+ ANC visit	
No	2628 (52.8)
Yes	2346 (47.2)
Religion	
Muslim	4568 (91.8)
Non-Muslim	406 (8.2)
Health care decision making	
Self or Jointly	3620 (72.8)
Others	1354 (27.2)
Child ever born	
1	1887 (37.9)
2-3	2477 (49.8)
4+	611 (12.3)
Place of residence	
Urban	1333 (26.8)
Rural	3641 (73.2)
Administrative division	
Barisal	282 (5.7)
Chattogram	1054 (21.2)
Dhaka	1275 (25.6)
Khulna	453 (9.1)
Mymensingh	424 (8.5)
Rajshahi	579 (11.6)
Rangpur	528 (10.6)
Sylhet	378 (7.6)
Institutional delivery	
Yes	2471 (49.7)
No	2503 (50.3)

The relationships between sociodemographic factors and the utilization of a healthcare facility during childbirth are depicted in **Table 2**. In this study, all covariates were significantly associated with institutional delivery except healthcare decision-making. Only significant variables were included in the multi-level model. The proportion of women who opted for

healthcare facility delivery was observed to be notably higher among the following demographic categories: older women (aged 20 years or older), women with higher levels of education, husbands with higher levels of education, women from wealthier backgrounds, those with exposure to mass media, women who attended more than four antenatal care (ANC) visits and individuals hailing from urban areas.

Table 2: Association between Bangladeshi women's socioeconomic characteristics and Institutional delivery

Variables	Institutional delivery		χ^2 value (P-value)
	Yes (%)	No (%)	
Age at first birth			
<20 years	43.9	56.1	197.16 **
≥ 20 years	65.7	34.3	(<.001)
Women's education			
No education	26.2	73.8	
Primary	32.8	67.2	525.10**
Secondary	53.3	46.7	(<.001)
Higher	78.7	21.3	
Husband's education			
No education	29.6	70.4	
Primary	38.3	61.7	515.27**
Secondary	55.5	44.5	(<.001)
Higher	78.0	22.0	
Wealth index			
Poor	31.8	68.2	
Middle	50.1	49.9	569.86**
Rich	69.5	30.5	(<.001)
Exposure to mass media			
Not exposed	32.5	67.5	327.88**
Exposed	59.6	40.4	(<.001)
4+ ANC visit			
Yes	66.1	33.9	442.41**
No	36.2	63.8	(<.001)
Religion			
Muslim	49.3	50.7	23.35**
			(<.001)

Non-Muslim	61.8	38.2	
Health care decision making			
Self or Jointly	50.9	49.1	1.52
Others	48.9	51.1	(.218)
Child ever born			
1	61.4	38.6	
2-3	47.5	52.5	226.52**
4+	27.5	72.5	(< .001)
Place of residence			
Urban	64.2	35.8	140.22**
Rural	45.3	54.7	(< .001)
Administrative division			
Barisal	39.4	60.6	
Chattogram	46.9	53.1	
Dhaka	58.3	41.7	114.93**
Khulna	60.9	39.1	(< .001)
Mymensingh	38.9	61.1	
Rajshahi	53.4	46.6	
Rangpur	48.6	51.4	
Sylhet	39.2	60.8	
$P < .05$, * $P < .001$			

Table 3 presents the factors influencing institutional delivery among women in Bangladesh. Women aged 20 and older were nearly 1.53 times more likely to access healthcare services during childbirth (OR 1.53, [1.31-1.78]) compared to younger women (under the age of 20). As the level of education of women increased, the likelihood of utilizing healthcare facilities for delivery also increased. Women with higher levels of education were almost twice as likely to opt for institutional delivery (OR 2.06, [1.40-3.02]) than uneducated women. Similarly, women with husbands who had a higher education had more than twice the chance of giving birth in a hospital. The socio-economic status of women had significant effects on the use of institutional delivery. Wealthy women had a 1.9-fold higher likelihood of it than economically disadvantaged women. The likelihood of giving birth in a health facility among middle-class women was 26%.

Delivery in a medical facility was significantly impacted by the mass media. Access to mass media increased a woman's likelihood of choosing institutional delivery by 36% compared to her counterpart (OR 1.36, [1.16-1.60]). With an increase in ANC visits, there was a higher chance of giving birth in a hospital. Non-Muslim women had a 1.31 times greater likelihood of giving birth in a medical facility than Muslim women. As a mother had more children, the likelihood of choosing an institutional birth decreased. Compared to urban women, rural women were 20% less likely to give birth in a hospital. Dhaka, Khulna and Rajshahi, three

divisions out of seven, were statistically significant. The Khulna divisions had the highest likelihood of delivering in a medical setting (OR 1.78, [1.16-2.74]).

Table 3: Multilevel binary logistic regression analysis of individual and community-level factors associated with institutional delivery services among women in Bangladesh.

Variables	Adjusted Odds Ratio (95% CI)	P-value
Age at first birth		
<20 years	Ref.	
≥ 20 years	1.53*** (1.31, 1.78)	< .001
Women's education		
No education	Ref.	
Primary	1.12 (0.81, 1.54)	.492
Secondary	1.50** (1.09, 2.07)	.012
Higher	2.06*** (1.40, 3.02)	< .001
Husband's education		
No education	Ref.	
Primary	1.06 (0.84, 1.33)	.627
Secondary	1.40** (1.11, 1.79)	.005
Higher	2.30*** (1.70, 3.10)	< .001
Wealth index		
Poor	Ref.	
Middle	1.26** (1.04, 1.53)	.018
Rich	1.90*** (1.55, 2.32)	< .001
Exposure to mass media		
Not exposed	Ref.	
Exposed	1.36*** (1.16, 1.60)	< .001
4+ ANC visit		
No	Ref.	
Yes	2.03*** (1.76, 2.33)	< .001
Religion		
Muslim	Ref.	
Non-Muslim	1.31* (0.99, 1.75)	.062

Child ever born		
1	Ref.	
2-3	0.69*** (0.60, 0.78)	< .001
4+	0.52*** (0.41, 0.80)	< .001
Place of residence		
Urban	Ref.	
Rural	0.80* (0.63, 1.02)	.067
Administrative division		
Barisal	Ref.	
Chattogram	1.13 (0.76, 1.68)	.552
Dhaka	1.45* (0.97, 2.17)	.072
Khulna	1.78*** (1.16, 2.74)	< .001
Mymensingh	0.95 (0.62, 1.46)	.819
Rajshahi	1.62** (1.07, 2.47)	.024
Rangpur	1.43 (0.92, 2.20)	.110
Sylhet	1.14 (0.73, 1.77)	.576
Random-effect variance		
Community	0.482***	< .001
ICC	0.13	

* $P < .1$, ** $P < .05$, *** $P < .001$

The study found significant community variations. The estimated intra-cluster correlation coefficient (ICC) for the model is 0.13, indicating that around 13% of the variation remained among community levels. Therefore, using multilevel modeling with these data is justified. This suggests that community-level factors play a notable role in explaining the differences observed in the utilization of institutional delivery services among mothers.

4. DISCUSSION

This study focused on investigating the prevalence of institutional delivery and the factors that significantly contributed to it in Bangladesh. According to the findings of this study, there was a higher rate of health facility delivery among women with higher levels of education. These highly educated women were more likely to be informed and knowledgeable about health-related issues. This finding is consistent with previous studies [20, 24-27]. Earlier research had also indicated that mothers with higher economic status were more inclined to choose hospital deliveries [28, 29]. This study yielded similar results. The poor's inability to pay for such services is one of the main barriers to the provision of health care. Rural women had a lower probability of delivering in a hospital compared to their urban counterparts. This discovery is consistent with the findings of a prior study conducted in Bangladesh [30]. Additionally, this study illustrated that the utilization of healthcare services during delivery was notably influenced by exposure to media. These findings were corroborated by earlier research [30, 31]. However, conflicting findings were found in another

study [24]. This study also found that women who had more than four ANC visits were more likely to deliver their babies in a hospital. Similar results were discovered in other studies [32, 33]. Non-Muslim women had a higher chance of delivering in a medical facility than non-Muslim women. This finding is consistent with prior studies [5, 13, 37]. Previous research indicated that women with low parity may be more motivated to choose health facility deliveries [24, 36,37]. This was in line with the study's findings. This trend could be attributed to the fact that first deliveries are often given more serious attention by family members. This study discovered an association between a woman's age at her first childbirth and her likelihood of delivering in a medical facility, which aligns with a similar finding from a study conducted in sub-Saharan Africa [4]. Women whose husbands have a high level of education were more inclined to give birth in a healthcare facility compared to those whose husbands had lower educational attainment. This pattern suggests that husbands with higher education levels tend to be more concerned about the health and well-being of their wives and children, which may lead to a higher likelihood of ensuring institutional delivery for their families. A comparable outcome was observed in a prior study conducted in Bhutan [36]. Geographical region was positively correlated with institutional delivery. According to this study, women who were from Dhaka, Khulna, and Rajshahi divisions had a higher likelihood of receiving institutional delivery. This finding is supported by the findings of earlier studies [13,18, 38]

The study faced a notable limitation in its reliance on BDHS data, primarily due to the frequently limited sample sizes available at the country level. Another significant constraint is that our research only included deliveries resulting in live births because BDHS data lacks information on pregnancy outcomes other than live births and does not specify their delivery locations, which introduces a level of complexity and potential bias in our findings.

5. CONCLUSION

A primary challenge impeding the achievement of Sustainable Development Goal 3 (SDG 3) in numerous developing countries is the limited access to healthcare facilities. The current study's findings indicated that several factors, including higher levels of education among women, favorable socio-economic conditions, exposure to media, age at first childbirth, and the educational background of husbands, were all influential factors contributing to a higher likelihood of giving birth in a healthcare facility. In Bangladesh, substantial barriers persist due to a lack of awareness regarding the importance of seeking delivery care from trained healthcare providers. Raising public awareness could be significantly enhanced through mass media, and policymakers should consider initiatives to increase the frequency and reach of media exposure. Low parity influences the utilization of healthcare facility delivery. The emphasis on low-parity women should be strengthened even though research and policy are trending in that direction. Improving access to healthcare facility delivery, especially in rural areas, is of utmost importance. The insufficient utilization of healthcare facilities during delivery was largely linked to women's socioeconomic status and educational level. As a result, when shaping health policies and strategies, the government of Bangladesh should place heightened attention on addressing these two key factors.

REFERENCES

1. World Health Organization (2023). Maternal Mortality. [Online] Who.int. Available at: <https://www.who.int/data/gho/indicator-metadata-registry/imr-details/4622> [Accessed 09 Sep. 2023].
2. World Health Organization (2023). Maternal Mortality. [Online] Who.int. Available at: <https://www.who.int/news-room/fact-sheets/detail/maternal-mortality> [Accessed 09 Sep. 2023].
3. Apanga, P.A. and Awoonor-Williams, J.K. (2018). Maternal Death in Rural Ghana: a Case Study in the Upper East Region of Ghana. *Frontiers in Public Health*, 6. doi:<https://doi.org/10.3389/fpubh.2018.00101>.
4. Benova, L., Macleod, D., Radovich, E., Lynch, C.A. and Campbell, (2017). Should I stay or should I go: consistency and switching of delivery locations among new mothers in 39 Sub-Saharan African and South/Southeast Asian countries. *Health Policy and Planning*, [online] 32, pp.1294–1308. doi:<https://doi.org/10.1093/heapol/czx087>.
5. Devkota, B., Maskey, J., Pandey, A.R., Karki, D., Godwin, P., Gartoulla, P., Mehata, S. and Aryal, K.K. (2020). Determinants of home delivery in Nepal – A disaggregated analysis of marginalised and non-marginalised women from the 2016 Nepal Demographic and Health Survey. *PLOS ONE*, 15(1), p. e0228440. doi:<https://doi.org/10.1371/journal.pone.0228440>.
6. Dahal RK. Central Department of Population Studies, Tribhuvan University, Kathmandu, Nepal. Factors influencing the choice of place of delivery among women in Eastern rural Nepal. *International Journal of Maternal and Child Health* 2013; 1:30–7.
7. Assarag B, Dujardin B, Delamou A, *et al.* Determinants of maternal near-miss in Morocco: too late, too far, too sloppy? *PLoS One* 2015;10: e0116675.
8. De Brouwere V, Richard F, Witter S. Access to maternal and perinatal health services: lessons from successful and less successful examples of improving access to safe delivery and care of the newborn. *Trop Med Int Health* 2010; 15:901–9.
9. Karkee R, Binns CW, Lee AH. Determinants of facility delivery after implementation of safer mother programme in Nepal: a prospective cohort study. *BMC Pregnancy Childbirth* 2013; 13:193.
10. Sharma, S.R., Poudyal, A.K., Devkota, B.M. and Singh, S. (2014). Factors associated with place of delivery in rural Nepal. *BMC Public Health*, 14(1). doi:<https://doi.org/10.1186/1471-2458-14-306>.
11. Das, S., Alcock, G., Azad, K., Kuddus, A., Manandhar, D.S., Shrestha, B.P., Nair, N., Rath, S., More, N.S., Saville, N., Houweling, T.A.J. and Osrin, D. (2016). Institutional delivery in public and private sectors in South Asia: a comparative analysis of prospective data from four demographic surveillance sites. *BMC Pregnancy and Childbirth*, 16(1). doi:<https://doi.org/10.1186/s12884-016-1069-7>.
12. Prata, N., Bell, S. and Quaiyum, M.A. (2014). Modeling maternal mortality in Bangladesh: the role of misoprostol in postpartum hemorrhage prevention. *BMC Pregnancy and Childbirth*, [online] 14(1). doi:<https://doi.org/10.1186/1471-2393-14-78>.

13. Kabir, S., Hasan, M.R., Hossain, M.I., Suraiya, S., Islam, F.B., Nayan, M.I.H., Haq, I. and Hossain, M.S. (2022). Determinants and Trends of Health Facility Delivery in Bangladesh: A Hierarchical Modeling Approach. *BioMed Research International*, 2022, p.e1359572. doi:<https://doi.org/10.1155/2022/1359572>.
14. Kane, T.T., Barkat-e-Khuda and Phillips, J.F. (1997). *Reproductive Health in Rural Bangladesh, Policy and Programmatic implications/ Thomas T Kane, Barkat-e-Khuda, James F Phillips*. [online] dspace.icddrb.org. International Centre for Diarrhoeal Disease Research, Bangladesh. Available at: <http://dspace.icddrb.org/jspui/handle/123456789/2565> [Accessed 13 Sep. 2023].
15. Kamal, S. M. (2013). Preference for institutional delivery and caesarean sections in Bangladesh. *Journal of health, population, and nutrition*, 31(1), 96.
16. Yaya, S., Bishwajit, G., & Ekholuenetale, M. (2017). Factors associated with the utilization of institutional delivery services in Bangladesh. *PloS one*, 12(2), e0171573.
17. Kamal, S. M., Hassan, C. H., & Alam, G. M. (2015). Determinants of institutional delivery among women in Bangladesh. *Asia Pacific Journal of Public Health*, 27(2), NP1372-NP1388.
18. Talukder, A., Anik, B. H., Hossain, M. I., Haq, I., & Habib, M. J. (2022). Socioeconomic and demographic factors for mothers' delivery at home: a comparative study among BDHS 2007, 2011 and 2014. *Asian Journal of Social Health and Behavior*, 5(1), 10.
19. C. Z. Olorunsaiye, L. B. Huber, S. B. Laditka, S. Kulkarni, and A. S. Boyd, "Factors associated with health facility delivery in west and central Africa: a multilevel analysis of seven countries," *Health Care for Women International*, vol. 41, pp. 3– 21, 2020.
20. Khan, J.R., Ara, T., Rahman, Md.M., Hossain, Md.B. and Muurlink, O. (2022). A multilevel assessment of the influence of education on women's uptake of institutional delivery services during childbirth in Bangladesh. *Midwifery*, 113, p.103425. doi:<https://doi.org/10.1016/j.midw.2022.103425>.
21. National Institute of Population Research and Training - NIPORT, Ministry of Health and Family Welfare, and ICF. 2020. Bangladesh Demographic and Health Survey 2017-18. Dhaka, Bangladesh: NIPORT/ICF. Available at <https://www.dhsprogram.com/pubs/pdf/FR344/FR344.pdf>.
22. H. Goldstein, *Multilevel Models in Education and Social Research*, Oxford University Press, Oxford, UK, 1987.
23. P. C. Austin and J. Merlo, "Intermediate and advanced topics in multilevel logistic regression analysis," *Statistics in Medicine*, vol. 36, no. 20, pp. 3257–3277, 2017.
24. Agha, S. and Carton, T.W. (2011). Determinants of institutional delivery in rural Jhang, Pakistan. *International Journal for Equity in Health*, 10(1), p.31. doi:<https://doi.org/10.1186/1475-9276-10-31>.
25. Chaka, E.E., Parsaeian, M. and Majdzadeh, R. (2019). Factors Associated with the Completion of the Continuum of Care for Maternal, Newborn, and Child Health Services in Ethiopia. Multilevel Model Analysis. *International Journal of Preventive Medicine*, [online] 10, p.136. doi:<https://doi.org/10.4103/ijpvm.IJPVM.26.19>.

26. Shibanuma, A., Yeji, F., Okawa, S., Mahama, E., Kikuchi, K., Narh, C., Enuameh, Y., Nanishi, K., Oduro, A., Owusu-Agyei, S., Gyapong, M., Asare, G.Q., Yasuoka, J., Ansah, E.K., Hodgson, A. and Jimba, M. (2018). The coverage of continuum of care in maternal, newborn and child health: a cross-sectional study of woman-child pairs in Ghana. *BMJ Global Health*, 3(4), p. e000786. doi:<https://doi.org/10.1136/bmjgh-2018-000786>.
27. Mitikie, K.A., Wassie, G.T. and Beyene, M.B. (2020). Institutional delivery services utilization and associated factors among mothers who gave birth in the last year in Mandura district, Northwest Ethiopia. *PLOS ONE*, 15(12), p.e0243466. doi:<https://doi.org/10.1371/journal.pone.0243466>.
28. Yaya, S., Bishwajit, G. and Gunawardena, N. (2019). Socioeconomic factors associated with choice of delivery place among mothers: a population-based cross-sectional study in Guinea-Bissau. *BMJ Global Health*, 4(2), p. e001341. doi:<https://doi.org/10.1136/bmjgh-2018-001341>.
29. Sarker, B.K., Rahman, M., Rahman, T., Hossain, J., Reichenbach, L. and Mitra, D.K. (2016). Reasons for Preference of Home Delivery with Traditional Birth Attendants (TBAs) in Rural Bangladesh: A Qualitative Exploration. *PLoS ONE*, [online] 11(1). doi:<https://doi.org/10.1371/journal.pone.0146161>.
30. Talukder, A., Anik, B., Hossain, M., Haq, I. and Habib, M. (2022). Socioeconomic and demographic factors for mothers' delivery at home: A comparative study among BDHS 2007, 2011 and 2014. *Asian Journal of Social Health and Behavior*, 5(1), p.10. doi:https://doi.org/10.4103/shb.shb_160_21.
31. Pandey, S. and Karki, S. (2014). Socio-economic and Demographic Determinants of Antenatal Care Services Utilization in Central Nepal. *International journal of MCH and AIDS*, [online] 2(2), pp.212–219. Available at: <https://pubmed.ncbi.nlm.nih.gov/27621975/> [Accessed 21 Apr. 2023].
32. Okonofua, F., Ogu, R., Agholor, K., Okike, O., Abdus-salam, R., Gana, M., Randawa, A., Abe, E., Durodola, A. and Galadanci, H. (2017). Qualitative assessment of women's satisfaction with maternal health care in referral hospitals in Nigeria. *Reproductive Health*, [online] 14(1). doi:<https://doi.org/10.1186/s12978-017-0305-6>.
33. Gebeyehu Workineh, Y. (2014). Factors Affecting Utilization of Postnatal Care Service in Amhara Region, Jabitena District, Ethiopia. *Science Journal of Public Health*, 2(3), p.169. doi:<https://doi.org/10.11648/j.sjph.20140203.15>.
34. Berhan, Y. and Berhan, A. (2014). A Meta-Analysis of Socio-Demographic Factors Predicting Birth in Health Facility. *Ethiopian Journal of Health Sciences*, 24(0), p.81. doi:<https://doi.org/10.4314/ejhs.v24i0.8s>.
35. Kebede, A., Hassen, K. and Teklehaymanot, A.N. (2016). Factors associated with institutional delivery service utilization in Ethiopia. *International Journal of Women's Health*, [online] 8, pp.463–475. doi:<https://doi.org/10.2147/IJWH.S109498>.
36. Wangdi, K., Gurung, M.S., Pelzom, D., Dema, T. and Wangdi, S. (2015). Preference of place of delivery and its determinants: a cross-sectional study in Chukha district, Bhutan. *Bhutan Health Journal*, 1(1), pp.23–29. doi:<https://doi.org/10.47811/bhj.05>.

37. J. K. Ganle, "Why Muslim women in northern Ghana do not use skilled maternal healthcare services at health facilities: a qualitative study," *BMC International Health and Human Rights*, vol. 15, no. 1, article 48, p. 10, 2015.

38. G. M. A. Kibria, S. Ghosh, S. Hossen, R. A. A. Barsha, A. Sharmeen, and S. M. I. Uddin, "Factors affecting deliveries attended by skilled birth attendants in Bangladesh," *Maternal Health, Neonatology and Perinatology.*, vol. 3, no. 1, article 46, p. 7, 2017.

UNDER PEER REVIEW