

***Original Research Article***  
**ESTIMATION OF SURFACE WATER VAPOUR  
DENSITY AND ITS VARIATION WITH  
METEOROLOGICAL PARAMETERS IN BENIN  
NIGERIA**

**ABSTRACT**

The present study investigated the monthly variation of Surface Water Vapour Density (SWVD) with meteorological parameters of monthly average daily mean temperature, relative humidity, surface pressure, cloud cover and sunshine hours during for the period of thirty eight (38) years (extending from 1979 to 2016) for Benin (Latitude: 6.32<sup>o</sup>N, Longitude: 5.10<sup>o</sup>E, 77.8m above sea level) were investigated. The daily variation of surface water vapour density SWVD for the two distinct seasons considering two typical months in each during the period of year 2014 was examined. The comparison assessment of the developed two variable SWVD based models was carried out using statistical indices of coefficient of determination (R<sup>2</sup>), Mean Bias Error (MBE), Root Mean Square Error (RMSE), Mean Percentage Error (MPE), Nash – Sutcliffe Equation (NSE) and Index of Agreement (IA). The results showed fluctuation in the amount of surface water vapour density SWVD in each day of the month for the period under investigation. The monthly average daily values indicated that the surface water vapour density SWVD is higher during the rainy season than in the dry season. It was observed that the maximum average value of surface water vapour density SWVD of (21.1448 gm<sup>-3</sup>) occurred in the month of August during the rainy season and minimum value of SWVD (17.7134 gm<sup>-3</sup>) occurred in the month of December during the dry season. The highest value of surface water vapour density SWVD was observed on 23<sup>rd</sup> May, 2014 with 27.57313 gm<sup>-3</sup> and the lowest on 26<sup>th</sup> December, 2014 with 9.656676 gm<sup>-3</sup>. The developed multivariate correlation regression model that relates pressure and precipitable water vapour with R<sup>2</sup> = 100%, MBE = -0.0204, RMSE = 0.0206, MPE = 0.1105, NSE = 99.9897% and IA = 99.9974% was found more suitable for surface water vapour density SWVD estimation with good fitting and therefore can be used for estimating surface water vapour density SWVD in Benin.

**Comment [A1]:** Specify seasons and months

**Comment [A2]:** What is the basis of selection of this particular year?

**Comment [A3]:** Round off is needed. Applicable for each similar case.

**Comment [A4]:** Good or best?

*Keywords: Correlation models, dry Season, ECMWF, rainy season, SWVD*

**1. INTRODUCTION**

Water vapor plays a very important role in climate change, hydrological processes, Earth's energy balance, and weather systems [1 – 4]. Water vapor is the most common greenhouse gas in the atmosphere, and it accounts for about 60% of the natural greenhouse effect [1, 5]. At the surface, water vapor (or humidity) is an important meteorological and climate variable that affects human comfort [6], surface evaporation and plants' transpiration. Surface specific ( $q$ ) and relative humidity (RH) is conventionally measured using wet and dry bulb

thermometers or RH sensors exposed in thermometer screens at a large number of weather and climate stations and on many marine platforms [7].

Water is the fundamental of the hydrological cycle, which is inter or intra movement of water, in the Earth's atmosphere, oceans, and continents. Transfer of heat and energy between the earth's surface and the atmosphere and within the planet takes place as a result of hydrological cycle. The primary and most effective greenhouse gas in the atmosphere is water vapour, as it absorbs long wave radiation and radiates it back to the surface, which contributes to warming [2, 8 – 11]. The atmosphere holds more water vapour as the temperature of the Earth's surface and atmosphere increases [2]. In the troposphere, water vapour molecules absorbed heat energy radiated from the Earth's surface [12].

As the Earth's surface temperature increases, the atmosphere tends to hold more water vapour. This atmospheric water vapour, acts as a greenhouse gas thus absorbing energy that would otherwise cause reduction of electromagnetic radiation travelling through the atmosphere, the penalty of these could be atmospheric or global warming. The proportion by volume of water vapour in the air at the ground level on the average changes from less than 0.001% in the arctic to more than 6% in the tropics [13 – 14]. This proportion decreases speedily with height [13].

Understanding the processes which manage the natural stability and variability of the climate system is one of the most intricate and challenging scientific problems faced by the climate science community today. This is due to the fact that human activities such as emission of greenhouse gases and land use change which result in external forcing are only partly predictable [15 - 16]. This is so because scientists lack the ability to actually do so since they cannot foretell population change, economic change, technological development and other relevant characteristic of future human activities [16].

The purpose of this study was to investigate the daily and monthly variation of surface water vapour density (SWVD) and to examine the monthly variation with meteorological parameters of mean temperature, relative humidity, surface pressure, cloud cover and sunshine hours for Benin located in South Western, Nigeria. The study also developed two variable correlation models for estimating surface water vapour density SWVD for the location under investigation.

This present study evaluated and compared six evapotranspiration models for estimating reference evapotranspiration ( $ET_o$ ) in Kano, Nigeria, using FAO – 56 PM method as standard. The reason for this comparison is to find out which of the six evaluated models is the most appropriate to be considered as an alternative to FAO – 56 PM model for estimating  $ET_o$  in Kano based on the four adopted statistical indicators.

**Comment [A5]:** Repetition. Same statement was found in the first paragraph. Try to write in a more compact way.

## 2. METHODOLOGY

The daily and monthly meteorological data such as average minimum temperature, maximum temperature, relative humidity, surface pressure, cloud cover and sunshine hours meteorological data used in this study was obtained from the European Centre for Medium-Range Weather Forecasts (ECMWF) at 2 m height for Benin, Nigeria. The period under study is thirty eight (38) years (1979 – 2016).

The surface water vapour density (SWVD), vapour pressure ( $e$ ) and mean temperature ( $T$ ) are related by the following expression [1, 14, 16] as:

$$SWVD = 216.7 \left( \frac{e}{T} \right) \quad (1)$$

They also obtained vapour pressure ( $e$ ) using the expression given by

$$e = RH \left( \frac{e_s}{100} \right) \quad (2)$$

where RH and  $e_s$  are the relative humidity and saturated vapour pressure, respectively. The saturated vapour pressure was evaluated determined using the Clausius Clapeyron equation defined as:

**Comment [A6]:** Write full form.

**Comment [A7]:** Mention reference

$$\log_{10} e_s = 9.4051 - \left(\frac{2353}{T}\right) \quad (3)$$

The mean temperature  $T$ , ~~is was also~~ obtained using [17]

$$T = \frac{T_{max} + T_{min}}{2} \quad (4)$$

where  $T_{max}$  and  $T_{min}$  are the maximum and minimum temperatures respectively. The SWVD is in  $gm^{-3}$  and  $e$  and  $e_s$  are in millibars (mb),  $T$  in Kelvin (K) and RH in percentage (%).

A correlation model between ambient Temperature  $T$  (Kelvin) and Partial Pressure  $P_s$  as reported by [1] is given by the semi empirical equation

$$P_s = \exp\left(26.23 - \frac{5416}{T}\right) \quad (5)$$

[18] presented the following formula, which expresses precipitable water in terms of relative humidity:

$$W = \frac{(0.493\phi_r P_s)}{T} \quad (6)$$

where  $\phi_r$  is relative humidity in fractions of one,  $T$  is ambient temperature in degrees Kelvin and  $p_s$  is the partial pressure of water vapour in saturated air.

Then dew point Temperature  $T_{dew}$  was obtained using [19]

$$T_{dew} = T - \frac{100 - RH}{5} \quad (7)$$

where  $T$  and  $RH$  are the Mean Temperature (Kelvin) and Relative Humidity in Percentage.

The virtual Temperature ( $T_{virtual}$ ) was also obtained using [20].

$$T_{virtual} = \frac{T}{1 - \frac{e}{p}(1 - \epsilon)} \quad (8)$$

where  $e$  is the vapour pressure and  $\epsilon$  is a constant denoted by 0.622

The potential temperature  $T_{potential}$  was obtained using [20]

$$T_{potential} = T_{mean} \left(\frac{p_o}{p}\right)^{\frac{R}{c_p}} \quad (9)$$

Equation (9) is called the Poisson's equation where  $p_o$  is the standard pressure generally taken as 1000hPa and  $\frac{R}{c_p} = 0.286$

## 2.1 Validation of the Models

The validation of the estimation of the estimated values was statistically tested by computing the Mean Bias Error (MBE), Root Mean Square Error (RMSE), Mean Percentage Error (MPE), Nash-Sutcliffe equation (NSE), ~~and the~~ Index of Agreement (IA), ~~similarly, also and~~ the coefficient of determination ( $R^2$ ) ~~was determined~~ for each of the models.

The expressions for the MBE, RMSE and MPE as stated according to [21 – 28], were given as follows.

$$MBE = \frac{1}{n} \sum_{i=1}^n (SWVD_{i,cal} - SWVD_{i,mea}) \quad (10)$$

$$RMSE = \left[ \frac{1}{n} \sum_{i=1}^n (SWVD_{i,cal} - SWVD_{i,mea})^2 \right] \quad (11)$$

$$MPE = \left( \frac{\frac{1}{n} \sum_{i=1}^n SWVD_{i,mea} - SWVD_{i,cal}}{SWVD_{i,mea}} \right) \cdot 100\% \quad (12)$$

$$NSE = 1 - \frac{\sum_{i=1}^n (SWVD_{i,mea} - SWVD_{i,cal})^2}{\sum_{i=1}^n \left( SWVD_{i,mea} - \frac{\sum_{i=1}^n SWVD_{i,mea}}{n} \right)^2} \quad (13)$$

$$IA = 1 - \frac{\sum_{i=1}^n (SWVD_{i,cal} - SWVD_{i,mea})^2}{\sum_{i=1}^n \left( |SWVD_{i,cal} - SWVD_{i,mea}| + \left| SWVD_{i,mea} - \frac{\sum_{i=1}^n SWVD_{i,mea}}{n} \right| \right)^2} \quad (14)$$

From equation (10) to (14)  $SWVD_{i,cal}$ ,  $SWVD_{i,mea}$  and  $n$  are the  $i^{th}$  calculated,  $i^{th}$  measured values of daily Surface Water Vapour Density and total number of observations, respectively. ~~also~~  $\frac{\sum_{i=1}^n SWVD_{i,mea}}{n}$  is the mean Surface Water Vapour Density.

Chen et al. [29] have recommended that a zero value for MBE is ideal and a low RMSE is desirable. Similarly, the smaller the value of the MBE and RMSE the better is the

Comment [A8]: Rephrase the statement.

Comment [A9]: Rephrase the statement.

performance of the model, a positive MPE and MBE values provide the averages amount of overestimation in the calculated values, while the negative values gives underestimation. A low value of MPE is desirable. The percentage error between  $-10\%$  and  $+10\%$  is considered acceptable [30 – 35]. High values of  $R^2$ , NSE and IA are desirable. The MBE and the RMSE are in  $\text{gm}^{-3}$ , while  $R^2$ , MPE, NSE and IA are in percentage (%) [36 – 39].

## 2.2 The Developed Two Variable Correlation Models

The proposed two variable correlation models are of the form:

$$SWVD = a + bP + cCC \quad (15)$$


$$SWVD = a + bP + cSSH \quad (16)$$

$$SWVD = a + bP + cPWV \quad (17)$$

where a, b and c are empirical constants and P, CC, SSH and PWV are Surface Pressure, Cloud Cover, Sunshine Hours and Precipitable Water Vapour, respectively.

## 3. RESULTS AND DISCUSSION


### 3.1 Surface Water Vapour Density and its Variation Meteorological Parameters for Benin



**Fig. 1. Monthly variations of SWVD for Benin, Nigeria**


Fig. 1 shows the monthly variation of Surface Water Vapour Density (SWVD) during the period under investigation for Benin. The result revealed that the SWVD during the rainy season is greater than that in the dry season. It was observed that the maximum and minimum values of SWVD of  $21.4753 \text{ gm}^{-3}$  and  $15.1798 \text{ gm}^{-3}$  in the months of June and January respectively. It was observed that the values of SWVD decreased in the month of July and August immediately after its maximum value in the month of June and later

increases in the month of September; this observation is in line with the result reported by Akpootu et al. [14] for Owerri located in the Southern zone of Nigeria.




**Fig. 2. Diurnal variation of SWVD in the dry season for Benin Nigeria**

Fig. 2 shows the diurnal variation of SWVD during the dry season for Benin. The result showed fluctuation in the amount of SWVD with the maximum and minimum values 27.5624 gm<sup>-3</sup> and 9.6567 gm<sup>-3</sup> on the 23<sup>rd</sup> March, 2014 and 26<sup>th</sup> December, 2014 respectively.



**Fig. 3. Diurnal variation of SWVD in the rainy season for Benin Nigeria**


Fig. 3 shows the diurnal variation of SWVD during the rainy season for Benin. The result shows that the maximum the minimum values of SWVD occurred with 27.5731  $\text{gm}^{-3}$  and 16.0151  $\text{gm}^{-3}$  on the 23<sup>rd</sup> May, 2014 and 10<sup>th</sup> August, 2014, respectively.



**Fig. 4. Variation of SWVD with Mean Temperature for Benin**

Fig. 4 shows the monthly variation of SWVD with mean temperature for Benin. The SWVD increased gradually from its minimum value of  $15.1798 \text{ gm}^{-3}$  in the month of January until it got to its maximum value of  $21.4753 \text{ gm}^{-3}$  in the month of June and decreased in August with a dip downward which suddenly increased to September and then dropped to December. The mean temperature increased with the SWVD from January and attained its maximum value of  $300.9395 \text{ K}$  in the month of March which then decreased continuously to its minimum value of  $297.6724 \text{ K}$  in the month of August and increased then to December. The drop in the SWVD as observed in the month of August may be as a result of August break, which is a period of short dryness; it is obvious that it corresponded to the period when the minimum temperature was observed in the study area. The results showed that high and low values of SWVD were observed during the rainy and dry seasons respectively; the reverse is the case for the mean temperature.


**Comment [A10]:** Split into short and simple sentences.



**Fig. 5. Variation surface water vapour density with relative humidity for Benin**


Fig. 5 shows the variation of SWVD with Relative humidity for Benin. The result shows that the relative humidity increases from its minimum value in January with 57.8482 % to its maximum value of 90.7671 % in July then drops down to August increases to September then drops down to December with 61.6150 % the variation with SWVD is in phase therefore there is a close similarities between their characteristics.

**Comment [A11]:** Split the sentence into simple sentences.




**Fig. 6. Variation of SWVD with Surface Pressure for Benin**

Fig. 6 is the variation of SWVD with the surface pressure; the result shows that the magnitude of surface pressure decreases from January to March then increases from April to its maximum value of 1000.9250 mbars in the month of July then continue dropping to December. The minimum value of 996.9097 mbars was observed in the month of March'




**Fig. 7. Variation of SWVD with Cloud Cover for Benin**

Fig. 7 shows the monthly variation of SWVD with cloud cover for Benin. The cloud cover increases with the SWVD from January to April with a little drop down in the month of May then increase to July then continue to its maximum value of 0.8886 in the month of September, then decreases from October to its minimum value of 0.5070 in the month of December.



**Fig. 8. Variation of SWVD with sunshine hours for Benin**


Fig. 8 shows the monthly variation of SWVD with sunshine hours for Benin. The sunshine hours decreases and increases at almost an equal interval from January to May and then decreases to its minimum value in the month of August at 6.4278 hours which corresponds to the August break observed for the SWVD. The sunshine hours increases from its minimum value in August to its maximum value of 7.8356 hours in December. The result revealed that high values of sunshine hours were observed during the dry season and low values during the rainy season which is the reverse case for the SWVD.



**Fig. 9. Variation of SWVD with Precipitable Water Vapour for Benin**

Fig. 9 shows the variation of SWVD with Precipitable Water Vapour (PWV) the result shows that the SWVD increase in the same pattern PWV they increase significantly from January until they reach their minimum value of  $21.0359\text{gm}^{-3}$  and  $4.6697\text{cm}$  in the month of July, the sudden drop in value of both the SWVD and the PWV in the month of August which is signified by the popular August break in the region.

**3.2. Variation of Mean Temperature, Virtual Temperature, Potential Temperature and Dew Point Temperature for Benin**



**Fig. 10. Variation of SWVD with Temperature ( $T_m$ ,  $T_p$ ,  $T_v$  and  $T_d$ ) for Benin**

Fig. 10 shows the monthly variation of mean temperature, dew point temperature, virtual temperature and potential temperature ( $T_d$ ,  $T_v$ ,  $T_p$  and  $T_m$ ) Benin. The result revealed that the mean temperature, the virtual temperature and potential temperature ~~are~~ were almost the same and at the same rate of variation with SWVD they all increased from January to their maximum values of 300.9395 K for mean temperature, 300.9292 K for virtual temperature and 301.2060 K for potential temperature. The temperatures continued to drop down from their maximum values to August where they also reached their minimum values of 297.6724 K for mean temperature, 297.6619 K for virtual temperature and 297.6096 for potential temperature. After the minimum level the temperatures rose to November then drop to December.

### 3.3. Two Variable Correlation Model for Benin

The two variable correlation model developed for Benin based on equation 15 to 17 are as follows:

$$SWVD = 295 - 0.288P + 16.3CC \quad 18a$$

$$SWVD = 1053 - 1.00P - 4.80SSH \quad 18b$$

$$SWVD = 0.363 - 0.000364P + 4.51PWV \quad 18c$$

**Table 1a. Statistical validation tests for Benin**


Models	R <sup>2</sup>	MBE	RMSE	MPE	NSE	IA
18a	92.5	0.1851	0.5714	-1.0531	96.5993	99.1695
18b	47.4	-0.0064	1.4271	-0.5888	78.7859	93.8017
18c	100	0.2882	0.2882	-1.4977	99.1346	99.7986

From table 1a above equation 18c has the highest R<sup>2</sup>, NSE and IA it also has lowest RMSE which shows that it is the best equation for estimating SWVD in the location, but equation 18b is the best in terms of MBE and MPE.

**Table 1b. ranking for the Models**

Models	R <sup>2</sup>	MBE	RMSE	MPE	NSE	IA	Ranking
18a	1	2	2	2	2	2	11
18b	3	1	3	1	3	3	14
18c	2	3	1	3	1	1	11

The ranking from table 1b shows that both equation 18a and 18c are suitable for estimation of SWVD for Benin and stations with similar weather conditions, as they has equal and lowest ranking.



**Fig. 11. Monthly variation of SWVD and developed models for Benin**

Figure 11 shows the monthly variation of SWVD with and the developed equations, the result shows that equation 18b overestimates SWVD in January and February, then August September and December. It also underestimates it from March to July.

#### 4. CONCLUSION

In this study, the question of estimating SWVD and its variation with other meteorological parameters during the period of thirty eight (38) years (1979 – 2016) and the daily variation of SWVD in each month for the year 2014 as well as the development of two variables correlation models has been addressed using monthly and daily average meteorological data obtained from the European Centre for Medium-Range Weather Forecasts (ECMWF) for Benin located in the Coastal region of Nigeria. The results of this study discovered that high values of SWVD are recorded during the rainy season and low values during the dry season for the location. The average value of SWVD was found to be  $21.14479 \text{ gm}^{-3}$ . The results of the diurnal variations of SWVD during the dry and rainy seasons show fluctuation in its values. The highest value of surface water vapour density was observed on 23<sup>rd</sup> May, 2014 with  $27.57313 \text{ gm}^{-3}$  and the lowest on 26<sup>th</sup> December, 2014 with  $9.656676 \text{ gm}^{-3}$ . Three simple two variable correlation models were developed and were statistically tested using statistical indices of coefficient of determination ( $R^2$ ), Mean Bias Error (MBE), Root Mean Square Error (RMSE), Mean Percentage Error (MPE), Nash-Sutcliff Equation (NSE) and Index of Agreement (IA) from which the model that relate Pressure (P) with Precipitable Water Vapour (PWV) was found to be the best (with the accuracy of about 100%) for estimating SWVD for the locations under investigation.

**Comment [A12]:** Should be deleted from this section and included in the result section.

#### REFERENCES

1. Akpootu DO, Iliyasu MI, Mustapha W, Salifu SI, Sulu HT, Arewa SP, Abubakar MB. Models for Estimating Precipitable Water Vapour and Variation of Dew Point Temperature with Other Parameters. *Journal of Water Resources and Ocean Science*. 2019a; 8(3), 28-36 doi: 10.11648/j.wros.20190803.11
2. Held IM, Soden BJ. Water vapor feedback and global warming, *Annual Review of Environment and Resources*. 2000; vol. 25, pp. 441- 475.
3. Trenberth KE, Fasullo J, Smith L. Trends and variability in column-integrated atmospheric water vapor. *Climate Dynamics*. 2005; vol. 24, no. 7-8, pp. 741-758.
4. Wang J, Carlson DJ, Parsons DB. Performance of operational radiosonde humidity sensors in direct comparison with a chilled mirror dew-point hygrometer and its climate implication. *Geophysical Research Letters*. 2003; vol. 30, no. 16.
5. Wagner T, Beirle S, Grzegorski M. Global trends (1996-2003) of total column precipitable water observed by Global Ozone Monitoring Experiment (GOME) on ERS 2 and their relation to near-surface temperature. *Journal of Geophysical Research: Atmospheres*. 2006; vol. 111, no. 12, Article ID D12102.
6. Changnon DM, Sandstrom, C, Schaffer C, Relating changes in agricultural practices to increasing dew points in extreme Chicago heat waves. *Climate Resources*. 2003; 24, 243–254.
7. Dai A. Recent climatology, variability, and trends in global surface humidity. *Journal of Climate Change*. 2006; vol. 19, no. 15: pp. 3589- 3606.

8. Zhao Y, Gong L, Zhou B, Huang Y, Liu C. Detecting tomatoes in greenhouse scenes by combining AdaBoost classifier and colour analysis. *Biosyst. Eng.* 2016; 148, 127–137.
9. Dai A, Wang J, Thorne P. A new approach to homogenize daily radiosonde humidity data. *Journal of Climate Change.* 2011; vol. 24, no. 4: pp. 965-991.
10. Khalameyda DD. Influence of fluctuations in the troposphere refractive index on the interferometer measurement accuracy of radio wave angles of arrival. *Telecommun. Radio Eng.* 70. 2011;
11. Khaniani AS, Nikraftar Z, Zakeri S. Evaluation of MODIS Near-IR water vapor product over Iran using groundbased GPS measurements. *Atmospheric Resources.* 2020; 231, 104657.
12. Israel E, David AK, Samuel OO, Gabriel AA, Omolara EG. Global Distribution of Surface Water Vapour Density Using in Situ and Reanalysis Data. *Journal of Water Resources and Ocean Science.* 2020; 9(3), 64-70. doi: 10.11648/j.wros.20200903.12
13. Ajayi GO. Physics of the tropospheric radiopropagation. *Proceedings of the ICTP College on Theoretical and Experimental Radio Propagation Physics.* 1989; 6–24, Trieste, Italy.
14. Akpootu DO, Mustapha W, Rabi AM, Iliyasu MI, Abubakar MB, Yusuf SO, Salifu SI. Estimation of surface water Vapour Density and its Variation with other Meteorological Parameters over Coastal region of Nigeria. *Hydrology.* 2019; 7(3), 46-55. DOI: 10.11648/j.hyd.20190703.12
15. IPCC. Intergovernmental Panel on Climate change. Third Assessment Report: Climate change 2001. WG1: The scientific basis, summary for policymakers, Geneva, Switzerland.
16. Adeyemi B, Ogolo EO. Diurnal and Seasonal Variation of Surface Water Vapour Density over some Meteorological Station in Nigeria. *Ife Journal of Science.* 2014; 16(2), 181-189.
17. Akpootu DO, Iliyasu MI, Nouhou I, Aina AO, Idris M, Mustapha W, Ohaji DE, Muhammad AD. Estimation and Variation of Saturation Mixing Ratio and Mixing Ratio over Potiskum, Nigeria. *Nigerian Journal of Basic and Applied Science.* 2022a; 30(1), 49-54. DOI: <http://dx.doi.org/10.4314/njbas.v30i1.7>
18. Leckner B. The spectral distribution of solar radiation at the earth's surface—elements of a model. *Sol Energy.* 1978; 20 (2), 143-150.
19. Lawrence MG. The relationship between relative humidity and the dewpoint temperature in moist air: A simple conversion and applications. *Bull. Amer. Meteor. Soc.*, 2005; 86, 225-233. doi: <http://dx.doi.org/10.1175/BAMS-86-2-225>.
20. Wallace JM, Hobbs PV. *Atmospheric Science, An Introductory Survey*, 2<sup>nd</sup> Edition, Elsevier, 2006; pp 66-82.
21. El-Sebaei A, Trabea A. Estimation of Global Solar Radiation on Horizontal Surfaces over Egypt, *Egypt. J. Solids.* 2005; 28(1): 163-175.
22. Gana NN, Akpootu DO. Angstrom Type Empirical Correlation for Estimating Global Solar Radiation in North- Eastern Nigeria. *The International Journal of Engineering And Science.* 2013a; 2(11), 58-78. ISSN: 2319-1813

23. Gana NN, Akpootu DO. Estimation of global solar radiation using four sunshine based models in Kebbi, North-Western, Nigeria. *Pelagia Research Library*. 2013b; **4**(5), 409-421. ISSN: 0976-8610.
24. Akpootu DO, Gana NN. Comparative Study of Global Solar Radiation between Nguru and Abuja. A paper presented at the 24<sup>th</sup> Annual Congress and Colloquium of the Nigerian Association of Mathematical Physics held at University of Benin, Benin City, Nigeria on the 25<sup>th</sup>-28<sup>th</sup> February, 2014. 2014;
25. Akpootu DO, Sanusi YA. A New Temperature-Based Model for Estimating Global Solar Radiation in Port-Harcourt, South-South Nigeria. *The International Journal of Engineering And Science*. 2015; **4**(1), 63-73.
26. Olomiyesan BM, Akpootu DO, Oyedum DO, Olubusade JE, Adebunmi SO. Evaluation of Global Solar Radiation Models Performance using Global Performance Indicator (GPI): A Case Study of Ado Ekiti, South West, Nigeria. A paper presented at the 43<sup>rd</sup> Annual Nigerian Institute of Physics, National Conference, held at the NnamdiAzikiwe University, Awka, Anambra State May 26-29, 2021. 2021;
27. Akpootu DO, Iliyasu MI. The Impact of some Meteorological Variables on the Estimation of Global Solar Radiation in Kano, North Western, Nigeria. *Journal of Natural Sciences Research*. 2015a; Vol.5, No.22, 1 – 13.
28. Akpootu DO, Iliyasu MI. A Comparative Study of some Meteorological Parameters for Predicting Global Solar Radiation in Kano, Nigeria Based on Three Variable Correlations. *Advances in Physics Theories and Applications*. 2015b; Vol.49, 1 – 9.
29. Chen R, Ersi K, Yang J. Validation of five global radiation Models with measured daily data in China. *Energy Conversion and Management*. 2004; **45**, 1759-1769.
30. Merges HO, Ertekin C, Sonmete MH. Evaluation of global solar radiation Models for Konya, Turkey. *Energy Conversion and Management*. 2006; **47**: 3149-3173.
31. Akpootu DO, Momoh M. Empirical Model for Estimating Global Solar Radiation in Makurdi, Benue State, North Central Nigeria. A paper presented at the 36<sup>th</sup> Annual Nigerian Institute of Physics, National Conference, held at the Department of Physics, University of Uyo, Nigeria on May 26-29, 2014. 2014;
32. Akpootu DO, Sulu HT. A Comparative Study of Various Sunshine Based Models for Estimating Global Solar Radiation in Zaria, North-Western, Nigeria. *International Journal of Technology Enhancements and Emerging Engineering Research*. 2015; **3**(12), 1 – 5.
33. Akpootu DO, Tijjani BI, Gana UM. Empirical models for predicting global solar radiation using meteorological parameters for Sokoto, Nigeria. *International Journal of Physical Research*. 2019c; **7**(2): 48 – 60. DOI: 10.14419/ijpr.v7i2.29160
34. Akpootu DO, Tijjani BI, Gana UM. Sunshine and Temperature Dependent Models for Estimating Global Solar Radiation Across the Guinea Savannah Climatic Zone of Nigeria. *American Journal of Physics and Applications*. 2019d; **7**(5): 125-135. doi: 10.11648/j.ajpa.20190705.15.

35. Akpootu DO, Tijjani BI, Gana UM. New temperature dependent models for estimating global solar radiation across the midland climatic zone of Nigeria. International Journal of Physical Research. 2019e; 7(2): 70 – 80. DOI: 10.14419/ijpr.v7i2.29214
36. Akpootu DO, Tijjani BI, Gana UM. New temperature dependent models for estimating global solar radiation across the coastal climatic zone of Nigeria. International Journal of Advances in Scientific Research and Engineering (ijasre). 2019f; 5(9): 126 – 141. DOI: 10.31695/IJASRE.2019.33523
37. Akpootu DO, Abdullahi Z. DEVELOPMENT OF SUNSHINE BASED MODELS FOR ESTIMATING GLOBAL SOLAR RADIATION OVER KANO AND IKEJA, NIGERIA. FUDMA Journal of Sciences (FJS). 2022; Vol. 6 No. 3, June, 2022, pp 290 – 300. DOI: <https://doi.org/10.33003/fjs-2022-0603-1001>
38. Akpootu DO, Iliyasu MI, Olomiyesan BM, Fagbemi SA, Sharafa SB, Idris M, Abdullahi Z, Meseke NO. MULTIVARIATE MODELS FOR ESTIMATING GLOBAL SOLAR RADIATION IN JOS, NIGERIA. Matrix Science Mathematic (MSMK). 2022b; 6(1) 05-12. DOI: <http://doi.org/10.26480/mkkm.01.2022.05.12>
39. Akpootu DO, Alaiyemola SR, Abdulsalam MK, Bello G, Umar M, Aruna S, Isah AK, Aminu Z, Abdullahi Z, Badmus TO. Sunshine and Temperature Based Models for Estimating Global Solar Radiation in Maiduguri, Nigeria. Saudi Journal of Engineering and Technology. 2023; 8(5): 82-90. DOI: 10.36348/sjet.2023.v08i05.001.