

# Blood Transfusion Request and Utilization: The Trend in a Tertiary Health Care Center in North Central Nigeria.

---

## ABSTRACT

**Background:** An effective blood transfusion service is anchored on an adequate supply of safe blood and its components as well as its accessibility and appropriate clinical use. There is a contrast in blood supplies and utilization between the high, middle and low-income countries necessitating the formulation of appropriate modalities aimed at ensuring judicious use of blood and minimizing wastage.

**Aims:** This study was aimed at assessing the blood transfusion request trend in our institution and its utilization with the intent of curtailing wastage of safe blood for transfusion.

**Study design:** It is a Retrospective study.

**Place and Duration of Study:** Jos University Teaching Hospital Blood Bank from January 2022 to December 2022.

**Methodology:** Appropriately filled blood transfusion request forms of the Jos University Teaching Hospital compared with the blood bank In-house records from January 2022 to December 2022 were reviewed. Request dates, requesting Department, Unit or Specialty, indications for request, Number of units requested and number eventually utilized were used to determine blood transfusion indices which are indicative of efficient and significant blood use.

**Results:** Eight thousand, Five hundred forty-eight (8,548) blood transfusion request forms from the blood bank and patients' folders were reviewed and compared with the Blood Bank's in-house records. Requests for blood transfusion for over 460 indications were received from twenty-six (26) departments and units. There was a general Crossmatch: Transfusion ratio (C: T) of 1.7 with a transfusion index (TI) of 1.2 and transfusion probability (TP) of 48.0%.

**Conclusion:** Transfusion requests were characterized by low utilization across departments and units. This calls for the establishment of transfusion protocol for the determination of blood volume required for every indication so that this limited commodity will be made available and accessible when needed.

*Keywords: Blood transfusion, Request, Trend, Utilization*

## 1. INTRODUCTION

Blood transfusion is a multi-specialist medical procedure that involves the infusion of whole blood or its components from a donor to a recipient. The estimated annual global blood donation is reported to be about 110 million units [1]. Apheresis-derived components which are rarely available in our health care centres' make up 12.4 % of all blood transfusions while whole blood, a source for several important components in blood transfusion, accounts

for 89.5% of all blood transfusions Worldwide [1]. The demand for blood transfusion cuts across departments and units such that no single department or unit can effectively function without this life-giving product. Increasing globalization and sophistication in patient care for haematologic disorders, obstetrics and gynaecology interventions and complications, surgical procedures, and several other medical conditions have created further demands for blood supply.

The average donation rate all across Africa is about 4.3 donations per 1000 population with an abysmal 0.2 donations per 1000 population in Nigeria [2]. Requests for blood transfusions are usually made in anticipation of bleeding that could occur during surgical interventions, blood loss from several other conditions or as a result of disorders arising from excessive blood cell destruction or due to inability of the bone marrow to produce blood cells to meet up the body's physiologic needs[3]. While requests during surgeries could be subjective, those due to an underlying disorder with evidently low haemoglobin concentration will require prompt supplies.

Several request and cross-match done in anticipation comes with minimal utilization and resultant wastage. It was estimated that in 2013 alone, about 1.8 million blood units were discarded globally [4]. Over the years donations have been grossly insufficient to meet the demand for transfusion, necessitating the need to closely conserve available supplies for judicious use where required.

To the best of our knowledge, no study of this magnitude has been carried out in our facility with the intent of determining blood transfusion requests, usage and utilization in departments and units of the hospital. This study is aimed at assessing the trend of blood transfusion requests in our institution and subsequent utilization towards identifying areas of wastage so these blood supplies can be appropriately channelled to where it is needed at all times.

## **2. MATERIAL AND METHODS**

### **2.1 Study Design**

This is a hospital-based retrospective study. Completed blood transfusion request forms from the Jos University Teaching Hospital Blood Bank were reviewed from January 2022 to December 2022.

### **2.2 Data Collection Method**

The Researchers reviewed the completed blood transfusion request forms and extracted request dates, requesting Departments and Units, indications for request, number of units requested and number eventually utilized. Blood Transfusion utilization indices for each department and unit were calculated using the following equations with their inferences.

- i. Cross-match to transfusion ratio (C: T ratio) = Number of units cross-matched/Number of units transfused.  
A ratio of less than 2.5 indicates efficient blood use[5].
- ii. Transfusion Index (TI) = Number of units transfused/Number of patients cross-matched.  
TI greater than 0.5 is indicative of significant blood utilization and blood will need to be cross-matched preoperatively [5].
- iii. Transfusion Probability (TP %) = Number of Patients transfused/Number of patients cross-matched x 100.  
A transfusion probability of  $\geq 30\%$  is considered significant blood usage[5].

### **2.3 Data analysis**

Microsoft Excel 2010 and Epi Info Version 7.2.5.0 software were used to analyze the data generated. The results were reported in proportions and percentages, and presented in tables.

## 2.4 Ethical Approval

Ethical approval was obtained from the Jos University Teaching Hospital Health Research Ethics Committee.

## 3. RESULTS

Eight thousand, Five hundred forty-eight (8,548) blood transfusion request forms obtained from the blood bank and patients' folders were reviewed and compared with the Blood Bank's in-house records. Units of blood requested for transfusion ranged from 1 unit to 10 units per request with 2 units accounting for 4668 (54.56%) requests. Three thousand eight hundred forty four (44.97%) requests made were never utilized, while the highest number of units transfused was 14 units (Table 1). Requests for blood transfusion were received from twenty-six (26) departments and units. The highest number of requests 2000 (23.4%) was received from the Department of Obstetrics and Gynaecology while the least request was from dermatology, a unit in the Department of Internal Medicine. Department of Haematology and the Paediatric haemato-oncology unit of the Department of Paediatrics had requests of 392 (4.6%) and 375 (4.4%) respectively (Table 2).

Seventeen thousand, eight hundred and twenty-four (17,824) units of blood were crossed match for the 8,548 requests we received within the study period but only 10,320 units were transfused. There was a general Crossmatch: Transfusion ratio (C: T) of 1.7 with a transfusion index (TI) of 1.2 and transfusion probability (TP) of 48.0%

Department of Obstetrics and Gynaecology which had the highest blood transfusion request, Department of General Surgery and the Department of Haematology had a C: T and TIs of 2.5, 1.7, 0.9, and 0.8, 1.2, 2.2 respectively (Table 2).

**Table 1: Blood units per request and blood units transfused (n=8548)**

Variables	Unit(s)	Frequency (n)	Percentage (%)
<b>Number of units requested for transfusion</b>	1	2000	23.40
	2	4668	54.61
	3	1264	14.79
	4	532	6.22
	5	20	0.23
	6	48	0.56
	7	0	0.00
	8	8	0.09
	9	0	0.00
	10	8	0.09
<b>Number of units not transfused/transfused</b>	0	3844	44.97
	1	1988	23.26
	2	1408	16.47
	3	596	6.97
	4	360	4.21
	5	140	1.64
	6	84	0.98
	7	52	0.61
	8	28	0.33
	9	16	0.19
10	20	0.23	

11	5	0.06
12	0	0.00
13	4	0.05
14	3	0.04

**Table 2: Number of transfusion requests, number of units' cross match, number of units transfused, C: T ratio, TI and TP by Departments and Units**

Department /Units transfusion request	Number of blood units	Number of blood units cross-matched	Number of transfused	C:T ratio	TI	TP(%)
Accident and Emergency	203	534	328	1.6	1.6	38.0
Cardiothoracic Surgery	263	696	464	1.5	1.8	37.8
Cardiology	141	385	301	1.3	2.1	36.6
Dermatology	4	8	8	1.0	2.0	50.0
Endocrinology	128	324	212	1.5	1.7	39.5
Oto-rhino-laryngology	138	246	38	6.5	0.3	56.1
Emergency Paediatric Unit	294	334	302	1.1	1.0	88.0
Gastroenterology	205	526	343	1.5	1.7	39.0
Haematology	392	766	850	0.9	2.2	51.2
Infectious disease	164	414	280	1.5	1.7	39.6
Maxillo Facial Surgery	182	332	98	3.4	0.5	54.8
Nephrology	420	1040	852	1.2	2.0	40.4
Neurology	100	244	172	1.4	1.7	41.0
Neuro-surgery	569	1480	504	2.9	0.9	38.5
Obstetrics &Gynaecology	2000	4123	1641	2.5	0.8	48.5
Oncology (Adult)	89	151	150	1.0	1.7	58.9
Orthopaedic Surgery	674	1438	620	2.3	0.9	46.9
Paediatric (General)	125	153	185	0.8	1.5	81.7
Plastic Surgery	300	608	348	1.8	1.2	49.3
Paediatric Haemato-oncology	375	488	502	1.0	1.3	76.8
Paediatric Surgery	256	369	146	2.5	0.6	69.4
Pulmonology	46	128	96	1.3	2.1	35.9
Rheumatology	82	170	124	1.4	1.5	48.2
Special Care Baby Unit	276	292	287	1.0	1.0	94.5
Surgery (General)	835	1904	1041	1.8	1.3	43.9
Urology	287	635	428	1.5	1.5	45.2
	<b>8,548</b>	<b>17,824</b>	<b>10,320</b>	<b>1.7</b>	<b>1.2</b>	<b>48.0</b>

C: T- Crossmatch: Transfusion, TI- Transfusion Index, TP- Transfusion Probability

#### 4. DISCUSSION

The first recorded successful blood transfusion occurred in 1665 in England when Doctor Richard Lower successfully kept a dog alive by transfusing it with blood from other dogs. By 1818, a British Obstetrician James Blundell successfully transfused human blood to a patient with postpartum haemorrhage marking the advent of human-to-human blood transfusion[6]. Little wonder that obstetrics and gynaecology continue to be one of the medical disciplines known to have a high demand for blood as demonstrated in this study. The first recorded blood transfusion in Africa was by an Arab physician in Egypt in 1242 and was later carried out in well-resourced medical centres in Uganda, Kenya and Senegal [7].

Humans realized how essential blood is from the ancient period hence the tag "*Vital force and essence of life*", however, it has become practically impossible to meet the world's safe blood transfusion demand despite efforts spanning decades of policies after policies[8]. Through well-formulated and supported policies, World Health Organization earmarked the year 2012 as the year for Nigeria and other Sub-Saharan African Countries to meet their demands for safe blood as well as sustain its availability[9, 10]. World Health Organization further recommended that the minimum blood donation by any given population should be 1% of that population but till date more than 70 countries globally have a blood donation rate of less than 1% [11]. Africa was only able to achieve 40% of that minimum requirement of blood donation in 2006 while Nigeria achieved only 33.3% [11]. These reports demonstrated the fact that this essential gift of life is grossly in short supply globally more so in the middle and low income countries, Nigeria inclusive.

Millions of people from different backgrounds experience one form of bleeding or the other in addition to obstetrics haemorrhages, paediatrics and adult anaemias caused by hundreds of benign and malignant disorders. This therefore calls for judicious utilization of this life-giving commodity to tackle the innumerable dire consequences associated with its scarcity. Obstetrics and Gynaecology (O&G) accounted for the highest transfusion request received in our centre and this is supported by several other reports in Nigeria and other Countries of the World[12-14]. Contrary to the predominance of O&G in blood transfusion requests in our study, Ajayi *et al*[15] in Ado-Ekiti, Nigeria reported the predominance of requests from the Department of Paediatrics. While it is true that more request was received from O&G in our centre, our findings demonstrated that less than half of the blood cross-matched were transfused. Several of these requests were made in anticipation of intra or post-operative bleeding while others arise from relative estimation of blood requirement for a procedure or a patient presenting with anaemia. There are several scientific methods of calculating blood requirement for patients based on packed cell volume (PCV) or haemoglobin (Hb) concentration at presentation relative to the desired blood level, patient weight in kilogram, and blood volume per kilogram body weight, but many health care personnel transfuse with the notion of 1 unit of blood increases PCV by 3% or Hb concentration by 10 gram per litre when it applies to red cell concentrate transfusion in an adult with 70-75 kilogram body weight [16,17]. This leads to overestimation and wastage of blood or underestimation of patients' blood requirement sometimes. This finding affirms the need for appropriate blood transfusion requests in quantity and component with the intent of maximizing utilization while minimizing wastage. A look at the individual statistics of the different departments and units captured in our study showed that a significant number of these departments and units rarely utilize half of the blood cross-matched based on several requests. However, worthy of note is the fact that the department and unit involved in managing haematologic disorders in adults and children as well as the paediatric emergency unit of the institution exceeded the number of units initially requested when compared with their transfusion rates. This may be linked to the fact that most patients in these categories present with clinically and laboratory-verifiable anaemia of varying degrees requiring instant blood transfusion. The surgical unit of Otorhinolaryngology had the lowest transfusion rate probably due to the predominant surgical procedures they are known to perform or their surgical precision which are not associated with significant blood loss. The general blood transfusion indices of the cross match: transfusion ratio (C: T), Transfusion index (TI) and transfusion probability (TP) determined in our study indicated significant blood usage and utilization. This cuts across all the departments and units identified to have requested blood for transfusion within the study period except for two surgical departments of Otorhinolaryngology and Neurosurgery despite having a significant transfusion probability. These slight differences may not be unconnected with the long-term surgical practice of blood grouping and cross-matching ahead of surgical procedures in anticipation and for prevention of complications that could occur in case of excessive bleeding during the surgery. The indices recorded in our study are slightly higher but within the range of significant blood usage and utilization when

compared to what was reported in studies from Calabar in Southern Nigeria and Sagamu in the South West region of Nigeria[18-19]. A report from Nepal, India showed a much higher transfusion probability indicating a very significant blood usage with minimal wastage [20]. There were over 460 indications for blood transfusion requests in our study cutting across the departments and units considered. Blood transfusion requests as assessed from this study indicated the fact that several requests are made in anticipation without quantifying the exact volume required resulting in keeping units of blood that may not be used or minimal quantity are utilized in comparison to what was requested. Scientific methods to determine blood volumes based on deficits incurred or anticipated are not utilized. Hospital transfusion committees could employ the use of schedules like the maximum surgical blood order schedule (MSBOS) using Mead's criterion where MSBOS= 1.5 x transfusion index or the Patient Specific Blood Ordering System (PSBOS) [21, 22]. These schedules and systems can aid the maintenance of inventory, adequate blood usage and utilization with resultant reduction in blood wastage thereby cutting cost and manpower required for blood safety. Blood transfusion indices by departments and units were used in our study, however, individual indications or yearly assessments can be carried out and these indices are determined for the same purpose as reported in several studies [23-25].

## **5. CONCLUSION**

Availability of blood and its components remains an integral part of any viable health care system but despite the glaring benefits of this product and the fact that our blood donation bases have continued to dwindle, blood transfusion requests are made without scientific quantification of volume required. This results in the unavailability of blood where required with resultant blood wastage.

## **6. RECOMMENDATION**

A robust system for improving blood utilization to curb blood wastage and channel excesses to where it is needed must be part of our health system planning. Health personnel working in surgical departments and units should be trained and encouraged to implore scientific methods of determining their patients' blood requirements during surgeries. Bloodless techniques and blood salvage during surgeries should also be encouraged.

## **10. ETHICAL APPROVAL**

Ethical approval was obtained from the Jos University Teaching Hospital Health Research Ethics Committee.

## **REFERENCES**

1. World Health Organization. Global status report on blood safety and availability 2021. Geneva: World Health Organization. 2022. 1-184. [www.who.int/bloodproducts](http://www.who.int/bloodproducts).
2. Tapko JB, Toure B, Sambo LG. Status of blood safety in the WHO African Region: Report of the 2010 Survey. Brazzaville, Republic of Congo: World Health Organization. Regional Office for Africa; 2014.
3. Outta AB. Transfusion practice. Clinical aspects and application in Blood banking and Transfusion. 1st ed. New Delhi. CBS Publishers; 2006: 213-321.
4. Enosolease ME, Imarengiaye C, Awodu AO. Donor blood procurement and utilization at the University of Benin Teaching Hospital. Benin City Afr J Reprod Health 2004; 8:59-63.
5. Feyisa T, Tesfaye Kiya G, Maleko WA. Assessment of recipients' characteristics, transfusion appropriateness, and utilization pattern of blood and blood products in Jimma Medical Center, Jimma, Ethiopia. PLoS ONE 2021; 16(4): e0250623. <https://doi.org/10.1371/journal.pone.0250623>.

6. Learoyd P. The history of blood transfusion prior to the 20<sup>th</sup> century-Part 1. *Transfus Med* 2012; 22: 308-314.
7. White L. The History of Blood Transfusion in Sub-Saharan Africa by William H, Schneider (Review). *Bull Hist Med* 2015; 89: 148-150.
8. Schroeder ML. Principles of Transfusion Medicine. In: Lee GR, Foerster J, Paraskevas F, Greer JP, Rogers GM, Eds. *Wintrobe's Clinical Haematology*; 10th ed. Baltimore, William and Wilkins, 1998: 817-872.
9. Aneke JC, Okocha CE, Blood Transfusion safety; Current Status and Challenges in Nigeria. *Asian J Transfus Sci* 2017; 11: 1-5.
10. Weimer A, Tagny CT, Tapko JB, et al. Blood transfusion safety in Sub-Saharan Africa: A literature review of changes and challenges in the 21<sup>st</sup> century. *Transfusion (Paris)* 2019; 59: 412-427.
11. Nwabuko OC, Nnaji TO. Challenges of Blood Transfusion in a Developing Country and the Ways Forward. *Biomed J Sci & Tech Res* 2021; 33(4):1-3. 10.26717/BJSTR.2021.33.005435.
12. Aliyu I, Michael G, Ibrahim H, Ibrahim ZF, Aliyu G, Isaiah AT. Blood transfusion request pattern in a medical centre in Northwestern Nigeria. *Glob J Transfus Med AATM* 2017; 2: 52-55.
13. Emeribe AO, Ejele AO, Attai EE, Usanga EA. Blood donation and patterns of use in Southeastern Nigeria. *Transfusion* 1993; 33: 330-332.
14. Kagu MB, Ahmed SG, Askira BH. Utilisation of blood transfusion service in North Eastern Nigeria. *Highland Med Res J* 2007; 5: 27-30.
15. Ibijola AA, Adegbamigbe OJ, Okunlola AI, Durowade KA, Adebara I, Fasakin KA. Pattern of blood component request and utilization in a tertiary hospital in Nigeria. *Niger J Med* 2022; 31: 87-91.
16. Adedoyin OT, Afolabi JK, Oyeyemi B. Proposed Formulae for Determining Blood Transfusion Requirements in Children with Severe Anaemia. *Nigerian Journal of Paediatrics* 2004; 31(1): 25-28.
17. Lozano M, Badawi M. Indications for transfusion of blood components. *ISBT Science Series* 2020; 15: 320-330. 10.1111/voxs.12605.
18. Okoroiwu HU, Okafor IM. Demographic characteristics of blood and blood components transfusion recipients and pattern of blood utilization in a tertiary health institution in southern Nigeria. *BMC Hematology* 2018; 18: 16-21. 10.1186/s12878-018-0112-5.
19. Akodu SO, Adekanmbi FA, Ogunlesi TA. A Review of Blood Products Requisition and Utilization in Post-neonatal Ward of a Tertiary Hospital in South Western Nigeria. *Journal of Epidemiological Society of Nigeria* 2021; 4(1): 97-107.
20. Shrestha AN, Aryal BB, Poudel A, Chhetri P, Poudel S, Adhikari A, Shrestha S. Blood transfusion practices in a tertiary care hospital in Nepal. *J Pathol Nep* 2020; 10: 1728-1732. 10.3126/jpn.v10i2.30424.
21. Guidelines on Blood and Blood Components Transfusion. Maximum Surgical Blood Order Schedule (MSBOS) For Elective Surgery. UKM Medical Center, 5<sup>th</sup> Edition 2019. 1-19.
22. Palmer T, Wahr JA, O'Reilly M, Greenfield ML. Reducing unnecessary cross-matching: a patient-specific blood ordering system is more accurate in predicting who will receive a blood transfusion than the maximum blood ordering system. *Anesthesia & Analgesia*. 2003;96 (2):369-375.

23. Akoko LO, Joseph AL. Blood utilization in elective surgery in a tertiary hospital in Dar es Salaam, Tanzania. *Tanzania Journal of Health Research* 2015; 17(4): 1-8. 10.4314/thrb.v17i4.5.
24. Quader MA, Islam QT. Analysis of Blood Request Form and Blood Utilization Pattern in a Tertiary Care Specialized Hospital. *Bangladesh J Medicine* 2022; 33: 259-262. 10.3329/bjm.v33i3.61371.
25. Murugesan M, Subbiah SP. Blood requisition and utilization practice in obstetric patients at tertiary care centre from South India: A descriptive study. *J Appl Hematol* 2018; 9:131-135. 10.4103/joah.joah\_42\_18.

UNDER PEER REVIEW