

Original Research Article

Antimicrobial Resistance profile of *Salmonella enterica* serovar *Typhi* isolated from human clinical samples in Ebonyi State

ABSTRACT

This study was designed to determine the antimicrobial Resistance profile of *Salmonella enterica* serovar *Typhi* isolated from human clinical samples in Ebonyi State. A non-duplicated Stool culture of *Salmonella enterica* serovar *Typhi* of patients diagnosed with typhoid fever at General Hospital Onicha Igboeze were collected from the hospital ward namely: A&E (n = 4), MS (n = 3), FS (n = 3), PD (n = 7), LW (n = 4), ORT (n = 1), LAB (n = 17), THE (n = 9), GOPD (n = 4), MM (n = 3). Antimicrobial studies of *Salmonella enterica* serovar *Typhi* was determined using the Kirby–Bauer disk diffusion method. The proportion of resistant ranges from 33 %-100% against colistin, Cefepime, Nalidixic acid, Cefoxitin, amikacin, Cefuroxime and piperacillin-tazobactam but isolates were only susceptible to meropenem 100%. The use of antimicrobial agent for treatment of *Salmonella enterica* serovar *Typhi* infection should be guided with antimicrobial susceptibility testing, Nonetheless, the diversity of the *Salmonella* isolates as a result of the dissemination of these resistant genes is a call for concern and emphasizes a need for an extensive investigation for the presence of these genes in Ebonyi State as well as the implementation of strict antimicrobial policies in a bid to restrict the spread of these resistance genes and prevent the emergence of new resistant strains.

Keywords: *Salmonella Typhi*, Resistant, human clinical, typhoid fever

1. INTRODUCTION

Salmonella enterica serovar *Typhi* (*S. typhi*) is a Gram-negative, rod-shaped, catalase positive flagellated bacterium [1, 2]. Moreover, the three *Salmonella* (*S. paratyphi*) types such as A, B and C are responsible for causing paratyphoid fever [1]. Typhoid fever being a systemic infection caused by *Salmonella enterica* serotype *Typhi* is a highly adapted human specific pathogen which possesses remarkable mechanism for persistence in host as reported by Onyenweet al. [3]. Earlier studies have shown that majority of enteric fevers are prevalent in most of the developing countries like India, Pakistan, Bangladesh, and several African nations [4, 5]. It is estimated to have approximately 5.4 million cases worldwide per year [3]. It can be transmitted via contaminated food, water and is mainly present in area with poor hygiene or low socioeconomic status [6]. According to the World Health Organization (WHO), they estimated that typhoid fever accounts for 222,000 deaths and 21 million infections annually on a global scale [7]. Africa is classified under region with medium incidence rate per year (10-100/100,000 cases/year) centered on a 22 community-based incidence study with only three African countries included in the analysis [7]. As of 2013, the annual mortality rate was reported at 2.8 per 100,000 persons in Africa with 2.5 per 100,000 persons reported for Nigeria [3, 8]. Appropriate treatment reduces the mortality rate as low as 0.5% [3]. Appropriate treatment reduces the mortality rate as low as 0.5% [3]. The emergence of resistance to Cotrimaxazole, chloramphenicol, fluoroquinolones and third-generation cephalosporins had led to the frequent use of azithromycin for empirical treatment of uncomplicated enteric fever. However, the recent emergence of cephalosporin resistant strains of *Salmonella enterica* serovar *Typhi* is a cause for concern in the management of enteric fever and has been noted in human patients as a serious problem in most healthcare settings. This incidence makes typhoid fever an infectious disease worth investigating for better treatment plan and management with the knowledge of local resistant profile.

2. METHODS

2.1 Culture processing

Prior to this research, ethical approval with reference No: SMOH/ERC/042/21 was obtained from the Research and Ethics Committee of Ebonyi State Ministry of Health, Abakaliki, Nigeria. A non-duplicated Stool culture of *Salmonella enterica* serovar *Typhi* of patients diagnosed with typhoid fever at General Hospital Onicha Igboeze located latitude 6° 6' 34" N and longitude 7° 49' 3" E were collected according to the hospital ward

namely: A&E (n = 4), MS(n = 3), FS(n = 3), PD (n = 7), LW (n = 4), ORT (n = 1), LAB (n = 17), THE (n = 9), GOPD (n = 4), MM (n = 3). The isolated *S. typhi* strains were genotypically confirmed by 16S rRNA sequencing at Bioinformatics Service laboratory, Ibadan, Nigeria, with universal specific primers.

2.3 Antibiotic susceptibility studies

Antibiotic susceptibility was performed by employing Kirby Bauer disk diffusion method using sterilized Mueller-Hinton agar in accordance with the guidelines of Clinical and Laboratory Standards Institute[9]. Bacteria suspension of the test isolate were prepared using 0.5 McFarland standards and seeded on solidified Mueller-Hinton agar. The plates were allowed to pre-diffuse for 15 min. Thereafter, the following antibiotics disc: Amoxicillin/clavulanic acid (30 µg), ceftazidime (30 µg), ceftriaxone (30 µg), ceftiofloxacin (30 µg), Colistin Sulphate (10µg), Ciprofloxacin (5 µg), imipenem (10 µg), Ertapenem (10 µg), Meropenem (10µg), Doripenem (10µg), tetracycline (30 µg), Nalidixic acid (30 µg), trimethoprim-sulfamethoxazole (25µg) was impregnated on the inoculated Mueller-Hinton (MH) agar plates and incubated at 37° C for 24 h. After overnight incubation, the diameters of zones of inhibition were measured and results was interpreted in accordance with the criteria of Clinical and Laboratory Standards Institute [9, 10, 11]

3. RESULT AND DISCUSSION

3.1 Result

Antimicrobial Resistant Pattern of *S. typhi*

S. typhi were resistance to Piperacillin-Tazobactam (33-100%), Nalidixic Acid (57-100 %), Cefepime (67-100 %) Cefuroxime (33-100%), Ofloxacin (100 %) colistin (33-100 %) as shown in Table 1. In table 2, the resistant pattern were colistin 50.0 %-100 %, Amoxicillin-Clavulanate acid 25 %-100 %, Cefoxitin 55-100%, Cefuroxime 65-100%. All *S. typhi* were susceptible to meropenem.

Table 1: Antimicrobial Resistant Pattern of *S. typhi* Isolated from fecal samples of patients at A&E, MS, FS, PD and LW

Antibiotic (μ g)	A&E (n = 4) R (%)	MS (n = 3) R (%)	FS (n = 3) R (%)	PD (n = 7) R (%)	LW (n = 4) R (%)
Colistin (30)	3 (75)	3 (100)	1 (33)	5 (71)	2 (50)
Amoxicillin-Clavulanate acid (30)	2 (50)	3 (100)	3 (100)	6 (86)	2 (50)
Piperacillin-Tazobactam (5)	4 (100)	1 (33)	3 (100)	5 (71)	4 (100)
Cefuroxime (20)	4 (100)	1 (33)	3 (100)	5 (71)	4 (100)
Cefepime (15)	3 (75)	3 (100)	3 (100)	7 (100)	4 (100)
Nalidixic Acid (20)	4 (100)	3 (100)	3 (100)	4 (57)	4 (100)
Meropenem (30)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Cefoxitin (30)	4 (100)	2 (67)	1 (33)	5 (71)	4 (100)
Ofloxacin (10)	4 (100)	2 (67)	3 (100)	7 (100)	4 (100)
Amikacin (5)	4 (100)	2 (67)	2 (67)	7 (100)	4 (100)

Key: R- Resistance, n=number of isolates, A&E = Accident and Emergency ward, MS = Male surgical ward, FS = Female surgical ward, PD = Paediatric ward, LW = Labour ward

Table 2: Antimicrobial Resistant Pattern of *S. typhi* Isolated from fecal samples of patients at ORT, LAB, THE, GOPD and MM

Antibiotic (μ g)	ORT (n = 1) R (%)	LAB (n = 17) R (%)	THE (n = 9) R (%)	GOPD (n = 4) R (%)	MM (n = 3) R (%)
Colistin (30)	1 (100)	14 (82)	9 (100)	2 (50)	0 (0)
Amoxicillin-Clavulanate acid (30)	1 (100)	10 (59)	4 (44)	1 (25)	3 (100)
Piperacillin-Tazobactam (5)	0 (0)	17 (100)	7 (77)	4 (100)	3 (100)
Cefuroxime (20)	0 (0)	11 (65)	7 (77)	4 (100)	3 (100)
Cefepime (15)	1 (100)	9 (53)	9 (100)	1 (25)	2 (67)
Nalidixic Acid (20)	1 (100)	12 (71)	9 (100)	1 (25)	2 (67)
Meropenem (30)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Cefoxitin (30)	1 (100)	15 (88)	5 (55)	4 (100)	3 (100)
Ofloxacin (10)	1 (100)	17 (100)	7 (77)	3 (75)	1 (33)
Amikacin (5)	1 (100)	17 (100)	9 (100)	3 (75)	1 (33)

Key: R- Resistance, n=number of isolates, ORT = Orthopaedic ward, LAB = Laboratory, THE = Theatre, GOPD = General outpatient department, MM = male medical ward

3.2 Discussion

Majority of these isolates were found resistant to first, second and third-generation cephalosporins, while 50% of them were resistant to fourth-generation cephalosporin. Susceptibility of the isolates to monobactams (tazobactam piperacillin), cefamycin (cefotaxime) and carbapenem (meropenem) antibiotics ranged between 10 % and 90%. The higher susceptibility of *Salmonella typhi* to some antibiotics may be due to the under use or over use of these antibiotics in animal rearing. In addition, low susceptibility (11% to 47%) was also observed against the common beta lactam and non-beta-lactam antibiotics. The fluoroquinolone, resistance was considered to be expressed by plasmid mediated quinolone (PMQR) determinants. These findings are in agreement with some recent studies which reported MDR *Salmonella* species and Non-typhoidal *Salmonella* species producing ESBL that were isolated from cattle [12], poultry [13], pigs [14], environment, [15] and humans [16]. The present study further reported a low susceptibility of ESBL *Salmonella* species and *Salmonella typhi* species towards other common non-beta-lactam antimicrobials such as aminoglycosides, fluoroquinolone, tetracycline, etc. This contrasts a Canadian survey, where 62.8% MDR *Salmonella* species isolates were collected from mastitis cows although they detected no ESBLs- producing *Salmonella* species [17].

Research has shown that the resistance to beta-lactam antimicrobials, such as ceftriaxone, is linearly correlated with the lactamase level over a period and resistance to beta-lactams and can be achieved by increasing enzyme levels [10]. Hence, the prolong use or misapplication of cephalosporins may lead to resistance over time.

For the treatment of non-typhoid *Salmonella* infections that are resistant to both ciprofloxacin and ceftriaxone, carbapenems may be the last drug of choice. However, in the present study, susceptibility to meropenem (100%) was also detected. Meanwhile for some of these isolates to be susceptible to a panel of antibiotics tested in the present study, the possibility of acquiring resistance due to horizontal gene transfer or mutation is highly possible.

The high level of tetracycline, amoxicillin resistance in the present study is not surprising as it is an over-the-counter medication used by most farmers in the study area to treat bacterial and tick-borne infections in livestock.. Interestingly, antimicrobial agents such as piperacillin/ tazobactam are not commonly used in animal health and production in Nigeria and this suggests the importance of other sources and mechanism of MDR *Salmonella* Spp. such as: the horizontal transfers of mobile resistance genetic element, imported *Salmonella*-contaminated feed, antimicrobials used in human treatment, livestock and foodstuff, and possibly cross-resistance among the related antimicrobials, the environment, or co-selection of resistances carried on the same DNA element[3, 11, 18].

This study showed high rates of resistant to ofloxacin and nalidixic acid was 33 % -100%. This agrees with several studies carried out where *Salmonella* isolates demonstrated 100% resistance to nalidixic acid [18,19, 20]. These antibiotics belong to the quinolone/fluoroquinolone drug class. Studies have shown quinolone resistance in salmonellae to be as a result of mutations in the DNA gyrase (*gyrA* and *gyrB*) and topoisomerase IV encoding (*parC* and *parE*) genes [3, 18, 19]. Other studies have also reported the presence of plasmid mediated quinolones resistant (PMQR) genes carried by the ESBL-producing plasmid, which facilitates the selection of higher-level resistance to quinolone drugs [3, 21, 22, 23, 24, 25].

Other studies have also suggested that the *in vitro* resistance to ofloxacin and nalidixic acid could be used as a pointer to the actual level of *in vitro* resistance to ciprofloxacin [26, 27] (Campioniet al., 2017; Klemm et al., 2018). According to the Clinical and Laboratory Standard Institute (CLSI) guidelines, the resistance to any antimicrobial agent in the fluoroquinolone drug class has an impact on the resistance of other antimicrobial agents within this drug class. This implies that the resistance observed by the *Salmonella typhi* to nalidixic acid in this study is a pointer to the development of resistance to other members of the fluoroquinolone class of antimicrobial agents such as ciprofloxacin, Levofloxacin and norfloxacin in humans in the Onicha region with time. This is important information given that fluoroquinolones are regarded as the antimicrobial class of first choice for the treatment of severe infections caused by *S. typhi* as well as other pathogenic Enterobacteriaceae in humans. For such patients, this study recommends the use of meropenem for case management as evidence with the 0.0 % resistant proportion reported in this study.

4. CONCLUSION

This study profiled antibiotic resistance, however, the trend of resistant from this study may emerge from injudicious use of antimicrobial agent in both human population, food and veterinary medicine and it is believed that this could be the most probable reason for the selective pressure and consequent resistance to

these drugs. Therefore, this study suggests strict selection and rotation of antimicrobial agents coupled with the continuous monitoring of susceptibility profiles of antimicrobial agents in order to manage the emergence and spread of antimicrobial resistance.

REFERENCES

1. Akram, J., Khan, S. A., Khan, H. A., Gilani, A. S., Akram, J. S., Ahmad, J. F and Mehboob, R (2020). Extensively Drug-Resistant (XDR) Typhoid: Evolution, Prevention and its Management. *BioMedical Research International*, **7**;23-45.
2. John, V and Ashurst, B. W (2018). *Salmonellatyphi*, NCBI, Ed, StatPearls Publishing.
3. Onyenwe, N. E., Nnamani, N. D., Okoro, J.C., Nwofor, C. N and Jesumirhewe, C (2020).Prevalence and gene sequencing of extended spectrum β -lactamases producing *Salmonella enterica* serovar. *Typhi* from South-East Nigeria. *African Journal of Pharmacy and Pharmacology*, **14**(7):192-202
4. Narasanna, R., Chavadi, M and Chandrakanth, K (2018). Prevalence of multidrug-resistant *Salmonella typhi* in typhoid patients and detection of blaCTX-M2 and blaCTX-M9 genes in cefetoxime-mediated extended spectrum β -lactamase-producing *Salmonella typhi* isolates. *Biomedical Research*, **29** (14): 3015-3021
5. Al- Mayahi, F. S and Jaber, S. M (2020) A preliminary study of multiple antibiotic resistance (MAR) and extensively drug-resistant (XDR) of bacterial causing typhoid fever isolated from stool specimens in Al-Diwaniya, Iraq. *Eurasia Journal of Bioscience*, **14**:2369-2378.
6. Mweu, E and English, M (2008). Typhoid fever in Children in Africa. *Tropical Medicine and International Health*, **13**(4):532-540
7. World Health Organization (WHO) (2015). Immunization, Vaccines and Biologicals: Typhoid. <http://www.who.int/immunization/diseases/typhoid/en/>
8. Health Grove (2018). Typhoid Fever in Nigeria: Statistics on Overall Impact and Specific Effect on Demographic Groups. <http://global202disease-burden.healthgrove.com/1/3644/Typhoid-Fever-in-Nigeria>
9. Clinical and Laboratory Standards Institute (CLSI). Performance standards for antimicrobial susceptibility testing; twenty-eighth edition (M100). Wayne, PA: Clinical and Laboratory Standards Institute; 2019.
10. Oke B, Iroha I R, Moses I B , Egwu I H, Elom E, Uzoh C V, Nwode V F, Okpada J O, Agbom J N, Peter I U, Ibeka G U. Killing Rate Kinetics of Commercially Available Brands of Ciprofloxacin and Cefotaxime on Clinical Bacterial Isolates Subjected to *in vitro* Antibiotic Treatments. *Int. J. Pharm. Sci. Rev. Res.* 2020; **64**(2):87-97
11. Uzoije U N, Moses I B, Nwakaeze E A, Uzoeto H O, Otu J O, Egbuna N R, Ngwu J N, Chukwunwejim C R, Mohammed D I, Peter I U, Oke B, Iroha I R. Prevalence of Multidrug-resistant Bacteria Isolates in Waste Water from Different Hospital Environment in Umuahia, Nigeria *Int. J. Pharm. Sci. Rev. Res.* 2021; **69**(2): 25-32.
12. Timofte, D., Maciucă, I. E., Evans, N. J., Williams, H., Wattret, A. and Fick, J. C. (2014). Detection and molecular characterization of *Salmonella* species CTX-M-15 and Non-typhoidal *Salmonella* SHV-12 beta-lactamases from bovine mastitis isolates in the United Kingdom. *Antimicrobial Agents and Chemotherapy*, **58**, 789-94.
13. Kar, D., Bandyopadhyay, S., Bhattacharyya, D., Samanta, I., Mahanti, A. and Nanda, P. K. (2015). Molecular and phylogenetic characterization of multidrug resistant extended spectrum beta-lactamase producing *Salmonella* species and Non-typhoidal *Salmonella* isolated from poultry and cattle in Odisha, India. Infection, genetics and evolution: *Journal of Molecular Epidemiology and Evolutionary Genetics in Infectious Diseases* **29**, 82-90.
14. Xu G., An, W., Wang, H. and Zhang X. (2015). Prevalence and characteristics of extended-spectrum beta-lactamase genes in *Salmonella* species and Non-typhoidal *Salmonella* isolated from piglets with post-weaning diarrhea in Heilongjiang province, China. *Frontier Microbiology*, **6**:1103

15. Gao, L., Hu, J., Zhang, X., Wei, L., Li, S. and Miao Z. (2015). Application of swine manure on agricultural fields contributes to extended-spectrum beta-lactamase-producing *Salmonella* species and Non-typhoidal *Salmonella* spread in Tai'an, China. *Front Microbiology*, 6, 313
16. Gu, B., Pan, S., Wang, T., Zhao, W. and Mei, Y. and Huang P. (2008). Novel cassette arrays of integrons in clinical strains of Enterobacteriaceae in China. *International Journal of Antimicrobial Agents*, 32, 529-33
17. Saini, V., McClure, J.T., Leger, D., Keefe, G. P., Scholl, D. T. and Morck, D. W. (2012). Antimicrobial resistance profiles of common mastitis pathogens on Canadian dairy farms. *Journal of Dairy Science*, 95, 4319-32.
18. Yhiler, N.Y., Basse, B. E., Paul, I., Francis, U. M., Anne, A and Okocha-Ejeko, A (2019). Antimicrobial resistance pattern in *Salmonella enterica* from clinical and poultry sources in Calabar, Nigeria. *Journal of Microbiology and Antimicrobials*, 11(2):5-10.
19. Ye, Q., Wu, Q., Zhang, S., Zhang, J., Yang, G., Wang, J., Xue, L and Chen, M (2018). Characterization of Extended-Spectrum β -Lactamase-Producing Enterobacteriaceae From Retail Food in China. *Frontiers in Microbiology*, 9:1709-1710.
20. Kownhar, H., Shankar, E. M., Rajan, R and Rao, U. A (2007). Emergence of Nalidixic acid-resistant *Salmonella enterica* serovar Typhi resistant to Ciprofloxacin in India. *Journal of Medical Microbiology*, 56:136-137.
21. Oghenevo, O., Basse, B., Yhiler, N., Francis, U and Angela, O (2016) Antibiotic Resistance in Extended Spectrum Beta-Lactamases (ESBLs) *Salmonella* Species Isolated from Patients with Diarrhoea in Calabar, Nigeria. *Journal of Clinical Infectious Disease Practice*, 1: 107-109.
22. Jacoby, G. A., Strahilevitz, J and Hooper, D. C (2014). Plasmid-mediated Quinolone Resistance. *Microbiology Spectrum*, 2(6):20-13.
23. Riyaz, A. A. A., Perera, V., Sivakumaran, S and de Silva, N (2018). Typhoid Fever due to Extended Spectrum β -Lactamase-Producing *Salmonella enterica* Serovar Typhi: A Case Report and Literature Review. *Case Reports in Infectious Diseases*, 12(23):34-66.
24. Kongsoi, S., Yokoyama, K., Suprasert, A., Utrarachkij, F., Nakajima, C., Suthienkul, O and Suzuki, Y (2015). Characterization of *Salmonella Typhimurium* DNA gyrase as a Target of Quinolones. *Drug Testing and Analysis*, 7:714-720.
25. Carfora, V., Alba, P., Leekitcharoenphon, P., Ballarò, D., Cordaro, G., Di Matteo, P., Donati, V., Ianzano, A., Iurescia, M., Stravino, F., Tagliaferri, T., Battisti, A and Franco, A (2018). Colistin Resistance Mediated by mcr-1 in ESBL-Producing, Multidrug Resistant *Salmonella* *Infantis* in Broiler Chicken Industry, Italy (2016–2017). *Frontiers in Microbiology*, 9:18-80.
26. Campioni, F., Souza, R. A., Martins, V. V., Stehling, E. G., Bergamini, A. M. M and Falcão, J. P (2017). Prevalence of gyrA Mutations in Nalidixic Acid-Resistant Strains of *Salmonella* *Enteritidis* Isolated from Humans, Food, Chickens, and the Farm Environment in Brazil. *Microbial Drug Resistance*, 23:421-428.
27. Klemm, E. J., Shakoor, S., Page, A. J., Qamar, F. N., Judge, K., Saeed, D. K., Wong, V. K., Dallman, T. J., Nair, S., Baker, S., Shaheen, G., Qureshi, S., Yousafzai, M. T., Saleem, M. K., Hasan, Z., Dougan, G and Hasan, R (2018). Emergence of an Extensively Drug-Resistant *Salmonella enterica* Serovar Typhi Clone Harboring a Promiscuous Plasmid Encoding Resistance to Fluoroquinolones and Third-Generation Cephalosporins. *Journal of Molecular Biology*, 9:105-18.

UNDER PEER REVIEW