

Monitoring and Assessment of Water Quality of Iril River, Manipur, India

Abstract:The present study can be concluded that the analysis of the physio chemical parameters of Iril River revealed that most of the values are within the permissible limit of water quality standards. Water quality is suitable as per Standards norm of BIS and CPCB based on the determined values of pH, EC, TDS, DO, BOD, Chloride, Total Hardness, Total Alkalinity and Sulphate was monitored during the period of January 2023 to May 2023. The value of water pH recorded between 7.4- 8.34, EC ranged between 179.6 $\mu\text{s/cm}$ -254 $\mu\text{s/cm}$, TDS ranged between 90.2 mg/l-126.8 mg/l, dissolved oxygen between 6.0-7.7 mg/l, BOD between 3.0 mg/l-3.9 mg/l, chloride between 28.9 mg/l-36.9 mg/l, total hardness between 112mg/l-170 mg/l, total alkalinity between 115 mg/l-174 mg/l and sulphate between 14.4 mg/l-39.3 mg/l. Hence, these rivers and hold to provide uninterrupted supply of drinking water to Iril River of Manipur without any major treatment.

Keywords: *Physio-chemical, Iril River, Water quality, Pollution*

1. Introduction:

Water is the prime natural resource. Water is the first need of all vital life processes, hence called “Liquid of Life.” According to Greek medieval philosophy, matter consists of four elementary substances namely water, air, stone and fire. Indian medieval also opines that the matter is composed of five Panch Mahabhut (Five elementary substances) namely water, air, light (fire), earth (stone) and sky (space). Among the all known planets only earth is blessed by these all five elements (Yadav *et al.*, 2010).

Iril river with its basin lying between the Latitude 24°40'N to 25°25'N and the Longitude 93°55'E to 94°20'E passes mainly through sub-urban areas of Imphal-east district suffer pollution due to urbanization and agricultural run off, domestic waste, municipal waste, sewage and over fishing into the river. The wastes running through the water may be hazardous and infectious (Jameel, 1998). The total length of the Iril river from its head to the point of its confluence with the Imphal

Comment [AM1]: I think $\mu\text{s/cm}$ instead of ms/cm ?

Formatted: Highlight

River is 144.5 km. The River originates from the south eastern slope of Hougdu Ching range in Ukhrul district at a height of 2473 m above MSL and flows through the hill tract of Ukhrul and Senapati districts and descends to the almost tracts of Imphal and Thoubal districts in the Manipur valley. It debounces its water into Manipur river (known as Imphal river in its upper part before it confluence with Khuga river) a tributary of the Chindwin-Irrawady system. (Romeo and Asha, 2009)

Many of the major problems that humanity is facing in the twenty-first century are related to water quantity and/or water quality issues. These problems are going to be more aggravated in the future by climate change, resulting in higher water temperatures, melting of glaciers, and an intensification of the water cycle, with potentially more floods and droughts. With respect to human health, the most direct and most severe impact is the lack of improved sanitation, and related to it is the lack of safe drinking water, which currently affects more than a third of the people in the world (Schwarznbach *et al.*, 2010).

Water quality is defined in terms of the chemical, physical and biological contents of water. The water quality of rivers and lakes changes with the seasons and geographic areas, even when there is no pollution present. Water quality guidelines provide basic scientific information about water quality parameters and ecologically relevant toxicological threshold values to protect specific water uses. Important physical and chemical parameters influencing the aquatic environment are temperature, rainfall, pH, salinity, dissolved oxygen, and carbon dioxide. Others are total suspended and dissolved solid, total alkalinity and acidity and heavy metal contaminates (Lawson, 2011).

The aim of the study is to analyze physio-chemical properties of water of Iril River of Manipur. Iril River is one of the main sources of water for the people of Manipur. It has a large population of endangered indigenous fish called Ngaton and Meitei Sareng.

2. Material And Methods:

The experiment was conducted during month of January to May 2023 at 10 different sites of Iril River, the name of sites are Sawombung(S1), Kangla(S2), Moirang Kampu(S3), Top Khongnangkong(S4), Naharup(S5), Bamon Kampu(S6),

Iribung(S7), Keirao(S8), Arapti(S9) and Lilong(S10) respectively are given in the Table: 1. The samples was collected thrice in a month and collected in the morning hours between 5 am to 11am, in clean plastic bottles without any air bubbles and closed tightly after collection and labeled in the field and preserved for physio-chemical analysis in laboratory. The water sample was immediately be brought into the laboratory for the estimation of all the physio chemical analysed in the laboratory of Directorate of Environment and Climate Change, Government of Manipur. All the parameters were analysed as per the standards method of APHA/AWWA (1991).

Sampling Sites:

The river water samples wascollected at different site starting fromSite1 at reference sites and covered upto to river stretch 30.2km and details of populationnear by the river stretch also given in the tabular form in Table 1.



Table 1: Location details of sampling site, distance and population at different site

| SITE Notation | SITE | Distance from S1 | Latitude & Longitude | Population as per 2021 census |
|---------------|--------------------|------------------|------------------------|-------------------------------|
| S1 | Sawombung | 0 | 24.8727° N, 94.0146° E | 111,287 |
| S2 | Kangla | 3.6 | 24.8459° N, 93.9934° E | 2075 |
| S3 | Moirang Kampu | 11.6 | 24.7959° N, 93.9808° E | 1626 |
| S4 | Top Khongnangkhong | 13.6 | 24.8097° N, 93.9591° E | 2474 |
| S5 | Naharup | 15.6 | 24.7584° N, 93.9687° E | 2394 |
| S6 | Bamon Kampu | 19.5 | 24.7706° N, 93.9772° E | 3401 |
| S7 | Iribung | 21.4 | 24.8063° N, 93.9481° E | 1014 |

| | | | | |
|-----|--------|------|---------------------------|--------|
| S8 | Keirao | 24.7 | 24.7479° N, 93.9796° E | 6489 |
| S9 | Arapti | 28.9 | 24.6908° N, 93.9483° E | 1824 |
| S10 | Lilong | 30.2 | 24.6821° N, 93.9452° E | 34,100 |

UNDER PEER REVIEW

Map Of Study Area



Comment [AM2]: Make a good location map with outline, scale etc...

Map 1: Map of the study area along with sampling sites.

3. Result and Discussion:

The results of physico-chemical parameters of the Iiril River are given in the Table No. 2. The value of water pH recorded between 7.4- 8.34, EC ranged between 179.6 ms/cm-254 ms/cm, TDS ranged between 90.2 mg/l-126.8 mg/l, dissolved oxygen between 6.0-7.7 mg/l, BOD between 3.0 mg/l-3.9 mg/l, chloride between 28.9 mg/l-36.9 mg/l, total hardness between 112mg/l-170 mg/l, total alkalinity between 115 mg/l-174 mg/l and sulphate between 14.4 mg/l-39.3 mg/l.

pH: pH is a measure of how acidic/basic water is, a measure of the hydrogen ion concentration. pH during study period ranged from 7.4- 8.34. The maximum value of 8.34 was recorded at Iirilbung(S7) during the month of February. The value recorded is within BIS/CPCB standard and which was alkaline in nature. The minimum value of 7.4 recorded at Keirao(S8) in January. The pH value ranges from 6.5 – 8.5 are prescribed for drinking and pH ranges from 6.5 – 9.0 are most suitable for fish production as proposed by ISI. (Surajel *et al.*, 2018)




Fig.1: pH value at different sites of Iiril River during sampling months

Electrical Conductivity: The EC is a measurement of capacity of water to transmit electric current in water bodies. EC during study period ranged between 179.6 ms/cm-254ms/cm. The maximum EC was 254 ms/cm recorded at Lilong(S10) during the month of March and minimum was recorded 179.6 ms/cm at Top Khongnangkhang(S4)

during the month of January 2023. The EC value of all the sites of Iril River were under the permissible limit given by the Standards. Other all the reference that the value of conductivity higher in the mouth of rainy season due to higher value of dissolved solids.(**Bhatt *et al.*, 1999;Romeo *et al.*, 2010**)




Fig.2: EC (µs/cm) of Iril River at different sites of month

Formatted: Highlight

Total Dissolved Solids:Total dissolved solids is the term used to describe the inorganic salts and small amounts of organic matter present in the water. TDS value during investigation period was ranged between 90.2 mg/l-126.8 mg/l. The maximum TDS was recorded at 126.8 mg/l at Lilong(S10) during month of March and minimum was observed at 90.2 mg/l at site Top Khongrangkhong(S4) during the month of January. The TDS value of all the sites of Iril River were under the permissible limit. The palatability of drinking water with a TDS level less than 600 mg/L is generally considered to be good.(**Bujar *et al.*,2022**)




Fig.3: TDS(mg/l) of Iril River at different sites of month

Dissolved oxygen:The DO content in the water body indicates the physical and biological processes prevailing in the water and is very important because it directly affects the survival and distribution of flora and fauna in an ecosystem. The dissolved oxygen value during the investigation period was observed between 6.0 to 7.7 mg/l. The maximum value was recorded at site Bamon Kampu (S6) during the month of March and the minimum value was observed at Moirang Kampu (S3) and Top Khongnangkong (S4) during the month of April. Other author reference that the highest value of dissolved oxygen in the river may be due to heavy rainfall and few of water. (Suraj, 2018).




Fig.4: DO of Iril River at different sites of month

Biochemical Oxygen Demand(BOD₃): BOD value was recorded between 3.0 mg/l-3.9 mg/l during the study periods. The maximum BOD₃ was observed 3.95 mg/l at site Moirang Kampu(S3) during the month of April, Bamon Kampu(S6) and Iribung(S7) during the month of May and minimum was 3.04 mg/l recorded at Kangla(S2) during the month of April. The BOD value of all the sites of Iril River were under the permissible limit. The variation of BOD value due to domestic waste water discharge into the river affect the quality of water. The similar result was found by the author **Singh *et al.* (2007)**




Fig.5: BOD of Iril River at different sites of month

Chloride: The value of chloride during estimation period ranged between 28.9 mg/l-36.9 mg/l. Chloride in water also act as an indicator of pollution by sewage. Higher value of chloride was recorded in Lilong(S10) during month of March and lower value of chloride was recorded in Bamon Kampu(S6) during the month of April, which may be due to high domestic sewage discharge at river water. The chloride concentration in the river water below 250mg/l and safe for uses and fall under the permissible limit. Similar finding was also reported by **(Patel and Sinha1998)**.




Fig.6:Chloride of Iril River at different sites of month

Total Hardness:Total hardness of water is generally caused by the nutrients like calcium and magnesium ions present in water. During study period the total hardness of water was estimated between 112mg/l-170 mg/l. The maximum value was recorded at Lilong(S10) during the month of March and minimum was recorded at Naharup(S5) during month of May. The value was non-uniform throughout the season. Similar result was conducted by other author(**Singh *et al.*,2008; Shrestha and Nath, 2019; Khan and Nath, 2014**)




Fig.7:Total Hardness of Iril River at different sites of month

Total Alkalinity: The total alkalinity of water neutralized a strong acid which may be due to the presence of carbonates, hydroxide and bicarbonates. The value of alkalinity during the study period was ranged between 115 mg/l-174 mg/l. The maximum value was recorded at Kangla(S2) in March may be due to sewage decomposition and use of detergent by the surrounding and minimum value was recorded at Naharup(S5) and Bamon Kampu(S6) itself in January. Higher Alkalinity was possible due to greater human activities including washing and bathing. (Sarita *et al.* 2019) revealed that the concentrations of total alkalinity were observed higher in the pre-monsoon.




Fig.8: Total Alkalinity of Iril River at different sites of month

Sulphate: The value of sulphate during study period was between 14.4 mg/l-39.3 mg/l. The maximum sulphate was 39.3 mg/l recorded at Sawombung(S1) during the month of May and the minimum was 14.4 mg/l recorded at Arapti(S9) during month of January. The sulphate value of all the sites of Iril River were under the permissible limit given by BIS and CPCB. (Singh *et al.* 2007 and Rajiv *et al.* 2017) found concentration of sulphate were low during monsoon and higher during premonsoon.




Fig.9: Sulphate of Iiril River at different sites of month

4. Conclusions:

All the parameters are under the acceptable limit. The present study can be concluded that the analysis of the physio chemical parameters of Iiril River revealed that most of the values are within the permissible limit of water quality standards. Water quality is suitable as per Standards norm of BIS and CPCB.

The water quality is suitable for drinking and aquacultures activities. Even though the water quality manageable for fish culture, the quality of the water is degrading gradually due to the anthropogenic activities of the human population. There is need to monitor periodically in order to maintain the changes in the water system as the anthropogenic activities is increasing day by day. So, the State Government and other NGOs must address the issue in time to prevent further degradation of the river.

References

1. **Albona, S., Berat, D., FS, S., and Hazir., S. (2020).** Assessment of Water Quality of Sitnica River by Using Water Quality Index. *Rasayan J. Chem.*13(1), 146-159(2020) Vol.13,pp.146 - 159
2. **Anita, B., Pooja, D., and M.P, G. (2016).** Impact of Mass Bathing and Activities on Water Quality Index of Prominent Water Bodies: A Multilocation Study in Haryana, India. *International Journal of Ecology.* Vol.2016, Article ID 2915905,pp.1-8
3. **Bhardwaj (2005)** Water quality monitoring in India- Achievements and Constraints, *Journal of Environmental Biology,* pp. 1-8.
4. **Bujar,H., Pajtim,B., Fidan,F. and Shkumbin, S.(2022).** Application of Water Quality Index for The Assessment the Water Quality in River Lepenci. *Ecological Engineering & Environmental Technology.*(2022), vol.23,pp. 188-200
5. **Dede, O.T., Telci, I.T. and Aral, M.M., (2013).** The use of water quality index models for the evaluation of surface for Kirmir basin, Ankara, Turkey. *Water qual expom health,*vol.8,pp:41-56.
6. **Dhanraj, K. and Asha, G. (2015).** Sediment Analysis from Imphal River, Manipur. *Bioglobia* <https://www.researchgate.net/publication>. Vol.2 (1): 2015, pp.49-55.
7. **Fakayode, S. O. (2005).**Assessment of the Risk of Toxicity of Fish in the River Sô by Contamination in Heavy Metals (Southern Benin, West Africa).*Journal of Geoscience and Environment Protection,* Vol.6 No.12, pp.202-215.
8. **Herojit, N., Maibam,P. and Sujata, S. (2021).** Physiochemical Assessment and Water Quality of Surface Water in Chandel and Tengnoupal Districts, Manipur for Domestic and Irrigational Uses.*International Research Journal of Engineering and Technology (IRJET)*Volume: 08, pp.732-738.
9. **Khan, S. and Nath, S. (2014)** Physio chemical Analysis of River Ganges at Mirzaur in Uttarpradesh, India, *IOSR Journal of Applied Chemistry (IOSR-JAC),* vol.7, Issue 12, pp.61-67.

10. **Kumar, R., Grover, A.S. and Wats, M. (2018)** Assessment of water quality of Lakes in Haryana, India, *International Journal of Recent Scientific Research*,vol. 9:(7), pp.27831-27835
11. **Lawson, E.O. (2011)** Physio-chemical parameters and heavy metal contents of water from the Mangrove swamps of Lagos Lagoon, Nigeria. *Advances in biological research*,vol.5(1),pp:08-21
12. **Liou, S.M., Lo, S.H., and Wang, S.H. (2003).** A generalized water quality index for Taiwan. *Environment monitoring and assessment*,vol. 96,pp:35-52.
13. **Mariana, M., Fachriah, W. and Arini,M. (2022).**Water Quality Index Assessment Methods for Surface Water: A Case Study of the Citarum River in Indonesia. www.cell.com/heliyon ,vol.8 pp.1-10.
14. **Patel, N. K. and Sinha, B.K. (1998)** Study of Pollution load in the ponds of Burla area near Hirakund Dam of Orrisa, *Journal of Environment & Pollution.*, Vol.5, pp. 761-771.
15. **Rajiv, D.K., Sarojini, D.B., Suganthi, K. and Munisamy, G. (2017)** Development of a water quality index (WQI) for the Loktak lake of India. *Appl water sci*,vol.7, pp: 2909-2918.
16. **Ram, P.S., Satyendra, N., Subhash, C.P. and Arvind, K.N. (2007)** Selection of suitable aggregation function for estimation of aggregate pollution index for river Ganges in India. *Journal of Environmental Engineering.* Vol.134, pp:689-701.
17. **Ravindra, G.J., Manik, A., Promod, P.N. and Utkash, M.(2022).** Analysis of season variation in surface water quality and water quality index (QWI) of Amba River from Dolvi Region, Maharashtra, India. *Arabian Journal of Geosciences.* Vol. (2022) 15:1261, pp.1-14
18. **Rizwan, R. and Gurdeep, S. (2010).** Assessment of river water quality status by using quality index (WQI) in industrial area of Orrisa. *International Journal of Applied Environmental Sciences.*Vol.5(2),pp.305-310.
19. **Romeo, S.M., and Asha.G. (2009).** Water Quality Status of Iril River, Manipur. *Journal of Current Sciences.* Vol.14 (1): 173-180.
20. **Romeo, S.M., and Asha. G. (2010).**Seasonal variations in certain Physio-chemical parameters of Imphal, Iril and Thoubal rivers from Manipur system, India. *Eco.Env. & Cons.*vol.16(2);pp 197-207.

Formatted: Indent: Left: 0", Hanging: 1.48 ch

21. **Rrishna, K. Y., Neha, G., Vinit, K. and Sudarshana, S.(2014).** Water Quality Assessment of Pahuj River using Water Quality Index at Unnao Balaji, M.P., India. *International Journal of Sciences: Basic and Applied Research (IJSBAR)*. Vol.19, No 1, pp 241-250.
22. **Sanjoy, S., Umesh, K. and Pankaj, M. (2019).** Water quality assessment of a tropical river using water quality index (WQI), multivariate statistical techniques and GIS. *Appl Water Sci*, vol. 9, 168, pp.1-21.
23. **Schwarzenbach, R.P., Thomas, E., Thomas, B.H., Gunten, U.V. and Wehrli, B. (2010)** Global water pollution and human health. *Annual review of environment and resource*, vol. 35:,pp.109-136.
24. **Sener, S., Sener, E. and Davaraz, A.(2017).** Evaluation of water quality index (WQI) methods and GIS in Akus river (SW-Turkey). *Science of the total environment*, 584-585, pp.131-144.
25. **Shrestha, S. and Satyendra, N. (2019).** Assessment of Gangariver water quality in Allahabad. *Indian Journal of Environmental Protection*, Vol. 39, pp. 770-775.
26. **Singh, M.R., Gupta, Asha and Beeteswari, Kh. (2010).** Physio-chemical Properties of Water Samples from Manipur River System, India. *Journal of Applied Sciences: Environmental and Management*.Vol.14(4),pp85-89.
27. **Sivaranjani, S., Amitava, R., and Samrath, S. (2015).** Water Quality Assessment with Water Quality Indices. *International Journal of Bio resource Science Vol (2)*,pp.85-94
28. **Suraj, D. W. and Rajmani, S.Kh. (2018).** Estimation of Water Quality of Imphal River, Manipur, India. *International Journal of Recent Scientific Research*. Vol.9, Issue, 5(J), pp. 27187-27190.
29. **Yadav, A.K., Khan, P. And Sharma, S.K., (2010).** Water quality index assessment of groundwater in Todaraisingh Tehsil of Rajasthan State, India- A green approach. *E-Journal of chemistry*, vol.7(S10), pp.5428-5432.
30. **Zayed, M., Fariha, T. A., Md, R. B.H. and Synthia, P. M. (2021).** Water Quality Index (WQI) for Evaluation of the Surface Water Quality of Bangladesh and Prediction of WQI from Limited Parameters. *Journal of Environment Protection and Sustainable Development*. Vol. 7, No. 2,pp.1-24.