

### Original Research Article

## Performance of integrated nutrient management and Foliar spray of micronutrients on growth, yield attribute and yield of wheat (*Triticum aestivum* L.)

### ABSTRACT

A field experiment was conducted during rabi seasons of 2021-22 at Students' Instructional Farm of Chandra Shekhar Azad University of Agriculture and Technology Kanpur, (U.P.) The experiment was carried out in SPD (Split Plot Design) with fifteen treatment combination were replication thrice. fifteen treatment treatments combination consisted of three organic manure + RDF nutrient source (Control (100% RDF), 100 % RDF + Vermicompost @ 2.5t/ha, and 100 % RDF + FYM @ 5t/ha) put under main plot and five foliar spray of micronutrient (control (100% RDF, Foliar application ZnSO<sub>4</sub> @ 0.25% at tillering and booting stage, spray of Nano Zinc @ 0.5% spray at tillering and booting stage, Foliar application of FeSO<sub>4</sub> @ 0.25% at tillering and booting stage and M<sub>5</sub>-Foliar application of ZnSO<sub>4</sub> @ 0.25% + FeSO<sub>4</sub> @ 0.25% at tillering and booting stage) allotted under sub plot. The study on yield attribute and yield of wheat. The result showed that the maximum increase growth parameters, yield attribute and yield of wheat viz plant height (cm), number of tillers/m<sup>2</sup>, and leaf area index and number of grain/spike, number spike/m<sup>2</sup>, total grain weight spike (g) and test weight (1000 grain weight) (g) and grain yield (t/ha) and straw yield (t/ha) recorded significantly higher with F<sub>2</sub>- 100 % RDF + Vermicompost @ 2.5t/ha treatments. Among the sub plot, M<sub>5</sub>- foliar application of ZnSO<sub>4</sub> @ 0.25% + FeSO<sub>4</sub> @ 0.25% at tillering and booting stage produced significantly maximum highest grain yield (6.11 t/ha) and straw yield (7.94 t/ha)

**Keywords:** *FYM, Iron sulphate, Nano-Zinc, Vermicompost, Wheat and Zinc sulphate*

### 1. Introduction

Wheat (*Triticum aestivum* L.) one of the most important cereal crops in the world. comes under the family "Poaceae" and genus "Triticum" It is a crop of temperate climate with winter season crop and hot summer being very favorable for its growth and maturity. Among the cereal wheat is the one of the most important sources of energy, carbohydrate and protein for the world population and wheat production is considerably important for food security. It's grown under different agro climate conditions. It occupies 220.29 million ha. area with would the production of 780.59 million tones and productivity is 3390 kg ha<sup>-1</sup>. rank second in production of wheat in the world after China. Important wheat producing state are Uttar Pradesh (30 million tons), Punjab (16.4 million tons), Haryana (11.6 million tons), with area 9.75 m/ha, 3.59 mha, 2.52 mha, respectively. Uttar Pradesh has first position in area and production of wheat in India. Although productivity of wheat (2661 kg ha<sup>-1</sup>) is still less than the national average (Anonymous, 2022-23). Due to continuous use of high doses of synthetic chemical fertilizer without use

Commented [tt1]: 2021-2022

Commented [tt2]: (Ministry of Agriculture & Farmers Welfare, 2023)

of organic source and intensive system of cultivation practices, there is change in soil physico-chemical condition and nutritional deficiencies are common in general and specifically with micronutrients, which are very essential for plants. In this endeavor proper blend of organic and inorganic fertilizer is important not only for increasing yield but also for sustaining soil health (Weber *et al.*, 2007 and Pullicinoa *et al.*, 2009). Judicious use of FYM improves soil physical, chemical and biological properties and improves crop productivity. It also helped in arresting the emerging deficiencies of macro and micro nutrients favorable and achieving economy and efficiency in fertilizers use. In view of shrinkage of land resources for cultivation, short supply and escalating cost of chemical fertilizers, environmental pollution and ill effects on soils, animals and human health, there is need to use farm yard manure with nano fertilizers for achieving the objective of environmentally and ecologically sustainable agriculture. The present investigation was, therefore, carried out to study the effect of FYM on yield attribute and yield, nutrient uptake and quality of wheat. (Kumar *et al.*, 2015). The vermin-compost is the excreta of the earth worms, and farm yard manure is the product of decomposition of the liquid and solid excreta of the livestock. As the source of nutrients, these materials may also reduce the cost of fertilizer input and improve the soil physio-chemical properties. It is imperative to develop strategy to use organic manures up to their maximum potential with proper technology to meet the shortage of fertilizers and for sustaining soil fertility on long term basis. Since organic manures are known to improve the soil health increase water retention and supply of most of the nutrients to the plants, hence the use of these in the form of farm yard manure and vermin composting will certainly boost up crop production from the soil. The organic material such as vermin compost and farm yard manure available in abundance at nominal cost in eastern Uttar Pradesh may be the alternative for the organic manures (Singh *et al.* 2020) which has worsened the quality of cereal staple in terms of Zn. Similar is the situation for Fe deficiency in cereals. In modern cultivated wheat, the seed concentration of Zn and Fe were found less than in the wild wheat (Cakmak *et al.*, 2004). Zinc and Fe are important minerals required for various metabolic functions. It is obvious for both animals and plants. Zinc is responsible for protein synthesis, gene expression, proper growth and immune system. Physically Zn deficiency is manifested as stunting, common health problem in children like diarrhea, low birth weight, high rate of infection, skin lesions and impaired wound healing (Mutangadura 2004). Fertilizers are necessary for enhancing productivity in crops especially in wheat, maximum use of macro nutrients and low use of micronutrients leading to an imbalance in soil fertility. A balance fertilization program with macro nutrients and micronutrients in plant nutrition is very essential in the high production of crops with good quality products, so there is a need balance use of fertilizers and agronomic procedures are needed to increase yield of this crop. The function of micronutrients is vital role in crop nutrition for improved yield attribute, growth and development and quality of wheat crop. (Saeed *et al.*, 2012).

Commented [tt3]: Does not appear in the references.

Commented [tt4]: (Weber *et al.*, 2007; Pullicinoa *et al.*, 2009)

Commented [tt5]: Does not appear in the references.

## 2. Materials and Methods

### 2.1. Experimental site:

The experiment was conducted during the rabi season 2021-22 at Students' Instructional Farm of Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, which is situated in the alluvial tract of Indo - Gangetic plains in central plain zone of Uttar Pradesh. A field experiment was conducted consecutive two season of 2021-2022-23 at Students' Instructional Farm Department of agronomy. College of agriculture, Chandra Shekhar Azad University of Agriculture and Technology Kanpur, (U.P.). The soil type and fertility status were determined by the mechanical and chemical analysis of the soil. In order to ascertain physico-chemical properties of the experimental soil, primary soil samples were drawn randomly up to 15cm depth from different spots of the entire experimental area. The soil of the experimental field was sandy loam in texture, well drained, plane topography, slightly saline in nature having initial values of pH (8.00), EC ( $0.29 \text{ dsm}^{-1}$ ), low in organic carbon (0.35%), low in available nitrogen ( $192.50 \text{ kg ha}^{-1}$ ), medium in phosphorus ( $15.75 \text{ kg ha}^{-1}$ ), Potash ( $173.51 \text{ kg ha}^{-1}$ ). Iron ( $1.75 \text{ mg kg}^{-1}$ ) and Zinc ( $61.15 \text{ mg kg}^{-1}$ ). respectively,

## 2.2. Experiment Details

A field experiment was laid out in Split Plot Design. There were fifteen treatments combinations consisting of three organic manure as a main treatment viz: F<sub>1</sub>- Control (100% RDF), F<sub>2</sub>- 100 % RDF + Vermicompost @ 2.5t/ha, and F<sub>3</sub>- 100 % RDF + FYM @ 5t/ha, and five Foliar application of micronutrient practices were applied as sub treatments viz: M<sub>1</sub>- control (100% RDF), M<sub>2</sub>- Foliar application @ 0.25% ZnSO<sub>4</sub> at tillering and booting stage, M<sub>3</sub>- Nano Zinc @ 0.5% spray at tillering and booting stage, M<sub>4</sub>- Foliar application @ 0.25% FeSO<sub>4</sub> at tillering and booting stage and M<sub>5</sub>- Foliar application @ 0.25% ZnSO<sub>4</sub> + 0.25% FeSO<sub>4</sub> at tillering and booting stage The treatments were replicated three times.

## 2.3. Geography and climate:

This zone has semi-arid climatic condition having alluvial fertile soil. The average relative humidity during the experimental season fluctuated between 44.4 % to 85.8 % and 47 % to 96% During 2021 to 2022 respectively. During the crop growing period, the mean weekly highest and lowest total rainfall recorded ranging from 0.0 mm to 23.5 mm and 0.0 mm to 39.2 mm and evaporation ranged from 7.6 to 26.0 mm/day<sup>-1</sup> and 8.8 to 25 mm/day<sup>-1</sup> during 2021 to 2022, respectively. The mean weekly total evaporation mm/day<sup>-1</sup> recorded ranging from 326.4 to 340.4 mm/day<sup>-1</sup> during 2021 to 2022, respectively. The weekly mean wind velocity during the year 2021-22 and 2022-23 experimental season was between 1.2 to 6.2 km and 1.1 to 6.0 km respectively.

## 2.4. Agronomic practices

The pre sowing irrigation was applied 8-10 days before sowing of wheat for optimum level

of moisture for field preparation for good germination, The seed of wheat crop variety K-1006 was selected for these studies. Clean and healthy seed @ 120 kg/ha were taken and sown at shallow furrows of 5 cm deep with 22.5 cm row spacing on both rabi seasons of 2021-22 and 2022-23.

Commented [tt6]: 2021-2022 and 2022-2023

#### 2.4.1 Manure and Fertilizers application

The experimental crop was uniformly fertilized-according to treatments, Farm yard manure (FYM) 5 t/ha, and Vermicompost (VC) 2.5 t/ha applied Before 15-20 days of sowing. The basal application of fertilizers @ 120 kg N and full doses  $P_2O_5$ , &  $K_2O$  was given and foliar application of micronutrients @ 0.25%  $ZnSO_4$ , Nano-Zinc @ 0.25%, @ 0.25%  $FeSO_4$  and foliar application @ 0.25%  $ZnSO_4$  + @ 0.25%  $FeSO_4$  at tillering and booting stage were done according to treatments and remaining nitrogen was top dressed in two splits.

#### 2.4.2 Irrigation: -

Frist irrigation (pre -sowing) was applied in experiment field to prepare the field and sow the crop at optimum soil moisture before the seed sowing. Irrigation in standing crop at different stage of crop were not applied due to abundant rainfall which provide water requirement of the crop. Irrigation was needed in the both years during the crop period. and after sowing of the crop, all five-six irrigations were given as per need of the crops coincide with critical stages of the crop during both the years.

#### 2.5. Observation Recorded:

The observations were recorded on plant height (cm), number of tillers/ $m^2$  and leaf area index as well as growth characters Plant height of the longest tiller was measured from five tagged plants from the base of the plant to highest terminal point by a meter scale. Average of five plant height at harvest was taken for analysis of plant height. Number of tillers was recorded by using meter row from three places in each plot at harvest and average of three places was taken for analysis of number of tillers  $m^{-2}$  and Plants from 22.5 cm row length removed from two places from the second row on either side in each plot were used for recording leaf area. The leaves were separated from the base of lamina. Leaf area was measured at 30 days intervals till the senescence of crop with the help of leaf area meter and the total number of grains per spike was counted from the spikes selected for measuring spike length; the grain was separated from spikelet's and the total numbers of grain/spike counted, A composite sample of grains was taken from the produce of each plot and thousand no of grains was counted with the help of automatic seed counter and the weight of 1000 grains expressed in gram, The grain yield per net plot and direction wise was weighed in kilograms. The yield in ton per hectare was calculated by multiplying the plot yield. The yield was used for statistical analysis and Straw yield was

worked out by subtracting the grain yield from total biological yield of net plot area and expressed in t/ha

## 2.6 Statistical analysis

using the F-test as per the procedure given by Gomez and Gomez (1984). Critical difference values at  $P=0.05$  were used to determine the significance of differences between treatment means.

## Result and discussions

### 3.1 Growth characters

The data plant height, number of tillers/m<sup>2</sup>, leaf area index during both the years of experimentation and data were pooled and presented in Table. 1. It is clear from the data the maximum tallest plant height, number of tillers/m<sup>2</sup> and leaf area index was recorded under the treatments F<sub>2</sub>- 100 % RDF + Vermicompost @ 2.5t/ha at 60 DAS, followed by F<sub>3</sub>- 100 % RDF + FYM @ 5t/ha. Both F<sub>2</sub> and F<sub>3</sub> treatments were significantly superior over the Control (100% RDF) during both the year as well as pooled basis data. Similar type of experimental finding reported by **Mohan et al., (2018)**, **Bairagya et al., (2019)** and **Tiwari et al., (2022)**. Among micronutrient foliar application M<sub>5</sub>- Foliar application of ZnSO<sub>4</sub> @ 0.25% + FeSO<sub>4</sub> @ 0.25% at tillering and booting stage recorded tallest plant height (cm), number of tillers/m<sup>2</sup> which was at par with M<sub>3</sub>- Nano Zinc @ 0.5% followed by, M<sub>2</sub>- foliar application of ZnSO<sub>4</sub> @ 0.25% M<sub>4</sub>- foliar application of FeSO<sub>4</sub> @ 0.25% at tillering and booting stage 60 DAS. Whereas minimum plant height (cm) and number of tillers/m<sup>2</sup> was recorded in control (No spray) for both the years as well as on pooled basis. Highest growth characters under these treatments may be due to micronutrient spray which gave better environment for crop growth and development. It confirms the conclusion drawn by **Solanki et al., (2020)** and **Akhtar et al., (2022)**. However, the maximum leaf area index recorded in M<sub>5</sub>- Foliar application of ZnSO<sub>4</sub> @ 0.25% + FeSO<sub>4</sub> @ 0.25% followed by M<sub>3</sub>- Spray of Nano Zinc @ 0.5% spray followed by M<sub>2</sub>- Foliar application of ZnSO<sub>4</sub> @ 0.25% at tillering and booting stage and M<sub>4</sub>- foliar application of FeSO<sub>4</sub> @ 0.25% at tillering and heading stage at harvest stages. The minimum leaf area index was recorded in control (No spray) 60 DAS, during both the years as well as pooled basis. These results corroborated with the findings of **Dimkpa et al., (2020)**.

Commented [tt7]: Mohan et al. (2018)

### 3.2 Yield attribute and yield

Data on Number of grains/spikes, Test weight (1000 grain wt.) (g) grain yield (ton/ha) and straw yield (ton/ha) was presented in Table. 2. Number of grains/spike, Test weight (1000 grain wt.) (g) grain yield (ton/ha) and straw yield (ton/ha) was recorded highest under the treatments F<sub>2</sub>- 100 % RDF + Vermicompost @ 2.5t/ha at harvest followed by F<sub>3</sub>- 100 % RDF + FYM @ 5t/ha. Both were significantly superior over the Control (100% RDF) during both the year as well as pooled basis data. These results corroborated with the findings of **Singh et al., (2020)**, **Fazily et al., (2021)** and **Loura et al., (2022)**. Among the foliar application of micronutrient, M<sub>5</sub>- Foliar application of ZnSO<sub>4</sub> @ 0.25% + FeSO<sub>4</sub> @ 0.25%

recorded highest Number of grains/spikes, Test weight (1000 grain wt.) (g) grain yield (ton/ha) and straw yield (ton/ha) was presented in Table.2. Number of grains/spikes, Test weight (1000 grain wt.) (g) grain yield (ton/ha) and straw yield (ton/ha) which was at par with M<sub>3</sub>- Spray of Nano Zinc @ 0.5% spray followed by, M<sub>2</sub>- Foliar application of ZnSO<sub>4</sub> @ 0.25%, and M<sub>4</sub>- foliar application of @ 0.25% FeSO<sub>4</sub> at tillering and booting stage respectively. It confirms the results corroborated with the findings of **Mahil et al., (2019)**, **Yadav and Tiwari (2022)** and **Yadav et al., (2023)**

UNDER PEER REVIEW

**Table 1. Effect of organic manure + RDF and foliar spray of micronutrients on plant height (cm), No. of tillers and LAI of wheat crop.**

Treatments	Plant height (cm) 60 DAS			Number of tiller/m <sup>2</sup> 60 DAS			LAI 60 DAS		
	2021-22	2022-23	Pooled	2021-22	2022-23	Pooled	2021-22	2022-23	Pooled
<b>Main plot (organic manure + RDF)</b>									
F <sub>1</sub> - Control (100% RDF)	43.83	44.41	44.12	313.60	315.40	314.50	2.10	2.17	2.14
F <sub>2</sub> - 100 % RDF + Vermicompost @ 2.5t/ha	51.44	51.71	51.58	399.80	401.40	400.60	3.94	4.03	3.99
F <sub>3</sub> - 100 % RDF + FYM @ 5t/ha	47.24	48.13	47.69	360.00	362.00	361.00	2.84	2.89	2.87
<b>S.Em, ±</b>	<b>0.065</b>	<b>0.128</b>	<b>0.150</b>	<b>1.500</b>	<b>0.919</b>	<b>1.187</b>	<b>0.005</b>	<b>0.005</b>	<b>0.014</b>
<b>C.D. (P=0.05)</b>	<b>0.263</b>	<b>0.517</b>	<b>0.606</b>	<b>6.046</b>	<b>3.706</b>	<b>4.787</b>	<b>0.021</b>	<b>0.019</b>	<b>0.057</b>
<b>Sub plot (foliar spray of micronutrient)</b>									
M <sub>1</sub> - Control (No spray)	45.73	46.25	45.99	338.33	340.00	339.17	2.54	2.62	2.58
M <sub>2</sub> - Foliar application of ZnSO <sub>4</sub> @ 0.25% at tillering and booting stage.	47.40	47.83	47.62	356.00	358.00	357.00	3.00	3.08	3.04
M <sub>3</sub> - Spray of Nano Zinc @ 0.5% spray at tillering and booting stage.	48.17	48.67	48.42	368.67	370.00	369.33	3.07	3.12	3.10
M <sub>4</sub> - Foliar application of FeSO <sub>4</sub> @ 0.25% at tillering and booting stage.	46.97	47.91	47.44	350.33	352.33	351.33	2.90	2.96	2.93
M <sub>5</sub> - Foliar application of ZnSO <sub>4</sub> @ 0.25% + @ 0.25% FeSO <sub>4</sub> at tillering and booting stage.	49.25	49.75	49.50	375.67	377.67	376.67	3.29	3.37	3.33
<b>S.Em, ±</b>	<b>0.379</b>	<b>0.399</b>	<b>0.283</b>	<b>3.012</b>	<b>1.789</b>	<b>2.234</b>	<b>0.016</b>	<b>0.018</b>	<b>0.022</b>
<b>C.D. (P=0.05)</b>	<b>1.112</b>	<b>1.170</b>	<b>0.831</b>	<b>8.845</b>	<b>5.254</b>	<b>6.558</b>	<b>0.047</b>	<b>0.053</b>	<b>0.065</b>
<b>Interactions</b>									
<b>S.Em, ±</b>	<b>0.591</b>	<b>0.631</b>	<b>0.463</b>	<b>4.902</b>	<b>2.921</b>	<b>3.658</b>	<b>0.026</b>	<b>0.028</b>	<b>0.037</b>
<b>C.D. (P=0.05)</b>	NS	NS	NS	14.895	8.892	11.164	0.076	0.084	0.114

**Note:** - RDF- (Recommended dose of fertilizers), FYM- (Farm yard manure), ZnSO<sub>4</sub> – (Zinc Sulphate) and FeSO<sub>4</sub> – (Ferrous Sulphate) and DAS (day after sowing)

**Table 2. Effect of organic manure + RDF and foliar spray of micronutrient on Spike length, no of spikelet/spike, harvest index and biological yield in wheat.**

Treatments	Test weight (g)			No of grain/spike			Grain yield (t/ha)			Straw yield (t/ha)		
	2021-22	2022-23	Pool ed	2021-22	2022-23	Pool ed	2021-22	2022-23	Pool ed	2021-22	2022-23	Pool ed
<b>Main plot (Organic manure +RDF)</b>												
F <sub>1</sub> - Control (100% RDF)	36.37	37.08	36.72	44.66	45.76	45.21	4.45	4.65	4.55	6.39	6.51	6.45
F <sub>2</sub> - 100 % RDF + Vermicompost @ 2.5t/ha.	41.68	43.04	42.36	59.68	60.73	60.20	5.80	6.21	6.01	7.53	7.76	7.65
F <sub>3</sub> - 100 % RDF + FYM @ 5t/ha.	39.58	40.61	40.10	54.45	55.61	55.03	5.01	5.42	5.22	6.93	7.24	7.09
<b>S.Em, ±</b>	<b>0.099</b>	<b>0.214</b>	<b>0.140</b>	<b>0.242</b>	<b>0.109</b>	<b>0.081</b>	<b>0.021</b>	<b>0.008</b>	<b>0.024</b>	<b>0.012</b>	<b>0.022</b>	<b>0.032</b>
<b>C.D. (P=0.05)</b>	<b>0.401</b>	<b>0.865</b>	<b>0.565</b>	<b>0.976</b>	<b>0.441</b>	<b>0.328</b>	<b>0.084</b>	<b>0.032</b>	<b>0.098</b>	<b>0.050</b>	<b>0.089</b>	<b>0.129</b>
<b>Sub plot (Foliar spray of micronutrient)</b>												
M <sub>1</sub> - Control (No spray)	37.81	38.92	38.36	46.68	47.87	47.27	4.24	4.47	4.35	5.85	6.00	5.92
M <sub>2</sub> - Foliar application of ZnSO <sub>4</sub> @ 0.25% at tillering and booting stage.	39.20	40.02	39.61	54.15	55.21	54.68	5.14	5.55	5.34	7.05	7.21	7.13
M <sub>3</sub> - Spray of Nano Zinc @ 0.5% spray at tillering and booting stage.	39.57	40.79	40.18	55.19	56.35	55.77	5.37	5.73	5.55	7.35	7.57	7.46
M <sub>4</sub> - Foliar application of FeSO <sub>4</sub> @ 0.25% at tillering and booting stage.	38.62	39.65	39.13	52.43	53.48	52.96	4.90	5.28	5.09	6.72	6.98	6.85
M <sub>5</sub> - Foliar application of ZnSO <sub>4</sub> @ 0.25% + @ 0.25% FeSO <sub>4</sub> at tillering and booting stage.	40.87	41.83	41.35	56.20	57.25	56.73	5.78	6.11	5.95	7.79	8.09	7.94
<b>S.Em, ±</b>	<b>0.397</b>	<b>0.393</b>	<b>0.340</b>	<b>0.426</b>	<b>0.330</b>	<b>0.484</b>	<b>0.042</b>	<b>0.045</b>	<b>0.044</b>	<b>0.062</b>	<b>0.062</b>	<b>0.039</b>
<b>C.D. (P=0.05)</b>	<b>1.164</b>	<b>1.153</b>	<b>0.999</b>	<b>1.252</b>	<b>0.968</b>	<b>1.422</b>	<b>0.122</b>	<b>0.132</b>	<b>0.128</b>	<b>0.181</b>	<b>0.181</b>	<b>0.115</b>
<b>Interactions</b>												
<b>S.Em, ±</b>	<b>0.622</b>	<b>0.645</b>	<b>0.545</b>	<b>0.703</b>	<b>0.522</b>	<b>0.755</b>	<b>0.068</b>	<b>0.070</b>	<b>0.072</b>	<b>0.096</b>	<b>0.098</b>	<b>0.069</b>
<b>C.D. (P=0.05)</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>2.156</b>	<b>1.559</b>	<b>2.225</b>	<b>0.215</b>	<b>0.221</b>	<b>0.227</b>	<b>0.303</b>	<b>0.310</b>	<b>0.218</b>

**Note:** - RDF- (Recommended dose of fertilizers), FYM- (Farm yard manure), ZnSO<sub>4</sub> – (Zinc Sulphate) and FeSO<sub>4</sub> – (Ferrous Sulphate) and DAS (day after sowing)

## Conclusion

In organic manure + RDF, application of 100 % RDF + Vermicompost @ 2.5t/ha are found significantly with respect of growth, yield attribute and yield as well as of wheat crop. Moreover, the combined foliar spray of ZnSO<sub>4</sub> @ 0.25% + FeSO<sub>4</sub> @ 0.25% at tillering and booting stage proved to be best for growth parameters, yield attribute as well as yield of wheat crop as compared to others treatment combinations respectively.

## REFERENCES

**Anonymous**, Annual report 2022-23, Department of Agriculture, Cooperation & Farmers' Welfare, *Ministry of Agriculture Govt. of India*, New Delhi. (2022-23)

**Akhtar, N. Jamil, M. Qazi, M. A., Muslim, N. Hayat Khan, M. U., Jamil, S. Bashir, F. and Iqbal, M. M.** Bio Fortification with Iron and Zinc Improves the Grain Quality and yield attributes of wheat crop. *Plant Cell Biotechnology and Molecular Biology*. 2022;23(15): 23-30.

**Bairagya M. D., Sarath Kumar D. Maitra S. Roja M. and Shankar T. (2019)**. Effect of FYM and Vermicompost on Yield and Economics of Rabi wheat (*Triticum aestivum*. L). *International Journal of Agriculture Sciences*. 2019;11 (7): 1-9.

**Cakmak I., Torun A., Millet E., Feldman M., Fahima T., Korol A., Nevo E., Braun H.J., Ozkan H. 2004.** An Important genetic resource for increasing zinc and iron concentration in modern cultivated wheat. *Soil Sci. Plant Nutrient*. 2004;50(7):1047-1054.

**Dimkpa, C. O., Andrews, J. Sanabria, J. Bindraban, P. S., Singh, U. Elmer, W. H., and White, J. C.** Interactive effects of drought, organic fertilizer, and zinc oxide nanoscale and bulk particles on wheat performance and grain nutrient accumulation. *Science of the Total Environment*. 2020;722, 137808.

**Fazily, T. Thakral, S. K., and Dhaka, A. K.** Effect of integrated nutrient management on growth, yield attributes and yield of wheat. *International Journal of Advances in Agricultural Science and Technology*. 2021;8(1):106-118.

**Kumar, Y. O., Singh, S. P., and Singh, V. P.** Effect of FYM and potassium on yield, nutrient uptake and economics of wheat in alluvial soil. *Annals of Plant and Soil Research*. 2015;17(1):100-3.

**Commented [tt8]:** Improve the reference writing format. The author is the Ministry of Agriculture & Farmers Welfare.

- Loura, D. Kumar, S. Kumar, P. Dhankar, A. Ahlawat, I. and Bedwal, S. (2022).** Comparative effect of organic and inorganic sources of nutrients on yield, soil properties, and economics of wheat under rice-wheat cropping system. *Journal of Sustainable Agriculture* 2022;32(1):37–50.
- Mohan, B. Kumar, P. and Yadav, R. A.** Effect of integrated nutrient management on yield attributes and yield of wheat (*Triticum aestivum* L.). *Journal of Pharmacognosy and Phytochemistry*, 2018;7(1):1545-1554
- Mahil, E. and Kumar, B. (2019).** Foliar application of Nano fertilizers in agricultural crops– A review. *Journal of Farm Science*, 2019;32(3):239-249.
- Saeed, B. Gul, H. Khan, A. Z., Badshah, N. L., Parveen, L. and Khan, A.** Rates and methods of nitrogen and sulphur application influence and cost benefit analysis of wheat. *Journal of Agricultural & Biological Science* 2012;7(2):81-85.
- Singh, H. Ingle, S. R., Pratap, T. Raizada, S. Singh, K. Singh, R. and Parihar, A. K. S.** Effect of integrated nutrient management on nitrogen content, uptake and quality of wheat (*Triticum aestivum* L.) under partially reclaimed sodic soil. *Pharma Innovation Journal*. 2020;11(2):2179-2183
- Solanki, D. Patel, B. T., Patel, D. K., Chaudhary, S** Effect of Iron and Zinc Enriched Organics on Growth, Yield Attributes and Yield of Wheat in Loamy Sand. *International Journal Current Microbiology Applied Science*. 2020;9(6):242-7.
- Tiwari, D. K., Chaturvedi, P. K., Kumar, R. and Yadav, D. S.** Effect of integrated nutrient management on growth and yield of wheat (*Triticum aestivum* L.). *The Pharma Innovation Journal*. 2022;11(12):5032-5034
- Yadav, L. K., and Tiwari, U. S. (2022).** Effect of integrated nutrient management on growth and productivity under rice-wheat cropping system under central plain zone of UP. *The Pharma Innovation Journal* 2022;11(6):307-311
- Yadav, A. Krishnan, P. Kundu, M. Bana, R. S., Shaloo, Choudhary, A. K., Shivay, Y. S., Meena, S. L., Begam, G. S., Ranjan, R. K., Yadav, N. S., Singh, T. Yadav, A. and Raj, R. (2023).** Zinc nano-fertilization enhances wheat productivity and biofortification. *International Lcense*. 2023;3(19):25-35

**Weber, J., Karczewska, A., Drozd, J., Licmar, M., Licznar, S. and Jarnroz, E. 2007.**

Agricultural and ecological aspects of a sandy soil as affected by the application of municipal solid waste composts. *Soil Biol. Biochemistry.*, 2007;39:1294-02.

UNDER PEER REVIEW