

Original Research Article

Salt Stress Responses in Early Seedling Growth Characteristics of Blackgram [*Vigna mungo* (L.) Hepper] Genotypes

ABSTRACT

In view of increasing salinity and intrusion of salt water into the cultivable land area under coastal saline region, germination and crop establishment is a major problem that influence on decrease in the production and productivity of blackgram. Hence, screening of 101 genotypes under different salt concentrations for growth characters paves way to identify the saline resistant genotypes. The upshot of saline stress on seedling growth characters of 101 genotypes which are collected from different eco geographical regions subjected to Half strength Hoagland and Arnon's solution (T₁), Half strength Hoagland Arnon's solution + NaCl treatment @ 100mM, 150mM, 200mM and 250mM concentration @ 7th, 10th, 13th, 16th days (T₂) respectively and Half strength Hoagland and Arnon's solution (control) (T₃) during January 2022. Highly significant variation exhibited among the various genotypes for different seedling growth characters viz., leaf area per plant, total number of leaves per plant, dry weight of leaves per plant, dry weight of shoot per plant, dry weight of roots per plant and dry weight of the whole plant. Salinity has the adverse effect on the growth characters. Among the genotypes studied, AUB-08-33(G81) and CBG-09-08(G18) showed maximum mean values under different NaCl concentrations (T₂), recommended as donor parent for hybridization as well as candidate genotype in crop improvement for saline soil.

Keywords: Blackgram, Salt stress, Seedling characters, NaCl

1. INTRODUCTION

Black gram is scientifically known as *Vigna mungo* (L.) Hepper, 2n=2x=22 and commonly known as urd in India. Black gram is mainly cultivated and distributed in tropical and subtropical regions of Africa, Asia, America, and Australia (Lewis *et al.*, 2005) whereas

majorly grown in South and Southeast Asian countries including Afghanistan, Bangladesh, Pakistan, and Myanmar. About 70 per cent of the world's black gram production comes from India. Worldwide it covers about 23.4 lakh tones of black gram annually from 46.7 lakh hectares of area with an average productivity of 501 kg per hectare in 2020-2021 (agricoop.nic.in). India identified as its primary origin (Fuller and Harvey, 2006) and domesticated from the wild species *V. mungo* var. *sylvestris* (Chandelet *et al.*, 1984). India is the largest producer and consumer of black gram with an area account for about 15.7 percent of total pulse acreage and contributes 9.09 per cent of total pulse production. In Kharif 2021-2022, black gram production accounts 20.5 lakh tones in an area of 39.43 lakh hectares (agricoop.nic.in).

Comment [EP1]: Not listed in References!

In India, states like Madhya Pradesh, Uttar Pradesh, Andhra Pradesh and Tamil Nadu shows upward production in black gram cultivation (Vennila *et al.*, 2018). The area and production of black gram in Tamil Nadu is about 4.2 lakh hectare and 2.78 lakh tons. (Department of Pulses Development, Bhopal, Annual Report; 2021-2022). There is a declining phase in production and productivity of black gram due to the influence of biotic, abiotic factors and urbanization. Its need of hour to increase the production and productivity to meet out the demands of increasing population (Farooq *et al.*, 2017). Among the abiotic factors, soil salinity plays an important role in declining the production and productivity of crop plants especially along the coastal region (Rahaman *et al.*, 2017). In India saline soil occupies about 6.73 million hectares (CSSRI -2022).

Comment [EP2]: Not listed in References!

Salt stress hampers the agricultural productivity by lowering the yield of various crops in arid and semi-arid regions of the world (Khan *et al.* 2019; Food and Agriculture Organization of the United Nations, 2022). Saline soils are present in the states of Gujarat, Bihar, Haryana, Rajasthan, Maharashtra, Odisha, Andhra Pradesh, Kerala, Tamil Nadu, Uttar Pradesh, and West Bengal (agricoop.nic.in).

The effect of salinity on the growth of black gram has been reported occasionally (Hasan *et al.*, 2018). Nirmala Sherawath *et al.* (2018) also opined the growth retarding effects of various salinity treatment in pulses. However, with increasing (0 to 180Mm of NaCl), decreases germination by 50% in the species of the genus *Phaseolus* reported by Bayuelo –Jimenez *et al.* (2002). Increase in salinity causes a decrease in fresh or dry weight of plants also it influences on

seedling growth, shoot height and root length which showed most sensitive to salt stress (Rajasekaret *al.*, 2022).

In the sight of increasing salinity stress, there is a necessary to find out the saline tolerance varieties which can be used to increase the black gram productivity under saline condition. Considering the above facts, the study was undertaken to identify the genotypes against salt stress by observing the growth characters.

2. MATERIALS AND METHODS

Location

To investigate the seedling response of black gram under various salinity treatment was conducted at the Department of Genetics and Plant Breeding, Faculty of Agriculture, Annamalai University, Annamalai Nagar, Tamil Nadu.

Plant materials and methods

Seeds of 101 black gram genotypes were collected from various geographical sources and the study was conducted. The list of the materials is presented in Table 1.

Experiment

To analyze the growth characters this experiment was conducted at Department of Genetics and Plant Breeding, Faculty of Agriculture, Annamalai University during January, 2022 using Factorial Completely Randomized Design (FCRD) described by Gomez and Gomez (1984) replicated twice.

Seeds of the 101 black gram genotypes were sown in plastic cup (9.0 x 6.5 cm) @ seven seeds per cup filled with sterilized coir pith. The cups are then placed in plastic trays (40 x 28 x 7 cm) @ 15 cups per tray which was irrigated with half –strength Hoagland and Arnon's nutrient solution (Hoagland and Arnon's, 1950) @ 2 liters per tray by following Asana Matsuura *et al.* (2005). Holes were made at the bottom of the cups to facilitate easy absorption of the nutrient solution by the coir pith. The details of the treatment is given below.

Chart 1. The details of the treatment

Code	Treatment	Observations recorded on

Comment [EP3]: I recommend also to state the growing conditions, for example the temperature, air humidity.

Comment [EP4]: evaluated genotypes

Comment [EP5]: are

Comment [EP6]: Table 1

T ₁	Half strength Hoagland and Arnon's solution	One day before the start of salinity treatment (6DAS) (Fig. 1)
T ₂	Half strength Hoagland Arnon's solution +NaCl treatment @100mM,150mM,200mM and 250mM concentration @7 th ,10 th ,13 th , &16 th days respectively	12days after the start of salinity treatment (18 DAS) (Fig. 2)
T ₃	Half strength Hoagland and Arnon's solution (control)	18 DAS

The observations were recorded on the growth characters viz., leaf area per plant, total number of leaves per plant, dry weight of leaves per plant, dry weight of shoots per plant, dry weight of roots per plant and dry weight of whole plant and the results were analyzed using TNAUSTAT.

Comment [EP7]: It is necessary to state at least the names of the methods used for analysis of for example measurement of leaf area per plant, ...

3.RESULT AND DISCUSSION

The analysis of variance indicated the existence of highly significant difference among all the 101 genotypes for all the traits studied (Table 2).

Total leaf area per plant

Diminution in leaf area were noted. The range for this trait was from 0.16 cm² to 2.53 cm² (G66) in T₁, 2.30 cm² to 25.97 cm² (G18) in T₂ and 5.60 cm² to 75.15 cm² (G59) in T₃. The genotype G18 recorded the maximum total leaf area of 25.975 cm² in T₂. It was found that out of 101 genotypes, 15 and 27 genotypes had significantly higher values than the general mean in T₂ and T₃ respectively. The leaf area was decreased to 97.54% Of the control respectively (Table 3). In general salinity stress reduces the leaf number, leaf area, shoot and root dry weight leading to low yield. (Li *et al.*, 2006; Sharifi *et al.* 2007; Dutta and Bera, 2014 and Udhaya Nandhini *et al.* 2018). Our results are consistence with Pankaj *et al.* (2016) who reported that NaCl highly reduced total leaf area.

Comment [EP8]: Not listed in References!

All cited authors/sources must be listed/mentioned in References and all references must be cited in manuscript!

Comment [EP9]: Not listed in References!

Total number of leaves per plant

The decline in the number of leaves were noted. The range for this trait was from 0.47 to 2.40 (G18) in T₁, 1.29 to 6.55 (G87) in T₂ and 3.00 to 10.81 (G47) in T₃. Genotype G87 recorded the maximum total number of leaves per pant of 6.55 in T₂. A total of 9 and 14 genotypes that significantly exceeded the general mean for this trait in T₂ and T₃ respectively. The leaf number also shown downward decrease to 95.09% over the control (Table 3). Leaf number was not affected by 25 and 50 mM NaCl concentrations so that there was no significant difference between those treatments in compare with control but increase of salinity to 100 mM significantly decreases leaf number. Similar reports were noticed in soybean (El Sabaghet *al.*, 2015 and Vennila, 2018). Reduced number of leaves in plant exposed to excessive salinity was also observed in Curly Kale (Salachna *et al.*, 2017).

Dry weight of leaves per plant

The drop in weight of leaves in black gram were observed. The range for this trait was from 0.0004 g to 0.0087 g (G65) in T₁, 0.002 g to 0.0553 g in T₂ (G18) and 0.0070 g to 0.0788 g in T₃ (G61). The genotype G18 was recorded the maximum dry weight of leaves per plant of 0.0553 g in T₂. One genotype in T₂ and 4 genotypes in T₄ stigmatically that exceeded the general mean. The dry weight of the leaves was decreased to 98.05% over the control (Table 4). Indeed, the plant dry weight was significantly decreased by increasing the NaCl levels (El Sabaghet *al.*, 2015 and Rajasekaret *al.*, 2022).

Dry weight of shoot per plant

Black gram genotypes remarkably reduced the shoot dry weight with increasing the salinity levels. The range for this trait was from 0.0011 g to 0.0285 g (G10) in T₁, 0.0050 g to 0.0495 g (G18) in T₂ and 0.0200g to 0.0626 g (G61) in T₃. It was found that the genotype G 18 (0.0495 g) had significantly higher values than the general mean in T₂. Genotypes *viz.*, G 24, G 61 and G 70 had significantly higher values than the general mean in T₃. Dry weight of the shoot decreased to 97.22% of the control respectively (Table 4). The shoot dry weight showed decreased with increasing salinity compared to control observed by El Sabagh *et al.* (2015), Hasan *et al.* (2017) and Subroto Kumar Podder *et al.* (2020).

Dry weight of roots per plant

The range for this trait was from 0.0002 g to 0.0321 g (G53) in T₁, 0.0041 g to 0.0476 g (G49) in T₂ and 0.0086 g to 0.0551 g (G49) in T₃. Genotype, G49 was recorded the maximum dry weight of roots per plant of 0.0476 g in T₂. It was found that out of 101 genotypes studied, 9, 18 and 14 genotypes had significantly higher values than the general mean in T₁, T₂ and T₃ respectively. Dry weight of the root also decreased to 96.74% (G1) of the control (Table 5). Velmaniet *al.* (2015) reported that the effect of NaCl on root dry weight decreased in black gram with increase in the concentration, compared to control. El-Kafafiet *al.* (2015) and Vennilaet *al.* (2018) observed that salinity significantly reduced the root dry weight and root was more affected than shoot.

Dry weight of whole plant

The fall of weight of the whole plant were observed. The range for this trait was from 0.0017 g to 0.0412 g (G70) in T₁, 0.0240 g to 0.1224 g (G18) in T₂ and 0.0506 g 0.1616 g (G61) in T₃. The genotype G18 was recorded the maximum dry weight of whole plant of 0.1224 g in T₂. Dry weight of the whole plant decreased to 97.22% over the control (Table 5). It is well documented that under saline condition, the ability of plant species to uptake water is reduced and this leads to rapid decrease in plant growth and productivity (Mahouachi, 2018 and Izadi–Darbandiet *al.* 2018).

4. CONCLUSION

Salt stress imperceptibly reduced the growth characters of black gram genotypes. The genotypes G87 and G49 recorded the maximum mean values for total number of leaves per plant and dry weight of roots per plant respectively. Genotype, G18 recorded maximum mean values for characters like leaf area per plant, dry weight of leaves per plant, dry weight of shoot per plant and dry weight of whole plant. Genotypes namely, G1, G18, G74, G81, G96 obtained from diversified regions which shows the maximum mean value under salt stress for almost all the growth characters studied (Fig. 4), it was deduced that these genotypes may be utilized as potential donor as well as taken as key candidate in crop improvement programmes specifically in coastal saline regions.

REFERENCES

Asana Matusuura, Shinobu Inanaga and Kenheri Murata. Differences in the vegetative growth between common and Tartary back wheat in saline hydroponic culture. *Plant Production Science*. 2005. 8(5): 533-538.

Central Soil Salinity Research Institute (CSSRI) -2022.

Chandel, K.P.S., Lester, R.N and Starling, R.J. The wild ancestors of mung beans (*Vigna mungo* (L.) Hepper and *V.radiata* (L.) Wilczek). *Botanical Journal of the Linnean Society*.1984. 89:85-96.

Department of Pulses Development, Bhopal, Annual Report, 2021-2022. DOI: 10.1007/s11356-015-5840-5.

Dutta P and Bera AK. Effect of NaCl salinity on seed germination and seedling growth of mung bean cultivars. *Legume Research*. 2014. 37:161 – 164.

El Sabagh, A., Omar, A., Saneoka, H. &Barutçular, C. Physiological performance of soybean germination and seedling growth under salinity stress. *Dicle University Journal of the Institute of Natural and Applied Science*.2015. 4(1): 6-15.

El-Kafafi, E.H., Helal, A.G., El Hafnawy S.F.M. &Flaah, R.F.E.L. Characterization and evaluation of some mung bean genotypes for salt tolerance. *World Applied Science Journal*. 2015.33 (3): 360-370.

Farooq M, Gogoi N, Hussain M, Barthakur S, Paul S, Bharadwaj N, Migdadi HM, Alghamdi SS, Siddique KHM. Effects, tolerance mechanisms and management of salt stress in grain legumes. *Plant Physiology and Biochemistry*. 2017. 118:199-217.

Food and Agriculture Organization of the United Nations (2022).<https://www.fao.org/soils-portal/soil-management>.

Fuller, D.Q. and Harvey, E.L. The archaeobotany of Indian pulses: identification processing and evidence for cultivation. *Environmental Archaeology*.2006. 11:241-268

Ghosh S, Mitra S and Paul A. Physiochemical studies of sodium chloride on mungbean (*Vigna radiata* L. Wilczek) and its possible recovery with spermine and gibberellic Acid. *The Scientific World Journal*.2015.doi: 10.1155/2015/858016.

Gomez KH and Gomez AA. *Statistical Procedures for Agricultural Research*. 1984; 381. Ref.: <https://goo.gl/BCaoZ3>.

Comment [EP10]: Many references are not cited in manuscript and not all cited authors/sources are listed in References! All References must be cited in article and all cited authors/sources must be listed in References. This problem must be solved!

Check and unification of reference format is needed (see instructions for authors <https://journalijpss.com/index.php/IJPSS/about/submissions#authorGuidelines>).

Comment [EP11]: Not cited in manuscript!

Hasan M.K, Islam M.S, Islam M.R, Ismaan H.N and Sabagh El A .Salinity tolerance of black gram cultivars during germination and early seedling growth. *CercetariAgronomice in Moldova*. 2018.1(3): 51-68.

Hasan, M.K., EL Sabagh, A., Sikdar, M.S.I., Alam, Md.J. Ratnasekera, D., Barutçular, C., Abdelaal, Kh.A.A. and Islam, M.S. Comparative adaptable agronomic traits of black gram and mung bean for saline lands. *Plant Archives*. 2017.17(1): 589-593.

Comment [EP12]: Not cited in manuscript!

Izadi-Darbandi, E., Mehdikhani, Hadi. Salinity effects on some of the morphological traits of three plantago species. *Scientia Horticulturae*. 2018. 236: 43-51.

Jeannette S Bayuelo- Jimenez, Richard Craig, Jonathan P Lynch. Salinity Tolerance of Phaseolus Species during Germination and Early Seedling Growth. *Crop Science*. 2002. 42(5): 1584-1594.

Khan .M.A.H, Baset .M.A, Ahmed. J.U, Karim. M.A and Saikat. M.M.H.Morpho –Anatomical appraisal of some pulse crops under salinity stress. *Bangladesh Journal of Agriculture research*: 2019. 44(3): 439-451.

Lewis, G. Schrire, B., Mackinder, B. and Lock, M. 2005. Legumes of the world, Kew Royal Botanical Gardens, Richmond Survey, p. 577.

Li X, An P, Inanaga S, Eneji AE, Tanabe K. Salinity and defoliation effects on soybean growth. *Journal of Plant Nutrition*. 2006. 29:1499-1508.

Mahouachi, J. Long term salt stress influence on vegetative growth and foliar nutrient changes in mango (*Mangifera indica* L.) seedlings. *Scientia Horticulturae*. 2018. 234: 95-100.

Nirmala Sehwat, Mukesh Yadav, Anil Kumar Sharma, Vikas Kumar and Kangila Venkataraman Bhat.Salt stress and mungbean (*Vigna radiata* (L.) effects, physiological perspective and management practices for alleviating salinity. *Archives of Agronomy and Soil Science*. 2018. 65 (9): 1287-1301.

Pankaj Awasthi, Himani Karki, Vibhuti, Kiran Bargali and Bargali.S.S. Germination and seedling growth of pulse crop (*Vigna Spp.*) as affected by soil salt stress: *Current Agriculture Research Journal*.2016. 4(2): 159-170.

Comment [EP13]: ?

ssp.

Salachna, P., Pirchocki, R. and Byczynska. A. Plant growth of curly kale under salinity stress. *Journal of Ecological Engineering*. 2017. 18(1): 119-124.

Rahman, Md.M. Zahan, F., Sikdar, S.I., ElSabagh, A., Ratnasekera, D., Barutçular, C. and Islam, M.S. Evaluation of salt tolerance mung bean genotypes and mitigation of salt stress through potassium nitrate fertilization. *Fresenius Environmental Bulltein*. 2017. 26 (12): 7218-7226.

Rajasekar, R., V. Babu Rajendra Prasad, M. Dianaguiraman, A. Sumathi and D. Jegadeeswari. Screening Blackgram (*Vigna mungo* (L.) Hepper) Genotypes for Salt Tolerance at Germination and Seedling Stages. International Journal of Environment and Climate Change. 2022. 12(11): 383-396.

Subroto Kumar Podder, Joyanti Ray, Debesh Das and BidhamChandro Sarker. Effect of salinity (NaCl) on germination and seedling growth of mung bean (*Vigna radiata* L.): Journal of Bioscience and Agriculture Research. 2020. 24(2):2012-2019.

Udhaya Nandhini D, Somasundaram E and Mohammed Amanullah M. Effect of rhizobial nod factors (lipo-chitooligosachharide) on seedling growth of black gram under salt stress: Legume Research, 2018. 41(1): 159-162.

Velmani, S., Murugesan, S. & Arulbalachandran, D. Growth and biochemical characteristics of black gram (*Vigna mungo* L., Hepper) under NaCl salinity. International of Current Research and Review. 2015.4(1):13-17.

Vennila, S. Effect of microbial inoculants on biometrical traits of blackgram (*Vigna mungo* L. Hepper). International journal of current research in life science. 2018.7(4):1487-1488.

Vennila, S., M Pushpakaran and K Palaniraja. Effect of presowing seed pelleting treatment using botanical leaf powders and biofertilizers on growth and yield characters in blackgram (*Vigna mungo* [L.] Hepper) variety VBN 5. Journal of Pharmacognosy and Phytochemistry. 2018. 7(4): 2923-2925.

Table 1. List of black gram genotypes used for the study

Comment [EP14]: List of evaluated black gram genotypes

Genotypes code	Genotypes	Source
G1	RU -8-701	Indian Institute of Pulse Research, Kanpur
G2	RU-8 703	Indian Institute of Pulse Research, Kanpur
G3	RU-8-704	Indian Institute of Pulse Research, Kanpur
G4	RU-8-705	Indian Institute of Pulse Research, Kanpur
G5	RU-8-706	Indian Institute of Pulse Research, Kanpur
G6	RU-8-707	Indian Institute of Pulse Research, Kanpur
G7	RU-8-711	Indian Institute of Pulse Research, Kanpur
G8	RU 8-708	Indian Institute of Pulse Research, Kanpur
G9	RU-8 -702	Indian Institute of Pulse Research, Kanpur
G10	PLU-621	NPRC, Vamban, Tamil Nadu, India
G11	RU -09-157	Indian Institute of Pulse Research, Kanpur

G12	RU 9-159	Indian Institute of Pulse Research, Kanpur
G13	RU 9- 158	Indian Institute of Pulse Research, Kanpur
G14	RU 9- 156	Indian Institute of Pulse Research, Kanpur
G15	RU 9-155	Indian Institute of Pulse Research, Kanpur
G16	RU 9-154	Indian Institute of Pulse Research, Kanpur
G17	RU 9-153	Indian Institute of Pulse Research, Kanpur
G18	CBG -09-08	Coimbatore , Tamil Nadu, India
G19	CBG- 09-06	Coimbatore , Tamil Nadu, India
G20	CBG-09-15	Coimbatore , Tamil Nadu, India
G21	CBG -09-10	Coimbatore , Tamil Nadu, India
G22	CBG- 09-04	Coimbatore , Tamil Nadu, India
G23	CBG -09-03	Coimbatore , Tamil Nadu, India
G24	CBG-09-02	Coimbatore , Tamil Nadu, India
G25	CBG-09-13	Coimbatore , Tamil Nadu, India
G26	CBG-09-01	Coimbatore , Tamil Nadu, India
G27	CBG-09-09	Coimbatore , Tamil Nadu, India
G28	VBN-5	NPRC, Tamil Nadu, India
G29	VBG-05-014	NPRC, Vamban, Tamil Nadu, India
G30	VBG-05-007(BR)	NPRC, Vamban, Tamil Nadu, India
G31	SD-2-143	NPRC, Vamban, Tamil Nadu, India
G32	RG RU- 448	NPRC, Vamban, Tamil Nadu, India
G33	H-21-40-41	NPRC, Vamban, Tamil Nadu, India
G34	PS-1	NPRC, Vamban, Tamil Nadu, India
G35	NTU -94-6	NPRC, Vamban, Tamil Nadu, India
G36	P-133/19	Punjab, India
G37	P-206	Punjab, India
G38	LU-272	NPRC, Vamban, Tamil Nadu, India
G39	MDU-99-2	Madurai, Tamil Nadu, India
G40	ADT-3	Aduthurai, Tamil Nadu, India
G41	PLU-227	NPRC, Vamban, Tamil Nadu, India
G42	Vanniyur local - 1	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G43	P-47	NPRC, Vamban, Tamil Nadu, India
G44	99-100	Taiwan
G45	Ariyapatti	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G46	Vanniyur local- 2	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G47	T9	NPRC, Vamban, Tamil Nadu, India
G48	VZM- 189/33	NPRC, Vamban, Tamil Nadu, India
G49	NUB-294	NPRC, Vamban, Tamil Nadu, India
G50	NP-7	NPRC, Vamban, Tamil Nadu, India
G51	U- 99-35	NPRC, Vamban, Tamil Nadu, India
G52	PLU-340	NPRC, Vamban, Tamil Nadu, India
G53	PLS 364/47	NPRC, Vamban, Tamil Nadu, India

G54	LO-217	NPRC, Vamban, Tamil Nadu, India
G55	Atthipatti local-1	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G56	PHM.8	Punjab, India
G57	PLV-554	NPRC, Vamban, Tamil Nadu, India
G58	Atthipatti local 2	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G59	TUS-27- Bold	NPRC, Vamban, Tamil Nadu, India
G60	Co-1	Coimbatore, Tamil Nadu, India
G61	Chidambaram local	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G62	PLU- 446	NPRC, Vamban, Tamil Nadu, India
G63	2KU-53	Uttar Pradesh India
G64	VBN-4	NPRC, Vamban, Tamil Nadu, India
G65	VBG05-02	NPRC, Vamban, Tamil Nadu, India
G66	PLS-254	NPRC, Vamban, Tamil Nadu, India
G67	P-13211	NPRC, Vamban, Tamil Nadu, India
G68	PLS – 364/42	NPRC, Vamban, Tamil Nadu, India
G69	IPU- 917	NPRC, Vamban, Tamil Nadu, India
G70	VBN-3	NPRC, Vamban, Tamil Nadu, India
G71	PLG-47	NPRC, Vamban, Tamil Nadu, India
G72	NLC-8183	NPRC, Vamban, Tamil Nadu, India
G73	Co-5	TNAU Tamil Nadu, India
G74	VBG-05-08	NPRC, Vamban, Tamil Nadu, India
G75	VBG-05-001	NPRC, Vamban, Tamil Nadu, India
G76	VBG-04-0012 (Br)	NPRC, Vamban, Tamil Nadu, India
G77	IPU- 232	NPRC, Vamban, Tamil Nadu, India
G78	VBG-05-0011	NPRC, Vamban, Tamil Nadu, India
G79	VBG-05-006	NPRC, Vamban, Tamil Nadu, India
G80	AUB-08-11	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G81	AUB-08-13	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G82	AUB-08-21	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G83	AUB-08-22	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G84	AUB-08-23	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G85	AUB-08-31	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G86	AUB-08-32	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G87	AUB-08-33	Plant Breeding Farm, Annamalai University, Tamil Nadu, India

G88	AUB-08-41	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G89	AUB-08-42	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G90	AUB-08-43	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G91	AUB-08-51	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G92	AUB-08-52	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G93	AUB-08-53	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G94	AUB-08-61	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G95	AUB-08-62	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G96	AUB-08-63	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G97	AUB-08-71	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G98	AUB-08-72	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G99	AUB-08-73	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G100	AUB-08-121	Plant Breeding Farm, Annamalai University, Tamil Nadu, India
G101	AUB-09-152	Plant Breeding Farm, Annamalai University, Tamil Nadu, India

Table 2. ANOVA for various growth characters of black gram genotypes

Source	df	MS					
		Total leaf area per plant	Total number of leaves per plant	Dry weight of leaves per plant	Dry weight of soot per plant	Dry weight of root per plant	Dry weight of whole plant
Genotypes	100	98.14**	2.92**	0.0002**	0.0006*	0.0008**	0.001**
Treatment	2	42662.95**	1066.48**	0.059	0.16**	0.0014**	0.18.**
Genotype x treatment	200	101.08**	3.12**	0.0002**	0.0006**	0.00086**	0.001**
Error	303	3.61	0.40	0.00006	0.0004	0.000021	0.0003
Total	605	192.48	5.24	0.0003	0.0006	0.00044	0.001
G x T	SED	1.9007	0.6359	0.0081	0.0223	0.0045	0.0174
	CD (0.05)	3.7403	1.2513	0.0163	0.4389	0.0089	0.0343
	CD (0.01)	4.9270	1.6483	0.0215	0.0578	0.0117	0.0452

Comment [EP15]: Results of Analysis of Variance of selected growth characters of black gram genotypes

*Significant @5% level

*Significant @1% level

Table 3. Mean performance of leaf area per plant and total number of leaves per plant for black gram genotypes

Genotypes	Total Leaf area per plant (cm ²)			Total number of leaves per plant		
	T ₁	T ₂	T ₃	T ₁	T ₂	T ₃
G1	1.4000	20.7500**	28.4950	1.4000	4.6000	7.6150*
G2	0.2500	10.0450	38.5250**	0.9000	2.8850	8.8000
G3	0.2500	10.0394	25.3000	0.9000	4.6500	6.9500

G4	0.3500	11.7600	26.4550	0.9000	2.6000	6.9250
G5	0.5000	9.0350	30.1750	1.9000	3.6500	4.5250
G6	1.0000	17.6250**	44.1400	1.9000	4.5500	10.3000**
G7	0.4450	14.6400	24.2850	0.9000	3.5500	5.5150
G8	0.2000	21.0250**	33.0550	0.9000	3.9000	6.3500
G9	0.3000	17.0600**	26.4000	0.9000	4.1000	5.6850
G10	0.6150	12.2350	29.2000	1.2500	3.3600	5.6650
G11	0.6000	21.6600**	27.1950	2.0000	4.6500	5.4350
G12	1.2000	15.1400	16.1750	1.7000	4.6000	4.8900
G13	1.1000	6.7100	38.9400**	1.9000	3.4900	7.5100
G14	0.8500	5.5150	40.8700**	1.9000	1.9250	8.2900**
G15	0.6000	10.0600	26.9000	1.9000	2.6500	6.1250
G16	0.6000	11.1100	29.9900	1.9000	3.7850	7.4100
G17	0.5500	3.6700	32.0700	0.9000	1.2950	5.3900
G18	0.9500	25.3350**	25.9750	2.4000	4.3600	5.6450
G19	1.4000	11.2800	15.1900	1.4000	3.6000	6.5250
G20	0.6000	19.4000	24.3050	1.9000	4.6000	6.7650
G21	0.2000	16.2500*	32.1850	0.9000	4.0500	6.5100
G22	0.4000	13.5150	32.0750	2.0500	4.0250	5.7900
G23	0.5000	16.1500*	31.9800	2.0500	3.0800	5.7750
G24	0.6000	10.4050	39.7250**	1.9000	1.6500	7.4900
G25	0.4300	3.0250	26.1900	1.4000	3.2150	6.6600

Con...

G26	0.6000	3.5000	5.6050	1.9000	2.3150	3.5100
G27	0.5000	19.9600**	27.1400	1.9000	4.6000	5.0500
G28	0.8000	16.4000**	23.2400	1.9000	3.3700	4.5100
G29	0.4000	4.3600	37.9750**	1.0000	2.0100	7.5050
G30	0.3500	10.2300	18.9500	0.9000	3.3500	4.6650
G31	0.4500	8.6050	46.5400**	1.4000	2.6500	7.5200
G32	0.7000	3.8600	20.9100	1.7000	1.7500	5.4900
G33	0.5000	6.3850	28.9400	1.9000	3.0500	6.8400
G34	0.9600	6.5350	14.1900	1.9000	2.7500	4.5150
G35	0.6000	10.2050	13.8850	1.9000	3.1500	4.5050
G36	0.6000	18.1450**	25.2200	1.9000	4.6000	6.0350
G37	1.0000	8.4950	15.1850	1.9000	2.9000	8.1200**
G38	1.0000	15.4400*	39.2000	1.9000	5.0000*	6.9500
G39	0.2800	11.3250	36.7850**	1.9000	2.9500	7.1600
G40	1.2250	2.8000	15.9350	1.9000	2.7100	5.4700
G41	0.3000	2.3000	31.6050	0.9000	2.2000	6.5250
G42	0.8000	2.3000	38.1150	1.9000	3.0500	6.8650
G43	0.4400	10.5890	11.6050	1.9000	3.1500	3.3050
G44	0.4000	13.3550	18.1800	1.9000	3.7900	4.0450
G45	0.4500	7.7650	10.0150	1.8750	2.2800	2.5600
G46	0.6500	3.6000	17.7350	1.9000	2.3800	3.6600
G47	0.6500	6.5050	17.7300	1.8750	2.6250	10.8100**
G48	0.5500	10.5250	26.2200	1.9000	3.8750	6.6550
G49	0.2000	15.7850*	33.0700	0.8500	4.2850	6.5100
G50	0.4000	12.1050	27.4300	1.4000	4.6000	5.2150
G51	0.4000	8.7750	19.9850	1.8750	3.2250	4.6550
G52	0.3500	16.5450**	44.8250**	1.9250	4.2700	4.4300
G53	0.3500	9.7150	36.9300	1.3750	3.6000	5.8200
G54	2.1300	9.8850	41.9750**	1.8250	4.1300	4.6650

G55	0.5000	13.8650	14.1200	1.8500	4.3350	4.6950
G56	0.1775	11.6500	32.8150	1.9000	5.0600*	5.1800
G57	0.1650	13.7600	32.0800	0.9000	3.9100	5.8250
G58	0.5500	5.8950	24.0950	1.9000	2.1850	4.8250
G59	0.2500	7.0650	75.1850**	1.9000	3.6000	8.5100**
G60	0.2500	11.6400	25.0100	1.9000	3.8500	5.4900
G61	0.3000	21.7750**	38.9600**	0.9000	4.8000	9.8700**
G62	1.1300	15.0300	24.3050	1.9000	5.5500**	7.0250
G63	0.7000	12.2900	18.4250	1.9000	5.6000**	5.8850
G64	0.7500	5.0000	46.8900**	1.9000	2.7500	6.4900
G65	1.1500	12.4250	41.2350**	1.8500	3.9500	7.4750
G66	2.5300	10.4400	25.4350	1.8500	4.1000	7.4900
G67	0.9000	7.3500	30.2500	1.9000	3.2250	5.9250
G68	0.4500	9.6700	36.5400**	1.9000	3.7000	5.8700
G69	0.4150	8.7850	13.6200	1.7000	4.2650	4.6300
G70	0.3750	11.3700	46.0550**	0.5600	3.3500	8.5150**
G71	0.7000	7.3000	42.3500**	0.9000	2.6500	5.6650
G72	1.4250	12.8400	29.7750	2.1500	5.6500**	6.5250
G73	0.9500	12.1750	17.3400	1.8500	2.9000	3.0050
G74	0.4000	15.9250*	26.9900	1.9000	6.2000**	6.5200
G75	0.6500	12.2850	17.4300	2.1850	3.4250	3.9650
G76	0.5500	12.8000	35.6750**	1.9000	4.0500	6.8150
G77	0.6500	12.8900	34.0250*	1.9000	3.6000	7.2000
G78	0.3500	14.7000	28.5600	1.9000	3.5500	6.5250
G79	0.6000	13.6500	40.8250**	1.2500	4.0500	7.6250*
G80	0.5000	13.0500	30.2150	1.9000	3.6000	5.6200
G81	0.4000	9.2200	31.5000	1.9000	3.2500	7.4900

Cont..

G82	0.7000	10.5550	34.3450*	1.4000	4.1000	7.5200
G83	0.5600	11.0050	28.4150	1.9000	3.1500	5.5200
G84	1.1000	12.9100	17.7200	1.9000	5.1000	5.8900
G85	1.1000	14.3450	52.9550**	1.9000	3.9500	8.5250**
G86	0.4500	8.2000	43.2400**	0.4700	3.9000	9.4900**
G87	1.9000	16.6400*	30.5400	1.9000	6.5500**	7.4900
G88	1.3000	12.0550	38.4050**	1.9000	4.0000	5.4900
G89	0.4500	12.2450	30.8750	0.9000	6.1150**	7.5300
G90	0.3300	7.3200	33.7750	0.9000	3.1500	6.1650
G91	1.0000	9.6200	34.2000	1.1000	3.9350	8.2150**
G92	0.6000	11.6300	36.6700**	0.6200	3.7250	6.6750
G93	0.7000	14.0050	51.9550**	0.9000	4.2000	7.7750*
G94	0.5000	11.4600	40.8250**	1.1000	4.6000	7.8750*
G95	0.1545	12.2750	38.9800	1.6000	5.1000*	6.7350
G96	0.4000	13.1150	29.3250	1.5000	3.6500	5.6350
G97	0.3800	13.3900	32.3000	1.6000	4.9000	6.0500
G98	0.1500	7.6650	37.2150**	1.9000	2.4750	7.4900
G99	0.5500	7.8200	26.5550	1.9000	3.8000	6.3000
G100	0.4000	11.6500	24.9500	1.9000	3.9500	10.7200**
G101	0.6500	12.4600	22.3850	1.9000	4.3500	4.6350
Mean	0.6442	11.4321	30.0619	1.6100	3.7157	6.3414
		CD(0.05)= 3.74 CD(0.01)= 4.92 SED= 0.63			CD(0.05)= 1.25 CD(0.01)= 1.64 SED= 1.9	

*Significant at 5% level

**Significant at 1%

Table 4. Mean performance of dry weight of leaves per plant and dry weight of shoot per plant for black gram genotypes

Genotypes	Dry weight of leaves per plant (g)			Dry weight of shoot per plant (g)		
	T ₁	T ₂	T ₃	T ₁	T ₂	T ₃
G1	0.0043	0.0417	0.0525	0.0247	0.0400	0.0485
G2	0.0022	0.0151	0.0436	0.0158	0.0216	0.0344
G3	0.0017	0.0301	0.0304	0.0195	0.0362	0.0383
G4	0.0010	0.0180	0.0309	0.0212	0.0305	0.0396
G5	0.0045	0.0180	0.0355	0.0135	0.0229	0.0302
G6	0.0028	0.0257	0.0425	0.0116	0.0293	0.0420
G7	0.0010	0.0275	0.0286	0.0011	0.0385	0.0400
G8	0.0010	0.0309	0.0400	0.0115	0.0405	0.0494
G9	0.0016	0.0322	0.0340	0.0223	0.0400	0.0432
G10	0.0028	0.0155	0.0165	0.0285	0.0328	0.0331
G11	0.0023	0.0200	0.0245	0.0120	0.0385	0.0412
G12	0.0050	0.0269	0.0309	0.0175	0.0295	0.0339
G13	0.0043	0.0068	0.0218	0.0164	0.0193	0.0278
G14	0.0027	0.0159	0.0300	0.0175	0.0204	0.0400
G15	0.0085	0.0310	0.0325	0.0200	0.0413	0.0506
G16	0.0015	0.0159	0.0354	0.0139	0.0293	0.0367
G17	0.0028	0.0204	0.0280	0.0208	0.0292	0.0375
G18	0.0060	0.0553*	0.0564	0.0235	0.0495*	0.0555
G19	0.0060	0.0075	0.0259	0.0170	0.0240	0.0470
G20	0.0055	0.0389	0.0400	0.0250	0.0391	0.0407
G21	0.0030	0.0316	0.0343	0.0170	0.0293	0.0315
G22	0.0020	0.0434	0.0598	0.0040	0.0162	0.0316
G23	0.0015	0.0199	0.0292	0.0079	0.0316	0.0409
G24	0.0022	0.0107	0.0490	0.0220	0.0285	0.0570*

G25	0.0048	0.0219	0.0284	0.0238	0.0298	0.0379
-----	--------	--------	--------	--------	--------	--------

G26	0.0045	0.0052	0.0114	0.0148	0.0152	0.0251
G27	0.0025	0.0384	0.0390	0.0117	0.0388	0.0470
G28	0.0014	0.0294	0.0330	0.0140	0.0375	0.0384
G29	0.0025	0.0247	0.0364	0.0175	0.0300	0.0360
G30	0.0085	0.0293	0.0305	0.0170	0.0319	0.0422
G31	0.0025	0.0169	0.0320	0.0194	0.0270	0.0343
G32	0.0013	0.0293	0.0300	0.0120	0.0353	0.0365
G33	0.0025	0.0219	0.0303	0.0200	0.0295	0.0404
G34	0.0035	0.0315	0.0362	0.0098	0.0396	0.0414
G35	0.0006	0.0223	0.0249	0.0015	0.0230	0.0271
G36	0.0050	0.0281	0.0301	0.0176	0.0299	0.0312
G37	0.0023	0.0108	0.0205	0.0166	0.0328	0.0425
G38	0.0034	0.0293	0.0305	0.0185	0.0312	0.0451
G39	0.0040	0.0229	0.0345	0.0086	0.0216	0.0546
G40	0.0005	0.0076	0.0365	0.0011	0.0159	0.0295
G41	0.0010	0.0022	0.0525	0.0068	0.0088	0.0315
G42	0.0044	0.0202	0.0395	0.0065	0.0196	0.0306
G43	0.0026	0.0160	0.0216	0.0152	0.0195	0.0362
G44	0.0075	0.0233	0.0293	0.0022	0.0161	0.0205
G45	0.0029	0.0241	0.0328	0.0020	0.0136	0.0201
G46	0.0057	0.0126	0.0309	0.0059	0.0098	0.0193
G47	0.0011	0.0297	0.0302	0.0157	0.0192	0.0288
G48	0.0022	0.0187	0.0220	0.0138	0.0321	0.0350
G49	0.0013	0.0303	0.0340	0.0098	0.0131	0.0235
G50	0.0042	0.0295	0.0305	0.0075	0.0243	0.0351
G51	0.0011	0.0097	0.0139	0.0033	0.0101	0.0273

G52	0.0005	0.0293	0.0617*	0.0033	0.0206	0.0525
G53	0.0005	0.0122	0.0224	0.0033	0.0200	0.0290

UNDER PEER REVIEW

G54	0.0041	0.0222	0.0410	0.0236	0.0310	0.0432
G55	0.0027	0.0107	0.0285	0.0186	0.0225	0.0246
G56	0.0071	0.0286	0.0386	0.0093	0.0225	0.0385
G57	0.0017	0.0025	0.0295	0.0060	0.0074	0.0317
G58	0.0020	0.0110	0.0275	0.0104	0.0239	0.0290
G59	0.0038	0.0078	0.0428	0.0075	0.0155	0.0388
G60	0.0042	0.0317	0.0345	0.0075	0.0132	0.0335
G61	0.0010	0.0239	0.0789*	0.0100	0.0210	0.0626*
G62	0.0064	0.0310	0.0320	0.0248	0.0290	0.0310
G63	0.0027	0.0299	0.0298	0.0028	0.0324	0.0355
G64	0.0043	0.0150	0.0501	0.0189	0.0295	0.0543
G65	0.0088	0.0166	0.0390	0.0082	0.0237	0.0305
G66	0.0055	0.0141	0.0201	0.0090	0.0111	0.0127
G67	0.0064	0.0225	0.0410	0.0216	0.0281	0.0526
G68	0.0031	0.0336	0.0454	0.0227	0.0284	0.0433
G69	0.0020	0.0243	0.0253	0.0140	0.0163	0.0253
G70	0.0039	0.0109	0.0619*	0.0219	0.0292	0.0600*
G71	0.0012	0.0212	0.0475	0.0084	0.0110	0.0414
G72	0.0028	0.0290	0.0435	0.0222	0.0229	0.0245
G73	0.0012	0.0171	0.0242	0.0017	0.0219	0.0230
G74	0.0020	0.0435	0.0452	0.0162	0.0255	0.0408
G75	0.0020	0.0230	0.0239	0.0136	0.0245	0.0385
G76	0.0014	0.0280	0.0386	0.0108	0.0289	0.0349
G77	0.0038	0.0292	0.0463	0.0099	0.0230	0.0528
G78	0.0020	0.0329	0.0342	0.0058	0.0327	0.0344
G79	0.0022	0.0245	0.0315	0.0106	0.0289	0.0299
G80	0.0015	0.0070	0.0070	0.0040	0.0050	0.0050
G81	0.0026	0.0245	0.0280	0.0225	0.0254	0.0327

G82	0.0025	0.0099	0.0388	0.0184	0.0237	0.0350
G83	0.0055	0.0175	0.0285	0.0116	0.0254	0.0262
G84	0.0032	0.0224	0.0333	0.0187	0.0295	0.0304
G85	0.0027	0.0279	0.0647*	0.0162	0.0186	0.0516
G86	0.0028	0.0113	0.0565	0.0227	0.0252	0.0439
G87	0.0050	0.0321	0.0350	0.0074	0.0245	0.0252
G88	0.0014	0.0214	0.0522	0.0028	0.0125	0.0407
G89	0.0059	0.0290	0.0424	0.0143	0.0317	0.0360
G90	0.0040	0.0232	0.0390	0.0053	0.0139	0.0159
G91	0.0025	0.0246	0.0400	0.0105	0.0169	0.0232
G92	0.0029	0.0332	0.0386	0.0225	0.0264	0.0321
G93	0.0021	0.0322	0.0439	0.0034	0.0241	0.0505
G94	0.0020	0.0198	0.0450	0.0155	0.0199	0.0362
G95	0.0022	0.0316	0.0361	0.0101	0.0235	0.0435
G96	0.0005	0.0380	0.0407	0.0009	0.0369	0.0450
G97	0.0015	0.0329	0.0392	0.0053	0.0304	0.0345
G98	0.0044	0.0113	0.0380	0.0076	0.0180	0.0352
G99	0.0033	0.0135	0.0151	0.0180	0.0293	0.0312
G100	0.0010	0.0283	0.0302	0.0070	0.0293	0.0303
G101	0.0024	0.0225	0.0284	0.0140	0.0244	0.0255
Mean	0.0031	0.0230	0.0354	0.0130	0.0255	0.0363
	CD(0.05)= 0.0266 CD(0.01)= 0.0347 SED= 0.0129			CD(0.05)= 0.0202 CD(0.01)= 0.0265 SED= 0.0098		

*Significant at 5% level

**Significant at 1% level

Table 5. Mean performance of dry weight of roots per plant and dry weight of whole plant for black gram genotypes

Genotypes	Dry weight of roots per plant (g)			Dry weight of whole plant (g)		
	T ₁	T ₂	T ₃	T ₁	T ₂	T ₃
G1	0.0106	0.0386**	0.0399**	0.0395	0.1203**	0.1409*
G2	0.0021	0.0319**	0.0320	0.0200	0.0686	0.1100
G3	0.0091	0.0297*	0.0327	0.0303	0.0959	0.1014
G4	0.0139	0.0249	0.0251	0.0360	0.0733	0.0956
G5	0.0085	0.0141	0.0224	0.0265	0.0550	0.0881
G6	0.0007	0.0336**	0.0365*	0.0151	0.0886	0.1210
G7	0.0005	0.0125	0.0130	0.0026	0.0785	0.0816
G8	0.0050	0.0205	0.0259	0.0175	0.0919	0.1153
G9	0.0174*	0.0223	0.0232	0.0412	0.0944	0.1003
G10	0.0043	0.0157	0.0164	0.0356	0.0640	0.0660
G11	0.0022	0.0248	0.0305	0.0165	0.0833	0.0962
G12	0.0052	0.0216	0.0230	0.0277	0.0780	0.0877
G13	0.0149	0.0175	0.0375*	0.0355	0.0435	0.0870
G14	0.0060	0.0105	0.0306	0.0262	0.0468	0.1006
G15	0.0050	0.0109	0.0146	0.0335	0.0831	0.0976
G16	0.0127	0.0159	0.0196	0.0280	0.0610	0.0916
G17	0.0025	0.0117	0.0145	0.0260	0.0612	0.0800
G18	0.0050	0.0177	0.0317	0.0345	0.1225**	0.1435**
G19	0.0106	0.0129	0.0321	0.0335	0.0444	0.1050
G20	0.0060	0.0141	0.0161	0.0365	0.0920	0.0968
G21	0.0075	0.0127	0.0139	0.0275	0.0735	0.0797
G22	0.0296**	0.0391**	0.0451**	0.0356	0.0987	0.1364*
G23	0.0029	0.0148	0.0226	0.0123	0.0663	0.0926
G24	0.0050	0.0137	0.0156	0.0292	0.0528	0.1216

G25	0.0030	0.0083	0.0487**	0.0316	0.0599	0.1149
-----	--------	--------	----------	--------	--------	--------

Con...

UNDER PEER REVIEW

G26	0.0082	0.0087	0.0229	0.0275	0.0291	0.0593
G27	0.0144	0.0205	0.0255	0.0285	0.0977	0.1114
G28	0.0011	0.0266	0.0272	0.0165	0.0935	0.0986
G29	0.0080	0.0175	0.0368	0.0280	0.0722	0.1092
G30	0.0055	0.0105	0.0129	0.0310	0.0716	0.0856
G31	0.0102	0.0132	0.0459**	0.0320	0.0570	0.1122
G32	0.0022	0.0299*	0.0310	0.0155	0.0945	0.0975
G33	0.0061	0.0106	0.0124	0.0285	0.0620	0.0830
G34	0.0153	0.0204	0.0259	0.0285	0.0915	0.1035
G35	0.0005	0.0332**	0.0424**	0.0025	0.0785	0.0943
G36	0.0044	0.0220	0.0246	0.0270	0.0800	0.0859
G37	0.0059	0.0148	0.0293	0.0248	0.0584	0.0923
G38	0.0046	0.0326**	0.0359*	0.0265	0.0931	0.1115
G39	0.0030	0.0266	0.0465**	0.0155	0.0711	0.1356*
G40	0.0002	0.0041	0.0145	0.0017	0.0275	0.0805
G41	0.0056	0.0061	0.0238	0.0133	0.0171	0.1078
G42	0.0041	0.0337**	0.0421**	0.0150	0.0735	0.1122
G43	0.0077	0.0100	0.0142	0.0255	0.0455	0.0720
G44	0.0149	0.0185	0.0236	0.0246	0.0578	0.0734
G45	0.0240**	0.0339**	0.0341	0.0289	0.0715	0.0869
G46	0.0108	0.0161	0.0237	0.0224	0.0385	0.0738
G47	0.0085	0.0195	0.0223	0.0253	0.0683	0.0812
G48	0.0086	0.0307*	0.0327	0.0245	0.0815	0.0897
G49	0.0146	0.0476**	0.0551**	0.0256	0.0909	0.1126
G50	0.0011	0.0125	0.0139	0.0127	0.0663	0.0795
G51	0.0243**	0.0276	0.0300	0.0287	0.0474	0.0711
G52	0.0013	0.0091	0.0295	0.0050	0.0589	0.1436**
G53	0.0321**	0.0364**	0.0448**	0.0359	0.0686	0.0962
G54	0.0081	0.0135	0.0341	0.0358	0.0667	0.1183

G55	0.0150	0.0173	0.0200	0.0363	0.0505	0.0731
G56	0.0031	0.0054	0.0242	0.0194	0.0565	0.1013
G57	0.0126	0.0142	0.0259	0.0203	0.0240	0.0871
G58	0.0009	0.0241	0.0299	0.0133	0.0590	0.0864
G59	0.0048	0.0068	0.0142	0.0160	0.0300	0.0957
G60	0.0098	0.0209	0.0247	0.0215	0.0657	0.0927
G61	0.0015	0.0144	0.0202	0.0125	0.0593	0.1617**
G62	0.0089	0.0204	0.0223	0.0401	0.0823	0.0834
G63	0.0199*	0.0244	0.0264	0.0254	0.0867	0.0917
G64	0.0012	0.0046	0.0358*	0.0243	0.0490	0.1402*
G65	0.0024	0.0103	0.0130	0.0193	0.0505	0.0825
G66	0.0258**	0.0301*	0.0305	0.0403	0.0553	0.0633
G67	0.0071	0.0195	0.0291	0.0350	0.0700	0.1226
G68	0.0030	0.0092	0.0231	0.0287	0.0712	0.1117
G69	0.0006	0.0149	0.0308	0.0166	0.0555	0.0814
G70	0.0155	0.0201	0.0215	0.0412	0.0601	0.1433*
G71	0.0080	0.0203	0.0296	0.0176	0.0525	0.1185
G72	0.0072	0.0147	0.0341	0.0321	0.0666	0.1021
G73	0.0228**	0.0306*	0.0331	0.0256	0.0696	0.0803
G74	0.0080	0.0146	0.0171	0.0262	0.0835	0.1031
G75	0.0010	0.0192	0.0203	0.0165	0.0667	0.0827
G76	0.0006	0.0201	0.0290	0.0128	0.0770	0.1025
G77	0.0023	0.0128	0.0194	0.0159	0.0650	0.1184
G78	0.0037	0.0182	0.0199	0.0115	0.0838	0.0885
G79	0.0122	0.0216	0.0266	0.0250	0.0750	0.0880
G80	0.0302**	0.0385**	0.0386*	0.0357	0.0505	0.0506
G81	0.0054	0.0069	0.0086	0.0305	0.0568	0.0692

G82	0.0125	0.0166	0.0199	0.0334	0.0501	0.0937
-----	--------	--------	--------	--------	--------	--------

Con...

UNDER PEER REVIEW

G83	0.0140	0.0179	0.0250	0.0310	0.0607	0.0797
G84	0.0073	0.0212	0.0249	0.0291	0.0730	0.0885
G85	0.0073	0.0160	0.0256	0.0262	0.0625	0.1418*
G86	0.0061	0.0140	0.0221	0.0315	0.0505	0.1225
G87	0.0107	0.0216	0.0311	0.0230	0.0782	0.0913
G88	0.0039	0.0149	0.0297	0.0080	0.0488	0.1225
G89	0.0029	0.0180	0.0198	0.0230	0.0787	0.0982
G90	0.0125	0.0446**	0.0485**	0.0218	0.0816	0.1034
G91	0.0155	0.0289	0.0302	0.0285	0.0704	0.0934
G92	0.0088	0.0114	0.0130	0.0342	0.0709	0.0836
G93	0.0050	0.0338**	0.0399	0.0105	0.0901	0.1343
G94	0.0059	0.0109	0.0294	0.0234	0.0505	0.1105
G95	0.0014	0.0180	0.0231	0.0136	0.0731	0.1027
G96	0.0002	0.0165	0.0207	0.0016	0.0914	0.1064
G97	0.0112	0.0207	0.0212	0.0180	0.0840	0.0949
G98	0.0051	0.0111	0.0258	0.0170	0.0404	0.0990
G99	0.0066	0.0195	0.0207	0.0279	0.0623	0.0670
G100	0.0091	0.0243	0.0272	0.0171	0.0819	0.0877
G101	0.0023	0.0151	0.0168	0.0187	0.0620	0.0707
Mean	0.0082	0.0195	0.0267	0.0243	0.0679	0.0984
		CD(0.05)= 0.0089 CD(0.01)= 0.0117 SED= 0.0045		CD(0.05)= 0.0343 CD(0.01)= 0.0452 SED= 0.0174		

*Significant at 5% level

**Significant at 1% level



Fig.1. Overview of the blackgram seedlings on 6DAS

UNDER PEER REVIEW



Fig. 2. Response of blackgram seedlings to T₃ and T₂ on 18 DAS



Fig.3. Response of the susceptible blackgramseedlings (RU-8-708) to salinity




Fig. 4. Performance of saline resistant blackgram genotypes for various seedling growth characters