

Original Research Article

Antioxidant Properties Comparison of Trunk's Barks of 10 Plants used in Traditional Medicine Against Hepatic Pathologies

ABSTRACT

Aims: Herbal remedies are known for their anti-inflammatory and antioxidant properties. The aim of this study was to compare the antioxidant activities of the aqueous extracts of the bark of the trunk of 10 medicinal plants used in traditional medicine for the management of hepatic pathologies. This is *Acacia nilotica* (L.) Willd. ex Delile (Fabaceae – Mimosoideae), *Adansonia digitata* L. (Bombacaceae), *Bombax costatum* Pellegr. & Vuill. (Bombacaceae), *Balanites aegyptiaca* (L.) Delile (Balanitaceae), *Cassia sieberiana* DC. (Fabaceae – Caesalpinioideae), *Piliostigma reticulatum* (DC.) Hochst. (Fabaceae – Caesalpinioideae), *Tamarindus indica* L. (Fabaceae – Caesalpinioideae), *Daniellia oliveri* (Rolfe) Hutch. & Dalziel (Fabaceae – Caesalpinioideae), *Khaya senegalensis* (Desr.) A.Juss. (Meliaceae) and *Gymnosporia senegalensis* (Lam.) Loes. (Celastraceae)..

Methodology: In this study, the total phenolic and flavonoid content of the aqueous extracts of the trunk bark of 10 plants were determined and their antioxidant activities by the DPPH and FRAP methods were evaluated. A correlation has been established between phenolic compounds and antioxidant activity.

Results: *Acacia nilotica* had the best contents of total phenolics and flavonoids respectively with 21.28 ± 0.18 g ETA / 100 g DM and 0.207 ± 0.003 g EQ / 100 g DM. For the evaluation of antioxidant activity, *Acacia nilotica* also gave the best activities by the DPPH method with a percent inhibition of 1.08 ± 0.03 AAE and a reducing capacity of Fe^{3+} to Fe^{2+} of 0.107 ± 0.03 AAE. A strong correlation was found between FRAP and total phenolics ($r^2 = 0.9559$).

Conclusion: *Acacia nilotica* having good antioxidant activity, it could be used for anti-inflammatory or even hepatoprotective properties

Keywords: Medicinal plants, total phenolics, flavonoids, DPPH, FRAP

1. INTRODUCTION

Liver disease (acute and chronic) is of global concern (1) and epidemiological studies have estimated that 3.5% of the world's population are chronically infected with viral hepatitis (2). Viral hepatitis B (HBV) has a prevalence in children in Africa of 3% (2). In Burkina Faso, its prevalence had already exceeded 12% since 2013 according to Lazarus et al. (3). More and more, other types of hepatitis are developing around the world (drugs, alcoholics, toxicants, etc.) (4). They are at the origin of a strong production of free radicals which can be at the origin of other pathologies or aggravate the hepatic consequences. Medical treatments for these diseases are often difficult to achieve and may have limited effectiveness. These are traditionally treated with medicinal plants especially in rural areas where they are widely available (1). These plants are widely used for their hepatoprotective activity through their

Comment [DN1]: Is this method or conclusion?

Comment [DN2]: Should be based on the problem statement or topic and not just a single plant.

Comment [DN3]: Who is being referred to as they here?

antioxidant, anti-inflammatory, etc. Herbal remedies provide a natural source of phytochemicals with biological activity, such as antioxidant properties (5). Among these phytoconstituents, phenolic compounds and flavonoids are the most well known for their antioxidant properties (6). In its program to combat hepatic diseases, the Institute for Research in Health Sciences carried out a survey in the Center-West region of Burkina Faso in order to identify the plants and parts of plants used for treatment of his pathologies. At the end of the survey, ten plants were selected according to the part used (trunk bark). The aim of this study was to compare the total phenolic and flavonoid contents as well as the antioxidant activity of the aqueous extracts of the trunk bark of the different plants selected.

Comment [DN4]: Antioxidant, antiinflammatory what?

Comment [DN5]: Why was the aim to compare the activities and not rather evaluate them. Its necessary to investigate and validate the traditional claims before thinking of comparisons.

2. MATERIAL AND METHODS

2.1. Chemicals and reagents used

1,1-diphenyl-2-picrylhydrazyl (DPPH), ascorbic acid and quercetin were purchased from Sigma Chemical Co. (St. Louis, MO, USA); potassium hexacyanoferrate [$K_3Fe(CN)_6$], trichloroacetic acid, trichloroferrate [$FeCl_3$] for evaluation of antioxidant activity, and Follin-ciocalteu and sodium carbonate, $AlCl_3$, for phenolics compounds

2.2. Plant material and extraction

The plant material consisted of the trunks bark of 10 plants collected in Ouagadougou in the province of Kadiogo in Burkina Faso: *Acacia nilotica* (L.) Willd. ex Delile (Fabaceae – Mimosoideae), *Adansonia digitata* L. (Bombacaceae), *Bombax costatum* Pellegr. & Vuill. (Bombacaceae), *Balanites aegyptiaca* (L.) Delile (Balanitaceae), *Cassia sieberiana* DC. (Fabaceae – Caesalpinioideae), *Piliostigma reticulatum* (DC.) Hochst. (Fabaceae – Caesalpinioideae) *Tamarindus indica* L. (Fabaceae – Caesalpinioideae), *Daniellia oliveri* (Rolfe) Hutch. & Dalziel (Fabaceae – Caesalpinioideae), *Khaya senegalensis* (Desr.) A.Juss. (Meliaceae) and *Gymnosporia senegalensis* (Lam.) Loes. (Celastraceae).

A maceration was used at 10%, ie 2.5 g of plant material in 2.5 ml of distilled water, for 24 hours at laboratory temperature. After 24 hours, the extract was first filtered with a nylon filter and then centrifuged at 3000 rpm for 10 min. The supernatant obtained was recovered and packaged in 4 Eppendorf tubes then placed in the freezer at $-4^{\circ}C$.

Comment [DN6]: What was the traditional mode of preparation from the survey? Is this method seeking to replicate it or modify it?

Comment [DN7]: Is this 10%

2.3. Determination of phenolic compounds in plant extracts

2.3.1. Total phenolics content

The total phenolic compounds were measured according to the Singleton method (7). These compounds react with the Folin Ciocalteu reagent (FCR) in an alkaline medium. The reaction mixture consisted of 1 ml of extract (1 mg / ml), 1 ml of 2N FCR and 3 ml of a 20% sodium carbonate solution in a test tube. After incubation for 40 min at room temperature, the absorbance of the mixture was measured at 760 nm on the spectrophotometer (Agilent 8453) against a tannic acid standard curve ($R^2 = 0.999$). In the blank control tube, the extract was replaced with distilled water. The tests were realized and the total phenol concentration of the extract, expressed in gram-equivalent tannic acid per 100 g dry matter (DM) (g ETA / 100 g DM), was calculated.

2.3.2. Total flavonoids content

The flavonoid dosage was realized according to the Kumaran method (8) adapted by Abdel-Hameed (9). Two (2) ml of extract (1 mg / ml in methanol) were mixed with 2 ml of aluminium trichloride (2%). After 40 min of incubation at ambient temperature, the absorbance was measured at 415 nm using a spectrophotometer (Agilent 8453) against a quercetin standard curve of $R^2 = 0.999$. The blank control tube consisted of 2 ml of methanol. The quantity of flavonoids in the plant extract was determined in gram-equivalent quercetin (EQ) per 100 g dry matter (DM) (g EQ / 100 g DM).

2.4. Evaluation of antioxidant activities

2.4.1. FRAP test (Ferric Reducing Antioxidant Power)

The FRAP (Ferric Reducing Antioxidant Power) method is based on the reduction of ferric ion (Fe^{3+}) to ferrous ion (Fe^{2+}). This method evaluates the reducing power of the compounds. The reducing power of the samples was evaluated according to the spectrophotometric method described by Apati et al., (10). A volume of 1 ml of the extract was mixed with 2.5 ml of phosphate buffer (0.2 M, pH 6.6) and 2.5 ml of potassium ferricyanide [$K_3Fe(CN)_6$] 1%. The mixture was incubated at 50 ° C for 20 min; 2.5 ml of 10% trichloroacetic acid (TCA) were then added before centrifuging the mixture at 3000 rpm for 10 min. A volume of 2.5 ml of the obtained supernatant was then taken and mixed with 2.5 ml of distilled water and 0.5 ml of 0.1% ferric chloride ($FeCl_3$) 0,1%. In the white control tube, the extract was replaced with 1 ml of the extraction solvent (distilled water or ethanol). The absorbance was read at 700 nm using the spectrophotometer (Agilent 8453) against a standard curve of ascorbic acid (0-0.2 mg / ml). For each plant, the measurements were carried out in quadruple from two independent tests. An increase in absorbance reflects an increase in the reducing power of the sample. The results are expressed in ascorbic acid equivalents (Ascorbate Acid Equivalent = AAE) per milliliter. When the AAE value is equal to 1 AAE.ml⁻¹, the reducing power of 1 ml of extract is equivalent to 1 μ mol of ascorbic acid. The value of the AAE is inversely proportional to the reducing power.

2.4.2. DPPH test (2,2-diphényl-1-picrylhydrazyl)

Free radical scavenging activity assessment (DPPH' assay): The antioxidant activity of the crude extracts and fractions was assessed by the mean of 2,2-diphenyl-1-picrylhydrazyl (DPPH') colorimetric method as described by Velazquez et al., (11), slightly modified. This method depends on the reduction of purple DPPH to a yellow-colored diphenyl picrylhydrazine and the remaining DPPH', which showed maximum absorption at 517 nm was measured (spectrophotometer Agilent 8453E). About 2 mL of a 20 mg mL⁻¹ DPPH' solution were added to 1 mL of a methanolic solution of each extract (1-100 μ g mL⁻¹). A mixture of 2 mL of DPPH' and 1 mL of methanol served as control. The mixture was shaken vigorously then incubated for 15 min in darkness at room temperature. Absorbance was measured at 517 nm. Methanol was used as blank. Ascorbic acid solution was used as positive controls. Each experiment was performed in triplicate. The DPPH' radical scavenging activity was calculated. Decreasing of the DPPH solution absorption indicates an increase of DPPH radical scavenging activity. For each plant, the measurements are carried out in quadruple from two independent tests. The results are expressed in mg content of antioxidants ascorbic acid equivalent (AAE).

2.5. Statistical analysis

All the assays were carried out in triplicate and each experiment was independently repeated at least three times, through which means and standard deviations (SD) were

generated. The results were analyzed with GraphPad Prism version 6, using one-way analysis of variance (ANOVA) followed by Tukey's multiple comparisons test with $P = 0.05$.

3. RESULTS AND DISCUSSION

3.1. Results

The Residual moisture content of the different powders varied from $4,29 \pm 0,24$ with *Tamarindus indica* to $7,43 \pm 0,18$ for *Gymnosporia senegalensis* (Table 1).

Comment [DN8]: This was not mentioned in the method. What is its relevance to the project under study?

Table 1 : Residual moisture content of different powder of different plants

Plants	Residual moisture content %
<i>Acacia nilotica</i>	$5,56 \pm 0,11^a$
<i>Adansonia digitata</i>	$7,07 \pm 0,84^b$
<i>Balanites aegyptiaca</i>	$4,37 \pm 0,17^c$
<i>Bombax costatum</i>	$5,02 \pm 0,13^a$
<i>Cassia sieberiana</i>	$5,38 \pm 0,05^a$
<i>Daniellia oliveri</i>	$7,03 \pm 0,03^b$
<i>Khaya senegalensis</i>	$6,60 \pm 0,07^d$
<i>Gymnosporia senegalensis</i>	$7,43 \pm 0,18^b$
<i>Piliostigma reticulatum</i>	$6,26 \pm 0,10^d$
<i>Tamarindus indica</i>	$4,29 \pm 0,24^c$

*Moisture content on oven dry weight basis

3.1.1. Phenolic compounds in plant extracts

The results of the total phenolic and flavonoid contents of the aqueous extracts of the trunk bark of the plants are shown in Figures 1 and 2. The total phenolic contents of the different extracts varied from $1,30 \pm 0,10$ g ETA/100 g DM for extracts of *Balanites aegyptiaca* to $21,28 \pm 0,18$ g ETA/100 g DM for extracts of *Acacia nilotica*. There is no statistically significant difference ($p < 0.05$) between the total phenolic contents of extracts of *Khaya senegalensis* ($14,91 \pm 0,11$ g ETA / 100 g DM) and *Piliostigma reticulatum* ($15,06 \pm 0,6$ g ETA / 100 g DM) and between those of *Adansonia digitata* ($1,64 \pm 0,4$ g ETA / 100 g DM) and *Balanites aegyptiaca* ($1,30 \pm 0,10$ g ETA / 100 g DM). As regards the flavonoid contents of the extracts of the trunk bark, they varied from $0,018 \pm 0,001$ g EQ / 100 g DM for extracts of *Balanites aegyptiaca* to $0,209 \pm 0,005$ g EQ / 100 g DM for extracts of *Cassia sieberiana*. There is no statistically significant difference ($p < 0.05$) between the flavonoid content of extracts of *Acacia nilotica* ($0,207 \pm 0,003$ g EQ / 100 g DM) and *Cassia sieberiana* ($0,209 \pm 0,005$ g EQ / 100 g DM), between *Gymnosporia senegalensis* ($0,069 \pm 0,0009$ g EQ / 100 g DM) and *Piliostigma reticulatum* ($0,07 \pm 0,007$ g EQ / 100 g DM).




Figure 1 : Total phenolic and flavonoid content of extracts from different plants
 *AD: *Adansonia digitata*, AN: *Acacia nilotica*, BA: *Balanites aegyptiaca*, BC: *Bombax costatum*, CS: *Cassia sieberiana*, DO: *Daniellia oliveri*, KS: *Khaya senegalensis*, GS: *Gymnosporia senegalensis*, PR: *Piliostigma reticulatum*, TI: *Tamarindus indica*
 Test drugs: significant from each drug, * $P < 0.05$
 Mean \pm S.E.M = Mean values \pm Standard error of means of 3 experiments

3.1.2. Antioxidant's activities in plant extracts

The antioxidant activity of the aqueous extracts of the bark of the trunks of different medicinal plants was evaluated using the DPPH free radical scavenging method and the method of the reduction of ferric ion (Fe^{3+}) to ferrous ion (Fe^{2+}). The results of the antioxidant activities are shown in Figure 2. The anti-free radical activity by the DPPH method of the different plants ranged from 0.069 ± 0.006 AAE for *Balanites aegyptiaca* to 1.08 ± 0.03 AAE for *Acacia nilotica*. There is no statistically significant difference ($p < 0.05$) between the anti-free radical activities of extracts from the trunk bark of *Bombax costatum* (0.518 ± 0.001 AAE), *Cassia sieberiana* (0.544 ± 0.05 AAE), *Daniellia oliveri* (0.503 ± 0.03 AAE), *Khaya senegalensis* (0.5 ± 0.02 AAE), and *Piliostigma reticulatum* (0.534 ± 0.09 AAE) and between *Tamarindus indica* (0.263 ± 0.08 AAE) and *Adansonia digitata* (0.27 ± 0.002 AAE). The reducing capacity of Fe^{3+} in Fe^{2+} of the extracts varied from 0.0023 ± 0.0006 AAE for *Balanites aegyptiaca* to 0.107 ± 0.03 AAE for *Acacia nilotica*. There is a statistically significant difference ($p < 0.05$) between extracts of *Daniellia oliveri* (0.077 ± 0.003 AAE), *Khaya senegalensis* (0.063 ± 0.002 AAE), *Gymnosporia senegalensis* (0.0704 ± 0.0009 AAE) and *Piliostigma reticulatum* (0.059 ± 0.007 AAE), between extracts of *Bombax costatum* (0.040 ± 0.001 AAE) and *Cassia sieberiana* (0.032 ± 0.005 AAE) and finally between extracts of *Tamarindus indica* (0.016 ± 0.007 AAE), *Adansonia digitata* (0.0063 ± 0.0002 AAE) and *Balanites aegyptiaca* (0.0023 ± 0.0006 AAE).




Figure 2 : Antioxidant activity by the FRAP and DPPH method of extracts from different plants

AD: *Adansonia digitata*, AN: *Acacia nilotica*, BA: *Balanites aegyptiaca*, BC: *Bombax costatum*, CS: *Cassia sieberiana*, DO: *Daniellia oliveri*, KS: *Khaya senegalensis*, GS: *Gymnosporia senegalensis*, PR: *Piliostigma reticulatum*, TI: *Tamarindus indica*

Test drugs: significant from each drug, * $P < 0.05$

Mean \pm S.E.M = Mean values \pm Standard error of means of 3 experiments

3.1.3. Correlations

The correlation between the contents of total phenolics, flavonoids and antioxidant activities by DPPH and FRAP is shown in Figure 3. A weak correlation was found between DPPH and total phenolics ($r^2 = 0.5054$) and between DPPH and flavonoids ($r^2 = 0.6245$). A strong correlation was found between FRAP and total phenolics ($r^2 = 0.9559$) and an $r^2 = 0.0495$ observed between FRAP and flavonoids.




Figure 3 : Correlation between FRAP and the total phenolics content (A) and between FRAP and total flavonoid content (B); Correlation between DPPH and the total phenolics content (C) and between DPPH and total flavonoid content (D)

3.2. DISCUSSION

The RMC of the powders of the trunk bark of the different plants was less than 10; this low humidity would indicate better storage and it helps prevent possible enzymatic reactions as well as the development of microorganisms (12). In fact, too high a water content could promote the deterioration of the microbiological quality, enzymatic reactions having consequences on the variation of secondary metabolites during the shelf life of the plant material powder. Thus, residual humidity would also promote the proliferation of microorganisms such as bacteria, yeasts and molds (13).

Comment [DN9]: What is meant by correlation in this sense? Why was it done?

Comment [DN10]: Isn't the moisture content dependent on the handling and storage techniques? This means the value is likely to change depending on how the samples are handled. Thus, not a major point.

Phenolic compounds are biologically active molecules which are widely used in therapy as vasoconstrictors, anti-inflammatories, enzymatic inhibitors, antioxidants and anti-radicals, antimicrobials, etc. Found in all parts of plants (roots, stems, leaves, flowers, pollens, fruits, grains and wood), phenolic compounds form the most important group of phytochemicals in plants (14). The levels of phenolic compounds vary from species to species, from part of the plant to another and depending on edaphic factors as well (15). A comparison of the contents of phenolic compounds, in particular the total phenolics and the flavonoids of the aqueous extracts of the bark of the trunks of different plants from the same family or not, is very important in the choice of the plant for possible pharmacological activities. The determination of total phenolics showed that *Acacia nilotica* had the highest levels followed by *Daniellia oliveri* and that the lowest levels were with *Balanites aegyptiaca* and *Adansonia digitata*. These results corroborate those of Boly et al. (16) who had found higher levels of phenolic compounds with extracts of *Acacia nilotica* and those of Traoré et al. (17) who after a comparative study on 5 plants found that *Balanites aegyptiaca* had the lowest levels. By comparing the flavonoid contents of the 10 species, it turns out that it is *Acacia nilotica* and *Cassia sieberiana* which gave the best levels and the lowest level always with *Balanites aegyptiaca*. These results always corroborate those of the two authors who in their respective studies had similar results. These phenolic compounds are the main compounds having their aromatic nucleus which allows them to stabilize and relocate unpaired electrons from their structures by giving up hydrogen atoms and electrons from their hydroxyl groups (15). They are known to act as antioxidants not only because of their ability to donate electrons, but also because of their stable radical intermediates, which can effectively prevent oxidation at the cellular and physiological level (18).

Phenolic compounds, in particular phenols, can adjust the concentration of ROS, thus activating a network of biochemical events to increase tolerance, hence the importance of studying the antioxidant activity of plant species (15). The two methods used make it possible to have a certain complementarity of the antioxidant activities of different plants. The results of the antioxidant activities by the two methods (DPPH and FRAP) showed that the aqueous extracts of *Acacia nilotica* had the best activities and the most activities were with the aqueous extracts of *Balanites aegyptiaca*. Several authors have shown that *Acacia nilotica* extracts have good antioxidant activity (16, 19). The work of Traoré et al. showed low antioxidant activity in aqueous extracts of *Balanites aegyptiaca* (17). Six plants had the same statistically significant activities by the DPPH method and among the six there are four which have the same activity by the FRAP method. The difference between the antioxidant activities of the different ones is probably due to their composition and their content of phenolic compounds and this reduction of the DPPH radical and the ferric ions to ferrous ions is generally not due to the action of a single compound but to the interactions that may exist from one extract to another (20). *Acacia nilotica* which had the high contents of total phenolics and flavonoids, gave the best antioxidant activities and *Balanites aegyptiaca* which had the lowest contents, also gave the lowest antioxidant activities. The high antioxidant capacities of the extracts of these plants would explain their ability to trap ROS. Several herbal studies have found a correlation between total phenolics and antioxidant power (21, 22). Phenolic acids, flavonoids and tannins are considered to be the main factors contributing to the antioxidant activity of medicinal plants. Indeed, the redox properties of these phenolic compounds would allow them to act as reducing agents, hydrogen donors, singlet oxygen deactivators and also metal chelators (23). Indeed, a good linear correlation between total phenolics and antioxidant activity would show that most of the antioxidant activity is due in part to the presence of a large part of total phenolics (8). Phenolic compounds could probably act, in synergy or in isolation, through their antioxidant properties to protect hepatocytes against free radicals and reactive oxygen forms generated by attacks in the body.

4. CONCLUSION

This study showed that the aqueous extracts of the bark of the trunks of the plants had higher or lower total phenolic and flavonoid contents and antioxidant activities. *Acacia nilotica* exhibited superior antioxidant activity than that of other medicinal plants, which is confirmed by its high content of phenolic compound also compared to that of other plants. *Acacia nilotica* could be a good choice for the evaluation of anti-inflammatory and even hepatoprotective activity.

Comment [DN11]: As in?

Comment [DN12]:

REFERENCES

1. Ali SA, Sharief NH, Mohamed YS. Hepatoprotective Activity of Some Medicinal Plants in Sudan. *Evid Based Complement Alternat Med.* 2019;2019:2196315.
2. Tibiri A, Boria S, Traoré TK, Ouédraogo N, Nikiéma A, Ganaba S, et al. Countrywide Survey of Plants Used for Liver Disease Management by Traditional Healers in Burkina Faso. *Frontiers in Pharmacology.* 2020;11(1750).
3. Lazarus JV, Sperle I, Safreed-Harmon K, Gore C, Cebolla B, Spina A. Associations between national viral hepatitis policies/programmes and country-level socioeconomic factors: a sub-analysis of data from the 2013 WHO viral hepatitis policy report. *BMC public health.* 2018;18(1):1-7.
4. Srivastava R, Srivastava P. Hepatotoxicity and the Role of Some Herbal Hepatoprotective Plants in Present Scenario 2018.
5. de Jesús Joaquín-Ramos A, López-Palestina CU, Pinedo-Espinoza JM, Altamirano-Romo SE, Santiago-Saenz YO, Aguirre-Mancilla CL, et al. Phenolic compounds, antioxidant properties and antifungal activity of jarilla (*Barkleyanthus salicifolius* [Kunth] H. Rob & Brettell). *CHILEAN JOURNAL OF AGRICULTURAL RESEARCH.* 2020;80(3):352-60.
6. Huyut Z, Beydemir Ş, Gülçin İ. Antioxidant and Antiradical Properties of Selected Flavonoids and Phenolic Compounds. *Biochemistry Research International.* 2017;2017:7616791.
7. Singleton VL, Orthofer R, Lamuela-Raventos RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in enzymology.* 1999;299:152-78.
8. Kumaran A, Joel Karunakaran R. In vitro antioxidant activities of methanol extracts of five *Phyllanthus* species from India. *LWT - Food Science and Technology.* 2007;40(2):344-52.
9. Abdel-Hameed E-SS. Total phenolic contents and free radical scavenging activity of certain Egyptian *Ficus* species leaf samples. *Food Chemistry.* 2009;114(4):1271-7.
10. Apáti P, Szentmihályi K, Kristó ST, Papp I, Vinkler P, Szoke E, et al. Herbal remedies of *Solidago*--correlation of phytochemical characteristics and antioxidative properties. *Journal of pharmaceutical and biomedical analysis.* 2003;32(4-5):1045-53.
11. Velázquez E, Tournier HA, Mordujovich de Buschiazio P, Saavedra G, Schinella GR. Antioxidant activity of Paraguayan plant extracts. *Fitoterapia.* 2003;74(1-2):91-7.
12. Ouédraogo S, Sombié BC, Ouédraogo JCW, Traoré TK, Traoré S, Nitiéma M. Standardization of Extracts from Trunks's Barks of *Lannea microcarpa* engl. and K. Krause (Anacardiaceae) and *Anogeissus leiocarpus* (DC) Guill. and Perr.(Combretaceae) for the Formulation of Antihypertensive Herbal Medicines.". *Int J Pharm Sci Rev Res.* 2018;48(1):92-7.

13. Chabrier JY. Plantes médicinales et formes d'utilisation en Phytothérapie. : Université Henri Poincare - Nancy 1; 2010.
14. Zhang L, Ravipati AS, Koyyalamudi SR, Jeong SC, Reddy N, Smith PT, et al. Antioxidant and anti-inflammatory activities of selected medicinal plants containing phenolic and flavonoid compounds. *Journal of agricultural and food chemistry*. 2011;59(23):12361-7.
15. Chaves N, Santiago A, Alias JC. Quantification of the Antioxidant Activity of Plant Extracts: Analysis of Sensitivity and Hierarchization Based on the Method Used. *Antioxidants*. 2020;9(1):76.
16. Abdoul GLB, Traore A, Ouedraogo M, Belemilga M, Traore TK, Belemnaba L, et al. Pharmacological study of trunk bark of *Acacia nilotica* var *adansonii* (Guill et Perr). o Ktze (Mimosaceae): Assays, antioxidant and antispasmodic activities. *Journal of Drug Delivery and Therapeutics*. 2019;9(3-s):524-30.
17. Tata Kadiatou T, André T, Noufou O, Nogma Ernest S, Jotham Yhi-pene Nd, Salfo O, et al. Ethnopharmacological plants used to treat hepatitis and their anti-oxidant activity of district of Bobo-Dioulasso (Burkina Faso). *International journal of Pharmacological Research*. 2018;8:15.
18. Ince I, Kayalar H, Elgin G, Koksall C, Yavasoglu NU. Antioxidant, anti-inflammatory activities and acute toxicity of the polyherbal formulation: Romix®. *Pharm Biol*. 2012;50(6):720-6.
19. Sadiq MB, Hanpithakpong W, Tarning J, Anal AK. Screening of phytochemicals and in vitro evaluation of antibacterial and antioxidant activities of leaves, pods and bark extracts of *Acacia nilotica* (L.) Del. *Industrial Crops and Products*. 2015;77:873-82.
20. Ouédraogo S, Traoré TK, Ouedraogo B, Atchadé BC, Coulibaly A, Lompo M, et al. Phytochemical Characteristics and Pharmacological Properties Studies of the Powders of *Punica granatum* L.(Lythraceae) Leaves from Seven Regions of Burkina Faso with a View to Standardizing. *Journal of Advances in Medical and Pharmaceutical Sciences*. 2021:1-11.
21. Boizot N, Charpentier J-P. Méthode rapide d'évaluation du contenu en composés phénoliques des organes d'un arbre forestier. *Le Cahier des Techniques de l'INRA*, Numéro spécial. 2006:79-82.
22. Marc F, Davin A, Deglene-Benbrahim L, Ferrand C, Baccaunaud M, Fritsch P. Méthodes d'évaluation du potentiel antioxydant dans les aliments. *M/S: médecine sciences*. 2004;20(4):458-63.
23. Rice-Evans CA, Miller NJ, Paganga G. Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free radical biology and medicine*. 1996;20(7):933-56.