

CORRELATION BETWEEN NUMBER OF HYSTEROSALPINGOGRAPHY ABNORMALITIES AND AGE AMONG WOMEN WITH INFERTILITY IN A TERTIARY HOSPITAL IN PORT HARCOURT

ABSTRACT

Background: The number of abnormalities detected on Hysterosalpingography (HSG) varies with age and the type of infertility. The more the numbers of abnormalities, the slimmer the chance of conception naturally and fecundity decreases with increasing age. HSG is still useful in evaluating the reproductive systems in developing countries.

Aim: The aim of this study is to determine the number of hysterosalpingography abnormalities and to determine if there is any relationship with age and type of infertility among women who had HSG in a tertiary hospital in Port Harcourt.

Methods: A retrospective study involving a total of 226 patient's data, carried out in a tertiary hospital in Port Harcourt. A *P*-value below 0.05 was considered statistically significant.

Results: Two-hundred and twenty-six women were studied. The mean age of the study population was 35.07 ± 5.45 years and ranges from 22-48 years. Most women were 30-39 years and have the highest number of abnormalities. Normal HSG findings were seen in 64, 28.3% and abnormal HSG (162, 71.7%). There was significant association with age and number of abnormalities on HSG, with *P*-value of 0.037 (scatter plot). Significant association was also seen

between number of HSG abnormalities and type of infertility with a P -value of 0.028 (multiple linear regression).

Conclusion: Hysterosalpingography (HSG) is a valuable tool in investigating women with infertility and most frequently encountered type in this study was secondary infertility. More than 71% of women had abnormal HSG. There was significant association between number of abnormalities in HSG with age and type of infertility.

Keywords: Correlation, Infertility, Age, Women, Hysterosalpingography, Abnormalities

INTRODUCTION

Infertility definition varies considerably. Often referred to as the inability of a couple to achieve gestation/conception within a time not less than a year of adequate/ enough unprotected sexual contact or coitus.[1] Basically two types of infertility have been identified; primary if the woman has never achieved pregnancy over a period of one year and not using birth control. Secondary infertility occurs when the woman has been able to achieve at least one prior pregnancy. In sub-Saharan Africa and in other parts of the world infertility is a foremost clinico-social concern.[2,3] Past records had shown that about 10-20% of females within the reproductive age bracket in sub-Saharan Africa has infertility predicament.[4] Female infertility is considered as a misfortune and is often the grounds for separation and divorce or polygamy as well as denial of rights in family functions in Nigeria.[5] Recently, in Nigeria fertility rate is just about 5.25 children per woman in year 2020.[6]

HSG is widely used to study the gross anatomy of the uterine cavity and establish the patency of fallopian tubes. The number of abnormalities detected in Hysterosalpingography (HSG) varies with age and the type of infertility. The more the numbers of abnormalities the slimmer the chance of conception naturally. As fecundity decreases with increasing age.

The aim of this study is to determine the number of hysterosalpingography abnormalities and to determine if there is any relationship with age and type of infertility among women who had HSG in a tertiary hospital in Port Harcourt. In this environment, there is a paucity of information on this hence the need for this present study and also to have a working document for the hospital.

MATERIALS AND METHODS:

The study was conducted at the Radiology and the Obstetrics and Gynaecology Departments of RSUTH, South-South Nigeria. Patients who presented for hysterosalpingography for infertility between 1st of January, 2020 to 1st of January 2022 were included in the study. Those with incomplete request forms were excluded from the study. The Biodata, indicated type of infertility under evaluation and the HSG findings were obtained and documented.

Ethical approval is not required in this kind of study.

Sample size calculation:

A sample size of 226 was used for this study while employing the formula for cross-sectional design. [7] Eduwem et al [8] recorded a prevalence rate of HSG abnormality as 85.8%. The sample size was gotten using desired precision of 0.05.

Data analysis:

Data obtained from the study proforma was entered into Microsoft Excel, and then exported to IBM Statistical Package for Social Sciences (SPSS) version 21 for statistical analysis. Tables and charts were used to display data as appropriate. Descriptive statistics employed means, standard deviation, median and ranges for numerical data, and absolute frequencies and percentages for categorical data.

Comparisons of number of abnormalities by infertility were performed using independent t-test to determine significant differences. A multiple linear regression analysis of age, infertility, and number of abnormalities in the study was also done. Statistical significance when *P*-value is less than 0.05.

RESULTS

Mean age of study population \pm S.D = 35.07 \pm 5.45 years, median age = 35.00 years. The range is 22-48 years.

Table: 1 Age distribution of women in the study

Variable	Frequency	Percentage
Age category		
20 – 29 years	40	17.7
30 – 39 years	130	57.5
40 – 49 years	56	24.8

Table 1 is showing the age category. Age 30-39 has the highest frequency/percentage (130, 57.5%), followed by 40-49 years and 20-29 years been the least.

Table2: Comparison of age category against number of abnormalities on HSG among the study population

Age category	Number of abnormalities				Total
	Normal	One	Two	Greater than two	
20 – 29 years	17 (42.5)	15 (37.5)	7 (17.5)	1 (2.5)	40 (100.0)
30 – 39 years	36 (27.7)	62 (47.7)	31 (23.8)	1 (0.8)	130 (100.0)
40 – 49 years	11 (19.6)	29 (51.8)	14 (25.0)	2 (3.6)	56 (100.0)
Total	64 (28.3)	106 (46.9)	52 (23.0)	4 (1.8)	226 (100.0)

Fisher's Exact = 8.199; p-value = 0.188

Table 2 revealed that the occurrence of at least one abnormality is seen more in 30-39 years category (n=62 or 47.7%) followed by 40-49years category and the 20-29years category. Similar pattern of occurrence is also seen with minimum of two abnormalities. However greater than two abnormalities is more common in 40-49 years category (n=2 or 3.6%) while 20-29 and 30-39 had equal numbers.

Table3: Comparison of age category against type of infertility among the study population

Age category	Primary infertility n (%)	Secondary infertility n (%)	Total n (%)
20 – 29 years	1 (2.5)	39 (97.5)	40 (100.0)
30 – 39 years	6 (4.6)	124 (95.4)	130 (100.0)
40 – 49 years	3 (5.4)	53 (94.6)	56 (100.0)
Total	10 (4.4)	216 (95.6)	226 (100.0)

Fisher's Exact= 0.432; p-value = 0.908

Table 3 revealed that secondary infertility was more common than primary infertility at all age category and it is not statistically significant with p -value=0.908. Age 30-39years had the highest occurrence rate for both primary (6, 4.6%) and secondary infertility (124, 95.4%) respectively.




Figure: 1 Scatter plot showing relationship between age (years) and number of abnormalities in the study.

Figure 1 showed that the relationship between age (years) and number of abnormalities are significant with p -value =0.037.

Table: 4 Classification of HSG among the study population

HSG Classification	Frequency	Percentage
Normal HSG	64	28.3
Abnormal HSG	162	71.7
Total	226	100.0

Table 4 is simply showing the numbers of normal HSG findings and abnormal HSG findings in the study population.

Table: 5 Number of abnormalities among those with abnormal HSG in the study population

Number of abnormality	Frequency	Percentage
None	64	28.3
One abnormality	106	46.9
Two abnormalities	52	23.0
Three abnormalities	4	1.8
Total	226	100.0

Table 5 is showing the number of abnormalities among the abnormal HSG findings. One abnormality is more prevalent than two or three abnormalities.

Table: 6 Comparison of number of abnormalities and age by type of infertility

	Primary	Secondary	T	p-value
	Mean \pm SD	Mean \pm SD		
Number of abnormalities	1.50 \pm 0.71	0.96 \pm 0.76	2.206	0.028*
Age	34.80 \pm 4.47	35.08 \pm 5.50	-0.160	0.873

*Statistically significant

Table 6 is showing the comparison number of abnormalities and age by type of infertility, which showed that there is an association between number of abnormalities and infertility type (P -value= 0.028). There is no association with age.

Table: 7 Multiple linear regression analysis of age, number of abnormalities and infertility in the study

Variables	Type of infertility			p-value
	B	95% Confidence Interval for β		
		Lower limit	Upper limit	
Age in years	0.001	-0.004	0.006	0.638
Number of abnormalities	-0.040	-0.076	-0.005	0.026*

*Statistical significance $p < 0.05$

Table 7 is displaying a multiple linear regression analysis of age, number of abnormalities and infertility in the study. Number of abnormalities and type of infertility showed significant association with P -value of 0.026 and age with type of infertility was not significant (p -value=0.638).



Figure 2: A spot film showing a capacious contrast opacified uterine cavity with irregular outline and depression at the fundal region. The fallopian tubes are not demonstrated. Features are in keeping with uterine myoma with bilateral tubal occlusion.



Figure 3: showed an elongated stretched contrast opacified uterine cavity angulated at the fundus and a well defined roundish calcified myoma on its right side. Both fallopian tubes are visualized and ends abruptly at the cornua.



Figure 4: showed a contrast opacified uterine cavity with irregular outline, both fallopian tubes were not demonstrated. Intravasation was noted.



Fig. 5a



Fig. 5b delayed image

Figure 5: showed contrast opacified uterine cavity with dilated and tortuous right fallopian tube, which is persistent in the delay image film.

DISCUSSION

Infertility is a big deal in Nigeria, female infertility is considered as a disaster and is often the rationale for separation and divorce or polygamy as well as denial of rights in family functions.[5]

In our study the mean age was 35.07 ± 5.45 years and the age range was 22-48 years. Age 30-39 years had the highest frequency of people presenting for HSG due to infertility. This is probably due to an increase in girl child education and more women in career pursuit.[9] This supports previous studies done.[10-15] The minimum age in our study was 22 years compared to studies done in Northern Nigeria with a minimum of 17 years. This could be linked to socio-cultural background in the North where early marriage is encouraged and common.[11,16]

In this study, secondary infertility (n=216, 95.6%) is commoner than primary infertility (n=10, 4.4%). Similar findings were made by other investigators.[2,10,13,16-23] The possible explanation for this may be as a result of pelvic inflammatory disease, sepsis, sexually transmitted diseases and unsafe abortion alongside post surgical procedures conducted in septic environs.[18,24] In contrast to our findings, others found out that primary infertility is a more frequent indication for HSG than secondary infertility.[12,25-26] This disparity in findings could be due to geographical location and methodology.

In this current study, 28.3% of the women had normal HSG findings. This observations corroborates with other studies[10,12-13,16,21,26] but disagrees with an earlier study done in Port Harcourt(44.2%) by Nwankwo et al[27] and in Northern Nigeria studies by Saidu et al-59% and Lawan et al-55% respectively.[19,28] A possible reason for this is early marriage, religious beliefs and/or premarital sex.

Of the two-hundred and twenty-six patients, 162 (71.7%) had abnormal HSG findings. This is similar to 73.1% and 75.2% reported by Udobi et al and Eze CU et al respectively.[21,26] Other studies [8,29-30] done within Nigeria also reported high percentage of abnormalities in infertility

work up. More often infertility cases are linked with problems resulting in incomplete or complete distortion of the uterine cavity and the occlusion of the fallopian tubes. HSG does clearly outline this reproductive pathway and thus an important tool in evaluation of infertility.[8] From this current study that an individual can have just one or more than one pathology is in tandem with earlier studies.[16-17,31-33]

Bello et al[17] also documented that pathology is associated with type of infertility and it is statistically significant with P -value <0.05 Similarly, Itanyi et al[31] reported that an association exist between abnormalities and type of infertility (P -value= <0.0001). Lash et al[34], established an association between secondary infertility and abnormality seen on HSG. These reports agree with the findings of this study using either multiple linear regression analysis (P -value= 0.026) or two by two table P -value= 0.028). Conversely, Aduayi et al[14] and Udobi et al[26] documented that there was no association between abnormalities and type of infertility.

With regards to age in this index study there is an association between abnormalities and age using scatter plot with P -value of 0.037 . This corroborates with Aduayi et al[14] (P -value= 0.002) and Akagbue et al[37, 38] (P -value= 0.007) On the other hand, some studies [26,31] disagree with this submission. Interestingly in our study there is no correlation between age and type of infertility and this agrees with Aduayi et al and Udobi et al respectively.[14,26]

CONCLUSION:

Hysterosalpingography (HSG) is a valuable tool in investigating women with infertility and most frequently encountered type in this study was secondary infertility and the commonest age category with abnormalities is 30-39 years. Abnormal HSG accounted for 71.7%. There was

significant association with age, type of infertility and number of abnormalities on HSG therefore women are encouraged to marry early once they reach adulthood..

LIMITATION OF THE STUDY: This is a retrospective study and the sample size is not representing the total population. Some patients still patronize unorthodox practitioners.

ETHICAL APPROVAL: Is not applicable in this study.

REFERENCES

1. Williams C.T, Giannopoulos and E.A Sheriff, 2003. Investigation of infertility with emphasis on laboratory testing and with reference to radiological imaging. J Clin Pathol 56:261-2267.
2. Ubeda B, M Paraiza, E Alert and RA Abuin 2010. Hysterosalpingography: Spectrum of normal variants and non-pathological findings. AJR, 177(1):131-135.
3. Steinkeler JA, AW Courtney, E Lazarus and MM Hillstrom, 2009. Female infertility: a systematic approach to radiologic imaging and diagnosis. Radiograph, 29:1353-13770.
4. Yu SL and Yap, 2003. Investigating the infertile couple. Ann Acad Med Singapore, 32:611-614.
5. Practice Committee of American Society for Reproductive Medicine, 2012. Diagnostic evaluation of infertile female: a committee opinion. Fertil Steril, 98:302.

6. Nigeria Fertility rate, 1950-2021-Knoema.com.Knoema.
Available:<https://knoema.com//atlas/Nigeria/topics/Demographics/Fertility/Fertility-rate>
rate
Accessed September 20, 2022.
7. Naing L, Winn TB, Rusli BN. Practical issues in calculating the sample size for prevalence studies. *Archives of orofacial Sciences*. 2006;1:9-14.
8. Eduwem, D. U., Akintomide, A. O., Bassey, D. E., & Ekott, M. I. (2016). Hysterosalpingographic patterns and relevance in the management of infertility in a Nigerian tertiary health Institution. *Asian Journal of Medical Sciences*, 7(5), 70–74.
<https://doi.org/10.3126/ajms.v7i5.15169>
9. Balen VF. Interpreting infertility, social science research on childlessness in a global perspective . Amsterdam. *Afr J Reprod Health* 2000;4:120-2.
10. Kiridi EK, Oriji PC, Ugwoegbu JU, Abasi IJ. Hysterosalpingography Findings among Women Presenting for Infertility Evaluation in Bayelsa State, South-South Nigeria. *J Adv Med Res*. 2022;34(5):7-17.DOI: 10.9734/JAMMR/2022/v34i531292
11. Danfulani M, Mohammed MS, Ahmed SS, Haruna YG. Hysterosalpingographic findings in women with infertility in Sokoto North-Western Nigeria. *Afr J Med Health Sci* 2014;13:19-23. DOI: 10.4103/2384-5589.139438
12. Okafor CO, Okafor CI, Okpala OC, Umeh E. The pattern of hysterosalpingographic findings in women being investigated for infertility in Nnewi, Nigeria. *Niger J Clin Pract* 2010;13(3):264-7.

13. Onwuchekwa CR, Oriji VK. Hysterosalpingographic(HSG) pattern of infertility in women of reproductive age. *J Hum Reprod Sci* 2017;10:178-84. DOI:10.4103/jhrs.JHRS_121_16
14. Aduayi OS, Akanbi GO, Akintayo AA, Aduayi VA. Hysterosalpingography findings among women presenting for gynaecological imaging in Ado-Ekiti, South Western Nigeria. *Int J Rreprod Contracept Obstet Gynecol* 2016;5:1906-11. DOI: 10.18203/2320-1770ijrcog20161688.
15. Eleje GU, Okaforcha EI, Umeononihu OS, Udegbonam OI, Etoniru IS, Okwuosa AO. Hysterosalpingographic Findings Among Infertile Women: Review at a Tertiary Health Care Institution in Nnewi, South-East Nigeria. *Afrimed J* 2012;3(2):20-23.
16. Bukar M, Mustapha Z, Takai UI Tahir A. Hysterosalpingographic findings in infertile women: A seven year review. *Niger J Clin Pract* 2011;14(2):168-70. DOI: 10.4103/1119-3077.84008.
17. Bello TO. Tubal abnormalities on hysterosalpingography in primary and secondary infertility. *West Afr J Med* 2006;25:130-3.
18. Kiguli-Malwade E, Byanyima RK. Structural findings at hysterosalpingography in patients with infertility at two private clinics in Kampala, Uganda. *Afr Health Sci* 1994;4(3):178-81.
19. Saidu HB, Mohammed MT, Abdullahi NA, El-Habeeb MA, Salihu FA, Ibrahim H, et al. evaluation of radiological pattern of HSG in female patients with infertility in Katsina Northwestern Nigeria. *Niger J Basic Clin Sci* 2022;19:15-19.

20. Tukur J, Zahradeen SU, Takai IU, Suwaid MA, Ibrahim UM, . evaluation of hysterosalpingographic findings of patients presenting with infertility in Kano, Northern Nigeria. Niger J Clin Res 2021;10:14-8.
21. Eze CU, Ohagwu CC, Abonyi LC, Njoku J, Iruhe N, Igbinedion FO. A spectrum of .hysterosalpingographic findings in infertile women in Benin City Nigeria. J Reprod infertile 2013;4:13-8. DOI: 10.5829/idosi.jri.2013.4.2.7692.
22. Botwe BO, Bamfo-Quaicoe K, Hunu E, Anim-Sampong S. Hysterosalpingographic findings among Ghanaian women undergoing infertility work-up: A study at the Korle-Bu Teaching hospital. Fertile Res Pract2015;1(9):1-6.
23. Larsen U 200. Primary and secondary infertility in Sub-Saharan Africa. Int J Epidemiol;29:285-291.
24. Bhartla N Infection as they infect individual organs. In; Jellcoates principles of gynaecology. 7th ed. Michigan, USA; Butterworths;2015,p:355-74.
25. Daniel A, Yewebdar N. Evaluation of the fallopian tubes in infertile women by hysterosalpingography in Tikur Anbessa Hospital, Addis Ababa, Ethiopia. Int J Nurs Midwifery 2011;3(11):178-81.
26. Udobi SI, Aronu ME. Hysterosalpingographic findings in women with infertility in Awka, Anambra State, South-East Nigeria. Niger J Surg Sci 2017;27:47-50. DOI: 10.4103/njss.njss_3_18.
27. Nwankwo NC, Akani CI. Pattern of hysterosalpingographic findings in infertility in Port Harcourt. West Afr J Radiol2005;12:15-19.

28. Lawan RO, Ibinaye PO, Onwuhafia P, Hamidu A. Evaluation of pattern of tubo-peritoneal abnormalities potentially responsible for infertility in Zaira , Nigeria.: Hysterosalpingohic assessment sub-Saharan Afr J Med 2015;2:110-6.
29. Oguntoyinbo AE, Aboyeji AP. Hysterosalpingographic (HSG) findings of infertile patients in Ilorin. Trop J ObstetGynaecol 2010;229(1):118-125.
30. Ibekwe PC, Udesi AM, Imo AO. Hysterosalpingographic findings in patients with infertility inSouthEasternNigeria. NigerJMed 2010;19(2):165-167.
31. Itanyi UD, Oluseyi HO. Spectrum of hysterosalpingographic findings among women presenting with infertility in Abuja, Nigeria's capital. Int J Reprod Contracept Obstet Gynecol 2017;6(4):1583-1589. DOI:10.18203/2320-1770.ijrcog20171433.
32. Walton SN, Mati JK. An evaluation of secondary infertility in Kenya. East Afr Med J 1976;53:310-4.
33. Dhaliwal LK, Gupta KR, Aggarwal N Is hysterosalpingography an important tool in modern gynaecological practice? Int J Fertil Women Med 1999;44:212-5.
34. Lash MM, Yaghamee W, Strohsnitter, Lalwani S. Association between secondary infertility and fallopian tube obstruction on hysterosalpingography.2008 J Reprod Med 2008;5(9): 677-680..
35. Akagbue VN, Nonye-Enyidah EI, Aderibigbe GJ Hysterosalpingographic Indications and Findings among Women Presenting to Radiology Department of Rivers State University Teaching Hospital.J Adv Med Pharma Sci 2022;24(9):14-24. DOI:10.9734/jamps/2022/v24i9576.
36. Akagbue V.N, Nonye-Enyidah EI, Aderibigbe GJ A Comparative Study on Hysterosalpingography Findings between Young-aged and Middle-aged Women at a

Tertiary Health Facility in Niger-Delta Region. Asian J Pregnan Childbirth2022;5(4):81-90. DOI:5(4):81-90,2022;Article no.AJPCB.94104

UNDER PEER REVIEW