

Original Research Article
**CHARACTERIZATION OF EXOTIC ACCESSIONS OF
SESAME (*Sesamum indicum*) THROUGH PRINCIPLE
COMPONENT ANALYSIS**

ABSTRACT

The present research conducted over 96 exotic sesame accession with five checks, to identify the minimum number of components, which can explain maximum variability out of the total variability using Principal Component Analysis (PCA); The investigation was performed under Project Co-ordinating Unit (Sesame and Niger) Research Farm, JNKVV, Jabalpur (M.P.) during kharif 2018 using Augmented block design. Among the studied traits, In this study, Component 1 had the contribution from the traits viz., number of primary branches per plant, number of capsules per plant, number of seeds per capsule, oil content and seed yield/plant. which accounted 30.71% to the total variability. Days to flower initiation and days to 50% flowering have contributed 17.11% to the total variability in component 2. The remaining variability of 11.26%, 9.94%, 7.48% and 6.73% was consolidated in PC3, PC4, PC5 and PC6 by various traits like number of secondary branches/plant, capsule length, days to maturity, thousand seed weight and plant height . The cumulative variance of 83.23% of total variation among 12 characters was explained by the first six axes. On the basis of PC scores PC1, PC3 and PC5 accounting mainly to yield and quality traits containing accesions viz., EC-334998, ES-38, EC-346426, EC-334958, EC-340538, RT-351 and GT-10 may be further utilized in breeding programme.

Keywords: [Principal component, Exotic sesame accession, Augmented block design, PC scores] (Arial, inclined, 10 font, justified)

(Note: 1. Case Reports should follow the structure of Abstract, Introduction, Presentation of Case, Discussion, Conclusion, Acknowledgements, Competing Interests, Authors' Contributions, Consent (where applicable), Ethical approval (where applicable), and References plus figures and/or tables. Abstract (not more than 250 words) of the Case reports should have the following sections: Aims, Presentation of Case, Discussion and Conclusion. Only Case Reports have word limits: Papers should not exceed 2000 words, 20 references or 5 figures. Other Type of papers have no word limits.

2. Review papers may have different headings of the sections and are exempted from following these suggestions.

3. Research Papers and Short Notes should follow the structure of Abstract, Introduction, Methodology, Results and Discussion, Conclusion, Acknowledgements, Competing Interests, Authors' Contributions, Consent (where applicable), Ethical approval (where applicable), and References plus figures and/or tables.)

1. INTRODUCTION

Sesame (*Sesamum indicum*L.) unremarkably known as til or benniseed from the family Pedaliaceae is a broadleaf, one of the most eminent annual oilseed crop in tropical and subtropical countries including India with chromosome number $2n = 26$. *Sesamum indicum*, the cultivated type, originated in India is tolerant to dry spell conditions in high heat, with residual moisture in soil. It can grow well in many ecological regions of tropical and subtropical climate,

and also grows in almost all kind of soils. Besides being drought tolerant and having acceptance of a range of different soils, sesame also stimulates a fauna of beneficial soil microbes and reduces the nematode populations.

It is an antediluvian oilseed, first noticed and recorded as a crop in Babylon and Assyria over 4,000 of years ago but naturalized well over 3000 years ago. Among *kharif* oilseed crops, sesame occupies an important place next to groundnut. It is one of the crops under cultivation from ancient times (Joshi, 1961 and Bistet *et al.*, 1998). Its production in India is boosting, as more than 30 % of world production is bestowed by India and India ranks top in total production (Pandey *et al.*, 2015). India is a major producer of sesame in the world and inhabits about 36% of the total acreage and contributes about 25% of the total output.

In India, sesame is cultivated in 1544.9 thousand hectare with production of 715.7 thousand tonnes and productivity of 463.26 kg/ha (Anonymous, 2017- 18). Madhya Pradesh contributes 18.8% and 24.03% share of country's area (424 thousand ha) and production (180 thousand tonne), respectively and ranked third in area and second in production. In India, major sesame producing states are Rajasthan, U.P., M.P., A.P., Orissa, Gujarat, Tamil Nadu and Karnataka.

Sesame, which is entitled as “the queen of oilseed crops”, is best known for caliber edible oil. Its seed contains 48-55% oil, protein (20-28%), carbohydrate (23 %), minerals, healthy polyunsaturated fatty acid and also few chemicals like lignans, pinosresinol and lariciresinol. Due to presence of potent antioxidants, sesame seeds are called as “the symbol of immortality” (Ayana. 2015). Sesame contributes in multifarious ways to human beings like culinary preparation, edible purposes, in industries like paints, cosmetics, insecticides, soap making, pharmaceuticals, etc. It is also used as a cattle supplement especially for milch cattle and as manure due to presence of 6-6.2% N, 2-2.2% P and 1-1.2% K in its cake (Singh *et al.*, 2009). Despite being largely self-sufficient in production, sesame productivity is declining mainly due to unavailability of high yielding varieties and it is a matter of great concern for scientists as well as the farmer's community. Though *Sesamum indicum* genotypes in Indian landmass are reported to represent a good diversity however, still no systematic efforts are created to characterize and document the native and exotic collections to reveal the genetic diversity (Kim *et al.*, 2002).

Principal component analysis measures the importance and contribution of each component to the total variance while each coefficient of proper vectors indicates the degree of contribution of every original variable with which each principal component is associated (Sanni *et al.*, 2012). Its presence of genetic diversity could be a fascinating demand for any breeding program. Both diversity analysis and PCA are useful for parent's choice in hybridization to develop high yield potential cultivars and to meet the diversified goals of plant breeding. A thorough knowledge on Principal component analysis will provide basis for selecting systematic breeding strategy to improve yield potential of the genotypes.

2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

The present experiment was conducted under Project Coordinating Unit (Sesame and Niger) Research Farm, JNKVV, Jabalpur (M.P.) during kharif 2018-19. The experimental material consisted of 96 *sesamum* germplasm along with 5 checks with distance between rows 0.40 m and plant to plant 0.12 m. The crop was raised under recommended package of practices along with prophylactic protection measures. PCA analysis was done using method given by Massay, 1965 and Jolliffe, 1986.

2.1 Subheading Subheading- second level heading.

2.1.1 Sub-subheading(ARIAL, BOLD, 10 FONT, LEFT ALIGNED, underlined)- third level heading.

2.1.1.1 Sub-sub-subheading- fourth level heading.]

3. RESULTS AND DISCUSSION

In PCA analysis, principal components (PCs) exhibited more than 0.5 Eigen value and showed the cumulative variance of 83.23% among the traits studied as shown in scree plot fig 1. Out of twelve, only six principal components were given due importance for further explanation. The PC1 had the highest variability (30.711 %) followed by PC2 (17.11%), PC3 (11.27%), PC4 (9.94%), PC5 (7.49%) and PC6 (6.73%) for all the traits under study as represented in Table no. 1. Almost similar findings were found by Baraki et al. (2015) and Shim et al. (2016) with total variance of 88.49% and 82.3%.

Table 1- Eigen values, % variance and cumulative eigen values of different traits in sesame genotypes

Characters	Principal component (PC)	Eigen value	Variability (%)	Cumulative %
DFI	PC1	3.685	30.711	30.711
DFE	PC2	2.054	17.114	47.824
NPB	PC3	1.352	11.267	59.091
NSB	PC4	1.193	9.943	69.035
PH	PC5	0.899	7.489	76.523
NCPP	PC6	0.808	6.733	83.257
CL	PC7	0.797	6.642	89.899
NSPC	PC8	0.563	4.693	94.591
DM	PC9	0.456	3.802	98.394
TSW	PC10	0.089	0.743	99.136
OC	PC11	0.066	0.551	99.687
SYPP	PC12	0.038	0.313	100.000

*Moisture content on oven dry weight basis

Results displayed in Table no. 2 and 3 revealed that the first principal component (PC1) was mostly related with yield traits such as Number of primary branches per plant, Number of capsules per plant, Number of seeds per capsule, Oil content (%) and Seed yield/plant. The second principal component (PC2) was dominated by traits number of Days to 50% flowering and Days to flower initiation and PC3 with Number of secondary branches per plant and Capsule length. On the basis of PCA, most of the important yield attributing and quality traits were present in PC1, PC2 and PC3. Similar results were reported by Saha et al. (2012) for days to 50% flowering, days to maturity and number of capsules per plant, Ismaila and Usman (2014) for number of capsules per plant and number of branches per plant, Shim et al. (2016) for capsule length and number of capsules per plant, days to maturity and plant height; Baraki and Berhe (2019) for seed yield, number of primary and secondary branches, number of capsule per plant and number of seeds per capsule. In contradiction Singh et al. (2018) reported that PC1 was mostly related with phenological traits viz., days to flower initiation, days to 50% flowering and days to maturity.

Table 2 Rotated component matrix for different traits of sesame genotypes

Traits	Principal Components					
	PC1	PC2	PC3	PC4	PC5	PC6

DFI	0.321	0.919	-0.039	-0.166	0.080	0.163
DFE	0.281	0.910	-0.021	-0.175	0.099	0.168
NPB	0.557	-0.172	0.214	-0.220	0.319	-0.126
NSB	0.459	-0.053	0.658	0.048	-0.297	0.005
PH	0.305	-0.378	0.289	-0.283	0.059	0.620
NCPP	0.943	-0.119	-0.048	0.046	0.023	-0.052
CL	-0.076	0.353	0.601	0.464	-0.353	-0.021
NSPC	0.899	-0.101	-0.135	0.013	0.030	-0.012
DM	0.063	-0.104	-0.381	0.626	-0.046	0.535
TSW	-0.139	0.055	0.371	0.496	0.740	-0.047
OC	0.542	0.166	-0.351	0.374	-0.113	-0.238
SYP	0.948	-0.123	-0.016	0.093	-0.018	-0.069

Table 3: Interpretation of rotated component matrix for the traits having maximum values in each PCs

	PC1	PC2	PC3	PC4	PC5	PC6
	Number of primary branches per plant	Days to flower initiation	Number of secondary branches per plant	Days to maturity	Thousand seed weight	Plant height
	Number of capsules per plant	Days to 50% flowering	Capsule length	-	-	-
Traits	Number of seeds per capsule		-	-	-	-
	Oil content (%)	-	-	-	-	-
	Seed yield/plant	-	-	-	-	-

Germplasm lines showing maximum positive PC scores and common in PC1, PC3 and PC5 for number of primary branches per plant, number of capsules per plant, number of seeds per capsule, oil content, seed yield per plant, number of secondary branches per plant, capsule length and thousand seed weight are EC-118571; and genotypes common in PC 1 and PC 3 are EC-334998, EC-335015, EC-346424, EC-346599, EC-346633 and EC-346681; genotypes common in PC1 and PC5 are RT-351, ES-38, EC-34625, EC-334958 and EC-340538. Genotypes showing maximum positive PC scores in PC1 for number of primary branches per plant, number of capsules per plant, number of seeds per capsule, oil content, seed yield per plant and negative scores in PC2 for traits days to flower initiation and days to 50% flowering are RT-351, EC-346424, EC-346400, EC-334958 and EC-346599. The result presented in table 4 and 5. Thus, selection of these lines can help in further development of new high yielding quality varieties.

*Test drugs: significant from normal control, * P < 0.05; ** P < 0.001*

Mean ± S.E.M = Mean values ± Standard error of means of six experiments

3.1 Subheading Subheading - second level heading.

3.1.1 Sub-subheading third level heading.

3.1.1.1 Sub-sub-subheading - fourth level heading.]

4. CONCLUSION

Principal component analysis measures the importance and contribution of each component to the total variance while each coefficient of proper vectors indicates the degree of contribution of every original variable with which each principal component is associated (Sanni et al., 2012). These results can now be used by breeders to develop high yielding sesame varieties and develop new breeding protocols for sesame improvement.

UNDER PEER REVIEW

CONSENT (WHEREEVER APPLICABLE)

No manuscripts will be peer-reviewed if a statement of patient consent is not presented during submission (wherever applicable).

This section is compulsory for medical journals. Other journals may require this section if found suitable. It should provide a statement to confirm that the patient has given their informed consent for the case report to be published. Journal editorial office may ask the copies of the consent documentation at any time.

Authors may use a form from their own institution or SDI Patient Consent Form 1.0. It is preferable that authors should send this form along with the submission. But if already not sent during submission, we may request to see a copy at any stages of pre and post publication.

If the person described in the case report has died, then consent for publication must be collected from their next of kin. If the individual described in the case report is a minor, or unable to provide consent, then consent must be sought from their parents or legal guardians.

Authors may use the following wordings for this section: "All authors declare that 'written informed consent was obtained from the patient (or other approved parties) for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editorial office/Chief Editor/Editorial Board members of this journal.'"

ETHICAL APPROVAL (WHEREEVER APPLICABLE)

This section is compulsory for medical journals. Other journals may require this section if found suitable. If human subjects are involved, informed consent, protection of privacy, and other human rights are further criteria against which the manuscript will be judged. It should provide a statement to confirm that the authors have obtained all necessary ethical approval from suitable Institutional or State or National or International Committee. This confirms either that this study is not against the public interest, or that the release of information is allowed by legislation.

All manuscripts which deal with animal subjects must be approved by an Institutional Review Board (IRB), Ethical Committee, or an Animal Utilization Study Committee. , and this statement, and approval number, must accompany the submission. If required, author should be ready to submit a scanned copy of the IRB or Ethical Committee Approval at any stage of publication (Pre or post publication stage). The manuscript should contain information about any post-operative care and pain management for the animals.

For manuscripts involving animal experiments, Authors may use the following wordings for this section"All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee"

All manuscripts which deal with the study of human subjects must be accompanied by Institutional Review Board (IRB) or Ethical Committee Approval, or the national or regional equivalent. The name of the Board or Committee giving approval and the study number assigned must accompany the submission. If required, author should be ready to submit a

scanned copy of the IRB or Ethical Committee Approval at any stage of publication (Pre of post publication stage).

For manuscripts involving human experiments, Authors may use the following wordings for this section: "All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki."

REFERENCES

Anonymous, 2017-18. 4th Advance Estimates, Agriculture Statistics Division, Directorate of Economics and Statistics, New Delhi.

Ayana NG. 2015. Status of production and marketing of ethiopian sesame seeds(*Sesamum indicum* L.): a review. *Agricultural and Biological Sciences Journal*1(5): 2017-223

Baraki F and Berhe M. 2019. Evaluating performance of sesame (*Sesamum indicum* L.) genotypes in different growing seasons in northern Ethiopia. *International Journal of Agronomy* 7:1-7.

Baraki F, Tsehaye Y and Abay F. 2015. Grain yield based cluster analysis and correlation of agronomic traits of sesame (*Sesamum indicum* L.) genotypes in Ethiopia. *Journal of Natural Sciences Research* 5(9): 11-17.

Bist IS, Mahajan RK, Loknathan TR and Agrawal RC. (1998). Diversity in Indian sesame collection and stratification of germplasm accessions in different diversity groups. *Genetic Resources and crop Evolution*,45: 325-335

Ismaila A and Usman A. 2014. Genetic variability for yield and yield components in sesame (*Sesamum indicum* L.). *International Journal of Science and Research* 3(9): 63-66.

Jolliffe I.T. (1986). *Principal Component Analysis*. Springer, New York.

Joshi AB. 1961. *Sesame - A Monograph*. Indian Central Oilseeds Committee, Hyderabad, India, p. 267

Kim DH, Zur G, Danin-Poleg Y, Lee S, Shim K, Kang C and Kashi Y. 2002. Genetic relationships of sesame germplasm collection as revealed by inter-simples sequence repeats. *Plant Breeding* 121:259–262.

Massay W.F. (1965). Principal components regression in exploratory statistical research. *Journal of the American Statistics Association*, 60: 234-246.

Pandey SK, Das A, Rai P and Dasgupta T. 2015. Morphological and genetic diversity assessment of sesame (*Sesamum indicum* L.) accessions differing in origin. *Physiological Molecular Biological Plants*. 21(4):519-29.

Saha, Sruba, Begumb T and Dasgupta T. 2012. Analysis of genotypic diversity in sesame based on morphological and agronomic traits. Conference on International Research on Food Security, Natural Resource Management and Rural Development organised by Georg-August Universität Göttingen and University of Kassel-Witzenhausen University of Calcutta,

Department of Genetics and Plant Breeding, 35 Ballygaunge Circular Road, 700019 Calcutta, India.

Shim KB, Shin SH, Shon JY, Shin GK, Yang WH, Heu SG. 2016. Classification of a collection of sesame germplasm using multivariate analysis. Journal Crop Science Biotechnology 19(2):151-155

Singh A, Bisen R and Tiwari A. 2018. Assessing inter: relationship of sesame genotypes and their traits using cluster analysis and principal component analysis. International Journal of Chemical Studies 6(1): 2151-2153.

Singh C, Singh P and Singh R. 2009. Modern techniques of raising field crops. Oxford&ibh publishing company. New Delhi. p311.

APPENDIX

FIG. 1: SCREE PLOT OF PRINCIPAL COMPONENT ANALYSIS OF SEASAMEGERMPLASM BETWEEN EIGEN VALUE AND PRINCIPAL COMPONENTS.




Table 4: PC scores of nigergermplasm having positive values >0.5 in each PCs.

S. No.	Genotypes	PC1 score	PC2 score	PC3 score	PC4 score	PC5 score	PC6 Score
1	TKG 22	1.235	2.668	-1.256	0.867	-0.184	0.36
2	GT 10	1.927	3.496	-2.099	1.033	0.079	-0.46
3	JTS 8	1.346	1.778	-1.235	0.787	0.172	-0.346
4	TKG 21	2.399	2.082	-0.347	0.243	-0.274	-1.743
5	RT-351	2.362	-1.537	-2.377	1.636	0.875	2.1
6	EC-38	1.269	0.862	-1.26	-0.906	0.579	-2.082
7	EC-177-A	-0.196	0.629	0.492	-0.112	-0.151	-0.047
8	EC-177-1	-0.885	-0.169	-0.107	-2.432	1.64	-0.535
9	EC-43708	-1.688	1.867	0.239	-0.264	-0.563	-0.274
10	EC-43760	-1.691	0.036	-0.627	-1.675	0.843	0.281
11	EC-89105	0.224	-0.059	-0.175	1.033	-0.525	-0.618
12	EC-89671	1.516	0.404	0.032	-0.928	0.146	0.063
13	EC-109693-A	-1.665	1.042	1.099	-0.091	-0.787	0.304
14	EC-118569	-2.134	1.077	0.753	-1.919	0.221	-0.611
15	EC-118571-B	-0.661	0.966	2.115	-0.773	-0.485	0.036
16	EC-118571	3.42	-1.146	0.927	-1.787	0.912	-0.995
17	EC-132827-1	0.278	0.39	-1.216	0.392	0.228	0.675
18	EC-34625	1.642	0.889	-0.196	-2.031	1.187	-0.935
19	EC-346347	-1.99	1.724	0.075	-0.646	0.048	0.955
20	EC-346377	0.906	0.653	-0.286	-1.49	0.445	0.474
21	EC346478	-2.039	0.918	-0.303	-0.177	-0.128	1.389
22	EC-346480	-0.41	2.933	2.338	-1.486	0.205	0.453
23	EC-346514	1.024	-1.096	0.049	-0.414	-0.192	0.989
24	EC-356526	0.776	-0.421	1.337	-0.919	0.005	0.465
25	EC-178365	2.054	0.412	0.757	-1.084	-0.082	-0.353
26	EC-133850-S	-2.116	-0.87	1.401	-0.868	-0.611	-0.211
27	EC-132828	0.715	2.912	1.36	-0.584	-0.196	0.822
28	EC-178374	4.048	2.432	0.3	-0.981	0.402	0.991
29	EC-334962	3.28	-0.922	-0.334	-0.039	0.595	-0.78
30	EC-334974	-2.548	-0.454	-0.516	-1.264	0.35	-1.337
31	EC-334982	-1.641	-2.529	-1.05	0.236	0.834	-0.039
32	EC-334993	-2.716	0.893	1.131	-0.805	0.129	0.148
33	EC-335005	1.51	-0.227	-1.025	-0.636	0.541	-0.258
34	EC-334998	3.607	1.274	2.605	-0.929	0.134	1.045
35	EC-335015	1.978	-0.082	1.027	0.095	-0.137	0.957
36	EC-334999	1.562	0.437	-0.4	-0.379	0.078	0.615
37	EC-335017	-1.758	0.914	0.408	1.006	-1.014	1.575
38	EC-342782	-1.794	0.086	1.073	0.95	-0.994	0.597
39	EC-346165	-3.428	0.868	1.191	-0.471	-0.111	0.469
40	EC-346192	-2.457	-1.59	-0.398	-1.555	0.378	-0.08
41	EC-346224	-1.791	-0.405	-0.436	0.196	-0.334	-0.396
42	EC-346300	-2.309	-0.836	0.564	-1.523	0.422	0.252
43	EC-346358	-2.152	-0.533	0.828	-0.217	-0.622	0.347
44	EC-346374	-2.442	-1.247	-0.674	-0.07	-0.273	-0.018
45	EC-346391	-1.834	-3.246	1.131	-0.537	-0.334	2.415
46	EC-346395	-2.59	0.287	-0.052	-1.564	-0.073	0.49
47	EC-346424	3.535	-3.247	3.619	-0.514	-0.459	0.273
48	EC-346448	-2.035	-2.813	-0.548	-1.442	0.751	0.03

S. No.	Genotypes	PC1 score	PC2 score	PC3 score	PC4 score	PC5 score	PC6 Score
49	EC-346453	2.054	0.412	0.757	-1.084	-0.082	-0.353
50	EC-346471	-2.944	0.632	3.781	5.196	7.189	-0.406
51	EC-346599	2.2	-1.22	0.846	1.135	-0.897	0.375
52	EC-346620	-1.461	0.037	0.494	-0.09	0.126	0.737
53	EC-346633	1.164	0.625	1.145	0.586	-0.659	0.973
54	EC-346698	-1.203	0.512	-0.054	-0.511	0.028	0.753
55	EC-346779	-0.009	-1.602	-0.11	0.946	-0.386	-0.094
56	EC-346926	2.409	-1.667	-0.104	0.904	-0.216	-0.684
57	EC-346999	-1.006	-1.288	0.033	1.102	-0.782	0.784
58	EC-347070	-2.12	-1.881	0.466	1.28	-0.827	-1.199
59	EC-347128	-0.719	0.274	0.169	0.746	-0.215	-0.891
60	EC-347142	0.091	0.932	1.069	-0.208	-0.623	-0.727
61	EC-347166	-0.89	-1.194	-0.622	0.949	-0.521	0.426
62	EC-350647	-0.341	-2.706	-0.027	0.604	-0.527	-0.057
63	EC-351826	0.872	0.403	-1.534	-0.236	0.479	-0.792
64	EC-351882	-0.512	-0.092	-1.132	1.633	-0.826	-0.025
65	EC-351887	-1.944	-2.218	-0.754	0.728	-0.707	-1.372
66	EC-351901	0.102	-1.194	-2.59	-0.59	0.929	0.971
67	EC-351904	0.49	-1.596	-1.309	-0.157	0.405	0.791
68	EC-355653	1.674	-1.549	-0.595	-0.02	0.14	-0.777
69	EC-355674	4.28	-1.253	0.278	1.159	0.154	1.162
70	EC-357021	1.089	-1.361	0.118	-0.409	0.249	0.543
71	EC-357313	-1.145	-0.38	0.428	0.651	-0.907	-0.645
72	EC-357327	-0.091	-1.876	-1.807	-0.443	0.522	0.11
73	EC-350401	-2.46	1.372	-0.146	1.323	-0.546	0.053
74	EC-376977	-2.944	0.632	3.781	5.196	7.189	-0.406
75	EC-346144	-0.825	0.345	-1.989	0.003	0.214	0.683
76	EC-346400	1.821	-0.817	0.804	1.703	-1.192	-0.23
77	EC-346411	2.638	0.218	0.487	1.209	-0.957	0.023
78	EC-346440	3.257	-0.439	0.04	-0.091	0.251	0.541
79	EC-346681	4.499	-1.338	2.046	0.037	-0.224	0.053
80	EC-346750	-0.991	-1.351	-1.044	-0.655	0.542	-0.343
81	EC-351880	0.833	0.458	0.132	-1.035	0.399	-0.024
82	EC-351888	-2.131	2.839	-0.539	0.079	-0.195	-1.984
83	EC-351903	-1.366	1.026	0.162	0.67	-0.588	0.072
84	EC-355670	-0.523	-0.677	-1.607	-0.281	1.379	0.251
85	EC-361719	-1.331	2.29	-0.998	-0.126	-0.122	1.003
86	EC-377321	2.661	0.733	-0.47	1.372	-0.796	-1.155
87	EC-377351	0.919	-0.799	0.143	-0.945	0.617	0.195
88	EC-0346210	0.437	0.531	-2.406	0.308	0.243	0.506
89	EC-0346341	1.297	2.638	0.436	0.271	-0.253	-0.053
90	EC-334956	-2.96	2.321	-0.743	1.428	-0.712	1.403
91	EC-334958	0.761	-0.803	-1.199	1.013	0.523	1.1
92	EC-340538	1.647	1.403	-1.092	-0.609	1.416	-0.835
93	EC-346173	-1.657	0.063	0.364	-1.095	-0.219	-1.121
94	EC-346174	2.343	0.788	-0.203	0.307	0.159	-2.447
95	EC-346175	0.743	0.295	-0.319	-1.155	0.368	1.301
96	EC-346186	-2.384	-0.742	1.188	0.925	-0.834	-1.137
97	EC-346197	0.042	0.2	-0.998	2.211	-0.858	0.4

S. No.	Genotypes	PC1 score	PC2 score	PC3 score	PC4 score	PC5 score	PC6 Score
98	EC-346226	0.135	-2.041	-1.504	0.011	0.215	-1.792
99	EC-346229	-0.825	0.345	-1.989	0.003	0.214	0.683
100	EC-346265	1.082	1.513	0.152	1.5	-1.722	-0.231
101	EC-346275	-1.931	-0.644	1.634	0.563	-0.934	-2.088

Table 5: List of selected genotypes in each principal component having positive values

PC1	PC2	PC3	PC4	PC5	PC6
TKG 22	TKG 22	EC-109693-A	GT 10	EC-177-1	RT-351
GT 10	GT 10	EC-118571-B	RT-351	EC-34625	EC346478
JTS 8	JTS 8	EC-346480	EC-89105	EC-346471	EC-334998
TKG 21	TKG 21	EC-356526	EC-335017	EC-355670	EC-335017
RT-351	EC-43708	EC-133850-S	EC-346471	EC-340538	EC-346391
EC-38	EC-109693-A	EC-132828	EC-346599	EC-355674	
EC-89671	EC-118569	EC-334993	EC-346999	EC-361719	
EC-118571	EC-346347	EC-334998	EC-347070	EC-334956	
EC-34625	EC-346480	EC-335015	EC-351882	EC-334958	
EC-346514	EC-132828	EC-342782	EC-355674	EC-346175	
EC-178365	EC-178374	EC-346165	EC-350401		
EC-178374	EC-334998	EC-346391	EC-346400		
EC-334962	EC-350401	EC-346424	EC-346411		
EC-335005	EC-376977	EC-346471	EC-377321		
EC-334998	EC-351888	EC-346633	EC-334956		
EC-335015	EC-351903	EC-347142	EC-334958		
EC-334999	EC-361719	EC-346681	EC-346197		
EC-346424	EC-0346341	EC-346186	EC-346265		
EC-346599	EC-334956	EC-346275			
EC-346633	EC-340538				
EC-346926	EC-346265				

EC-355653					
EC-355674					
EC-357021					
EC-346400					
EC-346411					
EC-346440					
EC-346681					
EC-377321					
EC-0346341					
EC-340538					
EC-346174					
EC-346229					
EC-346265					

UNDER PEER REVIEW