

Original Research Article

In-vitro assessment of different culture of media on the isolation and purification of cauliflower Alternaria Leaf spot disease caused by Alternaria brassicicola (Schwein)

Abstract

Alternaria leaf spot of cauliflower caused by *Alternaria brassicicola* (Schwein.) Wiltshire, is a significant disease of cauliflower in the Kanpur region. In the current study the pathogenic fungus was purified from infested cauliflower leaves on potato dextrose agar medium, Following inoculation on healthy cauliflower seedlings the pathogenicity of the causal agent was demonstrated. The results revealed that the fungus was *Alternaria brassicicola* based on common symptoms on foliage, microscopic findings, and culture characteristics of the fungus. Based on microscopic observation, the causal agent was identified as *Alternaria brassicicola* (Schwein.) Wiltshire. To proceed to the isolation and purification of the fungus, eight culture media were tested among which the potato dextrose agar medium was found most suitable and encouraged maximum radial mycelial growth (89.07mm) of *A. brassicicola*. The second best was Corn Meal Agar medium (67.46mm), followed by Potato carrot agar medium (54.83 mm), Rose Bengal agar (53.00 mm), Czapek dextrose agar (40.40 mm), Yeast Extract Agar (32.28 mm), and Nutrient Agar (24.28 mm), While minimum (22.16 mm) growth was recorded in Starch Agar. Thus, potato dextrose agar can therefore be used in the isolation, purification and pathogenicity of *A. brassicicola*.

Key words: Agriculture, Cauliflower, culture, growth, isolation and pathogenicity.

Introduction

Cauliflower (*Brassica oleracea* var. *botrytis*) is one of the most important winter vegetables grown in India. It is a cool-weather crop and closely related to broccoli and cabbage. however, cauliflower has more specific climatic requirements than any of the other cole crops. With the development of heat tolerant varieties, cauliflower is available from September to May in plains of India. The word cauliflower is derived from the Latin words 'caulis' meaning stalk and 'floris' meaning flower. It is grown for its white tender curd or head. Botanically, the edible part of cauliflower is hypocotyl branches or pre-floral fleshy epical meristem (Dhaliwal, 2017).

Cauliflower is grown in China, India, USA, Spain, Italy, Mexico and France, due to its higher nutritional value and widespread cultivation.

“In India, cauliflower was grown in 2022 in an area of about 452.59 thousand hectares with the production of 8668.22 thousand tonnes and the productivity was about 19.15 MT/ha” (Pavithra and Singh 2020).

In 2021-2022 growing season, cauliflower was grown in Pradesh in area of 22.85 million hectares and the production was 1.42 MT/h (**Horticulture statistics at a glance 2022**). “In India, cauliflower is cultivated in almost all the states, mainly Bihar, Uttar Pradesh, Orissa, Assam, Madhya Pradesh, Gujarat and Haryana”. (Gupta *et. al.* 2017). “Cauliflower plays an important role in the human diet due to its attractive appearance, good taste, and its nutritive value. It is a rich source of protein, carbohydrates, vitamin-B, and C as well as various minerals those are necessary for the human health. Cauliflower is grown for its edible flowering head and consume as a vegetable in curries, soups, and pickles” (Wani and Kaul, 2011).

Materials and Methods

Visual observations

To track the progression of the disease in a plant population under natural circumstances, visual observations of disease symptoms were made in the field and documented.

Collection of the diseased samples

“Leaves of cauliflower with dark brown spots having characteristic concentric circumferences, often with a yellowish chlorotic halo were collected from farmers field of nearby CSAUA&T, Kanpur and brought to the laboratory for isolation and further studies”. (Maude 1980, Huq 1999, Hossain 2005).

Microscopic study

“Spores of *A. brassicicola* were taken from the pure culture and mounted on the clear glass slide. Spores were mixed with thoroughly with lactophenol in order to obtain a uniform spread over which a cover slip was placed. The spores and hyphae of the fungus were observed for spore shape and size and photomicrographs were taken”.(Chadar *et al.* (2016)

Isolation

“For isolation of pathogen, small pieces of the leaves were cut from the diseased portion along with some healthy tissues and surface sterilized with 1% sodium hypochlorite solution for one minute followed by three consecutive washings with sterilized distilled water. The surface sterilized pieces were transferred to Petri plates containing Potato Dextrose Agar (PDA) and incubated at 25+1°C in BOD incubator. After seven days of incubation, the fungal growth was

transferred aseptically to PDA slants and purified following single spore technique". (Valvi *et al.* 2019 and Chatta *et al.* 2022)

Pathogenicity test

The pathogenicity was tested by following Koch's postulates. Cauliflower leaves were artificially inoculated with *A. brassicicola* by pin prick method (Pattanamahakul and Strange, 1999). To harvest conidia of the fungus, sterile distilled water was added to 10 days old culture growing on PDA Petri plates. The fungal spores were gently scrapped with the help of sterilized slide and conidial suspension was filtered through sterilized muslin cloth. The concentration of conidial suspension was adjusted to 4×10^5 spores/ml with the help of haemocytometer. Subsequently, conidial suspension was sprayed on 35 days old cauliflower seedlings grown in pots having dimensions 7×8.5 cm. Before inoculation the leaves were surface sterilized with 70 per cent ethanol with the help of a cotton swab. Then gentle pricking was done with the help of sterilized needle. After inoculum spraying, plants were covered with perforated and moistened transparent plastic cover at a temperature ranging from $20-25^\circ\text{C}$. Inoculated plants were labeled and kept under humid conditions to maintain proper moisture for disease development. High relative humidity was maintained with help of water spraying inside the polythene bags after every 12 hours. Leaves were observed for symptom development at regular intervals after inoculation. In case of control plants, sterile distilled water was sprayed. (Kashyap and Dhiman 2009)

Effect of culture media on growth and sporulation

Alternaria brassicicola isolates were grown on eight different media viz. Potato dextrose agar (PDA), Corn meal agar (CMA), Czapek's (dox) agar (CZA), Starch agar (SA), Nutrient agar (NA), Yeast extract agar (YEA), Potato carrot agar (PCA), and Rose Bengal agar (RBA) at $25 \pm 2^\circ\text{C}$ to identify the medium on which fungus grow faster than other. (Swati Deep *et al.* 2014)

1. Potato dextrose agar (PDA)

Potato	: 200 g
Dextrose	: 20 g
Agar-agar	: 20 g
Distilled water	: 1000 ml

2. Corn meal agar (CMA)

Corn meal	: 20 g
-----------	--------

Agar-agar	: 18 g
Distilled water	: 1000 ml

3. Czapek's dox medium

Sucrose	: 30.00 g
Sodium nitrate (NaNO ₃)	: 2.00 g
Potassium dihydrogen phosphate (KH ₂ PO ₄)	: 1.00 g
Magnesium sulphate (MgSO ₄ 7H ₂ O)	: 0.50 g
Ferric chloride (FeCl ₃ 6H ₂ O)	: 0.01 g
Potassium chloride (KCl)	: 0.50 g
Agar - agar	: 20.00 g
Distilled water	: 1000 ml

4. Starch's agar

Soluble starch	: 2.00 g
Beef extract	: 3.00 g
Bacto peptone	: 5.00 g
Agar agar	: 5.00 g
Distilled water	: 1000 ml.

5. Potato carrot agar (PCA)

Carrot	: 200. g
Potato	: 250. g
Agar	: 15. g
Distilled water	: 1000 ml.

6. Rose Bengal agar (RBA)

Soytone	: 5 g
Dextrose	: 10 g
KH ₂ PO ₄	: 1 g
Magnesium Sulfate	: 0.5 g
Rose Bengal	: 0.05 g
Agar	: 15 g
Distilled water	: 1000 ml

7. Yeast extract agar (YEA)

Soluble starch	: 10.00 g
Yeast extract	: 1.00 g
Agar agar	: 20.00 g
Distilled water	: 1000 ml

8. Nutrient agar (NA)

Yeast extract/Beef extract	: 3 g
Peptone	: 5 g
NaCl	: 5g
Agar	: 15 g
Distilled water	: 1000 ml

The media were prepared in a beaker where a quantity (grams) of the dehydrated powder or lab-prepared media was added to 1000 milliliters of distilled or deionized water. Suspension was then heated to boiling to dissolve the medium completely. The dissolved medium was then autoclaved at 15 lbs pressure (121°C) for 15 minutes. Once the autoclaving process was completed, the beaker was taken out and cooled to a temperature of about 40-45°C. If enrichment was desired, the addition of blood or biological fluids could be done following the autoclaving process. The media was then poured into sterile Petri plates under aseptic conditions. Once the media solidifies, the plates were placed in the hot air oven at a lower heat setting for a few minutes to remove any moisture present on the plates before further use.

Eight media were evaluated in the present study. Media were prepared with given composition. The initial pH of each medium was adjusted to 6.5 prior to autoclaving. The medium was prepared with given composition and dispensed in conical flask.

The flasks were plugged with non-absorbent cotton plugs and sterilized in an autoclave at 15 lbs. psi for 20 minutes. Petri plates were sterilized in hot air oven at 160 °C for 1 hour. Such sterilized Petri plates were poured with 20 ml of molten medium and allowed to solidify. To isolate the causal agent, five millimeter diameter disc of the test fungus was cut with the help of incinerated cork borer and inoculated at the center of Petri plates. The inoculated plates were then incubated at room temperature ($27 \pm 2^{\circ}\text{C}$) for 7 days. The compositions of all the media used were obtained from Ainsworth and Bisby's Dictionary of the fungi as mentioned above. Each treatment was replicated thrice. The measurement of the colony diameter was taken when the maximum (90mm) growth was achieved in any one of the media tested. The cultural

characters such as colony diameter, colony colour and degree of sporulation were recorded.

Statistical analysis

In the present investigation, lab experiment was conducted in complete randomize design. The data obtained from all the experiments were statically analyzed following the standard procedures (Gomez and Gomez 1984)

Results and Discussion

Visual observation

“Visual observations of disease symptoms were recorded in the field from all the aerial parts of the plants *viz.*, stem, leaves, fruits, pods and heads. Generally, in the beginning the spots remain small and circular or elliptical in shape with colour of spots varying from species to species” (Neeraj and Shilpi Verma 2010). These may be pale, brown, olivaceous brown, grayish or black *etc.* and typically surrounded by halo-chlorotic tissues. Later on these spots enlarged into gray to black lesions of 0.5 to 1 cm diameter. Then after the spots gradually increase in size (varying with species) in a concentric manner and often coalesce, leading to leaf spot appearance. (PLATE-1)

Microscopic examination

“Diseased leaves samples, exhibiting the characteristic of *Alternaria* leaf spot of cauliflower were examined visually followed by microscopic examination of sections of infected tissues. The pure culture isolates obtained from the diseased specimens of leaves were identified as *Alternaria brassicicola* (Schw.) Wiltsh” (Kumar et al., 2014). The morphological studies revealed that the mycelium of the fungal pathogen was septate, olive grey to greyish black in colour, conidiophores were olivaceous, septate and branched, conidia were dark cylindrical to oblong, muriform and produced in chains of 8-10. The conidia were devoid of any prominent beak and found to have 5-8 transverse and 0-4 longitudinal septation. The identification of the fungus was further reconfirmed as *Alternaria brassicicola* (Schw.) Wiltsh. (PLATE-2) (Kumar *et al.*, 2014).

Isolation

On potato dextrose agar medium, the pathogen was successfully isolated from diseased tissue displaying well-developed lesions together with healthy fraction that was brought to the lab from naturally infected cauliflower plants. The inoculation plates were incubated in the BOD incubator for 5 to 7 days at a temperature of 25 to 20C. The fungus culture was transplanted to

PDA in Petri plates and multiplied in the lab after being isolated from diseased tissue.

Purification of fungal culture

After seven days of incubation, the fungus generated greenish-gray to black-colored, fluffy, lanose to loose cottony growth on potato dextrose agar medium. The pure culture's slants were sealed with paraffin wax and kept in the lab's refrigerator for later use.

Pathogenicity test

The pure culture of *Alternaria brassicicola* obtained by single spore isolation method was used for pathogenicity test. The test was carried out by spore suspension spray inoculation on the foliage of 35 days old susceptible Pusa snowball variety of cauliflower. The symptoms appeared on inoculated leaves as brown, circular or oval necrotic spots with concentric rings and surrounded by yellow halo after twenty days after inoculation (Plate 2). The fungus was re-isolated and purified and compared for similarity to that of original culture. The plants which were not inoculated with the fungal spore suspension did not show any symptoms of the disease. Similar technique was followed by (Humpherson-Jones, 1992; Paul and Rawlinson, 1992; Verma and Saharan, 1994; Poapst *et al.*, 1979). The re-isolated *A. brassicicola* exhibited similar characters as in the originally isolated culture. (Akhtar *et al.* 2004)

Identification of the pathogen

The reisolated pure fungal culture was identified in the Department of Plant Pathology, Chandra Shekhar Azad University of Agriculture and Technology Kanpur by comparing its morphological characters with the information available on the standard websites for fungal identification as well as in the reviewed. In the current study, the causal agent was identified and confirmed as *A. brassicicola*. Earlier Ellis (1971) first time described *A. brassicicola* on cauliflower leaves producing dark brown to almost black circular 1-10 mm diameter zonate spots.



Plate 1: General view of field symptoms of leaf spot in cauliflower



Plate.2: Pure Culture of *Alternaria brassicicola* and its spores

Effect of culture media on growth and sporulation of *Alternaria brassicicola* (Schwein.) Wiltshire.


Radial growth of the pathogen


In order to find out the most effective culture medium for the growth of *A. brassicicola*.

eight culture media were evaluated against *A. brassicicola* under *in vitro* condition (Table 1). The result reveals that Potato dextrose agar medium was significantly superior over other tested media at 3, 5 and 7 days after inoculation. **(Fig-1)**

Seven days after inoculation maximum mycelial growth was recorded in Potato dextrose agar medium (89.07mm) followed by Corn meal agar (67.46mm), Potato carrot agar medium (54.83 mm), Rose Bengal agar (53.00 mm), Czapek dextrose agar (40.40 mm), Yeast Extract Agar (32.28 mm), and Nutrient Agar (24.28 mm), While minimum (22.16mm) growth was recorded in Starch Agar **(Plate-3)**.

The mycelial growth of *A. brassicicola* fully covered the plate on Potato dextrose agar medium at 7 days after inoculation and it was significantly superior over the remaining tested media. The next effective medium was Corn meal agar medium which was significantly superior over eight media. Excellent sporulation more than 20 spores/microscopic field) was observed on Carrot potato dextrose agar medium, Corn Meal Agar medium, Rose Bengal agar medium, and potato dextrose agar medium, whereas good sporulation was found in Yeast Extract Agar medium and Czapek dextrose medium. Poor sporulation was found in 2 culture media viz., Nutrient Agar and starch agar medium **(Table-1)**. Similar result was also revealed by (Kamei and Singh 2020 and Rex and Rajasekar 2021).





T₅

T₆

T₇

T₈

Plate.3: Effect of different culture media on growth of *Alternaria brassicicola*
Colour of the culture

Among all media tested for evaluation, colour of culture did not differ from each other.

Conclusion

It can be concluded that *Alternaria* leaf spot of cauliflower, which is brought on by *Alternaria*. The current study revealed that cauliflower was infected by *A. brassicicola* as the causal agent of *Alternaria* leaf spot of cauliflower. Among media tested PDA can be used because it makes the fungus grow quickly.

Reference

- Akhtar KP, Saleem MY, Asghar M, Haq MA (2004) New report of *Alternaria alternata* causing leaf blight of tomato in Pakistan. *Plant Pathology* **6**: 816
- Chadar LK, Singh RP, Singh RK, Yadav RR, Mishra MK, Pratap N, Vishnoi RK (2016) Studies on *Alternaria* in rapeseed mustard (*Brassica juncea* L.) caused by *Alternaria brassicae* (Berk.) Sacc. and its integrated management. *Plant Archives* **16**(2): 897-901
- Chatta, M. B., Muhammad. B., Razzaq, Shafique, S., Siddique, M. And Peerzada, H.H.E. (2022) Isolation, Characterization And Management Of *Alternaria* Leaf Blight Of Turnip through botanicals. *Pakistan Journal of Phytopathology*. **34**(2) 2305-0284
- Dhaliwal and S.N. Vishwakarma (1999). Morphological, and symptomatological variations in *Alternaria alternata* causing leaf blight in brinjal. *Indian Journal of Mycology, and Plant Pathology*, **29**(3) : 350- 354.
- Ellis, M.B. (1971). Dematiaceous hyphomycetes. Commonwealth Mycological Institute, Kew, Surrey, England. No-608; pp. 464-497.
- Gomez, K. A., & Gomez, A. A. (1984). *Statistical procedures for agricultural research*. John Wiley & sons.

- Gupta, V., Singh, D., Mishra, A. K., Singh, B. P., Kumar, R., & Pandey, R. K. (2017). A study on constraints faced by cauliflower growers in cauliflower cultivation in western Uttar Pradesh, India. *International Journal of Current Microbiology and Applied Sciences*, 6(7), 2646-2651.
- Hossain, Sakhawat, M. and Mian, I. H. (2005). Effect of planting date on Alternaria Blight and seed yield of cabbage. *Bangladesh Journal Plant Path.* 21:33-37.
- Humpherson-Jones, F. M., Dixon, G. R., Craig, M. A., & Ann, D. M. (1992). Control of clubroot using calcium cyanamide-a review. In *Brighton Crop Protection Conference, Pests and Diseases-1992. Volume 3.* (pp. 1147-1154). British Crop Protection Council.
- Huq, M. I., Malaker, P. K. and Nahar M. S. (1999). Economic assessment of yield loss of garlic due to leaf blotch disease. *Ann. Bangladesh Agric.* 9:11-17.
- Kamei, D. and Singh, A. U. (2020) In-Vitro Studies of Different Culture Media and Biocontrol Agents on Growth and Sporulation of *Alternaria Alternata* (Fr.) Keissler an Incitant of Broad bean (*Vicia Faba L.*) Leaf Blight Disease. *International Journal of Environmental & Agriculture Research*, 6 (11) 2454-1850
- Kashyap, P.L. and Dhiman, J. (2010). Ecofriendly strategies to suppress the development of Alternaria Blight and black rot of cauliflower. *World Appl. Sci. J.* 9:345-350.
- Kumar D, Maurya N, Bharati Y K, Kumar A, Kumar K, Srivastava K, Chand G, Kushwaha C, Singh S K and Mishra R K. 2014. Alternaria blight of oilseed brassicas: A comprehensive review. *African Journal of Microbiology Research* 8: 2816–2829.
- Maude, R. B., & Humpherson-Jones, F. M. (1980). Studies on the seed-borne phases of dark leaf spot *Alternaria brassicicola* and grey leaf spot *Alternaria brassicae* of brassicas. *Annals of Applied Biology*, 95(3), 311-319.
- Neeraj And Shilpi Verma., 2010, *Alternaria* diseases of Vegetable Crops and New Approaches for its Control. *Asian J. Exp. Biol. Sci.*, 1: 681-692
- Pattanamahakul, & Strange. (1999). Identification and toxicity of *Alternaria brassicicola*, the causal agent of dark leaf spot disease of Brassica species grown in Thailand. *Plant Pathology*, 48(6), 749-755.
- Paul, V. H., & Rawlinson, C. J. (1992). Diseases and Pests of Rape, Verlag Th. Mann Gelsenkirchen-Buer, Germany.
- Pavithra, S., & Singh, S. P. (2020). Economics of production and marketing of cauliflower in

Indore district of Madhya Pradesh. *The Pharma Innovation J*, 80-83.

Poapst, P. A., Ramsomair, B. A., & Gourley, C. O. (1979). On the promotion of senescence in *Brassica oleracea* var. capitata by *Alternaria brassicicola* and by *Botrytis cinerea*. *Canadian Journal of Botany*, 57(21), 2378-2386.

Rex, B. and Rajasekar, G. (2021) Influence Of Various Media And Nutrient Sources On *Alternaria Solani* Cause Early Blight Disease In Tomato. *Plant Archives Volume 21*, No 1, 2021 pp. 1470-1474

Swati Deep, Pratibha Sharma, Nirajian Behera And Pallem Chowdappa, 2014, Diversity in Indian Isolates of *Alternaria brassicicola* (Schwein) Wiltshire Causing Black leaf spot Disease in Cauliflower. *Pl. Pathol. J.*, ISSN 1812-5387.

Valvi HT, Kadam JJ, Bangar VR (2019) Isolation, pathogenicity and effect of different culture media on growth and sporulation of *Alternaria brassicae* (berk.) Sacc. causing *Alternaria* leaf spot disease of cauliflower. *International Journal of Current Microbiology and Applied Science* 8(4): 1900-1910.

Verma, P. R., & Saharan, G. S. (1994). *Monograph on Alternaria diseases of crucifers. Saskatoon Research Centre.*

Wani, Towseef A., Monika Sood, and Raj Kumari Kaul (2011) Nutritional and sensory properties of roasted wheat noodles supplemented with cauliflower leaf powder. *Annals. Food Science and Technology* 12.2: 102-107.

Table.1: Effect of culture media on growth and sporulation of *Alternaria brassicicola*

Treatments	Culture media	Average mycelial growth (mm)			Sporulation
		3 DAS	5 DAS	7 DAS	
T ₁	Czapek's Dox Agar	11.66	22.00	40.40	++
T ₂	Potato Carrot Agar	13.33	28.66	54.83	+++
T ₃	Potato Dextrose Agar (PDA)	31.00	62.99	89.07	+++
T ₄	Starch Agar	12.66	16.33	22.16	+
T ₅	Nutrient Agar	14.00	19.66	24.48	+
T ₆	Corn Meal Agar	18.33	51.00	67.46	+++
T ₇	Yeast Extract Agar	12.00	22.33	32.28	++
T ₈	Rose Bengal Agar	25.33	36.66	53.00	+++
	C.V (%)	2.41	5.86	6.01	
	S.Em.±	0.66	1.10	0.60	
	C.D. at 5 %	2.02	3.32	1.86	




Fig.1: Effect of culture media on growth of *Alternaria brassicicola*.

UNDER PEER