

Character Association and Path Analysis for yield and yield attributing traits in Bread Wheat [*Triticum aestivum* (L.) emThell] genotypes

ABSTRACT

The present investigation was carried out at Instructional Agronomy Farm, Rajasthan College of Agriculture, MPUAT, Udaipur during *Rabi*, 2021-2022 to estimate the association among yield components and their direct and indirect influence on grain yield of bread wheat. Significant genotypic differences were obtained for the entire examined trait depicting presence of huge amount of variation among studied genotypes. Grain yield per plant had positive and significant correlation with days to 50% heading, plant height, length of main spike, number of spikelets per spike, number of grains per spike, grain weight per spike and biological yield per plant at both genotypic and phenotypic levels. The maximum positive direct effect on grain yield per plant was exhibited by plant height followed by biological yield per plant, number of grains per spike, days to maturity and 1000-grain weight. Consequently, by considering specific traits *viz.*, plant height, biological yield per plant, number of grains per spike, it becomes possible to improve the grain yield per plant in bread wheat.

Keywords: Bread wheat, correlation, genotypes, variation, yield

1. INTRODUCTION

Bread wheat (*Triticum aestivum* (L.) em Thell) is an annual plant belongs to the family Poaceae, is one of the most important staple foods as well as the most consumed cereal amongst major cereals of the world. It is a monocotyledonous angiosperm originated in South West Asia. It stands second next to rice in production among cultivated crops and has been considered as “versatile cereal food”. In India, it is grown on an area of 31.82 million hectares with a production of 112.74 million tonnes and productivity of 3543 kg/ha (Anonymous, 2023). In addition to being a major source of starch and energy, it is good source of notably protein, vitamins (notably B vitamins), dietary fiber, and phytochemicals (Liu, 2011, Regvaret *al.*, 2011, and Lafiandraet *al.*, 2014). Wheat contains nearly 70% carbohydrate, 12% protein, 1.7% fat, 2.7% minerals, 2% dietary fiber, 12% moisture and considerable proportions of vitamins (Gillieset *al.*, 2012). The wheat grain contains 2-3% germ, 13-17% bran (outer layers of wheat grain) and 80-85% mealy endosperm (on a dry matter basis) (Belderoket *al.*, 2000). The demand for wheat is rising and it is expected that by 2050, the requirement of wheat would be 60% higher than the present year (FAO, 2018). Therefore, in the 21st century, meeting the food demand of a growing population necessitates significant enhancements in crop production and productivity (Poudel & Poudel, 2020).

Knowledge of the existing variation and the extent of association among yield contributing characters are essential for the development of high yielding cultivars. Correlation studies merely quantify the relationship between yield and its related attributes without providing the extent of yield dependence on the correlated factors. Path coefficient analysis measures the

direct effect of a predictor variable upon its response variable and the second component being the indirect effect(s) of a predictor variable (Dewey and Lu, 1959). To increase the yield, study of direct and indirect effects of yield components provides the basis for its successful breeding programme and hence the problem of yield increase can be more effectively tackled on the basis of performance of yield components and selection for closely related characters. So, the objective of this investigation was to study the inter-relationships, and direct and indirect effects of yield and yield attributing traits in the bread wheat genotypes.

2. MATERIALS AND METHODS

The experimental material comprised of 40 bread wheat genotypes obtained from AICRP on Wheat and Barley, Department of Genetics and Plant Breeding, Rajasthan College of Agriculture, MPUAT, Udaipur. All the 40 bread wheat genotypes were grown in Randomized Block Design with 3 replications. In each replication, genotypes were sown in 3 rows of 5m length with row to row and plant to plant to plant spacing of 20 cm and 5 cm, *respectively*. All the recommended package of practices was followed to raise a good and healthy crop. Five competitive plants from each genotype were selected randomly for recording observations for 13 traits under study *viz.*, plant height (cm), number of effective tillers per plant, length of main spike (cm), number of spikelets per spike, number of grains per spike (g), grain weight per spike (g), 1000-grain weight(g), biological yield per plant(g), grain yield per plant(g), harvest index(%) and protein content(%). The data on days to 50 percent heading and days to maturity were recorded on the whole plot basis. The analysis of variance was worked out per the method suggested by Panse and Sukhatme(1985). Genotypic and phenotypic correlation coefficient was computed as per the formula suggested by Al- Jibouriet *al.* (1958). The path coefficient was calculated by using the method of Dewey and Lu (1959). The protein content was estimated by using method of Linder (1944).

3. RESULTS AND DISCUSSION

The analysis of variance has unveiled noteworthy distinctions among the genotypes for the majority of the studied traits, signifying a significant level of variability in the materials. This substantiates the rationale behind selecting these experimental materials (Table 1).

3.1 Correlation coefficient analysis

The present investigation revealed that grain yield per plant had positive and significant correlation with days to 50 percent heading, plant height, length of main spike, number of spikelets per spike, number of grains per spike, grain weight per spike and biological yield per plant at both genotypic (fig. 1) and phenotypic levels (fig. 2) while it showed positive and significant correlation with harvest index at genotypic level only (Table 1 & fig.1). Similar results were also reported by Baye *et al.*(2020), Nagar *et al.*(2018). Days to maturity shows negative and significant correlation with harvest index while biological yield per plant shows negative and significant correlation with harvest index and protein content at both genotypic (fig.1) and phenotypic levels (fig. 2). While correlation analysis offers valuable insights, it is limited in that it does not shed light on the underlying causal factors driving the various inter-relationships. However, by conducting an analysis of path coefficients, we can gain a more comprehensive understanding of the causal basis behind these inter-relationships (Nukasani *et al.*, 2013).

3.2 Path coefficient analysis

3.2.1 Direct and indirect effects

The findings from the current investigation on path coefficient analysis are showcased in figure 3. The results indicate that several factors, plant height, biological yield per plant, number of grains per spike, days to maturity and 1000- grain weight had positive direct effect on grain yield. Consequently, focusing and selecting these specific traits would prove beneficial in effectively enhancing the overall productivity of bread wheat. However, traits like length of main spike, days to 50 percent heading, number of spikelets per spike, grain weight per spike, protein content and number of effective tillers per plant exhibited negative direct effect on grain yield per plant. Since the direct effect were negative, so the direct selection for these traits to improve yield will be undesirable. Similar results were in accordance with findings of Kumar *et al.*(2014),Ayer *et al.*(2017),Nagar *et al.*(2018). Number ofgrains per spike showed positive indirect effect *via* plant height, biological yield per plant.

3.2.2 Residual effect

Residual effect in the present study was computed ($R= 0.1037$) showing that 89.63 % of the variability in grain yield was demonstrated by the traits considered for path study. It also indicates that in addition to the studied characters, there are also other factors to justify grain yield per plant (AbdEl-Mohsen *et al.*, 2012)

Table 1: Analysis of variance for yield and its contributing traits in bread wheat

S.NO	Characters	Replication	Genotype	Error
		[2]	[39]	[78]
1.	Days to 50 percent heading	0.41	21.23**	2.79
2.	Days to maturity	0.91	27.28**	3.67
3.	Plant height (cm)	44.78	136.15**	28.45
4.	Number of effective tillers per plant	0.93	1.19**	0.40
5.	Length of main spike (cm)	0.93	2.98**	0.57
6.	Number of spikelets per spike	0.93	6.72**	1.04
7.	Number of grains per spike	35.21	376.65**	13.04
8.	Grain weight per spike (g)	0.66	0.89**	0.24
9.	1000-grain weight (g)	1.61	60.88**	3.20
10.	Biological yield per plant (g)	4.36	95.26**	9.55
11.	Grain yield per plant (g)	4.61	16.31**	2.78
12.	Harvest index (%)	15.56	34.83**	8.57
13.	Protein content (%)	0.51	6.67**	0.27

*, ** Significant at 5% and 1% respectively

Figure 1: Genotypic (rg) correlation coefficient for different characters in bread wheat

Character	Days to 50% heading	Days to maturity	Plant height	Number of effective tillers per plant	Length of main spike	Number of spikelets per spike	Number of grains per spike	Grain weight per spike	1000-grain weight	Biological yield per plant	Harvest index	Protein content	Grain yield per plant
Days to 50% heading		0.48**	0.56**	0.21	0.36*	0.16	0.28	0.25	0.03	0.43*	-0.12	-0.35*	0.39*
Days to maturity			-0.14	0.32	-0.01	0.06	0.00	-0.11	0.11	0.02	-0.37*	-0.11	-0.20
Plant height				-0.19	0.74**	0.63**	0.57*	0.67**	0.21	0.52**	0.31	0.06	0.73**
Number of effective tillers per plant					-0.44**	-0.32	-0.30	-0.30	-0.15	0.01	-0.19	-0.10	-0.06
Length of main spike						0.74**	0.78**	0.99**	0.41*	0.63**	-0.20	-0.01	0.51**
Number of spikelets per spike							0.77**	0.79**	0.22	0.49**	0.01	0.02	0.50**
Number of grains per spike								1.01**	0.17	0.57**	0.02	-0.15	0.60**
Grain weight per spike									0.46**	0.64**	-0.03	-0.06	0.63**
1000- grain weight										0.20	-0.09	0.09	0.17
Biological yield per plant											-0.37*	-0.35*	0.81**
Harvest index												0.15	0.41**
Protein content													-0.22
Grain yield per plant													

*, ** significant at 5% and 1% respectively, here 0.00 values indicate negligible correlation

Figure 2: Phenotypic (rp) correlation coefficient for different characters in bread wheat

Character	Days to 50 percent heading	Days to maturity	Plant height	Number of effective tillers per plant	Length of main spike	Number of spikelets per spike	Number of grains per spike	Grain weight per spike	1000-grain weight	Biological yield per plant	Harvest index	Protein content	Grain yield per plant
Days to 50percentheading		0.38**	0.31**	0.08	0.27*	0.19	0.24	0.13	0.03	0.35**	-0.03	-0.28*	0.32**
Days to maturity			-0.10	0.10	0.05	0.05	0.02	0.00	0.08	0.06	-0.25*	-0.07	-0.12
Plant height				0.02	0.54**	0.42**	0.43*	0.30**	0.13	0.37**	0.19	0.06	0.49**
Number of effective tillers per plant					-0.23*	-0.14	-0.23	-0.19	-0.09	0.02	-0.05	-0.01	-0.01
Length of main spike						0.49**	0.60**	0.56**	0.28	0.46**	-0.06	0.01	0.37**
Number of spikelets per spike							0.63**	0.49**	0.14	0.38**	-0.01	0.02	0.34**
Number of grains per spike								0.66**	0.15	0.49**	0.02	-0.13	0.46**
Grain weight per spike									0.27*	0.42**	-0.07	-0.08	0.34**
1000- grain weight										0.19	-0.08	0.08	0.14
Biological yield per plant											-0.29*	-0.26	0.75**
Harvest index												0.05	0.26
Protein content													-0.26
Grain yield per plant													

*, ** significant at 5% and 1% respectively

Figure 3: Estimate of direct effect (diagonal) and indirect effect (off diagonal) at genotypic level for grain yield/plant in bread wheat

S. No.	Character	Days to 50% heading	Days to maturity	Plant height	Number of effective tillers per plant	Length of main spike	Number of spikelets per spike	Number of grains per spike	Grain weight per spike	1000-grain weight	Biological yield per plant	Protein content	r with grain yield per plant
1	Days to 50% heading	-0.523	0.0889	0.6635	-0.0244	-0.2811	-0.0574	0.1936	-0.064	0.0044	0.3234	0.0646	0.3885*
2	Days to maturity	-0.2531	0.1837	-0.1626	-0.037	0.0067	-0.0213	-0.0026	0.0286	0.0166	0.0175	0.0211	-0.2023
3	Plant height	-0.2908	-0.025	1.1934	0.0212	-0.571	-0.2271	0.3943	-0.1731	0.0323	0.3901	-0.0116	0.7328**
4	Number of effective tillers per plant	-0.1119	0.0595	-0.2221	-0.1142	0.343	0.1145	-0.2101	0.078	-0.0219	0.0085	0.0181	-0.0586
5	Length of main spike	-0.1894	-0.0016	0.8781	0.0504	-0.7761	-0.268	0.5372	-0.2546	0.0614	0.4713	0.0025	0.5112**
6	Number of spikelets per spike	-0.0828	0.0108	0.7478	0.0361	-0.5738	-0.3624	0.53	-0.203	0.0338	0.369	-0.0029	0.5024**
7	Number of grains per spike	-0.1468	-0.0007	0.6821	0.0348	-0.6042	-0.2784	0.6899	-0.2606	0.026	0.428	0.0268	0.597**
8	Grain weight per spike	-0.1296	-0.0204	0.799	0.0344	-0.7646	-0.2846	0.6956	-0.2585	0.0693	0.4792	0.0103	0.6302**
9	1000-grain weight	-0.0151	0.0202	0.2548	0.0166	-0.3155	-0.081	0.1189	-0.1186	0.1511	0.1524	-0.0159	0.1679
10	Biological yield per plant	-0.2254	0.0043	0.6206	-0.0013	-0.4875	-0.1782	0.3936	-0.1651	0.0307	0.7503	0.0638	0.8056**
11	Protein content	0.1834	-0.0211	0.0753	0.0112	0.0107	-0.0057	-0.1006	0.0144	0.013	-0.2597	-0.1842	-0.2632

R= 0.1037

4. CONCLUSION

Based on the aforementioned findings, it may be concluded that Plant height, number of grains per spike, biological yield per plant exerted positive direct effect along with positive and significant correlation on grain yield per plant. These characters must be given preference in selection while selecting the superior genotypes. Therefore, these particular traits should be considered for enhancing yields in the wheat breeding program.

REFERENCES

- Anonymous, (2023). Director's Report of AICRP on Wheat and Barley 2022-23, Ed: Gyanendra Singh. ICAR-Indian P 90
- Liu, K. (2011). Comparison of lipid content and fatty acid composition and their distribution within seeds of 5 small grain species. *J. Food Sci.*, **76** (2): 334-342.
- Regvar, M., Eichert, D., Kaulich, B., Gianoncelli, A., Pongrac, P., Vogel-Mikus, K. and Kreft, I. (2011). New insights into globoids of protein storage vacuoles in wheat aleurone using synchrotron soft X-ray microscopy. *J. Exp. Bot.*, **62**(11): 3929-3939..s
- Lafiandra, D., Riccardi, G., Shewry, P. R. (2014). Improving cereal grain carbohydrates for diet and health. *Journal of Cereal Science*, **59**: 312- 326.
- Gillies, S.A., Futardo, A. and Henry, R.J. (2012). Gene expression in the developing aleurone and starchy endosperm of wheat. *Plant Biotech J.*, **10**(6): 668-679.
- Belderok, B., Mesdag, J., Mesdag, H. and Donner, D.A., (2000). Bread-making quality of wheat: a century of breeding in Europe. *Springer Science & Business Media*. pp 15-20
- FAO (2018) World food and agriculture statistical pocketbook (2018), Food and Agriculture Organization of the United Nations, Rome.
- Poudel PB & Poudel MR (2020) Heat Stress Effects and Tolerance in Wheat: A Review. *Journal of Biology and Today's World*, **9**(4): 217.
- Dewey D.R. and Lu K.H. (1959). A correlation and path coefficient analysis of components of crested wheat grass seed production. *Agronomy Journal*, **51**(9):515-518.
- Panse, V. G., and Sukhatme, P. V. (1985). Statistical Methods for Agricultural Workers, ICAR New Delhi.
- Al-Jibouri, H. A., Miller, P.A. and Robinson, H.F. (1958). Genotypic and environmental variances in upland cotton cross of interspecific origin. *Agronomy Journal*, **50**: 633-635.
- Lindner, R. C. (1944). Rapid analytical methods for some of the more common inorganic constituents of plant tissues. *Plant Physiology*, **19**(1), 76.
- Nagar, S. S., Kumar, P., Vishwakarma, S. R., Singh, G. and Tyagi, B. S. (2018). Assessment of genetic variability and character association for grain yield and its component traits in bread wheat (*Triticum aestivum* L.). *Journal of Applied and Natural Science*, **10**(2): 797-804.
- Nukasani, V., Potdukhe, N. R., Bharad, S., Deshmukh, S., & Shinde, S. M. (2013). Genetic variability, correlation and path analysis in wheat. *Journal of Cereal Research*, **5**(2).

- Kumar, R., Bhushan, B., Pal, R. and Gaurav, S. S. (2014). Correlation and path coefficient analysis for quantitative traits in wheat (*Triticumaestivum* L.) under Normal Condition. *Annals of Agri-BioResearch*, **19**(3): 447-450.
- Ayer, D.K., Sharma, A., Ojha, B. R., Paudel, A. and Dhakal, K. (2017). Correlation and path coefficient analysis in advanced wheat genotypes. *SAARC J. Agri.*, **15**(1): 1-12.
- Abd El-Mohsen, A. A., Abo Hegazy, S. R., & Taha, M. H. (2012). Genotypic and phenotypic interrelationships among yield and yield components in Egyptian bread wheat genotypes. *J. Plant Breed. Crop Sci*, **4**(1), 9-1

UNDER PEER REVIEW

UNDER PEER REVIEW