

Original Research Article **Modelling Count Variables: A Comparative Analysis of two Discretization Techniques**

ABSTRACT

Background: Different discretization methods have been proposed to provide a better fit to count observations with characteristics resembling a given continuous distribution. This study compares discretization through survival function and mixed Poisson process.

Methodology: The Ailamujia distribution is extended using the cubic rank transmutation map. The shapes of the continuous distribution are assessed, and two discretized versions of the distribution are unimodal and skewed, depicting characteristics of the continuous distribution. Parameters of the distributions are estimated using the method of maximum likelihood, and both AIC and chi-square are used for model comparison.

Results: Real-life assessment using five count data shows that the two propositions provide a better fit than the three competing distributions considered. Also, discretization through the mixed Poisson process offers a better fit than the survival function technique.

Conclusion: Various moment-based mathematical properties of the discretization through the mixed Poisson process are easily obtainable and hence, can be easily characterized.

Keywords: Discretization, survival function, mixed Poisson distribution, Ailamujia distribution.

1. INTRODUCTION

Some real lifetime data are discrete in observation even when they are primarily continuous in the real sense [1]. The discretization procedure was developed to improve efficiency in modelling count observations with shapes similar to a specific continuous distribution. The process involves using different mathematical concepts to derive discrete analogous to continuous distributions. Different approaches to discretizing a continuous distribution have been developed [2–4]. Among the prominent techniques for achieving this is the survival function of the continuous distribution, as was first used on the Weibull distribution [5–6]. A detailed survey of recent introductions in the discretization process has been reported [7].

If a continuous random variable has its CDF (distribution function) given as G_x and S_x is its survival function indexed with parameter vector θ , the PMF (probability mass function) of a new discrete random variable P_x is obtained [8–9] as:

$$P_x = S_x - S_{x+1}$$

where

$$S_x = 1 - G_x$$

An advantage of this technique is that the survival function for discretized count distributions has a functional form resembling its corresponding continuous distributions [10]. Leveraging on these advantages, many notable continuous distributions have been discretized. Among these are the discrete Weibull [5, 11, 12]; discrete Rayleigh [9]; the discrete Lindley distribution [13]; the discrete Lomax distribution [14]; the discrete generalized exponential distribution [15]; discrete Marshall-Olkin Weibull [16]; discrete normal [8]; and discrete Pareto and discrete Burr [17].

Another technique of obtaining new discrete distribution involves utilizing the mixed Poisson [18] concept. The process involves assuming a continuous distribution with positive supports for the Poisson parameter. In most cases, the newly obtained discrete distribution's shape mimics the continuous distribution assumed for the parameter. Other notable characteristics of this distribution are presented in [19–21]. The procedure has received patronage in modelling datasets from actuary science in particular and dispersed observations in general [22–27]. Among many of the obtained discrete distributions in this paradigm include the mixed Poisson Lindley [28] and its generalizations [24, 29]. Another very popular application is the mixed Poisson-gamma distribution [18] which turned out to be a form of the negative binomial distribution with $p = \left(\frac{1}{1+\beta}\right)$. Different extensions of this distribution pervade literature [30, 31] with applications in diverse fields of studies [32–34].

Suppose discrete random variable N has the Poisson distribution with parameter X . Also, if X is assumed to follow a continuous random distribution with positive supports $(0, \infty)$ with PDF denoted with g_x , a new discrete distribution is obtained in the mixed Poisson architecture by solving for the unconditional distribution for N in:

$$P_n = \int_0^{\infty} \frac{x^n e^{-x}}{n!} \cdot g_x dx$$

Different distributions have been proposed for the choice of g_x [35]. The shape of g_x has a resemblance with the shape of the obtained discrete distribution from the process [36, 37].

In this study, a new continuous distribution is obtained using the cubic rank transmutation map [38] to extend the Ailamujia distribution [39]. Both the survival function [9] and the mixed Poisson [18] approaches of discretization are compared using the obtained continuous distribution.

2. AILAMUJIA DISTRIBUTION

The Ailamujia distribution [39] has been used to model lifetime observations that are skewed and unimodal [40]. Several authors [41–44] have obtained an improved version of the distribution using different compounding techniques. The distribution function for the Ailamujia distribution is defined as:

$$F_x = 1 - (1 + \beta x)e^{-\beta x}, \quad \beta > 0 \quad (1)$$

Since introducing the quadratic transmutation map [45], many cubic transmutation maps that extend any baseline distribution pervade literature. The distribution function of the cubic rank transmutation (CRT) map of [38] is given as:

$$G_x = cF_x + (k - c)F_x^2 + (1 - k)F_x^3, \quad c \in [0,1], \quad k \in [-1,1], \quad (2)$$

2.1 Cubic Rank Transmuted Ailamujia Distribution

Some baseline distributions that have been extended using (2) include the inverse Rayleigh distribution [46], Gumbel distribution [47, 48], modified Burr III Pareto distribution [49], inverseWeibul distribution [50], Gompertz and Frechet distributions [48].

Inserting (1) into (2) gives the cubic rank transmuted Ailamujia(CRTA) distribution with CDF, PDF, and survival function respectively obtained as:

$$G_x = c(1 - (1 + \beta x)e^{-\beta x}) + (k - c)(1 - (1 + \beta x)e^{-\beta x})^2 + (1 - k)(1 - (1 + \beta x)e^{-\beta x})^3 \quad (3)$$

$$g_x = \beta^2 x e^{-\beta x} (3 - c - k + 2(c + 2k - 3)(1 + \beta x)e^{-\beta x} - 3(k - 1)(1 + \beta x)^2(e^{-2\beta x})) \quad (4)$$

$$S_x = 1 - c(1 - (1 + \beta x)e^{-\beta x}) - (k - c)(1 - (1 + \beta x)e^{-\beta x})^2 - (1 - k)(1 - (1 + \beta x)e^{-\beta x})^3 \quad (5)$$

Figure 1 shows different shapes of the PDF for the CRTA distribution for different parameter combinations. The figure reveals that the distribution is unimodal with positive skewness.




Figure 1. Shapes of PDF for CRTA distribution

2.2 Moments of the CRTA Distribution

Proposition 1. If a random variable X has a CRTA distribution, then the r^{th} moment is obtained as:

$$E(x^r) = (3 - c - k)(r + 1)r! + \frac{2(c + 2k - 3)(r + 1)r!}{4(2\beta)^r} \left(1 + \frac{\beta(r + 2)}{2\beta}\right) - \frac{3(k - 1)(r + 1)r!}{9(3\beta)^r} \left(1 + \frac{2\beta(r + 2)}{3\beta} + \frac{(r + 3)(r + 2)}{(3\beta)^2}\right) \quad (6)$$

Proof:

$$\begin{aligned} E(x^r) &= \int_0^{\infty} x^r g_x dx \\ &= \int_0^{\infty} x^r \left(\beta^2 x e^{-\beta x} (3 - c - k + 2(c + 2k - 3)(1 + \beta x)e^{-\beta x} - 3(k - 1)(1 + \beta x)^2(e^{-2\beta x})) \right) dx \\ &= \int_0^{\infty} \beta^2 x^{r+1} e^{-\beta x} (3 - c - k + 2(c + 2k - 3)(1 + \beta x)e^{-\beta x} - 3(k - 1)(1 + \beta x)^2(e^{-2\beta x})) dx \\ &= \beta^2 \int_0^{\infty} (3 - c - k)x^{r+1} e^{-\beta x} + 2(c + 2k - 3)(x^{r+1} e^{-2\beta x} + \beta x^{r+2} e^{-2\beta x}) \\ &\quad - 3(k - 1)(x^{r+1} e^{-3\beta x} + 2\beta x^{r+2} e^{-3\beta x} + x^{r+3} e^{-3\beta x}) dx \\ &= \beta^2 \left[\int_0^{\infty} (3 - c - k)x^{r+1} e^{-\beta x} dx + \int_0^{\infty} 2(c + 2k - 3)(x^{r+1} e^{-2\beta x} + \beta x^{r+2} e^{-2\beta x}) dx \right. \\ &\quad \left. - \int_0^{\infty} 3(k - 1)(x^{r+1} e^{-3\beta x} + 2\beta x^{r+2} e^{-3\beta x} + x^{r+3} e^{-3\beta x}) dx \right] \\ &= \beta^2 \left[(3 - c - k) \frac{(r + 1)r!}{\beta^2} + \frac{2(c + 2k - 3)(r + 1)r!}{(2\beta)^{r+2}} \left(1 + \frac{\beta(r + 2)}{2\beta}\right) \right. \\ &\quad \left. - \frac{3(k - 1)(r + 1)r!}{(3\beta)^{r+2}} \left(1 + \frac{2\beta(r + 2)}{3\beta} + \frac{(r + 3)(r + 2)}{(3\beta)^2}\right) \right] \\ &= (3 - c - k)(r + 1)r! + \frac{2(c + 2k - 3)(r + 1)r!}{4(2\beta)^r} \left(1 + \frac{\beta(r + 2)}{2\beta}\right) \\ &\quad - \frac{3(k - 1)(r + 1)r!}{9(3\beta)^r} \left(1 + \frac{2\beta(r + 2)}{3\beta} + \frac{(r + 3)(r + 2)}{(3\beta)^2}\right) \end{aligned}$$

Hence, the first four moments of the CRTA distribution are obtained as:

$$m_1 = \frac{32(1-k) + 9\beta^2(15c + 22k - 37) + 216\beta^3(3-c-k)}{108\beta^3} \quad (7)$$

$$m_2 = \frac{160(1-k) + 3\beta^2(243c + 398k - 641) + 1944\beta^3(3-c-k)}{324\beta^4} \quad (8)$$

$$m_3 = \frac{320(1-k) + \beta^2(1701c + 2986k - 4687) + 7776\beta^5(3-c-k)}{324\beta^5} \quad (9)$$

$$m_4 = \frac{560(1-k) + 15\beta^2(243c + 446k - 689) + 29160\beta^6(3-c-k)}{243\beta^6} \quad (10)$$

3.0 DISCRETIZED TRANSMUTED AILAMUJIA DISTRIBUTION

Proposition 2: With the distribution function of the CRTA distribution obtained in (3) and the corresponding survival function obtained in (5), the discretized CRTA distribution (DCTA) is obtained as:

$$P_x = S_x - S_{x+1}, \quad x = 0, 1, 2, \dots$$

Hence,

$$P_x = c \left((1 + \beta x)e^{-\beta x} - 1 \right) - (k - c) \left(1 - (1 + \beta x)e^{-\beta x} \right)^2 - (1 - k) \left(1 - (1 + \beta x)e^{-\beta x} \right)^3 + c \left(1 - (1 + \beta(x+1))e^{-\beta(x+1)} \right) + (k - c) \left(1 - (1 + \beta(x+1))e^{-\beta(x+1)} \right)^2 + (1 - k) \left(1 - (1 + \beta(x+1))e^{-\beta(x+1)} \right)^3, \quad x = 0, 1, 2, \dots \quad (11)$$

Special cases:

i. When $k = 1$, equation (11) becomes the DCTA I:

$$P_x = c \left((1 + \beta x)e^{-\beta x} - 1 \right) - (1 - c) \left(1 - (1 + \beta x)e^{-\beta x} \right)^2 + c \left(1 - (1 + \beta(x+1))e^{-\beta(x+1)} \right) + (1 - c) \left(1 - (1 + \beta(x+1))e^{-\beta(x+1)} \right)^2, \quad x = 0, 1, 2, \dots \quad (12)$$

ii. When $k = c$, equation (11) becomes the DCTA II

$$P_x = c \left((1 + \beta x)e^{-\beta x} - 1 \right) - (1 - c) \left(1 - (1 + \beta x)e^{-\beta x} \right)^3 + c \left(1 - (1 + \beta(x+1))e^{-\beta(x+1)} \right) + (1 - c) \left(1 - (1 + \beta(x+1))e^{-\beta(x+1)} \right)^3, \quad x = 0, 1, 2, \dots \quad (13)$$

iii. When $c = 0$, equation (11) becomes the DCTA III:

$$P_x = -k \left(1 - (1 + \beta x)e^{-\beta x} \right)^2 - (1 - k) \left(1 - (1 + \beta x)e^{-\beta x} \right)^3 + k \left(1 - (1 + \beta(x+1))e^{-\beta(x+1)} \right)^2 + (1 - k) \left(1 - (1 + \beta(x+1))e^{-\beta(x+1)} \right)^3, \quad x = 0, 1, 2, \dots \quad (14)$$




Figure 2. Shapes of PMF for DCTA distribution

The PMF of the DCTA distribution for different combinations of parameters show positive skewness, unimodality and resembles the shapes of the PDF of the CRTA distribution in figure 1.

If S_x is the survival function of the CRTA distribution, the distribution function (CDF) and the survival function for the DCTA distribution [10, 16] are obtained from:

$$F(x) = 1 - S_x + P_x$$

$$S(x) = 1 - F(x) + P_x$$

Hence, the CDF and survival functions are obtained as:

$$F(x) = c(1 - (1 + \beta(x+1))e^{-\beta(x+1)}) + (k-c)(1 - (1 + \beta(x+1))e^{-\beta(x+1)})^2 + (1-k)(1 - (1 + \beta(x+1))e^{-\beta(x+1)})^3, \quad x = 0,1,2, \dots \quad (15)$$

$$S(x) = 1 - c(1 - (1 + \beta x)e^{-\beta x}) - (k-c)(1 - (1 + \beta x)e^{-\beta x})^2 - (1-k)(1 - (1 + \beta x)e^{-\beta x})^3, \quad x = 0,1,2, \dots \quad (16)$$

3.1 Moments of the DCTA Distribution

Proposition 3. If a random variable X has a CRTA distribution, then the r^{th} moment of the DCTA distribution is obtained as:

$$E(x^r) = \sum_{x=0}^{\infty} x^r \left[c \left((1 + \beta x)e^{-\beta x} - 1 \right) - (k-c)(1 - (1 + \beta x)e^{-\beta x})^2 - (1-k)(1 - (1 + \beta x)e^{-\beta x})^3 + c(1 - (1 + \beta(x+1))e^{-\beta(x+1)}) + (k-c)(1 - (1 + \beta(x+1))e^{-\beta(x+1)})^2 + (1-k)(1 - (1 + \beta(x+1))e^{-\beta(x+1)})^3 \right] \quad (15)$$

Proof:

$$\begin{aligned} E(x^r) &= \mu_r' = \sum_{x=0}^{\infty} x^r P_x \\ &= \sum_{x=0}^{\infty} x^r \left[c \left((1 + \beta x)e^{-\beta x} - 1 \right) - (k-c)(1 - (1 + \beta x)e^{-\beta x})^2 - (1-k)(1 - (1 + \beta x)e^{-\beta x})^3 + c(1 - (1 + \beta(x+1))e^{-\beta(x+1)}) + (k-c)(1 - (1 + \beta(x+1))e^{-\beta(x+1)})^2 + (1-k)(1 - (1 + \beta(x+1))e^{-\beta(x+1)})^3 \right], \quad r = 1,2, \dots \end{aligned}$$

In particular, the mean of the distribution is obtained from:

$$\begin{aligned} \mu'_1 = & \left[c \sum_{x=0}^{\infty} x \left((1 + \beta x) e^{-\beta x} - 1 \right) - (k - c) \sum_{x=0}^{\infty} x \left(1 - (1 + \beta x) e^{-\beta x} \right)^2 - (1 - k) \sum_{x=0}^{\infty} x \left(1 - (1 + \beta x) e^{-\beta x} \right)^3 \right. \\ & + c \sum_{x=0}^{\infty} x \left(1 - (1 + \beta(x + 1)) e^{-\beta(x + 1)} \right) + (k - c) \sum_{x=0}^{\infty} x \left(1 - (1 + \beta(x + 1)) e^{-\beta(x + 1)} \right)^2 \\ & \left. + (1 - k) \sum_{x=0}^{\infty} x \left(1 - (1 + \beta(x + 1)) e^{-\beta(x + 1)} \right)^3 \right] \end{aligned}$$

In general, there is no close form for the moments of the DCTA distribution.

3.2 MLE of the DCTA distribution

Proposition 4: Given that x_1, x_2, \dots, x_n are random samples of size n drawn from the DCTA distribution, the log-likelihood function for the distribution is obtained as

$$\begin{aligned} \mathcal{L} = \prod_{i=1}^n P_{x_i} &= \prod_{i=1}^n \left(c \left((1 + \beta x) e^{-\beta x} - 1 \right) - (k - c) \left(1 - (1 + \beta x) e^{-\beta x} \right)^2 - (1 - k) \left(1 - (1 + \beta x) e^{-\beta x} \right)^3 \right. \\ & \quad \left. + c \left(1 - (1 + \beta(x + 1)) e^{-\beta(x + 1)} \right) + (k - c) \left(1 - (1 + \beta(x + 1)) e^{-\beta(x + 1)} \right)^2 \right. \\ & \quad \left. + (1 - k) \left(1 - (1 + \beta(x + 1)) e^{-\beta(x + 1)} \right)^3 \right) \\ \ell = \log \mathcal{L} &= \sum_{i=1}^n \log \left(c \left((1 + \beta x) e^{-\beta x} - 1 \right) - (k - c) \left(1 - (1 + \beta x) e^{-\beta x} \right)^2 - (1 - k) \left(1 - (1 + \beta x) e^{-\beta x} \right)^3 \right. \\ & \quad \left. + c \left(1 - (1 + \beta(x + 1)) e^{-\beta(x + 1)} \right) + (k - c) \left(1 - (1 + \beta(x + 1)) e^{-\beta(x + 1)} \right)^2 \right. \\ & \quad \left. + (1 - k) \left(1 - (1 + \beta(x + 1)) e^{-\beta(x + 1)} \right)^3 \right) \end{aligned}$$

Estimating the parameters (c, k, β) denoted with $(\hat{c}, \hat{k}, \hat{\beta})$ involves solving a system of non-linear equations. In this study, the *optimr* package [51] in the *R-language*[52] is used to obtain the estimates. A similar approach of parameter estimation was used in similar propositions [16].

4.0 MIXED POISSON CRTA DISTRIBUTION

Proposition 5. If $N \sim \text{Poisson}(X)$, where the PDF of X is given in equation (4), then the probability mass function (PMF) of the mixed Poisson CRTA distribution (MCTA) is obtained as:

$$\begin{aligned} P_n = \beta^2 & \left[\frac{(3 - c - k)(n + 1)}{(1 + \beta)^{n+2}} + \frac{2(c + 2k - 3)(n + 1)(1 + 4\beta + n\beta)}{(1 + 2\beta)^{n+3}} \right. \\ & \quad \left. - \frac{3(k - 1)(n + 1)}{(1 + 3\beta)^{n+4}} \left((1 + 3\beta)^2 + 2\beta(1 + 3\beta)(n + 2) + \beta^2(n + 3)(n + 2) \right) \right] \end{aligned} \quad (16)$$

Proof:

$$\begin{aligned} P_n &= \int_0^{\infty} \frac{x^n e^{-x}}{n!} \cdot g_x dx \\ &= \int_0^{\infty} \frac{x^n e^{-x}}{n!} \beta^2 x e^{-\beta x} \left(3 - c - k + 2(c + 2k - 3)(1 + \beta x) e^{-\beta x} - 3(k - 1)(1 + \beta x)^2 (e^{-2\beta x}) \right) dx \\ &= \frac{\beta^2}{n!} \left[(3 - c - k) \int_0^{\infty} x^{n+1} e^{-(1+\beta)x} dx + 2(c + 2k - 3) \int_0^{\infty} x^{n+1} e^{-(1+2\beta)x} (1 + \beta x) dx - 3(k \right. \\ & \quad \left. - 1) \int_0^{\infty} x^{n+1} e^{-(1+3\beta)x} (1 + 2\beta x + \beta^2 x^2) dx \right] \\ &= \frac{\beta^2}{n!} \left[\frac{(3 - c - k)(n + 1)n!}{(1 + \beta)^{n+2}} + \frac{2(c + 2k - 3)(n + 1)n!}{(1 + 2\beta)^{n+2}} \left(1 + \frac{\beta(n + 2)}{1 + 2\beta} \right) \right. \\ & \quad \left. - \frac{3(k - 1)(n + 1)n!}{(1 + 3\beta)^{n+2}} \left(1 + \frac{2\beta(n + 2)}{1 + 3\beta} + \frac{\beta^2(n + 3)(n + 2)}{(1 + 3\beta)^2} \right) \right] \end{aligned}$$

$$= \beta^2 \left[\frac{(3-c-k)(n+1)}{(1+\beta)^{n+2}} + \frac{2(c+2k-3)(n+1)(1+4\beta+n\beta)}{(1+2\beta)^{n+3}} - \frac{3(k-1)(n+1)}{(1+3\beta)^{n+4}} ((1+3\beta)^2 + 2\beta(1+3\beta)(n+2) + \beta^2(n+3)(n+2)) \right]$$

Special cases:

i. When $k = 1$, equation (16) becomes the MCTA I:

$$P_n = \beta^2 \left[\frac{(2-c)(n+1)}{(1+\beta)^{n+2}} + \frac{2(c-1)(n+1)(1+4\beta+n\beta)}{(1+2\beta)^{n+3}} \right], \quad x = 0,1,2, \dots \quad (17)$$

ii. When $k = c$, equation (16) becomes the MCTA II:

$$P_n = \beta^2 \left[\frac{(3-2c)(n+1)}{(1+\beta)^{n+2}} + \frac{6(c-1)(n+1)(1+4\beta+n\beta)}{(1+2\beta)^{n+3}} - \frac{3(c-1)(n+1)}{(1+3\beta)^{n+4}} ((1+3\beta)^2 + 2\beta(1+3\beta)(n+2) + \beta^2(n+3)(n+2)) \right], \quad (18)$$

$x = 0,1,2, \dots$

iii. When $c = 0$, equation (11) becomes the MCTA III:

$$P_n = \beta^2 \left[\frac{(3-c-k)(n+1)}{(1+\beta)^{n+2}} + \frac{2(c+2k-3)(n+1)(1+4\beta+n\beta)}{(1+2\beta)^{n+3}} - \frac{3(k-1)(n+1)}{(1+3\beta)^{n+4}} ((1+3\beta)^2 + 2\beta(1+3\beta)(n+2) + \beta^2(n+3)(n+2)) \right], \quad (19)$$

$x = 0,1,2, \dots$




Figure 3. Shapes of PMF for the MCTA distribution

Figure 3 shows that the shapes of the MCTA distribution resemble the shapes of the PDF of the CRTA distribution. The shapes suggest that the distribution can model unimodal and positively skewed count observations.

4.1 Moment-Generating Function of the MCTA Distribution

Proposition 6. Given that g_n is the mixing distribution of a random variable N with the CRTA distribution, the probability generating function (PGF) of MCTA distribution is defined as:

$$P_n(z) = \int_0^{\infty} e^{n(z-1)} g_n dn$$

$$= \int_0^{\infty} e^{n(z-1)} \beta^2 n e^{-\beta n} \left(3-c-k + 2(c+2k-3)(1+\beta n) e^{-\beta n} - 3(k-1)(1+\beta n)^2 (e^{-2\beta n}) \right) dn$$

$$\begin{aligned}
&= \beta^2 \left[(3-c-k) \int_0^{\infty} ne^{-(1+\beta-z)n} dn + 2(c+2k-3) \int_0^{\infty} ne^{-(1+2\beta-z)n} (1+\beta n) dn - 3(k-1) \int_0^{\infty} ne^{-(1+3\beta-z)n} (1+\beta n)^2 dn \right] \\
&= \beta^2 \left[\frac{(3-c-k)}{(1+\beta-z)^2} + 2(c+2k-3) \int_0^{\infty} (ne^{-(1+2\beta-z)n} + \beta n^2 e^{-(1+2\beta-z)n}) dn - 3(k-1) \int_0^{\infty} ne^{-(1+3\beta-z)n} (1+2\beta n + \beta^2 n^2) dn \right] \\
&= \beta^2 \left[\frac{(3-c-k)}{(1+\beta-z)^2} + 2(c+2k-3) \left(\frac{1}{(1+2\beta-z)^2} + \frac{2}{(1+2\beta-z)^3} \right) - 3(k-1) \left(\frac{1}{(1+3\beta-z)^2} + \frac{4\beta}{(1+3\beta-z)^3} + \frac{6\beta^2}{(1+3\beta-z)^4} \right) \right]
\end{aligned}$$

Hence the PGF of the MCTA distribution is:

$$\begin{aligned}
P_x(z) = \beta^2 \left[\frac{(3-c-k)}{(1+\beta-z)^2} + 2(c+2k-3) \left(\frac{1}{(1+2\beta-z)^2} + \frac{2}{(1+2\beta-z)^3} \right) - 3(k-1) \left(\frac{1}{(1+3\beta-z)^2} + \frac{4\beta}{(1+3\beta-z)^3} + \frac{6\beta^2}{(1+3\beta-z)^4} \right) \right] \quad (20)
\end{aligned}$$

Also, the moment generating function for the PMF in (16) is obtained by replacing z with e^t in (20). This is given as:

$$\begin{aligned}
M_x(t) = \beta^2 \left[\frac{(3-c-k)}{(1+\beta-e^t)^2} + 2(c+2k-3) \left(\frac{1}{(1+2\beta-e^t)^2} + \frac{2}{(1+2\beta-e^t)^3} \right) - 3(k-1) \left(\frac{1}{(1+3\beta-e^t)^2} + \frac{4\beta}{(1+3\beta-e^t)^3} + \frac{6\beta^2}{(1+3\beta-e^t)^4} \right) \right] \quad (21)
\end{aligned}$$

From (21), the first four raw moments for the MCTA distribution are obtained as:

$$\begin{aligned}
m_1 &= \frac{2\beta(295-81c-106k) + 81(c+2k-3)}{108\beta^2} \\
m_2 &= \frac{3\beta^2(295-81c-106k) + \beta(2399-729c-698k) + 243(c+2k-3)}{162\beta^3} \\
m_3 &= \frac{6\beta^3(295-81c-106k) + 12\beta^2(1321-405c-430k) + 8\beta(2279-729c-578k) + 1215(c+2k-3)}{324\beta^4} \\
m_4 &= \frac{18\beta^4(295-81c-106k) + 24\beta^3(4745-1458c-1586k) + 18\beta^2(20905-6723c-6406k) + 5\beta(55603-18225c-14050k) + 1215(c+2k-3)}{972\beta^5}
\end{aligned}$$

Hence, the variance for the distribution can be obtained from:

$$Var = m_2 - (m_1)^2$$

The index of dispersion, skewness and kurtosis for the MCTA distribution can be obtained using the moment-based relationships [53] respectively as:

$$DI = \frac{Var}{m_1}$$

$$S_k = \frac{m_3 - 3m_2m_1 + 2(m_1)^3}{(Var)^{\frac{3}{2}}}$$

$$Ku = \frac{m_4 - 4m_3m_1 + 6m_2(m_1)^2 - 3(m_1)^4}{(Var)^2}$$

4.2 MLE of the MCTA distribution

Proposition 7: Given that n_1, n_2, \dots, n_k are random samples of size k drawn from the MCTA distribution, the log-likelihood function for the distribution is obtained as

$$\mathcal{L} = \prod_{i=1}^k P_{n_i} = \prod_{i=1}^k \left(\beta^2 \left[\frac{(3-c-k)(n+1)}{(1+\beta)^{n+2}} + \frac{2(c+2k-3)(n+1)(1+4\beta+n\beta)}{(1+2\beta)^{n+3}} \right. \right. \\ \left. \left. - \frac{3(k-1)(n+1)}{(1+3\beta)^{n+4}} ((1+3\beta)^2 + 2\beta(1+3\beta)(n+2) + \beta^2(n+3)(n+2)) \right] \right) \\ \ell = \log \mathcal{L} = \sum_{i=1}^k \log \left(\beta^2 \left[\frac{(3-c-k)(n+1)}{(1+\beta)^{n+2}} + \frac{2(c+2k-3)(n+1)(1+4\beta+n\beta)}{(1+2\beta)^{n+3}} \right. \right. \\ \left. \left. - \frac{3(k-1)(n+1)}{(1+3\beta)^{n+4}} ((1+3\beta)^2 + 2\beta(1+3\beta)(n+2) + \beta^2(n+3)(n+2)) \right] \right)$$

The parameter estimates of $(\hat{\beta}, \hat{c}, \hat{k})$ is obtained using the *optimr* package [51] in the *R-language* [52].

5.0 APPLICATIONS

The proposed distributions in this study are compared with (EDW) the exponentiated discrete Weibull distribution [17], (DMOG) the discrete Marshall-Olkin generalized exponential distribution [54], and (DBX) the discrete Bur XII distribution [55].

Table 1. PMF of the competing distribution

Distribution	PMF
EDW	$P_x = (1 - \beta^{(x+1)^k})^c - (1 - \beta^{x^k})^c$
DMOG	$P_x = \frac{k(1 - (1 - \beta^x)^c)}{k + (1 - k)(1 - \beta^x)^c} - \frac{k(1 - (1 - \beta^{(x+1)})^c)}{k + (1 - k)(1 - \beta^{(x+1)})^c}$
DBX	$P_x = \beta^{\log(1+x^c)} - \beta^{\log(1+(x+1)^c)}$

Five real-life datasets are utilized to compare the new propositions and other competing distributions. The first dataset represents yeast cell counts per square, while the second dataset is the counts of the European red mites on apple leaves. Both data have been previously used in new propositions involving count distributions [16, 26, 56]. The third dataset is the frequency of epileptic seizures previously used on other discrete distributions [25, 57]. The fourth dataset on the number of mistakes in copying groups of random digits has been used to model various count distributions [25, 26, 28, 58]. The last dataset represents the number of strikes in a UK coal mining industries from 1948-1959 [16, 59].

RESULTS AND DISCUSSION

The maximum likelihood estimation technique using various non-linear algorithms that come with the *optimr* package in the *R-language* is used to obtain the estimates. The results obtained are presented in Tables 2 – 6. The Akaike Information Criterion (AIC) and the chi-square goodness of fits are used for model comparisons.

Table 2. Results of data on yeast cell counts per square

X	Freq.	DCTA	DCTA I	DCTA II	DCTA III	MCTA	MCTA I	MCTA II	MCTA III	EDW	DMOG	DBX
0	128	127.9	120.6	118.7	114.7	126.8	124.1	123.7	121.8	143.0	127.3	127.8
1	37	37.3	51.8	54.2	62.3	40.3	45.7	46.1	48.9	26.5	41.0	42.5
2	18	17.3	11.5	11.4	8.7	14.5	12.9	12.9	12.8	10.1	13.0	10.2
3	3	3.7	2.5	2.2	1.1	4.2	3.3	3.2	2.8	4.2	3.9	3.4
4	1	0.6	0.5	0.4	0.1	1.0	0.8	0.8	0.6	1.8	1.2	1.4
β		2.06	1.79	1.93	2.33	5.73	3.82	4.21	5.14	0.56	0.30	0.19
c		2.80	1.44	1.15	2.05	4.14	1.32	1.05	1.84	0.47	0.74	1.72
k		-1.44				-5.54				1.44	1.58	
χ^2		0.32	9.05	11.72	25.32	1.44	3.69	3.90	5.81	9.23	2.14	4.36
AIC		343.04	349.48	351.68	365.78	344.39	344.74	344.85	346.34	344.62	345.82	349.61

Table 3. Results of data on the European red mites on Apple leaves

X	Freq.	DCTA	DCTA I	DCTA II	DCTA III	MCTA	MCTA I	MCTA II	MCTA III	EDW	DMOG	DBX
0	38	38.2	28.8	27.5	22.1	38.2	32.7	32.3	30.1	44.3	37.2	38.1
1	17	16.6	28.0	28.5	36.9	16.3	23.4	23.5	25.5	13.2	19.1	22.0
2	10	10.9	13.4	14.2	14.7	11.5	12.6	12.9	13.9	7.6	11.1	8.5
3	9	7.7	5.7	6.0	4.5	7.4	6.2	6.3	6.3	4.8	6.1	4.0
4	3	3.9	2.4	2.4	1.3	3.8	2.9	2.9	2.6	3.2	3.2	2.2
5	2	1.7	1.0	0.9	0.4	1.7	1.3	1.3	1.0	2.1	1.6	1.3

6	1	0.7	0.4	0.3	0.1	0.7	0.6	0.5	0.4	1.5	0.8	0.9
β		1.13	1.06	1.16	1.46	2.38	1.54	1.70	2.08	0.78	0.50	0.39
c		2.36	1.35	1.07	1.87	4.34	1.31	1.04	1.81	0.49	0.61	1.79
k		-0.89				-6.01				1.48	2.16	
χ^2		0.72	12.71	14.69	37.52	0.93	5.16	5.52	9.26	4.63	1.45	5.01
AIC		240.89	249.73	251.51	273.43	241.09	243.34	243.55	246.78	241.96	242.77	250.21

Table 4. Results of data on the number of epileptic seizures

X	Freq.	DCTA	DCTA I	DCTA II	DCTA III	MCTA	MCTA I	MCTA II	MCTA III	EDW	DMOG	DBX
0	126	124.9	94.5	89.9	61.1	125.6	113.3	111.8	99.7	151.4	121.0	128.9
1	80	81.9	114.9	114.9	148.3	81.0	97.1	97.4	105.1	64.4	93.1	109.0
2	59	56.6	69.7	72.9	86.0	58.8	62.7	63.7	70.8	40.8	58.3	45.3
3	42	42.0	36.4	38.5	35.1	39.7	36.3	37.0	39.2	27.5	34.3	21.9
4	24	24.3	18.2	18.8	13.2	23.3	19.9	20.1	19.6	19.1	19.6	12.3
5	8	12.0	9.0	8.8	4.8	12.2	10.5	10.5	9.2	13.5	11.0	7.6
6	5	5.4	4.4	4.0	1.7	5.8	5.5	5.4	4.2	9.6	6.1	5.1
7	4	2.3	2.1	1.8	0.6	2.6	2.8	2.7	1.9	6.9	3.4	3.6
8	3	1.0	1.0	0.8	0.2	1.1	1.4	1.3	0.8	4.9	1.9	2.7
β		1.01	0.87	0.95	1.20	1.78	1.20	1.30	1.54	0.80	0.85	0.52
c		1.85	1.32	1.05	1.82	2.42	1.20	1.00	1.83	0.65	1.20	2.19
k		-0.48				-2.92				1.44	1.17	
χ^2		0.16	27.16	30.21	110.00	0.15	6.64	6.98	16.71	19.81	4.57	29.11
AIC		1192.43	1215.20	1219.95	1317.17	1191.94	1195.45	1195.73	1206.71	1192.83	1196.78	1248.64

Table 5. Results of data on the number of mistakes in copying groups of random digits

X	Freq.	DCTA	DCTA I	DCTA II	DCTA III	MCTA	MCTA I	MCTA II	MCTA III	EDW	DMOG	DBX
0	35	35.0	29.1	28.1	25.4	34.9	31.2	31.0	29.8	38.9	34.5	34.9
1	11	10.9	20.4	21.2	26.2	11.4	17.2	17.4	18.8	9.8	13.9	14.8
2	8	8.4	7.1	7.5	6.6	7.9	7.2	7.3	7.7	4.9	6.3	5.0
3	4	3.9	2.3	2.3	1.4	3.8	2.7	2.8	2.6	2.7	2.9	2.1
4	2	1.3	0.7	0.7	0.3	1.4	1.0	1.0	0.8	1.5	1.3	1.1
β		1.47	1.36	1.47	1.83	3.87	2.26	2.50	3.06	0.69	0.45	0.29
c		2.55	1.38	1.10	1.92	6.95	1.31	1.04	1.78	0.48	0.82	1.61
k		-1.16				-10.88				1.40	1.17	
χ^2		0.29	10.68	12.54	27.11	0.23	4.91	5.17	7.48	2.30	1.69	3.68
AIC		149.95	156.16	157.45	169.06	149.93	152.02	152.14	153.77	151.50	152.50	155.26

Table 6. Results of data on the number of strikes in UK coal mining industries

X	Freq.	DCTA	DCTA I	DCTA II	DCTA III	MCTA	MCTA I	MCTA II	MCTA III	EDW	DMOG	DBX
0	46	46.0	46.1	46.2	46.3	46.0	46.0	46.0	46.0	76.3	55.1	53.2
1	76	74.8	75.0	74.2	74.6	75.2	75.2	75.0	74.7	45.1	62.8	72.8
2	24	27.1	26.9	27.8	26.9	26.6	26.5	26.8	27.2	20.1	26.7	17.9
3	9	6.4	6.4	6.3	6.5	6.5	6.5	6.5	6.5	8.5	8.2	5.9
4	1	1.3	1.3	1.2	1.4	1.4	1.4	1.3	1.3	3.5	2.3	2.5
β		1.86	1.84	1.93	1.84	5.82	5.76	6.24	6.86	0.59	0.27	0.55
c		0.03	-0.03	0.27	0.96	-3.82	-3.97	-2.46	-7.79	1.34	2.38	3.86
k		0.84				0.64				1.66	1.66	
χ^2		1.27	1.20	1.52	1.22	1.08	1.08	1.12	1.22	39.72	6.14	6.18
AIC		380.96	378.93	379.14	378.94	377.70	378.73	378.77	378.80	381.58	386.09	390.01

The parameter estimates, goodness of fit statistics, observed frequencies and expected frequencies when each proposition and competing distributions are assumed are presented in Tables 2 – 6. For dataset I and II, presented in Tables 2 and 3, the discretized cubic rank transmuted Ailamujia distribution (DCTA) has the least chi-square and AIC while the mixed Poisson cubic rank transmuted distribution (MCTA) provides the second best fit for both datasets.

The MCTA gives the best fit for datasets III, IV, and V, as shown in Tables 4 – 6. The second best fit for the three datasets is obtained when the DCTA is assumed. The MCTA and the DCTA provide a better fit than the three considered competing distributions. In most cases, the two-parameter special cases of the MCTA provide a relatively better fit to the dataset when compared with the special cases of the DCTA.

CONCLUSION

This study introduces two discrete versions of the continuous cubic rank transmuted Ailamujia distribution. The first version is obtained using the survival function of the continuous distribution. For the second version, the parameter of the classical Poisson distribution is assumed to follow the cubic rank transmuted Ailamujia distribution in the mixed Poisson architecture.

Both proposed distributions are unimodal and positively skewed. Five real-life count datasets are used to assess the flexibility of the new propositions. Comparisons are made between the two discretization techniques and three other discrete distributions. Parameters of the distributions are estimated using the method of maximum likelihood, and both AIC and chi-square are used for model comparison.

In all cases, the two propositions provide a better fit than the three competing distributions considered. Also, from the five real-life applications, the discretization through the mixed Poisson process provides

a better fit than the survival function technique. Also, moment-based mathematical properties of the discretization through the mixed Poisson process are easily obtainable and hence, can be easily characterized.

REFERENCES

- [1] Yari G, Tondpour Z. Some New Discretization Methods with Application in Reliability. *Appl Appl Math*. 2018;13(2):664–76.
- [2] Ghosh T, Roy D, Chandra NM. Discretization of Random Variables with Applications. In: *Special 7-th Triennial Proceedings Volume of Calcutta Statistical Association Bulletin*. 2011. p. 249–52.
- [3] Drezner Z, Zerom D. A Simple and Effective Discretization of a Continuous Random Variable. 2015.
- [4] Tovissodé CF, Kakaï RG, Honfo SH, Doumatè JT. On the Discretization of Continuous Probability Distributions using a Probabilistic Rounding Mechanism. *Mathematics*. 2021;9(5):1–17.
- [5] Nakagawa T, Osaki S. The Discrete Weibull Distribution. *IEEE Trans Reliab*. 1975;24(5):300–1.
- [6] Al-Wakeel AA `S. On Discrete Weibull Distribution. *J Econ Adm Sci*. 2014;20(79):1–8.
- [7] Chakraborty S. Generating discrete analogues of continuous probability distributions-A survey of methods and constructions. *J Stat Distrib Appl [Internet]*. 2015;2(6):1–30. Available from: <http://dx.doi.org/10.1186/s40488-015-0028-6>
- [8] Roy D. The Discrete Normal Distribution. *Commun Stat Methods*. 2003;32(10):1871–83.
- [9] Roy D. Discrete Rayleigh Distribution. *IEEE Trans Reliab*. 2004;53(2):255–60.
- [10] EL-Helbawy AA, Hegazy MA, AL-Dayian GR, EL-Kader A. A Discrete Analog of the Inverted Kumaraswamy Distribution: Properties and Estimation with Application to COVID-19 Data. *Pakistan J Stat Oper Res*. 2022;18(1):297–328.
- [11] Khan MSA, Khalique A, Abouammoh AM. On Estimating Parameters in a Discrete Weibull Distribution. *IEEE Trans Reliab*. 1989;38(3):348–50.
- [12] Johnson NL, Kotz S, Kemp AW. *Univariate Discrete Distributions*. 3rd ed. John Wiley and Sons Inc. New Jersey: John Wiley and Sons Inc.; 2005.
- [13] Gómez-Déniz E, Calderín-Ojeda E. The Discrete Lindley Distribution: Properties and Applications. *J Stat Comput Simul*. 2011;81(11):1405–16.
- [14] Para BA, Jan TR. On Discrete Three Parameter Burr Type XII and Discrete Lomax Distributions and their Applications to Model Count Data from Medical Science. *BiomBiostat Int J*. 2016;4(2):1–15.
- [15] Nekoukhou V, Alamatsaz MH, Bidram H. Discrete Generalized Exponential Distribution of a Second Type. *Statistics (Ber)*. 2013;47(4):876–87.
- [16] Opone FC, Izekor EA, Akata IU, Osagiede FEU. A Discrete Analogue of the Continuous Marshall-Olkin Weibull Distribution with Application to Count Data. *Earthline J Math Sci*. 2021;5(2):415–28.

- [17] Krishna H, Pundir PS. Discrete Burr and Discrete Pareto Distributions. *Stat Methodol.* 2009;6(2):177–88.
- [18] Greenwood M, Yule GU. An Inquiry into the Nature of Frequency Distributions Representative of Multiple Happenings with Particular Reference to the Occurrence of Multiple Attacks of Disease or of Repeated Accidents. *J R Stat Soc.* 1920;83(2):255–79.
- [19] Willmot GE. Mixed Compound Poisson Distributions. *ASTIN Bull.* 1986;16(S1):59–79.
- [20] Maceda EC. On generalized Poisson Distribution. *Ann Math Stat.* 1948;19:414–6.
- [21] Holgate P. The modality of some Compound Poisson Distributions. *Biometrika.* 1970;87(3):666–7.
- [22] Simeunović I, Balaban M, Bodroža D. Pricing Automobile Insurance using Mixed Poisson Distributions. *Industrija.* 2018;46(1):61–78.
- [23] Iyer-Biswas S, Jayaprakash C. Mixed Poisson distributions in exact solutions of stochastic autoregulation models. *Phys Rev E - Stat Nonlinear, Soft Matter Phys.* 2014;90(5):052712.
- [24] Das KK, Ahmed I, Bhattacharjee S. A New Three-Parameter Poisson-Lindley Distribution for Modelling Over-dispersed Count Data. *Int J Appl Eng Res [Internet].* 2018;13(23):16468–77. Available from: <http://www.ripublication.com>
- [25] Adetunji AA, Sabri SRM. An Alternative Count Distribution for Modeling Dispersed Observations. *Pertanika J Sci Technol.* 2023;31(3):1587–603.
- [26] Adetunji AA, Sabri SRM. On zero-inflated mixed Poisson Transmuted Exponential Distribution: Properties and Applications to observation with excess zeros. *Maejo Int J Sci Technol.* 2023;17(01):68–80.
- [27] Adetunji AA, Sabri SRM. A New Three-Parameter Mixed Poisson Transmuted Weighted Exponential Distribution with Applications to Insurance Data. *Sci Technol Indones.* 2023;8(2):235–44.
- [28] Sankaran M. The Discrete Poisson-Lindley Distribution: 275 . Note. *Biometrics.* 1970;26(1):145–9.
- [29] Gómez-Déniz E, Sarabia JM, Balakrishnan N. A Multivariate Discrete Poisson-Lindley Distribution: Extensions and Actuarial Applications. *ASTIN Bull J IAA.* 2012;42(2):655–78.
- [30] Willmot GE. On Recursive Evaluation of Mixed Poisson Probabilities and Related Quantities. *Scand Actuar J.* 1993;1993(2):144–133.
- [31] Sastry DVS, Bhati D, Rattihalli RN, Gómez-Déniz E. Zero Distorted Generalized Geometric Distribution. *Commun Stat - Theory Methods.* 2016;45(18):5427–42.
- [32] Dzipire NC, Ngare P, Odongo L. A Poisson-Gamma Model for Zero Inflated Rainfall Data. *J Probab Stat.* 2018;2018.
- [33] Wu S. Poisson-Gamma Mixture Processes and Applications to Premium Calculation. *Commun Stat - Theory Methods [Internet].* 2020;0(0):1–29. Available from: <https://doi.org/10.1080/03610926.2020.1850791>
- [34] Çakmakyapan S, Özel G. The Poisson Gamma Distribution for Wind Speed Data. *AIP Conf Proc.* 2016;1726.

- [35] Ong SH, Low YC, Toh KK. Recent Developments in Mixed Poisson Distributions. *ASM Sci J*. 2021;14(3):1–10.
- [36] Willmot GE. Asymptotic Tail Behaviour of Poisson Mixtures with Applications. *Adv Appl Probab*. 1990;22(1):147–159.
- [37] Rémillard B, Theodorescu R. Inference Based on the Empirical Probability Generating Function for Mixtures of Poisson Distributions. *Stat Decis*. 2000;18(4):349–66.
- [38] Granzotto DCT, Louzada F, Balakrishnan N. Cubic Rank Transmuted Distributions: Inferential Issues and Applications. *J Stat Comput Simul*. 2017;87(14):2760–78.
- [39] Lv HQ, Gao LH, Chen CL. Ailamujia Distribution and its Application in Supportability Data Analysis. *J Acad Armored Force Eng*. 2002;16(3):48–52.
- [40] Jamal F, Chesneau C, Aidi K, Ali A. Theory and Application of the Power Ailamujia Distribution. *J Math Model*. 2021;9(3):391–413.
- [41] Jan U, Fatima K, Ahmad SP. On Weighted Ailamujia Distribution and its Applications to Lifetime data. *J Stat Appl Probab*. 2017;6:619–33.
- [42] Haq MA, Usman RM, Hashmi S, Al-Omeri AI. The Marshall-Olkin Length-Biased Exponential Distribution and its Applications. *J King Saud Univ - Sci*. 2019;31(2):246–51.
- [43] Ekhosuehi N, Kenneth GE, U.K. K. The Weibull Length Biased Exponential Distribution: Statistical Properties and Applications. *J Stat Econom Methods*. 2020;9:15–30.
- [44] Adetunji AA. Transmuted Ailamujia Distribution with Applications to Lifetime Observations. *Asian J Probab Stat*. 2023;21(1):1–11.
- [45] Shaw WT, Buckley IRC. The alchemy of probability distributions: beyond Gram-Charlier expansions, and a skew-kurtotic-normal distribution from a rank transmutation map. *Res Rep*. 2007;
- [46] Taniş C, Saraçoğlu B. Cubic rank transmuted inverse Rayleigh distribution: Properties and applications. *Sigma J Eng Nat Sci*. 2022;40(2):421–32.
- [47] Elhertaniy DA, Rahim A, Hamid B. The Cubic Rank Transmuted Gumbel Distribution. *J Al-QadisiyahComput Sci Math*. 2023;15(1):1–18.
- [48] Celik N. Some Cubic Rank Transmuted Distributions. *J Appl Math Stat Informatics*. 2018;14(2):27–43.
- [49] Bhatti FA, Hamedani GG, Sheng W, Ahmad M. Cubic Rank Transmuted Modified Burr III Pareto Distribution: Development, Properties, Characterizations and Applications. *Int J Stat Probab*. 2018;8(1):94.
- [50] Ogunde AA, Chuckwu UA, Agwuegbo O-NS. The Characterization of the Cubic Rank Inverse Weibull Distribution. *Asian Res J Math*. 2020;16(7):20–33.
- [51] Nash JC, Varadhan R, Grothendieck G. optimr Package: A Replacement and Extension of the “optim” Function. 2019.
- [52] R-Core Team. R: A language and Environment for Statistical Computing. [Internet]. R Foundation for Statistical Computing, Vienna, Austria.; 2020. Available from: <https://www.r-project.org/>

- [53] De Jong P, Heller GZ. Generalized Linear Models for Insurance Data. London: Cambridge University Press; 2008. 29 p.
- [54] Almetwally EM, Abdo DA, Hafez EH, Jawa TM, Sayed-Ahmed N, Almongy HM. The new discrete distribution with application to COVID-19 Data. Results Phys [Internet]. 2022;32(November 2021):104987. Available from: <https://doi.org/10.1016/j.rinp.2021.104987>
- [55] Nekoukhou V, Bidram H. The Exponentiated Discrete Weibull distribution. SORT. 2015;39:127–46.
- [56] Shanker R, Fesshaye H. On Poisson-Lindley distribution and its Applications to Biological Sciences. Biometrics Biostat Int J. 2015;2(5):1–5.
- [57] Samutwachirawong S. Poisson-Exponential and Gamma Distribution: Properties and Applications. J Appl Stat Inf Technol. 2021;6(2):17–24.
- [58] Sah BK, Mishra A. A Generalised Exponential-Lindley Mixture of Poisson Distribution. Nepal J Stat. 2019;3:11–20.
- [59] Hussain T, Aslam M, Ahmad M. A two parameter Discrete Lindley distribution. Rev Colomb Estadística. 2016;39:45–61.