

## **Original Research Article**

# **Impact of weed control measures on soybean (*Glycine max* L.) in kymore plateau and satpura hills**

### **ABSTRACT**

A field study was carried out at college farm of JNKVV, Jabalpur, M.P. to investigate the effect of different weed control treatments on weed flora and soybean. The experiment was laid out in randomized block design with three replications. Nine treatments were comprised of chlorimuron-ethyl 12, 24, 36, 48 and 72 g ha<sup>-1</sup>, hand weeding, mechanical weeding, combined application of weeding and chlorimuron and lastly weedy check. Application of herbicide and weeding effectively controlled weeds flora and their density rather than weedy check. Among all the treatments hand weeding recorded higher yield attributes, yield (1.69 t ha<sup>-1</sup> grain) and net monetary returns (20011.7 ha<sup>-1</sup>), but it was at par with chlorimuron-ethyl 24 g/ha along with mechanical weeding, that recorded 1.69 t ha<sup>-1</sup> grain yield and 20023.8 ha<sup>-1</sup> NMR.

*Keywords* : Chlorimuron-ethyl, NMR, Soybean, Weeding and Yield

### **1. INTRODUCTION**

Among oilseeds, soybean is one of the important crops of India. Soybean produced 23% of total vegetable oil of India (National Food Security Mission, 2018). It has also been termed as miracle bean and golden bean because, it contains about 40 per cent protein, 20 per cent oil which is rich in poly-unsaturated fatty acids and anti-oxidants. India recorded 11.33 million ha area, 13.79 million tones production and around 12.17 q ha<sup>-1</sup> productivity of soybean (Agricultural Statistics at a Glance, 2019).

Being a *kharif* season crop, it suffers from severe infestation of weeds, which rob it of essential nutrients, space and moisture, causing substantial loss in yield as it reduces the quality of seeds, it depends on weed flora and density [(Bhan and Kewat, 2003), (Tehulie et al., 2021)]. The excessive occurrence of weeds, limit the full expression of yield potential of crop, thus an early control of weeds in soybean is very essential and if it is not done, the yield losses may reach up to 25-80 per cent (depending upon type of weed flora and density). Hand weeding is one of the most efficient mean to control weeds in soybean, but it is time consuming and difficult due to unavailability of labours during peak period of demand. Weeding during critical growth stages is sometimes not possible due to continuous heavy downpours (Kumar et al., 2022). Use of mechanical weeding by hand hoe is popular in the Malwa tract of the state.

Thus, weeds can be controlled manually by pulling, mechanically, and chemically by application of herbicides or by a combination of these methods. But, application of suitable herbicide appears to be an effective alternative option to minimize the weed problems timely (Meseldzija et al., 2020). Chlorimuron-ethyl has been found very effective post emergence herbicide, for controlling both sedge and broad leaves weed but less in grassy weeds in soybean. Therefore, present investigation was carried out to assess the impact of weed control measures on soybean.

### **2. MATERIAL AND METHODS**

A field experiment was executed at product testing unit, Adhartal farm, J.N.K.V.V., Jabalpur, M.P. in *kharif* season of 2012. Jabalpur is situated at 23° 9' North latitude and 79° 58' East longitudes with an altitude of 411.78 meters above the mean sea level. It is classified as "Kymore Plateau and Satpura Hills" agro climatic zone as per norms of National Agriculture Research Project (NARP), New Delhi. The total rainfall received during the crop season was 1542 mm, which was equally distributed in 54 rainy days from June to third week of November.

The soil of the experimental field was clayey in texture, neutral in reaction (pH 7.2), medium in organic carbon (0.60 per cent), available N (372 kg ha<sup>-1</sup>), available P (16.40 kg P<sub>2</sub>O<sub>5</sub>ha<sup>-1</sup>) and high in available K (293 kg K<sub>2</sub>O ha<sup>-1</sup>).

The experiment involved nine treatments such as chlorimuron-ethyl 12, 24, 36, 48, 72 g ha<sup>-1</sup>, hand weeding, mechanical weeding, combined application of chlorimuron 24 g ha<sup>-1</sup> + hand hoeing and last one is weedy check, which were arranged in a randomized block design with three replications. The herbicide spray solution was prepared by mixing the required quantity of herbicide in water at 500 litreha<sup>-1</sup> for each plot. Before sowing, the seeds were treated with carbendazim at 2.0 g kg<sup>-1</sup> of seeds followed by inoculation with *Rhizobiumjaponicum* culture at 5 g kg<sup>-1</sup> of seeds. Seed rate was 70 kg ha<sup>-1</sup> of JS 97-52 variety of soybean. Fertilizer application (20 kg N + 60 kg P<sub>2</sub>O<sub>5</sub> + 20 kg K<sub>2</sub>O ha<sup>-1</sup>) was applied as basal application through urea, single super phosphate and murate of potash.

Crop management was done by using the recommended package of practices. The observations on population of major weeds and other associated weeds were recorded at 45 DAS and harvest by quadrat count method. The percentage composition of weed flora was estimated from weedy check plot. Formula of relative density proposed by Mishra (1968).

Number of individuals of the same species

$$\text{Relative Density (\%)} = \frac{\text{Number of individuals of the same species}}{\text{Number of individuals of all species}} \times 100$$

Weed index measures the reduction in crop yield due to weed competition as against weed free treatment is expressed in percentage and it was determined with the help of following formula (suggested by Gill and Kumar (1969). Here X is seed yield from weed free plot (hand weeding), Y is seed yield from the treated plot.

$$\text{WI (\%)} = \frac{X - Y}{X} \times 100$$

Data of plant parameters as plant height, dry weight, nodulation, number of pods per plant, seeds per pod, seed index is directly measured from the representative samples from each plot. The seed yield per net plot was recorded after winnowing the seed with the help of double pan balance. Finally, seed yield of each plot was converted in to seed yield per hectare. Haulm yield was determined by subtracting seed yield (economical yield) of each plot from biological yield (bundle weight) of the same plot. The net monetary returns (NMR) per hectare under each treatment was determined by subtracting the cost of cultivation of a particular treatment from the GMR of the same treatment. The collected data was analysed statistically using RBD as delineated by Panse and Sukhatme (1967) with the help of ANOVA.

### 3. RESULTS AND DISCUSSION

#### 3.1 Weed flora and Relative density of weeds

After the application of the herbicide weeds are controlled. But, in weedy check plots highest intensity of monocot were recorded at 45 DAS and harvest of soybean. Among monocot *Echinochloacolona* (26.27 and 21.18 per cent) was the most dominant weed followed by *Cyperus iria* (11.59 and 14.70 per cent) at 45 DAS and harvest respectively, whereas dicot weeds like *Eclipta alba* (9.34 and 12.58 per cent), *Commelinabenghalensis* (4.82 and 8.91 per cent), *Alternanthera philoxiroides* (16.37 and 14.89 per cent) and other weeds (31.61 and 23.65 per cent) were present in lesser number in soybean ecosystem (Table 1). Almost similar

weed flora associated with soybean was reported by Halvankar et al., (2005), Girothia et al., (2006), Pratap Singh and Raj Kumar (2008).

### 3.2 Weed Index

Results revealed that maximum reduction in yield (36 per cent) occurred in weedy check plots where weeds were not controlled throughout the crop season. Combined application of Chlorimuron-ethyl 24 g ha<sup>-1</sup> + mechanical weeding curbed the weed menace substantially bringing down WI to the tune of 5.3 per cent (Table 2). But in the hand weeding plots zero per cent yield reduction was observed as compared to other treatments of the investigation. The findings correlate with the findings of Bali et al., (2016).

### 3.3 Yield attributes and yield

Yield attributes, namely number of pods per plant and number of seeds per pod were significantly superior in the weed free treatment (154.06 and 2.67, respectively) than other treatments. Treatment chlorimuron-ethyl 24 g ha<sup>-1</sup> + mechanical weeding produced higher number of pods per plant and seeds per pod as (151.64 and 2.61, respectively) as compared to other treatments on account of maximum reduction in weed growth (Table 2). Excellent growth and development of soybean plants under weed free environment during critical period of crop growth might have resulted in higher number of pods per plant and seeds per pod under weed free treatment as compared to weedy check, which had severe weed competition right from early growth stages and ultimately resulted in most inferior yield attributes. Almost similar results were obtained by Raghuvanshi et al., (2005) and Singh et al., (2006).

Minimum seed index (9.02 g) was recorded in weedy check plot. Among the herbicidal treatments combined application of Chlorimuron-ethyl 24 g ha<sup>-1</sup> + mechanical weeding (9.83 g) produced higher seed index as compared to other treatments (Table 2). It may be because of maximum reduction in weed growth, coupled with no inhibitory effects on soybean plants and maximum transformation of nutrients as assimilates towards seeds resulted in higher seed index in these treatments. Results are in close conformity with the findings of Raghuvanshi et al., (2005), Singh et al., (2006) and Shete et al., (2008).

All the herbicidal treated plots produced significantly higher seed and stover yield than weedy check (10.86 q ha<sup>-1</sup> and 27.01 q ha<sup>-1</sup>, respectively). Yield was increased with the increased application rates of chlorimuron-ethyl 12, 24, 36, 48, 72 g ha<sup>-1</sup> and when combined application of Chlorimuron-ethyl 24 g ha<sup>-1</sup> + mechanical weeding was applied yield was increased at higher magnitude (16.08 q ha<sup>-1</sup> seed yield and 34.0 q ha<sup>-1</sup> stover yield). However, hand weeded plots (20 and 40 DAS) registered the highest seed and stover yield (16.98 q ha<sup>-1</sup> and 34.96 q ha<sup>-1</sup>, respectively) and proved significantly superior to all the treatments (Table 2), except chlorimuron-ethyl 24 g + mechanical weeding (40 DAS). Dhaker et al., (2015) also supported the findings. The crop under weed free plots attained lush growth due to elimination of weeds from inter and intra row spaces besides better aeration due to manipulation of surface soil and thus, more space, water, light and nutrients were available for the better growth and development, which resulted into superior yield attributes and development, and consequently the highest yield. Halvankar et al., (2005) and Jhadav and Kashid (2019) also reported the similar results. that, hand weeding as an effective method of weed control for achieving the maximum yield of soybean.

### 3.4 Economics

It is obvious from the data that there was very less net monetary returns Rs 9071.6 ha<sup>-1</sup> when crop was not weeded (Weedy check) throughout the crop season while highest NMR was gained from combined application of chlorimuron-ethyl 24 g ha<sup>-1</sup> + mechanical weeding (Rs

20023.8ha<sup>-1</sup>), closely followed by T<sub>6</sub>(Hand weeding) Rs 20011.7 ha<sup>-1</sup> (Figure 1). Pandya *et al*(2005) and Bali *et al.*, (2016).

#### 4. CONCLUSION

Among herbicidal treatments, combined application of Chlorimuron-ethyl 24 g ha<sup>-1</sup> + mechanical weeding, was most effective to reduced most of weed flora. Application of chlorimuron-ethyl herbicide at 24 g ha<sup>-1</sup> as early post-emergence along with mechanical weeding was significant superior for yield attributes and seed yield (16.08 q ha<sup>-1</sup>) of soybean than rest of the treatments without any phytotoxicity on soybean plants. It was also found more remunerative in terms of NMR (Rs 20023.8 ha<sup>-1</sup>) than application of chlorimuron-ethyl at 12 to 72 g ha<sup>-1</sup> as early post-emergence.

**Table 1. Weed flora of soybean during *kharif* season in weedy check plots at 45 DAS and at harvest**

S. No.	Weed flora	Density m <sup>-2</sup>		Relative density (%)	
		45 DAS	Harvest	45 DAS	Harvest
	<b>Monocot</b>				
1	<i>Echinochloacolona</i>	58.67	65.56	26.27	21.18
2	<i>Cyperus iria</i>	25.89	45.50	11.59	14.70
	<b>Dicot</b>				
3	<i>Alternanthera philoxaroides</i>	36.57	46.07	16.37	14.89
4	<i>Eclipta alba</i>	20.88	38.93	9.34	12.58
5	<i>Commmelinabenghalensis</i>	10.78	27.58	4.82	8.91
6	<b>Others</b>	70.62	85.86	31.61	27.65
	Total	223.40	309.5	100.00	100.00

**Table 2. Influence of weed control treatments on yield attributes, yield, weed index and economics of soybean**

Treatments	Pods plant <sup>-1</sup>	Seeds pod <sup>-1</sup>	Seed Index	Weed index	Seed yield (q ha <sup>-1</sup> )	Stover yield (q ha <sup>-1</sup> )	GMR (Rs ha <sup>-1</sup> )	NMR (Rs ha <sup>-1</sup> )
T <sub>1</sub> - Chlorimuron-ethyl 12g/ha	130.88	2.20	9.17	30.2	11.85	29.17	28990.7	11135.7
T <sub>2</sub> - Chlorimuron-ethyl 24g/ha	146.65	2.33	9.35	12.4	14.88	31.95	35752.8	17872.8
T <sub>3</sub> - Chlorimuron-ethyl 36g/ha	150.41	2.37	9.42	11	15.12	32.35	36435.1	18530.1
T <sub>4</sub> - Chlorimuron-ethyl 48g/ha	151.67	2.38	9.48	10.4	15.22	32.41	36715.9	18785.9
T <sub>5</sub> - Chlorimuron-ethyl 72g/ha	153.07	2.40	9.52	9.7	15.34	32.65	37046.1	19065.1
T <sub>6</sub> - Hand weeding (20 & 40 DAS)	154.06	2.67	9.95	0	16.98	34.96	41141.7	20011.7
T <sub>7</sub> - Mechanical weeding (20 DAS)	147.05	2.45	9.49	12.2	14.91	32.52	35848.6	15918.6
T <sub>8</sub> - Chlorimuron-ethyl 24g/ha + MW (40 DAS)	151.64	2.61	9.83	5.3	16.08	34.00	38863.8	20023.8
T <sub>9</sub> - Weedy check	130.90	2.00	9.02	36	10.86	27.01	26601.6	9071.6
SEm±	0.19	0.15	0.06	–	0.49	0.27		
CD at 5%	0.58	0.46	NS	–	1.47	0.82		

GMR : Gross Monetary Returns, NMR : Net Monetary Returns

## 5. REFERENCES

- Agricultural Statistics at a Glance, Ministry of Agriculture and Farmers' welfare. Government of India, 2019;p 74.
- Bali A, Bazaya BR, Chand L and Swami S. Weed management in Soybean (*Glycine max* L.). *The Bio-Scan (Supplement on Agronomy)*. 2016; 11(1): 255-257.
- Bhan M and Kewat ML. Activity and persistence of pendimethalin applied pre emergence to soybean in Vertisol. *Annals of Agricultural Research*. 2003; 24(4):970-972.
- Dhaker SC, Munda SL, Dhaker RC and Sumeriya HK. Effect of weed management and Sulphur on nutrient content and uptake by weeds and soybean. *Legume Research- An International Journal*. 2015; 38(3): 411-414.
- Gill GS and Kumar V. Weed index, a new method for reporting weed control trials. *Indian Journal of Agronomy*, 1969; 14(2): 96-98.
- Girothia OP and Thakur HS. Efficacy of post emergence herbicides for weed management in soybean. *Soybean Research*, 2006; 4(1/6):20-23.
- Halvankar GB, Varghese P, Taware SP and Raut VM. Effects of herbicides on weed dynamics and yield of soybean. *Journal of Maharashtra Agricultural Universities*, 2005; 30(1):35-37.
- Jadhav V T and Kashid NV. Integrated weed management in soybean., *Indian Journal of Crop and Weed*. 2019; 51(1): 81-83.
- Kumar S, Rana SS and Ramesh. Weed management strategies in soybean (*Glycine max*)- A Review. *Indian Journal of Agricultural Sciences*. 2022; 92(4): 438-44.
- Meseldzija M, Rajkovi M, Dudi M, Vranesevi M, Bezdani A, Jurisi A and Ljevnaić-Masi B. Economic feasibility of chemical weed control in soybean production in Serbia. *Agronomy*. 2020; 10: 291.
- Mishra, R. Community structure ecology work book. Oxford IBH Publ. Co. New Delhi, 1968; pp. 31-34.
- National Food Security Mission, 2018. Present status of oilseed crops and vegetable oils in India. <https://www.nfsm.gov.in/statusPaper/NMOOP2018.pdf>
- Pandya N, Chauhan GS and Nepalia. Influence of integrated weed management on yield and economic viability of soybean [*Glycine max* (L.) Merrill] grown at different crop geometries. *Indian Journal of Agriculture Sciences*. 2005; 75(8): 510-512.
- Panse VG and Sukhatme PV. Statistical methods for Agricultural Workers, ICAR, Publication New Delhi. 1967.
- Pratap Singh and Rajkumar. Agro-Economic Feasibility of weed management in soybean grown in South-Eastern Rajasthan. *Indian Journal of Weed Science*. 2008; 40(1&2):62-64.
- Raghuwanshi OPS, Deshmukh SC and Raghuwanshi SRS. Effect of some new post emergence herbicides on weed parameters and seed yield of soybean. *Research on Crops*. 2005; 6(3):448-451.
- Shete BT, Patil HM and Ilhe SS. Effect of cultural practices and post emergence herbicides against weeds control in soybean. *Journal of Maharashtra Agricultural Universities*. 2008; 33(1):118-119.

Singh VP, Mishra JS, Dixit A and Singh PK. Comparative efficacy of herbicides against spurge (*Euphorbia geniculata*) in soybean. *Indian Journal of Agricultural Sciences*.2006; 76(7):420-422.

Tehulie NS, Misgan T and Awoke T. Review on weeds and weed controlling methods in soybean (*Glycine max L.*). *Journal of Current Research in Food Science*.2021; 2(1): 01-06.




Figure 1. Economics of different weed control treatment in soybean