

Impact of industrial effluent on physico-chemical properties of soil in Unnao District of Uttar Pradesh

Abstract

A study was carried out during 2021-22 considering various vegetable growing farmers in Unnao District (U.P.) to find out the effect of using industrial effluents as the only source of irrigation to their crops. 250 surface soil samples collected from farmer's field were analyzed for physical as well as chemical parameters. The physical properties of soils improved due to industrial effluent irrigation. Among the physical properties bulk density ranged from 1.15 to 1.55 Mg m⁻³ and particle density ranged between 2.61 to 2.86 Mg m⁻³. Chemical Parameters of soil such as pH, EC, OC, Available N, P & K were also analyzed and the results indicated that soils had a mean pH of 8.1, E.C. of 1.25, O.C. of 49.80. Available nitrogen ranged from 377 – 1035 kg ha⁻¹ with mean of 658.26 kg ha⁻¹. The Phosphorus availability was found ranging from 20.5 – 48.80 kg ha⁻¹ with a mean of 35.97 kg ha⁻¹, while Potassium was found with a mean of 204.12 kg ha⁻¹ in range of 163 – 302.2 kg ha⁻¹.

Key words: Bulk density, Particle density, pH, Nitrogen and Potassium

INTRODUCTION

Water and nutrients are the major inputs for crop production. With the rapid growth in industrialization in different countries, the pollution problem is also on the increase, which have resulted in utilization of significant quantities of fresh water available for agriculture. It is predicted that most of the Asian countries will face severe problem related to water availability by 2025 (Singh, 1999). This has resulted in the application of waste water/effluents to agricultural soil are a useful source of plant nutrients, particularly nitrogen and phosphorous and also organic matter that can potentially improve soil fertility and physical properties. However, in addition to these beneficial effects, effluents often contain appreciable amounts of both organic and inorganic toxic materials. Many organic pollutants, being biodegradable, are less persistent, and presumably have transient and less serious effects as they are eventually metabolized to carbon dioxide and other inorganic substances.

In many areas of developing countries, untreated wastewater flows through channels into rivers where it is diverted by subsistence farmers to small plots of vegetables including

tomato, cabbage, beetroot and others which are easily consumed as salad. The public health risks of using such contaminated streams for irrigation are obvious (Mead and Griffin, 1998; WHO 2004). However, treated effluents can be used for irrigation under controlled conditions to minimize the transfer of pathogenic and toxic contaminants into agricultural products, soil surfaces and groundwater (Batarseh *et al.* 1989).

Since, industrial effluent is a pool of various elements, it may be beneficial in enhancing the fertility status of soil as it contains appreciable amount of essential plant nutrients that are taken up by crops during plant growth and development. Amount of available nitrogen, phosphorus and other nutrients also increase in the soil but excess of them can leach and pollute groundwater under continuous effluent use for long periods (Chaney, 1990).

Materials and methods

Unnao is the second township across the river Ganga, located between the rivers Ganga and Sai, roughly 22 km from Kanpur. The District is located between Latitudes 26°8' and 27°2' North and 80°3' and 81°3' east. It is divided from Kanpur and Fatehpur by the Ganga on the west, Lucknow on the east, Rae Bareli on the south, and Hardoi on the north. It covers 4589 sq. km of land. The Ganga, which forms the district's western and southern boundaries, and the Sai, which forms the district's northern and eastern boundaries, are its two principal rivers. About twenty or so tanneries are located in the district of Unnao, which is adjacent to Kanpur.

In the present investigation total 250 soil samples (0-20cm) were collected from farmer's field with the help of tube auger randomly from field during rabi season nearby effluent outlet who engage in irrigating their crops with industrial effluents.

Result and discussion:

Particle density was not changed in effluent irrigated soil. This can be attributed to improvement in total porosity and aggregate stability in the effluent irrigated soils due to addition of organic matter which plays an important role in improving soil physical environment. Rattan *et al.* (2001) observed enhanced available water content in the soils due to continuous application of effluents and sewage. In case of industrial effluent irrigated soils; the bulk density increase in comparison to sewage irrigated soil; the reason being abundant load of metal in soil causes deflocculation of soil aggregates by the activity of Na. Similar conclusions were suggested by Rai *et al.* (2011).

1.15	1.38	1.25	1.33	1.35	2.62	2.61	2.68	2.65	2.68	44.28	49.54	51.48	47.18	47.39
1.33	1.45	1.33	1.48	1.44	2.66	2.65	2.75	2.72	2.83	47.55	52.69	55.33	49.55	51.25
1.27	1.53	1.41	1.35	1.39	2.63	2.78	2.63	2.68	2.75	45.83	50.35	52.76	48.23	48.93
1.19	1.49	1.29	1.44	1.42	2.67	2.63	2.71	2.63	2.84	46.39	51.78	54.89	47.95	50.76
1.35	1.35	1.35	1.39	1.38	2.72	2.72	2.65	2.81	2.79	45.27	53.42	53.25	48.69	49.44
1.22	1.47	1.28	1.41	1.51	2.65	2.68	2.74	2.75	2.81	47.83	49.35	54.62	49.83	51.86
1.18	1.51	1.42	1.37	1.47	2.68	2.75	2.62	2.69	2.69	48.65	50.72	53.85	50.25	47.69
1.31	1.48	1.39	1.45	1.43	2.64	2.71	2.73	2.83	2.77	44.89	53.28	55.44	49.72	50.23
1.24	1.55	1.27	1.38	1.38	2.69	2.64	2.69	2.77	2.86	48.23	51.63	51.97	48.98	49.78
1.32	1.39	1.43	1.42	1.52	2.67	2.74	2.65	2.74	2.72	46.78	52.59	54.33	47.68	51.85

Table-2: Effect of industrial effluents on chemical properties of soil

pH					Electrical Conductivity (dS/m)					Organic Carbon (%)				
E ₁	E ₂	E ₃	E ₄	E ₅	E ₁	E ₂	E ₃	E ₄	E ₅	E ₁	E ₂	E ₃	E ₄	E ₅
7.95	8.02	8.05	8.02	7.93	1.07	1.24	1.05	1.13	1.35	44.72	49.54	51.37	47.34	46.82
8.08	8.08	8.08	8.06	8.05	1.11	1.37	1.22	1.27	1.42	48.55	53.85	55.82	49.65	49.55
7.97	8.05	8.15	8.11	7.99	1.09	1.28	1.08	1.15	1.39	46.23	49.78	52.65	48.89	47.63
8.12	8.03	8.11	8.08	8.07	1.15	1.33	1.17	1.21	1.47	47.89	51.33	50.73	49.23	51.95
8.07	8.11	8.07	8.15	7.94	1.13	1.29	1.25	1.19	1.33	45.61	50.87	53.68	47.98	48.36
7.99	8.07	8.04	8.07	8.11	1.08	1.32	1.18	1.32	1.47	48.38	54.49	52.74	48.15	50.09
8.05	8.15	8.12	8.16	7.97	1.14	1.35	1.09	1.28	1.35	45.87	52.38	51.88	49.76	46.25
8.02	8.12	8.09	8.13	8.03	1.12	1.41	1.16	1.17	1.44	46.52	50.15	54.35	48.55	49.38
8.14	8.09	8.11	8.18	8.08	1.17	1.39	1.13	1.24	1.37	48.91	53.62	53.89	50.83	48.79
8.09	8.14	8.16	8.14	8.15	1.15	1.27	1.27	1.29	1.41	44.83	49.95	52.57	47.49	50.45

Table-3: Effect of industrial effluents on chemical properties of soil

Available Nitrogen (kg/ha)					Available Phosphorus (kg/ha)					Available potassium (kg/ha)				
E ₁	E ₂	E ₃	E ₄	E ₅	E ₁	E ₂	E ₃	E ₄	E ₅	E ₁	E ₂	E ₃	E ₄	E ₅
383	738	930	377	653	32.4	34.3	20.5	40.2	27.5	215	206	168	163	185
394	745	948	385	678	39.7	38.5	26.2	46.8	31.8	258	211	173	175	193
387	763	967	394	694	45.2	37.9	24.7	45.3	29.4	293	217	175	168	188
403	774	984	408	686	48.8	35.2	23.4	41.9	33.9	245	208	169	173	197
398	812	1035	423	702	37.5	36.7	29.2	43.5	32.7	302	215	174	169	194
425	795	992	389	679	43.9	39.4	25.9	44.6	35.3	277	212	171	175	191
443	784	1021	411	713	42.2	41.2	21.5	47.4	34.5	285	219	178	177	189
457	833	975	447	708	38.7	38.8	28.8	42.9	28.5	269	214	165	184	195
413	825	1013	435	725	46.4	40.5	27.3	43.7	30.8	234	213	177	178	192
468	847	989	397	668	33.9	37.3	22.9	46.5	32.6	298	218	182	181	198

Conclusion

Continuous use of industrial effluent irrigation recorded improvement in soil physical properties like bulk density, water retention, organic carbon and build-up of soil available N, P, K, status. The industrial effluent irrigated soils had higher electrical conductivity and organic carbon as compared to the ground water irrigated soils. The electrical conductivity although increased due to effluent, it was within the tolerance limit to cause any soil salinity hazard. Available macronutrients including Nitrogen, Phosphorus and Potassium were also found to be higher than the normal level which lead to enhanced growth of vegetable crops thereby benefiting the farmers.

References:

Akan, J.C., Abdulrahman, F.I., Sodipo, O.A. and Lange, A.G. (2010). Physico-chemical Parameters in Soil and Vegetable Samples from Gongulon Agricultural Site, Maiduguri, Borno State, Nigeria., Journal of American Science 6(12).

- Amusan, A.A., Ige, D.V. and Olawale, R. (2005). Characteristics of soils and crops uptake of metals in municipal waste dump sites in Nigeria. *Journal of Human Ecology*, 17 : 167-171.
- Awomesoa, J.A., Ufoegbunea, G.C., Oluwasanyaa, G.O. and Ademola-Aremua, O.O. (2009). Impact of industrial effluents on water, soils and plants in the Alakia industrial area of Ibadan, South West Nigeria. 91 : 5-15.
- Batarseh, L. I. O. A., Rimavi, E. Salameh, 1989. Treated wastewater reuse in agriculture. Part1. Hussein Medical Center Project. The Water Research and Study Center, University of Jordan, Issue no. 12.
- Bhise, P. M., Cholla, O. and Venugopalan, M. V. (2007). Effect of waste water irrigation on soil properties under different land system. *Journal of the Indian Society Soil Science*, 55: 254-258.
- Brar, M.S., Malti, S.S., Singh, A.P., Arora, C.L. and Gill, K.S. (2000). Sewage water irrigation effects on some potentially toxic trace elements in soil and potato plants in Norton western India. *Canadian Journal of Soil Science*, 80 : 465-471.
- Chaney, R. L., 1990. Twenty years of land application research. *Biocycle*, 31:54-59.
- Chojnacka, K., Witek-Krowiak, A., Moustakas, k., Skrzypczak, D., Mikula, K., Loizidou, M. (2020). A transition from conventional irrigation to fertigation with reclaimed wastewater: Prospects and challenges. *Renewable and Sustainable Energy Reviews* 130 109959. <http://www.elsevier.com/locate/rser>.
- Gupta, A.P., Narwal, R.P., Antil, R.S. (1999). Sewer water composition and its effect on soil properties. *Bioresource Technology*, 65 : 171-173.
- Jackson, M. (1973). "Soil chemical analysis Prentice hall Inc. Englewood Cliffs" New Jersey, U.S.S.
- Kharche, V.K., Desai, V.N. and Pharande, A.L. (2011). Effect of Sewage Irrigation on Soil Properties, Essential Nutrient and Pollutant Element Status of Soils and Plants in a Vegetable Growing Area around Ahmednagar City in Maharashtra. *Journal of the Indian Society of Soil Science*, 59 :177-184.

- Kumar, Vipin and Nagar, Nidhi .(2012). Effect of industrial effluent on content of nutrient and heavy metals in plant and soil. Proc. International Conference on Chemical, Bio-chemical and Environmental Sciences (ICBEE'2012) December 14-15, 2012 Singapore.
- Leila El-Bassi, Ahmed Amine Azzaz, Salah Jellali, Hanene Akrouf, Evan A.N. Marks, Camélia Matei Ghimbeu, Mejdijeguirim (2020). Application of olive mill waste-based biochars in agriculture: Impact on soil properties, enzymatic activities and tomato growth, *Science of the Total Environment*
- Malla, R. and Totawat, K. L. (2006). Effect of municipal sewage on soil properties and chemical build up in vegetable crops grown on Haplusteps of sub humid southern plains of Rajasthan. *Journal of the Indian society of Soil Science*, 54 : 226-231.
- Mead, P. S., and P. M. Griffin, 1998. Escherichia coli 0157: H7. *Lancet* 352, 1207-1212.
- Mitra, A. and Gupta, S.K. (1999). Effect of sewage irrigation on essential plant nutrient and pollutant status in a vegetable growing area around Calcutta. *Journal of the Indian Society of Soil Science*, 47: 99-105.
- Rattan. R. K., Datta, S. P. Chonkar, P. K. Suribabu, K. and Singh, A. K. (2005). Long term impact of irrigation with sewage effluents on heavy metal content in soils, crops and ground water - a case study, *Journal of Agric. Ecosystems Environ.*, 109 : 310-322.
- Singh, P.K., Deshbhratar, P.B., Ramteke, D.S. (2012) Effects of sewage wastewater irrigation on soil properties, crop yield and environment. *Agricultural Water Management*, 103 : 100-104.
- Singh, R. Sand, R. P. Singh, 1999. Distribution of DTPA extractable Cd, Pb, Cr, Cu, Zn, Mn and Fe in soil profiles contaminated by sewage and industrial effluent. *J. Indian Soc. Soil. Sci.* 42(3):466-468.

UNDER PEER REVIEW