

Efficacy of Scabdel and GoldenCopper Fungicides on PhytopathogenicFungi Isolated from Guava Leaves

Abstract

The objectives of the study identify the fungal species from guava leaves symptomatic with leaf spots. Six fungal species (*Alternariaatra*, *Botryotrichiumverrucosum*, *Cochliobolusspecifer*, *Drechsleralalodes*, *Humicolagrisea* and *Stachybotryschartarum*) were isolated from infected guava leaves by the direct plate method on PDA medium at 28 °C. Pathogenicity test illustrated that only two species were pathogenic to guava leaves, and *B. verrucosum* exhibited moderate virulent ability, but *C. specifer* was weakly virulent. Half of the fungal species were pathogenic to apple fruits indicating that the species were non-host-specific to the guava plant. Scabdel fungicide completely inhibited the growth of *B. verrucosum* at 50 ppm, but in the case of *C. specifer* the inhibition ranged 18.8- 33.3 % with dose increases. Golden Copper was stimulatory to the growth of tested fungi at 50-1000 ppm doses.

Keywords: Guava leaves, identification, pathogenicity, fungicides.

1. INTRODUCTION

Guava (*Psidium guajava* L.) is an important fruit cultivated in several countries around the world [1]. It contains carbohydrates, protein, and adequate amounts of vitamin C and some minerals [2,3]. Due to its high nutritional value and low price, guava is considered one of the most popular fruits. In addition, it is used in folk medicine like the treatment of coughs, dental pain and diarrhoea [4]. Guava trees are susceptible to attack by more than 170 pathogens [5]. Several studies documented different diseases on guava trees e.g.; damping off and blight of seedlings, anthracnose, stem canker, leaf spots and blight, smooty mold and rots of fruits [6,7,8]. Among these diseases, leaf spots were considered mainly fungal infections incited by *Alternariaalternata* [9], *Alternariatenuissima* [10], *Cercosporapsidii*, and *Cercosporasawadae* [11] and *Pestalotiapsidii* [4]; or algal infection incited by *Cephaleuros virescens* [12].

There are different methods used for the management of phytopathogenic fungi on cultivated crops, including the use of fungicides [13]. The fungicides showed variable effects on different fungal species. Özer et al. [14] reported that seven of nine fungicides had an inhibitory influence on the mycelial growth of *Botrytis cinerea* isolates. Also, Equation Pro and Kema Zed showed inhibitory effects on mycelial growth and hydrolytic enzymes produced by broad bean leaf pathogens [15]. Recently; Abdel-Wahed [16] studied the effect of fungicides (Topsin M 70% and Vitavax 200 75%) and other compounds on the growth of *F. oxysporum*, *R. solani* and *F. solani*, and he concluded that the fungicides were the superior treatment in controlling the pathogens as compared with other compounds.

This work aimed to identify the fungal species from infected guava leaves, evaluate their virulence on guava leaves and apple fruits and control them with two fungicide applications.

2. MATERIAL METHODS

2.1 Collection of Guava Leaves

Fifteen samples of infected guava leaves symptomatic with leaf spots were collected from a South Valley University farm in May 2023. Each infected sample was put in a plastic bag transferred to the mycological lab and stored at 5 °C until further analysis.

2.2 Isolation of mycobiota from infected guava leaves.

The direct plate method was employed for fungal species isolation on potato dextrose agar (PDA) medium for 5-7 days at 28°C as described by Abdel-Fatthah et al. [17]. The developed colonies were purified and stored on PDA slant for further study. The isolated fungi were grown on PDA medium and morphological identification was performed based on macro (colony shape, colour, reverse and diameter) and microscopic features (conidial shape, color and size).

2.3 Pathogenicity Test

The virulence of fungal species (*Alternaria atra*, *Botryotrichum verrucosum*, *Cochliobolus specifer*, *Drechslera halodes*, *Humicola grisea* and *Stachybotrys chartarum*) were tested on detached guava leaves as described by Li et al. [18], with some modifications. Healthy appearance and freshly guava leaves were washed under tap water, then sterilized by 70% ethanol. Leaves were cut into pieces and each two pieces placed in a petri dish containing three filter papers. Leave pieces were wounded using a sterile needle and inoculated by fungal discs cut

from the margin of 1-week-old cultures. The control was inoculated with an agar plug without the fungus. Ten ml of distilled water was added to each plate, and then the plates were incubated at 28±1 °C for seven days[18]. The developed lesions were determined. Fungal re-isolation was performed from developed lesions according to Koch's postulates.

To determine whether the guava plant is a host specific to isolated fungi or not, the aggressiveness of isolated fungi was evaluated on apple fruits as follows; healthy and uniform apple fruits were surface sterilized by sodium hypochlorite (1%) for 3-5 min, then rinsed in sterilized distilled water twice and let dry under sterilized conditions. Active-growing fungal discs were inserted into holes made in sterilized apple fruits, as described by Baiyewu et al.[19] and Chukwuka et al.[20]. The control was carried out using sterile PDA. The inoculated fruits were put in polyethylene bags and incubated for 2 weeks at 28 °C. After the incubation period, the virulence of the tested fungi was estimated by measuring the diameter of the lesion. All experiments were conducted in triplicate.

2.4 Effect of Fungicides on *Botryotrichum verrucosum* and *Cochliobolus specifer* growth.

The efficacy of two fungicides, Scabdel 70% and Golden Copper 50% were tested against *B. verrucosum* and *C. specifer* growth. Their chemical names, active ingredients, manufactures and agricultural uses are shown in Table (1). For this purpose, different doses of fungicides (50, 100, 500 and 1000 ppm) were incorporated into autoclaved and cooled (about 50 °C) PDA medium, then active growing discs were extracted from 7 day colonies and placed in the center of PDA medium plates containing the fungicide dose. PDA medium without fungicides was used as a control. The plates were placed in an incubator at 28°C[21]. The radial fungal growth was determined after 7 days of incubation and the percentage of inhibition was determined by Equation $\% I_x = [X_c - X_i / X_c] \times 100$ [22]. where: % I_x: percentage of radial inhibition., X_c: mean radius (mm) of the control colony and X_i: mean radius (mm) of colonies in media with fungicide.

2.5 Statistical Analysis

Data were analyzed using one-way ANOVA and statistically significant values were considered at P < 0.05.

3. RESULTS

3.1 Mycobiota recovered from infected guava leaves.

Six fungal species attributed to six fungal genera namely; *Alternariaatra*, *Botryotrichiumverrucosum*, *Cochliobolusspecifer*, *Drechslerahalodes*, *Humicolagrisea* and *Stachybotryschartarum* were collected from guava diseased leaves on plates of Potato Dextrose Agar medium. From this, *Botryotrichiumverrucosum*, *Cochliobolusspecifer* and *Humicolagrisea* were recovered from 6.6%, 6.6% and 13.3% of the matching 16.6% of total fungi for each (Table 2).

3.2 The Pathogenicity of Isolated Fungi

The virulence of *Alternariaatra*, *Botryotrichiumverrucosum*, *Cochliobolusspecifer*, *Drechslerahalodes*, *Humicolagrisea* and *Stachybotryschartarum* was evaluated on guava leaves and apple fruits. The obtained data indicated that *B. verrucosum* and *C. specifer* were successful in lesion production on guava leaves after 7 days of incubation. *B. verrucosum* showed moderate virulence capacity with lesions less than 50% from leaf area, but in case *C. specifer*, the virulence capacity was low with lesions less than 25% from leaf area. The remaining four species failed to show any detectable disease symptoms on guava leaves (Table 3 and Fig.1).

Exactly 50% of the tested fungi (*B.verrucosum*, *C.specifer* and *D. halodes*) exhibited virulence capacity on apple fruits with different degrees. The highest fruit infection (30 mm) was observed on *B.verrucosum* followed by *C. specifer* (20 mm) and the lowest necrosis was achieved by *D. halodes* (12 mm). The success of some tested fungi to produce necrosis symptoms on apple fruits means these species are not host-specific to the guava plant (Table 3 and Fig. 2).

3.3 Effect of Fungicides on *Botryotrichiumverrucosum* and *Cochliobolusspecifer* Growth.

Four doses (50, 100, 500 and 1000 ppm, active ingredients) from two fungicides (Scabdel 70% and Golden copper 50%) were used to estimate the effect of fungicides on *B. verrucosum* and *C. specifer* growth.

Scabdel 70% exhibited strong inhibition activity on *B. verrucosum* growth; the fungicide at a concentration of 50 ppm completely stopped the fungal growth. The growth of *C. specifer* was greatly inhibited by Scabdel doses and the inhibition rate increased with increasing fungicide concentrations from 50-1000 ppm. *B. verrucosum* greatly influenced by scabdel than *C. specifer* (Table 4 and Figs. 3, 4).

The data illustrated that golden copper 50% has **no** inhibitory effect on the growth of both tested fungi, on the contrary the growth of *B. verrucosum* and *C. specifer* were significantly increased by increasing the fungicide dose from 50-1000 ppm (Table 4 and Figs. 3, 4).

4. DISCUSSION

Six fungal species were isolated from infected guava leaves symptomatic of leaf spots and the highest fungal count was achieved by *Botryotrichum verrucosum*, *Cochliobolus specifer* and *Humicola grisea*. Pandey et al. [23] studied the mycoflora of guava from the bud stage to the leaf fall stage in different seasons and they found that *Colletotrichum gloeosporioides*, *Fusarium oxysporum* f. sp. *psidii*, *Pestalotiopsis* and *Phoma psidii* were the most potent pathogenic species during all seasons. In Egypt, Youssef et al. [24] isolated *Alternaria alternata*, *Lasiodiplodia theobromae*, *Fusarium semitectum* and *Pestalotiopsis* from the dropped guava.

Our results showed that *B. verrucosum* and *C. specifer* succeeded in producing of disease symptoms on guava leaves and the *B. verrucosum* was more virulent than *C. specifer*. Youssef et al. [24] evaluated disease intensity on petioles and fruits of guava after artificial inoculation with *Alternaria alternata*, *Lasiodiplodia theobromae*, *Fusarium semitectum* and *Pestalotiopsis*, they concluded that the highest disease incidence on both petioles and fruits was achieved by *L. theobromae*. The inoculation of guava leaves with *Pestalotiopsis* resulted in typical leaf spot symptoms [4]. Song et al. [10] noticed that wilt and necrosis symptoms appeared on guava seedlings after wound inoculation with *Alternaria tenuissima*.

Scabdel 70% strongly affected the radial growth of *B. verrucosum* and *C. specifer* but in case of Golden Cooper 50%, the radial growth of tested fungi increases with the doses of fungicide. This finding is fully agreement with Özer et al. [14] he evaluated the effect of nine fungicides on the mycelial growth of *Botrytis cinerea* isolates and they concluded that all tested fungicides except Triadimentol and Tebuconazol showed inhibitory effects to different degrees. Abdel-Kader et al. [25] illustrated that, Euparen fungicide had no effect on the growth of *Alternaria alternata*, *Fusarium moniliforme*, *Myrothecium verrucaria* and *Thermoascus aurantiacus* at 4.8, 23.8 and 47.5 ppm concentrations. Also, Vacomilplus 50% (Copper oxychloride + Metalaxyl) at concentrations 50-200 ppm did not exhibit any detectable effects on

Lasiodiplodiatheobromae and *Fusarium semitectum* growth [24]. Abdul Wahid [26] studied the effect of two fungicides (Rovral and Sumislex) on the growth of five isolates of *Colletotrichum gloeosporioides* causing guava anthracnose in Egypt, he claimed that all tested isolates were strongly affected by the two fungicides used. The efficacies of fungicides toward pathogenic fungi were varied and affected by fungal species and fungicide dose [27]. Saleem et al. [15] found that, the mycelial growth of *A. alternata* was stimulated at lower doses (50 ppm) of Aquation Pro and Kema Zed fungicides and reduced at higher concentrations.

5. CONCLUSION

Third of the isolated fungi were able to infect guava leaves, Scabdelfungicide had inhibitory effect against *Botryotrichium verrucosum* and *Cochliobolus specifer*, on contrary Golden Copper showed positive effect on tested species.

ACKNOWLEDGMENTS

The experimental work was funded by project no. 609550-EPP-1-2019-1-BG-EPPKA2-CBHE-JP (2019-I943/001-001) new professional diploma in plant clinic and phytosanitary technologies (Program Erasmus+ capacity building in the field of higher education).

REFERENCES

1. Atawia AAR, El-Gendy FMA, Bakry Kh AI, Abd El-Ghany NA, Singer MAA. Physiological studies on flowering and fruiting of guava trees. Middle East J. Agric. Res. 2017;6(1):143-151.
2. Bardi, E. 1975. Tropical fruits: Guava. Abster Trop Agric. 1975;1:9-16.
3. Valentino MJG, Pineda FG, Fandialan MF. Phytopathogenicity of fungi associated with crown rot of Guava (*Psidium guajava*). Plant Pathol Quarantine. 2015;5(1):7-13.
4. Moustafa MSH, El-Dakar HAM, Alkolaly AMA. Pestalotia leaf spot a new disease affect guava trees in Egypt. International Journal of Scientific & Engineering Research. 2015;6(10):1306-1312.
5. Misra AK. Wilt of guava - a disease of national importance. Indian Phytopathol 2006;59(3):269-280.

6. Younis M, Mehmood K, Rashid A, Ashiq A. Effect of carbon, nitrogen sources and ascorbic acid on the colony growth and acervulus production of *Pestalotiapsidii*. Inter J Agric Biol. 2004;6(6):1110-1112.
7. Ismail OM, Abd El-Moniem EAA, Abd-Allah ASE, El-Naggar MAA. Influence of some post-harvest treatments on guava fruits. Agric Biol J NorthAmer. 2010;1(6):1309-1318.
8. Dwivedi SK, Neetu D. Antifungal activity of some plant extracts against guava wilt pathogen. Internat Environ Sci. 2012;3(1):412-420.
9. Perumal P, Sellamuthub K, Vanitha K, and Manavalasundaram V.K. Guava Leaf Disease classification using support vector machine. Turkish Journal of Computer and Mathematics Education.2021; 12(7):1177-1183.
10. Song XB , Cui YP , Peng AT, Ling JF, Chen X. First report of brown spot disease in *Psidiumguajava* caused by *Alternariatenuissima* in China. J Plant Pathol. (2020);102:1309.
11. Al-Sherif AAH, Khalil FAA, Mourad MY. Planting and production of guava in Egypt. Agriculture Research Center, Extension Publication. 2007; 640.
12. Abd El-Wahab FM, Shehata ASF, and Kamhawy MAM. First record of algal fruit and leaf spots of guava caused by *Cephaleuros virescens* in Egypt. Egypt J Phytopathol.2013;217-218.
13. Orina A, Gavrilova O, Gagkaeva T. The effect of fungicides on growth of *Fusarium* fungi in vitro. BIO Web of Conferences. 2020;18: 00022.
14. Özer N, Köycü ND, Delen N. Analysis of polygalacturonase enzyme production in isolates of *Botrytis cinerea* sensitive to different fungicides used in vineyards. J Phytopathol. 2010;158 (11-12): 744-749.
15. Saleem A, El-Said AHM, Magraby TA, Hussein MA. Effect of equation pro and kema zed fungicides on cellulase and pectinase production by some phytopathogenic fungi of broad bean. Afr J Biotechnol. 2014;13(46):4330-4337
16. Abdel-Wahed GA. Application of some fungicides alternatives for management root rot and wilt fungal diseases of marigold (*Calendula officinalis* L.). Scientific Journal of Agricultural Sciences. 2020;2(1):31-41.
17. Abdel-Fattah HM, Moubasher AH, Abdel-Hafez SI. Fungus flora of root and leaf surface of broadbean cultivated in oases, Egypt. Naturaliamonspeliensia. Seriebotanique. 1979;27:167-177.

18. Li H, Bian R, Liu Q, Sun L. Identification of a novel hypovirulence-inducing hypovirus from *Alternaria alternata*. *Front Microbiol.* 2019;10:1076.
19. Baiyewu RA, Amusa NA, Ayoola OA, Babalola OO. Survey of the postharvest diseases and aflatoxin contamination of marketed Pawpaw fruit (*Carica papaya L.*) in South Western Nigeria. *Afr J Agr Res.* 2007;2(4):178-181.
20. Chukwuka KS, Okonko IO, Adekunle AA. Microbial ecology of organisms causing Pawpaw (*Carica papaya L.*) fruit decay in Oyo state, Nigeria. *Am Eu J Tox Sci.* 2010;2(1):43-50.
21. Nisa T, Bhat MY, Pala SA, Mir RA. 2011 In vitro inhibitory effect of fungicides and botanicals on mycelial growth and spore germination of *Fusarium oxysporum*. *J Biopest.* 2011;4(1):53-56.
22. Plascencia M, Viniegra G, Olayo R, Castillo M, Shirai K. Effect of chitosan and temperature on spore germination of *Aspergillus niger*. *Macromol Biosci* 2003;3(10):582-586.
23. Pandey RR. Succession of microfungi on leaves of *Psidium guajava L.* *Bulletin of the Torrey Botanical Club.* 1990;117(2):153-162.
24. Youssef KYA, Mustafa ZMM, Mounir GA, Abo Rehab MEA. Preliminary studies on fungal species associated with guava fruit drop disease and possible management. *Egypt J Phytopathol.* 2015;43(1-2):11-23.
25. Abdel-Kader MIA, Moubasher AH, Abdel-Mallek AY. Studies on the effect of fungicide Euparen on cellulose-decomposing fungi in Egyptian soil. *J Basic Microbiol.* 1989;29:329-335.
26. Abdul Wahid OA. Comparative study on five isolates of *Colletotrichum gloeosporioides* causing guava anthracnose in Egypt and its control. *Microbiol Res.* 1999;154:63-69
27. Moharram AH, Abdel-Hafez SII, El-Said AHM, Saleem A. Effect of amistar and mocut fungicides on fungi faba bean plant and on amylase activity. *Afr J Microbiol Res.* 2004;5(26):4492-4507.

Table1. Fungicides used, their chemical names, active ingredients, manufacturers and agricultural uses.

	Scabdel 70%	Golden copper 50%
Chemical name	Dimethyl 4,4-(o-phenylene) bis (3 thioallophanate)	Copper oxychloride
Active ingredient	Thiophanate Methyl 70%	Copper oxychloride 50%, Cymoxanil 4%.
Manufactures	Wuxi Xinan, China	Aristo, India
Agriculture uses	Systemic fungicide used to control many plant diseases including: Powderly mildews, scab, leaf spots and rot of fruits and roots.	Systemic fungicide used to control many plantdiseases including: Early, late blight, downy mildews, Scab rot and fall of fruits.

Table 2. Total counts (TC), percentage count (%C) and percentage frequency of isolated fungi on Potato Dextrose Agar medium at 28 °C.

Fungal species	TC	% C	%F
<i>Alternariaatra</i>	1	8.3	6.6
<i>Botryotrichiumverrucosum</i>	2	16.6	6.6
<i>Cochliobolusspecifer</i>	2	16.6	6.6
<i>Drechslerahalodes</i>	1	8.3	6.6
<i>Humicolagrisea</i>	2	16.6	13.3
<i>Stachybotryschartarum</i>	1	8.3	6.6
<i>Sterile mycelia</i>	3	25	
Total count	12		

Table 3. Pathogenicity level of tested fungi on detached guava leaves and apple fruits.

Fungal species	Pathogenicity rate	
	Guava leaves	Apple fruits (in mm)
<i>Alternariaatra</i>	ND	ND
<i>Botryotrichiumverrucosum</i>	++	30.5
<i>Cochliobolusspecifer</i>	+	20
<i>Drechslerahalodes</i>	ND	12
<i>Humicolagrisea</i>	ND	ND
<i>Stachybotryschartarum</i>	ND	ND

+: Weak virulent (lesion less than 25% of leaves area); ++: moderate virulent (lesion 25-50% of leaves area)

Table 4. Effect of Scaddel70% and Golden Cooper 50% fungicides on the growth (in mm) of *Botryotrichiumverrucosum* and *Cochliobolusspecifer*.

Fungicide dose	Scabdel 70%		Golden copper 50%	
	<i>B. verrucosum</i>	<i>C. specifer</i>	<i>B. verrucosum</i>	<i>C. specifer</i>
Control	43	45	43	45
50	0*	36.5*	52*	46
100	0*	35*	57*	65*
500	0*	31.5*	65*	70*
1000	0*	30*	71*	73.5*
Inhibition (%)	100	18.8 - 33.3	-20.9-(-65.1)	-2.2-(-63.3)

* *P* value < 0.05 was considered statistically significant from the control.




Fig. 1. Disease symptoms on guava leaves after 7 days of incubation; A: control; B: weak virulent (*C.specifer*)and C: moderate virulent (*B.verrucosum*).



Fig. 2. External symptoms of infection resulted from tested fungi on apple fruits after 15 days of incubation, A: control, B: *A. atra* (nonpathogenic), C: *D. halodes*, D: *C. specifer*and E: *B.verrucosum*




Fig. 3. Colony diameter of *B.verrucosum* and *C.specifer* on different concentration of fungicides, A: *B.verrucosum* on Scabdel, B: *B.verrucosum* on Golden Copper, C: *C.specifer* on Scabdel and D: *C.specifer* on Golden Copper.




Fig. 4. Effect of different concentration of Scabdel and Golden Copper fungicides on *B. verrucosum* and *C. specifer* growth.