

EVALUATION OF DIFFERENT JUICE VARIETIES OF GRAPE(*Vitis* sp.) FOR QUALITATIVE CHARACTERISTICS

Abstract

The present investigation was conducted at College of Horticulture, Rajendranagar, during the year 2022. The experiment was laid out in completely randomized design. The research experiment was conducted to evaluate qualitative parameters of different juice varieties of grape. The results revealed that maximum TSS (21.05 °B) and brix/acid ratio (42.96) were recorded in T₁ – H-516. Maximum pH was recorded in T₆ – Manjari Medika (4.03) and highest titrable acidity was recorded in T₅ – Gulabi X Bangalore Purple (0.71 %). Reducing sugars were highest in T₁ – H-516 (17.24 %), whereas total sugars were maximum in T₄ – Concord (28.56 %).

Introduction

Grape is one of the important fruit crops that belongs to the family Vitaceae. The family contains about 1000 species belonging to 17 genera that are typically shrubs or woody lianas which climb by means of their tendrils (Keller, 2015). Grape is a refreshing fruit, rich in sugars, acids, minerals, vitamins and tannins. Major constituents of fruits are carbohydrates (15 %), minerals (0.2-0.6 %), organic acids (0.3-1.5 %), nitrogenous compounds (0.03-0.7 %), iron (0.003-0.017 %/100 g), calcium (0.004-0.025 %), potassium (0.15-0.25 %), vitamin A (1-80 microgram), vitamin B complex (391-636 mg/100 g), and vitamin C (1-1.25mg/100 mg) (Winkler, 1965). As fresh fruit, grapes are very delicate and extremely perishable and have a very high rate of loss during harvest and distribution. Therefore, grapes are processed into different products like wine, raisins, juice, crush, jelly, canned grapes etc to reduce waste and improve the marketability and profit from grape cultivation. (ChandrasekarVenkitasamy *et al.*, 2019). Grape juice contains 81 to 86 percent of water, in which nutrient elements, sugar and natural acids are present in readily available forms. Hence grape juice assimilates in body fluid immediately after consumption, and act as an excellent health drink with instant source of energy

(Gurak *et al.*, 2010). The present study aims at evaluating the different juice varieties of grape for their qualitative parameters.

Material And Methods

The experiment was conducted at College of Horticulture, Rajendranagar, during the year 2022. Eight grape varieties (T₁ – H-516, T₂ – Arka Shyam, T₃ – Black Cornechen, T₄ – Concord, T₅ – Gulabi X Bangalore Purple, T₆ – Manjari Medika, T₇ – Pusa Navrang, T₈ – Bangalore Blue) were collected from grape vines planted at Grape Research Station, Rajendranagar.

Total soluble solids (TSS) (°Brix)

The TSS of grape juice was measured with the help of digital refractometer (HI 96801, Hanna, Romania). Data was expressed as degree Brix (°B).

pH

The pH was determined using pH meter. pH meter was calibrated with the help of standard buffer solutions (pH 4.0 and 7.0). The juice sample was taken in 100 ml beaker and electrode was kept in the sample and read on pH meter.

Titration acidity (%)

Titration acidity in juice of different varieties of grape was estimated by adopting the procedure suggested by Lunke and Hashizume (2014).

$$\text{Acidity (\%)} = \frac{\text{TV} \times \text{Normality of alkali} \times \text{Eqwt of acid} \times \text{Vol made (ml)} \times 100}{\text{Volume of aliquot (ml)} \times \text{Weight of sample (g)} \times 1000}$$

Brix/Acid ratio

Brix-acid ratio was calculated by dividing the TSS value by the acid value.

TSS value

$$\text{Brix/Acid ratio} = \frac{\text{-----}}{\text{Acid value}}$$

Reducing sugars (%)

The reducing sugars were analysed by Lane and Eynon method suggested by Ranganna (1986).

Factor X Volume made up X 100

Reducing sugars (%) = -----

Titre value X Weight or volume of sample

Total sugars (%)

Total sugars in juice of different varieties were estimated by adopting the Lane and Eynon method suggested by Ranganna (1986).

Factor X Volume made up X 100

Total sugars (%) = -----

Titre value X Weight or volume of sample

Results and Discussion

Trials were conducted to evaluate different juice varieties of grape for qualitative parameters. The data recorded were statistically analysed and the results obtained are presented in the Table 1.

Total soluble solids (^oBrix)

The evaluation of TSS content established significant differences between different grape juice varieties. Among the different grape juice varieties, total soluble solids were significantly highest in T₁ – H-516 (21.05 ^oBrix) which was statistically on par with T₄ – Concord (20.87 ^oBrix), whereas significantly lowest total soluble solids were observed in T₅ – Gulabi X Bangalore Purple (17.47 ^oBrix). The variation in TSS may be attributed to changes in site, locality, topography and environment (Soni *et al.*, 2019). The variation in TSS of grape grown under same environment could be because of experimental conditions (Vijaya *et al.*, 2018). The results from current investigation are in accordance with the findings reported by Mehan *et al.* (2006), Patil *et al.* (2008), Gill *et al.* (2009), Ratnacharyulu (2010) and Vijaya *et al.* (2018) in different varieties of grape.

pH

pH was found significant in all treatments. Of all the varieties evaluated, T₆ – Manjari Medika had significantly highest pH (4.03), followed by T₁ – H-516 (4.00), while T₅ – Gulabi X Bangalore Purple recorded significantly lowest pH (3.22). The variation in pH of berry juice depends on genotypes, cultivars and environmental condition (Eshghi *et al.*, 2014). Karibasappa *et al.* (2008), Ratnacharyulu (2010), Sahoo *et al.* (2018) and Akram *et al.* (2020) reported similar findings in different grape varieties.

Titration acidity (%)

Significantly maximum titration acidity was obtained in T₅ – Gulabi X Bangalore Purple (0.71 %) which was statistically on par with T₃ – Black Cornechen (0.68 %), whereas the minimum titration acidity was obtained in T₆ – Manjari Medika (0.48 %). The difference in acidity among different varieties might be due to varietal difference and other factors like storage conditions (Akram *et al.*, 2020). The reduction in acidity at the time of harvest is also due to dilution effect caused by increased fruit size (Khan *et al.*, 2011). Similar observations in different varieties of grape were reported by Ghosh *et al.* (2006), Kumar and Rajan (2008), Patil *et al.* (2008), Gill *et al.* (2009) and Soni *et al.* (2019).

Brix/Acid ratio

Among the different grape juice varieties, T₁ – H-516 recorded significantly highest brix/acid ratio (42.96) which was statistically on par with T₆ – Manjari Medika (41.52), while T₅ – Gulabi X Bangalore Purple recorded the lowest brix/acid ratio (24.61). The brix/acid ratio varies in different varieties of grape because amount of TSS/TA ratio is governed by genetical constitution, phenotypical factors and also the day and night temperature generally favours the accumulation of solutes (Sahoo *et al.*, 2018). Mehan *et al.* (2006), Ghosh *et al.* (2006) and Gill *et al.* (2009) reported similar findings in various grape varieties.

Reducing sugars (%)

All the varieties exhibited significant differences in the reducing sugars with significantly highest being T₁ – H-516 (17.24 %) which was followed by T₄ –

Concord (16.39 %), whereas T₇ – Pusa Navrang recorded significantly lowest reducing sugars (10.86 %). The difference in reducing sugars among the grape varieties might be because of the fact that the sugars in grapes are greatly influenced by varietal difference and environmental condition (Yinshan *et al.*, 2017; Akram *et al.*, 2020). The results of present investigation are supported by findings of Ratnacharyulu (2010) and Bahkshet *al.* (2022) in different grape varieties.

Total sugars (%)

Total sugars were found significant in all treatments. Of all the varieties evaluated, significantly highest total sugars were recorded in T₄ – Concord (28.56 %) which was followed by T₁ – H-516 (25.60 %), while lowest total sugars were recorded in T₂ – Arka Shyam (14.21 %). The variation in the total sugars might be due to genetic makeup of the variety and environmental condition (Shiraishi *et al.*, 2010; Khan *et al.*, 2011). Similar findings were reported by Ghosh *et al.* (2006), Ratnacharyulu (2010) and Akram *et al.* (2020) in different grape varieties.

Conclusion

The results from the present investigation revealed that TSS was recorded maximum in T₁ – H-516 (21.05 °B) which was statistically on par with T₄ – Concord (20.87 °B). T₆ – Manjari Medika recorded maximum pH (4.03) followed by T₁ – H-516 (4.00). The maximum titrable acidity was found in T₅ – Gulabi X Bangalore Purple (0.71 %) which was statistically on par with T₃ – Black Cornechen (0.68 %). Highest brix/acid ratio was recorded in T₁ – H-516 (42.96) which was statistically on par with T₆ – Manjari Medika (41.52). T₁ – H-516 recorded maximum reducing sugars (17.24 %) followed by T₄ – Concord (16.39 %), whereas maximum total sugars were recorded in T₄ – Concord (28.56 %) followed by T₁ – H-516 (25.60 %).

References

- Akram, M.T., Qadri, R.W.K., Jaskani, M.J. and Awan, F.S. 2020. Phenological and physicochemical evaluation of table grapes germplasm growing under arid subtropical climate of Pakistan. *Pak. J. Bot.* 52(3): 1011-1018.
- Bahksh, A., Riaz, S., Kanwal, M.A.J., Nasir, M., Abbas, M.M., Hameed, A., Mahreen, N. and Aziz, M.M. 2022. Performance of various grapes cultivars under agro-climatic conditions of Central Punjab, Pakistan. *JPAA.* 7(3).
- Eshghi, S., Salehi, L. and Karami, M.J. 2014. Antioxidant activity, total phenolic compounds and anthocyanin contents in 35 different grapevine (*Vitis vinifera* L.) cultivars grown in Fars Province. *International Journal of Horticultural Science and Technology.* 1(2): 151-161.
- Ghosh, S.N. 2006, February. Performance of eight grape cultivars in laterite soil of West Bengal. In *International Symposium on Grape Production and Processing.* 785: 73-78.
- Gill, M.I.S. and Arora, N.K. 2009. Performance of different grape varieties under north Indian conditions. *Indian Journal of Ecology.* 36(1): 15-17.
- Gurak, P.D., Cabral, L.M., Rocha-Leao, M.H.M., Matta, V.M. and Freitas, S.P. 2010. Quality evaluation of grape juice concentrated by reverse osmosis. *Journal of food engineering.* 96(3): 421-426.
- Karibasappa, G.S. and Adsule, P.G. 2006, February. Evaluation of wine grape genotypes by National Research Centre for Grapes at their farm at Pune, Maharashtra, India. In *International Symposium on Grape Production and Processing.* 785: 497-504.
- Keller, M. 2020. *The science of grapevines.* Academic press.
- Khan, A.S., Ahmad, N., Malik, A.U., Saleem, B.A. and Rajwana, I.A. 2011. Phenophysiological revelation of grapes germplasm grown in Faisalabad, Pakistan. *Int. J. Agric. Biol.* 13(5): 791-95.

- Kumar, R. and Rajan, S. 2006, February. Grape cultivars flame seedless and Pusa Navrang can be commercially grown in north Indian conditions. In *International Symposium on Grape Production and Processing*. 785: 69-72.
- Lunkes, L.B.F. and Hashizume, L.N. 2014. Evaluation of the pH and titratable acidity of teas commercially available in Brazilian market. *RGO-Revista Gaucha de Odontologia*. 62: 59-64.
- Mehan, V.K., Gill, M.I.S. and Sharma, J.K. 2006, February. H-516: A Promising Grape Hybrid under N-Indian Conditions. In *International Symposium on Grape Production and Processing*. 785: 109-112.
- Patil, S.G., Karkamkar, S.P. and Rao, V.S. 2008. Selection of New Promising Grape Varieties for Their Commercial Cultivation. *Actahorticulturae*. 785: 105.
- Ranganna, S. 1986. Handbook of Analysis and Quality Control for Fruit and Vegetable Products. Tata McGraw Hill Publishing Company, New Delhi, 623-624.
- Ratnacharyulu, S.V. 2010. *Evaluation of coloured grape varieties for yield, juice recovery and quality* (Doctoral dissertation, Msc Thesis. Andhra Pradesh Horticultural University, Rajendranagar, Hyderabad, AP).
- Sahoo, T., Verma, M.K., Singh, S.K., Thakre, R.R.S., Jaiswal, S. and Arora, A. 2018. Morphological and biochemical characterization of newly evolved grape (*Vitis vinifera*) hybrids. *Indian Journal of Agricultural Sciences*. 88(8): 1280-6.
- Shiraishi, M., Fujishima, H. and Chijiwa, H. 2010. Evaluation of table grape genetic resources for sugar, organic acid, and amino acid composition of berries. *Euphytica*. 174: 1-13.
- Soni, N., Patil, P., Meena, K.C., Haldar, A., Patidar, D.K. and Tiwari, R. 2019. Evaluation of different coloured varieties of grapes under nontraditional area of Malwa Plateau: A thin line tool for doubling the farmer income. *Int. J. Curr. Microbiol. App. Sci*. 8(3): 1968-1976.

Venkitasamy, C., Zhao, L., Zhang, R. and Pan, Z. 2019. Grapes. In *Integrated processing technologies for food and agricultural by-products*. 133-163. Academic Press.

Vijaya, D., Reddy, R.G., Veena, J. and Mamatha, K.A. 2018. Evaluation of juice and wine varieties of grapes (*Vitis* spp) for petiole nutrient content, bud break, yield and yield components. *IJCS*. 6(6): 2739-2745.

Winkler, A.J. 1965. *General viticulture*. Univ of California Press.

Yinshan, G., N. Zaozhu, S. Kai, Z. Jia, R. Zhihua, Z. Yuhui, G. Quan, G. Hongyan and G. Xiuwu. 2017. Composition and content analysis of sugars and organic acids for 45 grape cultivars from northeast region of China. *Pak. J. Bot.* 49(1): 155-160.

Table 1 : Evaluation of different juice varieties of grape for qualitative parameters

Treatments	Total soluble solids (°Brix)	pH	Titration acidity (%)	Brix/Acid ratio	Reducing sugars (%)	Total sugars (%)
T ₁ : H - 516	21.05	4.00	0.49	42.96	17.24	25.60
T ₂ : Arka Shyam	18.57	3.44	0.54	34.38	12.82	14.21
T ₃ : Black		3.38				

Cornechen	19.13		0.68	28.13	11.36	15.16
T₄ : Concord	20.87	3.64	0.56	37.27	16.39	28.56
T₅ : Gulabi X Bangalore Purple	17.47	3.22	0.71	24.61	13.51	19.18
T₆ : Manjari Medika	19.93	4.03	0.48	41.52	14.49	22.53
T₇ : Pusa Navrang	18.84	3.99	0.51	36.94	10.86	16.34
T₈ : Bangalore Blue	20.67	3.34	0.58	35.64	12.65	15.92
SEm ±	0.09	0.01	0.01	0.60	0.26	0.29
CD at 5%	0.26	0.02	0.03	1.79	0.78	0.86
CV	0.76	0.27	2.74	2.94	3.28	2.52