

**Assessment of Physical and Chemical Properties of Continuously Cultivated Rice-Fallow and Uncultivated Soils of Nalbari District, Assam**

**ABSTRACT**

The present study was undertaken to assess the changes in physical and chemical properties of soils due to long term continuous cultivation of rice under rice –fallow system and that was compared with adjacent uncultivated soils. Soil samples were collected from respective sites and the soil physical (BD, WHC, MC) and chemical properties (pH, EC, organic carbon, available N, available P, available K, exchangeable Ca, exchangeable Mg, available S, available Zn and available B) analyzed were evaluated using descriptive statistics. The results indicated that the soils under cultivation with rice- fallow were significantly higher in bulk density ( $1.40 \text{ Mg m}^{-3}$ ) and lower in WHC (41.34%) than the adjacent uncultivated soils (BD= $1.34 \text{ Mg m}^{-3}$  & WHC=42.26%). The values of soil chemical properties (pH=5.51, organic carbon=0.63%, available N= $245.56 \text{ kg ha}^{-1}$ , available P=  $31.27 \text{ kg ha}^{-1}$ , available K= $120.98 \text{ kg ha}^{-1}$ , exchangeable Ca= $5.35 \text{ C mol (p+) kg}^{-1}$ , available S= $53.42 \text{ kg ha}^{-1}$ , available Zn= $0.31 \text{ mg kg}^{-1}$  and available B= $0.41 \text{ mg kg}^{-1}$ ) were found lower in cultivated rice- fallow soils than the uncultivated soils (pH=5.93, organic carbon=0.89%, available N= $418.37 \text{ kg ha}^{-1}$ , available P=  $42.62 \text{ kg ha}^{-1}$ , available

**K=145.90 kg ha<sup>-1</sup>, exchangeable Ca=5.93 C mol (p+) kg<sup>-1</sup>, available S=61.14 kg ha<sup>-1</sup>, available Zn=0.39 mg kg<sup>-1</sup> and available B=0.48 mg kg<sup>-1</sup>) and differed significantly. It indicated that continuous cultivation of crops without adopting proper management practices leads to decline in soil physical and chemical properties and proper management practices has to be adopted to sustain the properties of soil and crop productivity.**

**Key words:** Rice –fallow; uncultivated; physical properties; chemical properties; productivity.

## **1. INTRODUCTION**

Soil physical and chemical properties play vital role in maintaining soil fertility that has been threatened by various improper agricultural practices resulting deterioration of soil health and productivity . Continuous cultivation of crops with improper management practices on long term deteriorates soil physical and chemical properties and decline crop yield [1]. A better understanding of impact of continuous cultivation on soil properties is, therefore, essential for improvement of soil quality and sustaining soil health and productivity. Comparison of soil under natural vegetation and adjoining cultivated soils has revealed that prolonged agricultural land use alters some soil properties, mostly those related to fertility [2]. Continuous cultivation with frequent tillage results in a rapid loss of organic matter through increased microbial activity [3]. Changes in soil properties due to continuous cultivation and management practices and their consequences on crop productivity is an important area of research for soils of Nalbari district, Assam.

Nalbari district is located in central Western part of Assam is one of the agriculturally important district in Lower Brahmaputra Valley Zone of Assam lies between 26<sup>o</sup> N Latitude and 91<sup>o</sup> E Longitude with mean elevation 89 m above msl and comes under AESR 15.2 Q8B8 and 16.1C10A9. Though the district comprises 2.6% of state's geographical area, it has 5.46% and 4.96% of the state's net and gross cropped area, respectively. The district of Nalbari is primarily agrarian, with about 80% of the population directly or indirectly dependent on agriculture and allied activities. Rice is the major crop and rice based cropping systems are predominantly practiced by the farmers in the district and majority of the area comes under rice

-fallow system. Farmers cultivate rice crop with improper management practices that involves imbalance and injudicious use of nutrients, low farm input, and removal of residues from field may lead to decline the soil organic carbon resulting decline in the productivity and sustainability of intensive rice based cropping system.

The present study was undertaken with an objective to assess the impact of long term continuous cultivation of rice crop under rice –fallow system on physical and chemical properties of soils comparing with adjacent uncultivated soils.

## 2. MATERIALS AND METHODS

Geo-referenced (N:26°31.882' to 26°18.224' and E:091°30.536' to 091°15.750') soil samples (0-15cm) from rice -fallow cultivated soils were collected after harvest of rice crop during the year 2014-16. For comparison, the soil samples from adjacent uncultivated sites were also collected. The representativeness and uniformity of the fields were taken into consideration while collecting the soil samples. The sampling is focused on the plough layer because; this is where most soil quality changes are expected to occur due to long -term land use and soil management practices. All total 120 soil samples, 60 from cultivated rice-fallow and 60 from uncultivated soils, were collected covering 23 villages of the district. At the time of collection of soil samples, the crop history including management practices was recorded from respective farmer. The soil samples were mixed uniformly, air dried in shade and sieved with 2mm sieve and thereafter kept in zipped poly pouches with proper labelling inside and outside the packet for analysis of physical and chemical properties.

The physical properties like bulk density (BD), soil moisture content, soil texture and water holding capacity of soils were determined for soil samples collected from cultivated rice-fallow as well as adjacent uncultivated soils. For determination of bulk density undisturbed soil samples were collected from the field in natural condition using a tube core sampler (5.2cm diameter and 9cm length) following the standard method [4]. The soil moisture content (MC) was determined using the gravimetric method for field moist soils by drying at 105°C for 24 hours [4]. The texture of the soil samples were determined by International Pipette

Method [4]. Maximum water holding capacity (WHC) in percentage was determined by using Keen Rackzowski Box [4]. The chemical properties such as soil pH, electrical conductivity, organic carbon, CEC, available N, P, K, exchangeable Ca and Mg, available S, micronutrients such as available Zinc and boron were estimated following standard procedure [5].

Descriptive statistics was used for calculating the characteristics like mean, standard deviation (SD), minimum and maximum of analyzed soil physical and chemical properties. To compare the means of each individual soil parameters between cultivated rice-fallow and uncultivated soils, the paired t test was carried out. SPSS software (version 16.0) was used for analysis of the entire dataset.

### **3. RESULTS AND DISCUSSION**

#### **1. Crop History and soil management practices under rice-fallow system**

High yielding rice variety Ranjit, has been grown for past 10 years along with few other varieties like *Bahadur* and *Mahsuri* and other traditional varieties of *Joha* and glutinous rice-*Bora*, *Baismuthi* under rain fed conditions as mono crop. Average yield of rice is recorded as (4.3 t ha<sup>-1</sup>) for HYV and 2.5 to 3.0 t ha<sup>-1</sup> for traditional varieties. Use of organic manure was low and very irregular. Nutrient management conjoint with unscientific use of urea and SSP and a very little use of MOP leading to imbalanced nutrition. Tillage operation was carried out by power tiller and /or tractor. Farmers notice the yield decline over the years. After harvest of rice the field remains fallow.

#### **2. Soil physical properties**

##### **2.1. Soil texture**

The soil textures of the selected sites of rice-fallow system were recorded as sandy, loamy sand, sandy loam, loam, silt loam and clay (Table 1). The textural distribution of soils varied widely depending upon the parent materials. This was in conformity with the findings of extensive research works carried out on 107 soil series of Assam by Vadiveluet *al.* [6] including the soil series of Maroa, Tihu and Nalbari of Nalbari district.

##### **2.2. Bulk Density (BD)**

The mean value of bulk density (BD) of the soils under rice-fallow system was  $1.40 \text{ Mg m}^{-3}$  ( $\pm 0.06$ ) in cultivated sites compared to  $1.34 \text{ Mg m}^{-3}$  ( $\pm 0.07$ ) in adjacent uncultivated sites and differed significantly (Table 1). In the cultivated rice-fallow soils, 91.67% soils showed a higher BD ( $>1.3 \text{ Mg m}^{-3}$ ) values and in case of uncultivated soils 36.67% was found to have higher BD. The higher BD in ricegrowing sites might be due to soil compaction arising from puddling operation during rice cultivation. Singh *et al.*[7] reported that excessive tillage and wet tillage (puddling) under rice-wheat cropping system in general, resulted in gradual compaction of soils thus increases the BD. Both wetting drying and freezing-thawing cycles after tillage may also cause the bulk density to increase because of natural soil reconsolidation as reported by Assouline, [8] and Hu *et al.*[9]. The farm mechanization in tillage operation with continuous use of power tiller / tractor and less incorporation of organic inputs in soils also contributed towards higher value of BD.

### **2.3. Moisture Content (MC)**

The mean soil moisture content (MC) of the cultivated rice-fallow soils [26.58% ( $\pm 6.03$ )] after harvest of rice was non-significant (t test value 1.86) with uncultivated soils [27.01% ( $\pm 5.09$ )] (Table 1). In both cultivated and uncultivated soils showed a higher soil MC ( $>15\%$ ) in 98.33% and 100% of selected sites, respectively. The higher soil moisture content might be due to collection of soil samples immediately after harvest of rice and characterized by poorly drained characteristic of rice soils. Janssen *et al.*[10] reported that puddling over many years have potential to increase the plant available water capacity in the puddled layer, but reduces the small coarse and meso-pores (50 to  $0.2 \mu\text{m}$ ) in the plough pan. In case of uncultivated soil it was attributed due to higher content of organic matter which helps in retention of water.

### **2.4. Water Holding Capacity (WHC)**

The mean value of maximum water holding capacity (WHC) of cultivated rice-fallow [41.34% ( $\pm 11.17$ )] and uncultivated [42.26% ( $\pm 11.13$ )] soils varied significantly (Table 1) in which  $>80\%$  sites showed high WHC ( $>30\%$ ). The soil texture in the study sites varies from sandy loam, loam, silt loam and clay and WHC invariably controlled by soil texture.

Table: 1. Descriptive statistics of soil physical properties

Parameters	Cultivated			Uncultivated			Paired t-test value
	Minimum	Maximum	Mean ( $\pm$ SD)	Minimum	Maximum	Mean ( $\pm$ SD)	
BD ( $\text{Mg m}^{-3}$ )	1.25	1.6	1.40 ( $\pm$ 0.06)	1.2	1.50	1.34( $\pm$ 0.07)	8.07*
MC (%)	13.60	34.80	26.58( $\pm$ 6.03)	15.40	35.30	27.01( $\pm$ 5.09)	1.86 <i>ns</i>
WHC (%)	19.80	63.10	41.34( $\pm$ 11.17)	21.20	63.70	42.26( $\pm$ 11.13)	6.80*

*ns*- non significant; \* Significant at < 0.05 probability level.

### 3. Soil chemical properties

#### 3.1. Soil organic carbon (OC)

In the present investigation, the results illustrated that the mean value of OC of cultivated rice-fallow [0.63% ( $\pm$ 0.12)] and uncultivated [0.89% ( $\pm$ 0.22)] soils varied significantly (Table 2, Fig1). Of the total cultivated rice-fallow soils only 15% soils were in the high range (>0.75%) of OC compared to 76.67% in uncultivated soils. In the present investigation, inappropriate soil management with low inputs, excessive tillage and continuous mono cropping primarily with rice for longer period might have declined OC in cultivated rice soils compared to uncultivated soils (where no tillage operation were done). Maia *et al*[11] reported that conventional tillage caused a huge soil C loss by soil erosion and increasing soil OC decomposition rate while no-tillage increased soil OC pool in the soil by improving soil structure, reducing erosion, enhancing aggregates stability and providing physical protection.

#### 3.2. Soil pH

In the investigation the mean value of soil pH of cultivated rice-fallow [5.51 ( $\pm$ 0.51)] and uncultivated [5.93 ( $\pm$ 0.68)] soils remained non-significant (Table 2 & Fig 1), in which 60% and 83.33% sites showed moderately (4.5-6.0) acidic in cultivated and uncultivated soils respectively. Earlier Reza *et al*. [12] reported that the cultivated soils were moderately acidic in Assam which is in conformity with the present findings.

#### 3.3. Cation Exchange Capacity (CEC)

The low status ( $<10 \text{ C mol (p+ kg}^{-1})$ ) of CEC in the cultivated rice-fallow and uncultivated soils with the mean value of  $6.72 \text{ C mol (p+ kg}^{-1} (\pm 1.79)$  and  $6.88 \text{ C mol (p+ kg}^{-1} (\pm 1.83)$ , respectively were recorded (Table 2). Moral and Borah [13] also reported lower value of CEC in the soils of LBVZ of Assam. Moreover, the results were in conformity with the findings of extensive research works carried out on Nalbari soil series of Assam by Vadivelu *et al.* [6].

### **3.4. Electrical Conductivity (EC)**

The electrical conductivity (EC) of the soils under rice cultivation and uncultivated soils ranged between  $0.01$  and  $0.08 \text{ dSm}^{-1}$  and  $0.01$  and  $0.06 \text{ dSm}^{-1}$  with mean values of  $0.03 \text{ dSm}^{-1} (\pm 0.01)$  and  $0.03 \text{ dSm}^{-1} (\pm 0.01)$ , respectively exhibiting no significant difference of the mean value of EC in between the cultivated and uncultivated soils (Table 2). Karmakar [14]; Baruah *et al.* [15] reported that EC of Assam soil is low ( $<1 \text{ dSm}^{-1}$ ) indicating no salinity problem in Assam soils and has no adverse effect on crop growth.

### **3.5. Soil available Nitrogen (N)**

Conventional rice agro ecosystem, especially in Assam, have been characterized by low input of chemical fertilizers and organic amendments, leading to decline of available nutrient status. The mean value of available N content of soils under rice cultivation was  $245.56 \text{ kg ha}^{-1} (\pm 86.07)$  compared to the value of  $418.37 \text{ kg ha}^{-1} (\pm 85.45)$  in uncultivated soils and varied significantly (Table 2 & Fig 1). In the cultivated rice soils 66.67% of the samples were low in available N content ( $<272 \text{ kg ha}^{-1}$ ) compared to 3.33% in uncultivated soils. N is highly mobile in the soil and has highest probability, among the major nutrients, to be lost from rice eco system through volatilization and leaching and especially in high rain fall areas like Assam. Higher organic matter content in uncultivated soils attributed to higher level of available N in soils as compared to cultivated rice-fallow soils. Moreover, use of low input of chemical fertilizers and organic amendments are the characteristic of crops growing system of Assam, leading to decline in available nutrients status. The cultivated soils in the subtropical region coupled with preponderance of tillage practices and low external inputs are rarely sufficient in N as reported by Sanyal, *et al.* [16] which is in conformity with the present study.

### **3.6. Soil available Phosphorus (P)**

The mean value of available P content of the soils under rice cultivation was  $31.27 \text{ kg ha}^{-1} (\pm 5.87)$  compared to the mean value of  $42.62 \text{ kg ha}^{-1} (\pm 8.2)$  in uncultivated soils and varied significantly (Table 2 & Fig 1). The soil under rice cultivation showed 1.67% of soils as low ( $< 22.5 \text{ kg ha}^{-1}$ ), in available P and 98.33% in medium range ( $22.5\text{-}56.0 \text{ kg ha}^{-1}$ ) and 3.33% were high ( $> 56.0 \text{ kg ha}^{-1}$ ) in available P content. Since the soils of Nalbari district of Assam are acidic with low CEC, the low to medium range of available P in the soils under cultivated soils might be primarily due to the soil acidity that fixes the P into unavailable form as Al and/or Fe phosphate as well as low rate of application of phosphatic fertilizers as evidenced during soil samples collection. This was also reported that the average P fertility index is low to medium in Al and Fe- rich alluvial soils of Assam by Vadivelu *et al.*[6].

### **3.7. Soil available Potassium (K)**

The estimation of available K was widely differed across the location of the surveyed sites with a mean value of  $120.98 \text{ kg ha}^{-1} (\pm 37.62)$  in cultivated rice-fallow soils as compared to the mean value of  $145.90 \text{ kg ha}^{-1} (\pm 54.32)$  in uncultivated soils and exhibited significant differences (Table 2 & Fig 1). In the cultivated rice soils 70.00% of the total samples were low ( $< 136 \text{ kg ha}^{-1}$ ) in available K content, whereas 46.67% of the soils were found to be low ( $< 136 \text{ kg ha}^{-1}$ ) in case of uncultivated soils. Under low input situation, research carried out by Talukdar and Das [17] in Brahmaputra Valley Zone of Assam, demonstrated varied K supply power of soil including K deficiency.

### **3.8. Exchangeable Calcium (Ca)**

Nutritional responses to Ca can occur with sensitive crops in acid soils with low CEC in high rain fall zones. The estimation of exchangeable Ca was wide differed across the location and the mean exchangeable Ca content in cultivated sites of rice-fallow was found to be of  $5.35 (\pm 2.97) \text{ C mol (p+) kg}^{-1}$  compared to the  $5.93 (\pm 2.66) \text{ C mol (p+) kg}^{-1}$  in uncultivated soils respectively (Table 2 & Fig 1). The results illustrated that more than 50% of soils belonged to low [ $2\text{-}5 \text{ C mol (p+) kg}^{-1}$ ] to medium [ $5\text{-}10 \text{ C mol (p+) kg}^{-1}$ ] in exchangeable Ca content. Similar status of exchangeable Ca was also observed in a pedological study carried out in Assam soils by Dey and Sehgal,[18].

### **3.9. Exchangeable Magnesium (Mg)**

The mean exchangeable Mg content in cultivated sites of rice-fallow was found to be of 2.13 ( $\pm 1.02$ ) C mol (p+)  $\text{kg}^{-1}$  compared to the 2.38 ( $\pm 0.93$ ) C mol (p+)  $\text{kg}^{-1}$  in uncultivated sites (Table 2 & Fig 1). The results illustrated that more than 50% of soils under rice-fallow system belonged to the low [0.3-1 C mol (p+)  $\text{kg}^{-1}$ ] to medium [1-3 C mol (p+)  $\text{kg}^{-1}$ ] category in exchangeable Mg content. Extensive research works carried out by Vadivelu *et al.* [6] in Nalbari soil series of Assam reported 3.81 C mol (p+)  $\text{kg}^{-1}$  Ca+Mg which were in conformity with the present investigation.

### **3.10. Soil available Sulphur (S)**

In the present study, the mean available S content in cultivated sites of rice-fallow was found to be of 53.42 ( $\pm 23.22$ )  $\text{kg ha}^{-1}$  compared to the 61.14 ( $\pm 20.52$ )  $\text{kg ha}^{-1}$  in uncultivated sites (Table 2 & Fig 1). The results illustrated that more than 60% of soils under rice-fallow system belonged to high ( $> 44.8 \text{ kg ha}^{-1}$ ) category in available S content. In the present study, high level of S is due to continuous indiscriminate use of S containing fertilizers. Moreover, higher content of S in uncultivated soils might be attributed due to high level of organic matter as compared to cultivated rice-fallow soils.

### **3.11. Soil available Zinc (Zn)**

The mean DTPA-Zn content in cultivated sites of rice-fallow, was found to be of 0.31 ( $\pm 0.12$ ),  $\text{mg kg}^{-1}$  compared to the 0.39 ( $\pm 0.09$ ),  $\text{mg kg}^{-1}$  in uncultivated sites (Table 2 & Fig 1) and differed significantly. The results illustrated that more than 90% of soils under rice-fallow system belonged to deficient ( $< 0.6 \text{ mg kg}^{-1}$ ) in DTPA-Zn content. Similar findings of Zn deficiencies in soils of Assam was also observed by Bhuyan, *et al.* [19].

### **3.12. Soil available Boron (B)**

The estimated mean available B content in cultivated sites of rice-fallow, found to be of 0.41 ( $\pm 0.15$ )  $\text{mg kg}^{-1}$  compared to the 0.48 ( $\pm 0.11$ ),  $\text{mg kg}^{-1}$  in uncultivated sites (Table 2 & Fig 1). The results illustrated that more than 70% of soils under rice-fallow system belonged to deficient ( $< 0.5 \text{ mg kg}^{-1}$ ) category in available B content. B deficiencies in Assam soils was reported by Baruah *et al.* [15] which were attributed due to leaching of B during monsoon rain and less use of B in soil.

UNDER PEER REVIEW

Table 2. Descriptive statistics of soil chemical properties

Soil chemical Parameters	Cultivated			Uncultivated			Paired t- test value
	Minimum	Maximum	Mean ( $\pm$ SD)	Minimum	Maximum	Mean ( $\pm$ SD)	
pH	4.6	7.0	5.51( $\pm$ 0.51)	5.0	7.4	5.93 ( $\pm$ 0.67)	4.3*
EC (dSm <sup>-1</sup> )	0.01	0.08	0.03 ( $\pm$ 0.01)	0.01	0.06	0.03 ( $\pm$ 0.01)	0.38 $ns$
CEC [C mol (p+) kg <sup>-1</sup> ]	4.42	12.31	6.72 ( $\pm$ 1.79)	4.51	12.41	6.88 ( $\pm$ 1.83)	0.62 $ns$
OC (%)	0.45	1.03	0.63 ( $\pm$ 0.12)	0.54	1.70	0.89 ( $\pm$ 0.22)	12.86*
N (kg ha <sup>-1</sup> )	110.50	510.4	245.56 ( $\pm$ 86.07)	255.4	545.0	418.37( $\pm$ 85.45)	13.71*
P (kg ha <sup>-1</sup> )	20.57	47.57	31.27( $\pm$ 5.87)	30.5	57.2	42.62 ( $\pm$ 6.82)	13.64*
K (kg ha <sup>-1</sup> )	64.8	240.33	120.98 ( $\pm$ 37.62)	10.0	300.10	145.90 ( $\pm$ 54.32)	4.28*
Ca [C mol (p+) kg <sup>-1</sup> ]	1.40	13.20	5.35 ( $\pm$ 2.97)	1.7	13.2	5.93 ( $\pm$ 2.66)	3.05*
Mg [C mol (p+) kg <sup>-1</sup> ]	0.4	4.6	2.13 ( $\pm$ 1.02)	0.80	5.0	2.38 ( $\pm$ 0.93)	1.53 $ns$
S (kg ha <sup>-1</sup> )	3.67	112.9	53.42 ( $\pm$ 23.22)	28.8	121.2	61.14 ( $\pm$ 20.52)	6.49*
Zn (mg kg <sup>-1</sup> )	0.13	0.86	0.31 ( $\pm$ 0.12)	0.25	0.66	0.39 ( $\pm$ 0.09)	7.63*
B (mg kg <sup>-1</sup> )	0.11	0.89	0.41 (0.15)	0.30	0.78	0.48 (0.11)	4.68*

$ns$ - non significant; \* Significant at < 0.05 probability level.




Fig. 1 Soil chemical properties under rice-fallow and uncultivated soils

#### 4. CONCLUSION

The present investigation indicated that due to continuous cultivation of rice crop there was increase of BD value of soils as compared to the adjacent uncultivated soils. Similarly, nutrient status of soil also showed a declining trend. It can be concluded from the above study that continuous cultivation of rice crop on long term decline the soil physical and chemical properties. Therefore, cultivation of crop with proper soil management practices like balanced use of chemical fertilizers on the basis of soil test values, use of organic manures and inclusion of legume crops in crop sequence etc has to be adopted for sustaining soil health and crop productivity.

#### REFERENCES:

1. Prasad D, Verma A K, Jain V, Agrwal S, Tedia K. Effect of different rice-based cropping systems on Physico-chemical properties of soil. *Agronomy Journal*.2022; 114(3): 1642-1654.
2. Pez-Gonzalez A., Vieira SR, Taboada Castro MT. The effect of cultivation on the spatial variability of selected properties of an umbric horizon. *Geoderma*.2000; 97:273–292.
3. Shepherd TG, Sagggar S, Newman RH, Ross CW, Dando JL. Tillage-induced changes in soil structure and soil organic matter fractions. *Australian Journal of Soil Research*. 2001; 39:465–489.
4. Baruah TC, and Barthakur H P. A text book of Soil Analysis. Vikas Publishing House, New Delhi, India. 1997; pp 28-31.
5. Jackson, M. L. *In: Soil Chemical Analysis*. Prentice Hall of India Pvt. Ltd., New Delhi. 1973.
6. Vadivelu S, Sen TK., Bhaskar BP, Baruah U, Sarkar D, Maji AK., Gajbhiya K S. Soil Series of Assam, NBSS Publication No. 101, NBSS & LUP (ICAR) Nagpur. 2004; pp 22-27.
7. Singh K B, Jalota S K, Sharma, BD. Effect of continuous rice wheat rotation on soil properties from four agro ecosystems of Indian Punjab. *Communications in soil science and plant analysis*. 2009; 40: 2945-2958.
8. Assouline S. Bulk density of soils and impact on their hydraulic properties. In Glinski, J., Horabac, J. and Lipiec, J. Eds *Encyclopedia of agrophysics* springer, Dordrecht, The Netherlands.2011; pp.95-100.
9. Hu W, Shao MA, Si B. Seasonal changes in surface bulk density and saturated hydraulic conductivity of natural landscapes. *European Journal of Soil Science*. 2012; 63: 820-830.

10. Janssen I, Peng X, Horn R. Physical soil properties of paddy fields as a function of cultivation history and texture. *Advances in Geo Ecology*. 2006; 38: 446-455.
11. Maia SM F, Ogle SM, Cerri CC, Cerri CEP. Changes in soil organic carbon storage under different agricultural management systems in the South West Amazon Region of Brazil. *Soil and Tillage Research*. 2010; 106:177-184.
12. Reza SK., Baruah U, Bandyopadhyay S, Sarkar D, Dutta D P. Characterization of soil acidity under different land uses in Assam. *Agropedology*. 2012;22 (2):123-127.
13. Moral G, Borah D K. Salient Research Findings, Department of Soil Science, Assam Agricultural University, Jorhat-13. 1992; pp 35
14. Karmakar RM. Characterization and potentiality evaluation of soils developed in different land forms of North Bank Plain Zone of Assam. *Agropedology*. 2014; 24 (01): 52-63.
15. Baruah B K, Hague A, Das B, Medhi C, Mishra AK. Boron in soil and water samples in tea garden belt of Golaghat district, Assam. *Advances in Applied Science and Research*. 2011; 2(3): 298-305.
16. Sanyal SK, Majumdar K, Singh VK. Nutrient management in Indian Agriculture with Special reference to Nutrient mining. A Relook. *Journal of Indian Society of Soil Science*. 2014; 62 (4): 307-325.
17. Talukdar M C, Das M. Potassium supplying power of soil of Brahmaputra Valley of Assam. *Journal of Potassium Research*. 1994; 10 (3):285-289.
18. Dey JK, Sehgal JL. Characteristics and classification of some alluvium derived paddy and associated non paddy soils of Assam. *Agropedology*. 1997; 7:22-31.

19. Bhuyan N, Barua NG, Borah DK, Bhattacharyya D, Basumatary A. Georeferenced Micro nutrient status in soils of Lakhimpur district of Assam. Journal of the Indian Society of Soil Science. 2014; 62(2): 102-107.

UNDER PEER REVIEW