

Original Research Article

Influence of Diaminazene Aceturate (Berenil®) on the Haematology of Yankasa Sheep Experimentally Infected with *Trypanosoma evansi*

Comment [o1]: rams

Comment [o2]: delete

ABSTRACT

Haematological profiles were determined in Yankasa sheep experimentally infected with *Trypanosoma evansi* (*T. evansi*) and treated with diaminazene aceturate (berenil®). A total of 30 animals were divided into 6 groups (A to F) (n=5). Animals from each group were either uninfected or infected with *T. evansi*, and treated with Berenil®. Infection of the infected groups (A, C and E) was done via intravenous inoculation of *T. evansi*, while the infected group C and E were treated with berenil® at 3.5 and 7 mg/kg BW, respectively, by day 16 post infection (PI). The infected groups had pre-patent period of 8 days, with similar levels of parasitaemia (4.7 ± 0.27). In group A, the mean parasite count rose significantly ($p < 0.05$) to 72.8 ± 1.07 by day 12 PI and continue to a peak value of 250.6 ± 1.98 by day 28 PI. In group C and E, the initial parasitaemia rose significantly ($p < 0.05$) to 80.8 ± 1.12 and 78.2 ± 1.11 by day 12 PI, following treatment with 3.5 and 7.0 mg/kg BW of berenil®, by day 20 PI, respectively, and was completely eliminated by day 9 and 5 post treatment (PT), respectively. As parasitaemia increased, PCV, RBC, Hb, WBC, platelets count, absolute lymphocytes and monocytes, significantly declined ($p < 0.05$) in group A, leading to anaemia at day 16 PT. It is therefore,

demonstrated that both the two doses of berenil[®] were effective in the treatment of the disease under experimental conditions but 7.0 mg/kg cleared the parasitaemia faster.

KEYWORD: Diaminazene aceturate, Experimental, Haematology, *Trypanosoma evansi*, Yankasa breed of sheep

1. INTRODUCTION

Trypanosoma evansi, known as “surra”, is a disease of most domestic and wild animals. The pathogenic effect of the disease varies according to the virulence of the trypanosome strain, the host, species, general stress and other concurrent infections. [1]. *Trypanosoma evansi* is the most widely distributed of the pathogenic animal trypanosomes affecting camel, cattle, buffalo, horses and donkeys [2], causing economic losses including; severe weakness, abortion, infertility, reduced milk yield, weight loss and reduced drought capabilities that later lead to neuropathy and immune suppression coupled with anemia and eventually death of the animal [3]. *Trypanosoma evansi* is wide spread in different ecological regions in Africa, Americas, some parts of Europe and Asia [4]. Various species of domestic livestock can be infected with *T. evansi*, including horses, camels, buffaloes, sheep, goats, cattle and pigs. It is primarily transmitted mechanically by biting flies from the species like *Tabanus*, *Stomoxys*, *Haematopota*, *lyperosia*, and *Chrysops* spp [5]. In South-America, next to flies, vampire bats, during certain seasons can transmit the disease [6]. The diagnosis of surra is usually based on the demonstration of the parasite in blood, supplemented by haematological, biochemical and serological tests [7]. In general, clinical sign of *T. evansi* infections include; pyrexia directly associated with parasitaemia together with a progressive anaemia, and loss of condition [8].

The field control of animal trypanosomosis has, over the years, relied on two broad strategies: using chemotherapeutic agents on infected animals and vector control. The chemotherapeutic approach is used much more widely acceptable than vector control because it is easier to eliminate the trypanosomes than the flies [9]. Chemotherapy, exerts its action by stopping the multiplication of the trypanosomes, thereby helping the immune system to overcome the infection [10]. Chemotherapeutic drugs are also known to be toxic and often have a similar

disruptive effect on the cells of the host [11], and are therefore always used with care and at the recommended dose level only [12]. The most widely used and preferred curative trypanocide is diaminazene aceturate [13]. Other trypanocidal drugs that can be used include isometamidium chloride (curative and preventive), cymelarsan (curative and used only in camels), suramin, and quinapyramine (curative and preventive) [14]. In Nigeria, extensive work has been done on other animal trypanosomosis, but there remains paucity of information on infection due to *T. evansi* especially in Yankasa breed of sheep. Therefore, this research was designed to evaluate the haematological changes in Yankasa breed of sheep experimentally infected with *T. evansi* and with different doses of diaminazene aceturate.

2. MATERIALS AND METHOD

2.1 STUDY LOCATION

The study was conducted at the Department of Veterinary Medicine, University of Maiduguri, Maiduguri, Borno State capital, which lies in the northeastern geopolitical zone of Nigeria. The state is situated within the semi-arid zone of west Africa. It lies on latitude 11° 5'N and 13° 5'E. It has a total area of 72,609 square km, with temperature ranging from 35-40°C for most of the year

2.2 EXPERIMENTAL ANIMALS

A total of thirty (30) adult Yankasa breeds of sheep between 12-18 months of age with average weight of 30.22 ± 25 kg were procured from Maiduguri livestock cattle market and used for this study. The animals were kept in fly-proof pens and acclimatized for a period of two weeks. During this period, they were screened for haemoparasites and helminths by collecting blood and fecal samples for standard clinical examination procedure, respectively. They were fed twice a day with hay and concentrate supplement while salt lick and water were given *ad libitum*.

2.3 EXPERIMENTAL DESIGN

Comment [a3]: Delete s

The 30 adults Yankasa breed of sheep were randomly divided into six groups (A to F) of five animals each and labelled as follows:

Group A infected with 0.5ml of *Trypanosoma evansi*, but untreated control.

Group B was uninfected and untreated control.

Group C was infected with 0.5ml of *Trypanosoma evansi* and treated with diaminazeneaceturate (berenil[®]) at a single dose of 3.5mg/kg body weight at day 16 post infection (PI).

Group D was uninfected, treated with diaminazene aceturate at a single dose of 3.5mg/kg body weight at day 16 (PI).

Group E was infected with 0.5ml of *Trypanosoma evansi* and treated with diaminazene aceturate at a single dose of 7.0mg/kg body weight by day 16 (PI).

Group F was uninfected, treated with diaminazene aceturate at a single dose of 7.0mg/kg body weight by day 16 (PI).

2.4 INFECTIONS OF THE EXPERIMENTAL ANIMALS

Trypanosoma evansi used in this study was obtained from the Department of Veterinary Parasitology and Entomology, Ahmadu Bello University Zaria, Kaduna State, Nigeria. The organism was initially isolated from the blood of naturally infected Camel in Kano State Abattoir and maintained by serial passage in albino rats. Six (6) adult albino rats were used as donors. After patency, the donor rats were bled via the tail vein into a petri dish diluted with phosphate buffered saline glucose (pH 7.4). Each sheep in group A, C and E were inoculated through the jugular intravenous (IV) route with 0.5ml of blood containing 1.0×10^6 *Trypanosoma evansi*. The estimation of the number of infective trypanosomes was determined using the rapid matching method of Herbert and Lumsden [15].

2.5 ESTIMATION OF PARASITEMIA AND CLINICAL SIGNS

One milliliters of blood were collected from the jugular vein every 4 days into sample bottles containing EDTA. This blood was used to estimate parasitemia. Parasitemia was determined using wet mount and hematocrit centrifugation techniques for the detection of trypanosomes as described by Adeyeye et al. [16] and Desquesnes et al. [17]. The number of parasites was estimated following the method described by Herbert and Lumsden [18]. Post inoculation the animals were closely monitored for clinical signs of trypanosomosis such as pyrexia, palor of visible mucous membranes, anorexia, depression, lacrimation, nasal discharge, enlargement of lymph node.

2.6 SOURCE OF DRUG AND TREATMENT

Diaminazeneaceturateberenil[®] used in this study was manufactured by InterchemieWerken “De Adelaar” B.V. Metaalweg and venry, Holland. It was administered at a single dose of 3.5 mg/kg body weight to groups C and D, and 7.0mg/kg body weight to groups E and F, by deep intramuscular (IM) route on day 16th post infection at the peak of parasitaemia ($>45 \times 10^3/\text{uL}$).

2.7 BLOOD SAMPLE COLLECTION

About 1 ml of blood sample were obtained from the experimental animals via jugular vein, every 4 days (day 0, 4, 8, 12, 16, 20, 24 and 28) and dispensed into EDTA sample bottles. Blood samples collected were properly labelled according to group and used for determination of haematological parameters.

Comment [o4]: s

2.8 DETERMINATION OF HAEMATOLOGICAL PARAMETERS

Determination of packed cell volume (PCV) was carried out using microhaematocrit centrifuge as described by Coles, [19], Haemoglobin concentration (HB) were determined using cyanmethaemoglobin method [20], Red blood cell counts (RBC) and Total white blood cell counts (TWBC) were determined using Neubauer haemocytometer as described by Coles, [21], while thin blood smears stained with Giemsa stain were used to determine differential white blood cell counts. The Platelet counts were determined by the use of commercial kit (Randox lab Ltd, Ardmore, U.K.) according to the method of Cheesbrough, [22], via improved Neubauer counting chamber.

2.9 STATISTICAL ANALYSES

Data generated were expressed as mean \pm standard deviation (S.D) using Analysis of variance (ANOVA). Turkey – Kramer multiple comparison test was used to compare within and between group means and $P < 0.5$ was considered significant [23].

3. RESULTS

3.1 Mean Parasite Counts ($\times 10^3/\text{ul}$)

The mean parasite counts of the sheep experimentally infected with *T. evansi* and treated with diaminazene aceturate and their controls are presented in Figure 1. All the infected groups (A, C, and E) had a pre-patent period of 8 days post infection (PI) with a uniform mean parasitaemia of 4.7 ± 0.27 . In group A, the mean parasitaemia count of 4.7 ± 0.27 by day 8 (PI), rose significantly ($p < 0.05$) to $100.2 \pm$ on day 16 PI and continued without abatement to a peak value of 250.6 ± 1.98 by day 28 PI. At these points, the mucous (ocular and buccal) membranes of the animal in this group became pale. They became extremely weak, disinclined to move, anorectic and recumbent. They had to be humanly euthanized at this point (day 28 PI) to avoid painful death in accordance with international guidelines of using uninfected/uninfected animals for biomedical research. At this point, (day 28 PI), it was deemed that all the infected untreated animals had died of the infection. In the infected treated groups C and E, the initial parasitaemia of 4.7 ± 0.27 , which occurred on 8 PI, reached a significant ($p < 0.05$) peak count of 80.8 ± 1.12 and 78.2 ± 1.11 on day 12 (PI), respectively. Following treatment, the parasitaemia dropped significantly ($p < 0.05$) to 20 ± 0.56 on day 9 post treatment and subsequently cleared from the blood stream. In group E the parasite was completely cleared from blood stream on day 5 post treatment. No death or relapse parasitaemia was encountered for 90 days after monitoring parasitaemia in the infected, treated groups (C and E) respectively.




Fig. 1. Mean parasite counts ($\times 10^3/\mu\text{L}$) of Yankasa breed of sheep experimentally infected with *T. evansi* and treated with two different doses of berenil[®] and their controls.

3.2 Packed cell volume (PCV)

The mean PCV changes of the sheep experimentally infected with *T. evansi* and their controls are presented in Figure 2. In group A, the pre-infection PCV value of 38.7 ± 0.78 experienced a gradual but significant decline ($p < 0.05$) from day 8 PI, when parasitaemia became patent. By day 16 PI, parasitaemia had declined to 17.4 ± 0.52 and it continued unabatedly to 15.0 ± 0.48 on day 28 PI. In group B, D, and F, the pre-infection PCV of 38.8 ± 0.78 , 38.5 ± 0.78 and 38.2 ± 0.77 all remained fairly constant ($p < 0.05$) all throughout the period of the experiment. Meanwhile in group C, the pre-infected value of 38.4 ± 0.77 rose to a peak count of 20.7 ± 0.57 . Following treatment with 3.5mg/kg BW of berenil[®] on day 16 PI, the declined values began to rise significantly ($p < 0.05$), until its pre-infection PCV of 38.4 ± 0.77 was attained on day 28 PI (day 13 PT). Similarly, in group E, the pre-infection PCV of 38.4 ± 0.78 also declined significantly ($p < 0.05$) to 19.4 ± 0.56 . Following treatment with 7.0mg/kg BW of berenil[®], the values rose sharply ($p < 0.05$), thereby attaining its pre-infection value on day 24 PI (day 9 PT).




Fig. 2. Mean packed cell volume (%) of Yankasa breed of sheep experimentally infected with *T. evansi* and treated with two different doses of berenil[®] and their controls.

3.3 Red blood cell count

The mean total red blood cell counts of the sheep experimentally infected with *T. evansi* and their controls are presented in Figure 3. In group A, the pre-infection value of 11.84 ± 0.43 , experienced a significant ($p < 0.05$) decline which continued without abatement to 5.0 ± 0.28 on day 28 PI. In group B, D and F, their pre-infection values of 11.82 ± 0.43 , 11.84 ± 0.4 and 11.80 ± 0.43 all remained fairly constant ($p < 0.05$) throughout the study. However, in group C, the pre-infection value of 11.80 ± 0.43 declined significantly ($p < 0.05$) to 7.0 ± 0.34 on day 16 PI. Following treatment with 3.5mg/kg BW of berenil[®] on day 16 PI, the declined values began to rise significantly ($p > 0.05$) thereby, attaining its pre-infection value on day 28 PI or on day 13 PT. Similarly, in group E, the pre-infection value of 11.81 ± 0.43 declined significantly ($p < 0.05$) to 7.0 ± 0.33 on day 16 PI. Following treatment with 7.0mg/kg BW of berenil[®] on day 16 PI, the declined values appreciated significantly ($p < 0.05$) to its pre-infection value on day 24 PI or on

day 9 PT.




Fig. 3. Mean total red blood cell counts ($\times 10^6/\text{mm}^3$) of Yankasa breed of sheep experimentally infected with *T. evansi* and treated with two different doses of berenil[®] and their controls.

3.4 Haemoglobin concentration

The mean haemoglobin concentrations of the sheep experimentally infected with *T. evansi* and treated with two different doses of berenil[®] and their controls are presented in Figure 4. In group A, the pre-infected value of 9.5 ± 0.39 experienced a significant decline ($p < 0.05$) to 6.0 ± 0.31 on day 16 PI. Thereafter, the value began to decline further to 4.0 ± 0.25 on day 28 PI. Meanwhile in group B, D, and F, the pre-infected values of 9.6 ± 0.39 , 9.5 ± 0.39 and 9.7 ± 0.39 respectively, remained fairly constant ($p > 0.05$) throughout the study. However, in group C the pre-infection value of 9.7 ± 0.39 declined significantly ($p < 0.05$) to 5.8 ± 0.30 on day 16 PI. Following this treatment on day 16 PI, the declined values began to rise significantly ($p < 0.05$),

thereby attaining its pre-infection value on day 28 PI or on day 13 PT. Similarly, in group E, the pre-infection value of 9.7 ± 0.39 declined significantly ($p < 0.05$) to 7.0 ± 0.33 on day 16 PI. Following treatment with 7.0 mg/kg BW of berenil[®], on day 16 PI, the value rose significantly ($p < 0.05$), thereby attaining its pre-infection value on day 24 PI or on day 9 PT.




Fig.4. Mean haemoglobin concentrations (g/dl) of Yankasa breed of sheep experimentally infected with *T. evansi* and treated with two different doses of berenil[®] and their controls.

3.5 Total white blood cell count

The mean white blood cell counts (WBC) of the sheep experimentally infected with *T. evansi* and their controls is presented in Figure 5. In group A the mean pre-infection value of 10.0 ± 0.39 experienced significant ($p < 0.05$) decline to 7.2 ± 0.34 on day 16 PI. This decline in values continued unabatedly to 5.0 ± 0.28 on day 28 PI. For group B, D, and F, the pre-infection WBC counts of 10.2 ± 0.39 , 10.0 ± 0.39 , and 10.3 ± 0.39 , respectively, remained fairly constant ($P > 0.05$) throughout the study. Meanwhile, in group C, the pre-infection value of 10.3 ± 0.39 , declined significantly ($p < 0.05$) to 7.0 ± 0.33 on day 16 PI. Following treatment with 3.5 mg/kg

BW of berenil[®], by day 16 PI, the pre-infection value was attained on day 28 PI (day 13 PT). Similarly, in group E, the pre-infection value of 10.2 ± 0.39 declined significantly ($p < 0.05$) to 7.4 ± 0.35 on day 16 PI. Following treatment with 7.0mg/kg BW of berenil[®], on day 16 PI, the values began to increase significantly ($P < 0.05$), until its pre-infection value was attained on day 24 PI (day 9 PT).




Fig.5. Mean white blood cell counts ($10^3/\text{mm}^3$) of Yankasa breed of sheep experimentally infected with *T. evansi* and treated with two different doses of berenil[®] and their controls.

3.6 Absolute platelet counts

The absolute platelet counts of sheep infected with *T. evansi* and their controls are presented in Figure 6. In group A, the pre-infection value of 151.6 ± 1.54 experienced a significantly ($p < 0.05$) decline in value to 60.0 ± 0.97 on day 16 PI. Thereafter, it fluctuated on day 20 PI and continued to decline further, to 45.6 ± 0.84 on day 28 PI. In group B, D, and F, with pre-infection values of 152.6 ± 1.54 , 152.9 ± 1.55 and 150.3 ± 1.53 all remained fairly constant respectively ($p > 0.05$).

throughout the study. For group C, the pre-infection value of 151.8 ± 1.54 declined significantly ($p < 0.05$) to 70.0 ± 1.05 on day 16 PI. Following treatment with 3.5mg/kg BW of berenil[®] on day 16 PI, the pre-infection value was attained on day 28 PI (day 13 PT). Similarly, in group E, the declined in values of 75.0 ± 1.08 on day 16 PI increased significantly ($p < 0.05$) following treatment with 7.0mg/kg BW of berenil[®] on day 16 PI to its pre-infection value on day 24 PI (day 9 PT).




Fig.6. Mean absolute platelet counts ($\times 10^9/L$) of Yankasa breed of sheep experimentally infected with *T. evansi* and treated with two different doses of berenil[®] and their controls.

3.7 Absolute Neutrophil counts

The absolute neutrophil counts of the sheep experimentally infected with *T. evansi* and their controls are presented in Figure 7. In group A with a pre-infection value of 7.36 ± 0.34 experienced an unabated but significant ($p < 0.05$) decline of values to 4.3 ± 0.26 on day 16 PI.

Thereafter, a further decline continued unabated to 3.0 ± 0.22 on day 28 PI. In groups B, D and F, the pre-infection values of 7.80 ± 0.34 , 7.80 ± 0.35 and 7.08 ± 0.33 remained fairly constant ($p > 0.05$) throughout the period of the experiment respectively. In group C, the pre-infection value of 7.82 ± 0.35 continued to decline ($P < 0.05$) without abatement to 4.0 ± 0.25 on day 16 PI. Following treatment with 3.5mg/kg BW of berenil[®], on day 16 PI the values increased gradually until the pre-infection value was attained on day 28 PI (day 13 PT). In group E, the pre-infection value of 7.06 ± 0.33 experienced a significant decline ($p < 0.05$) to 4.2 ± 0.28 on day 16 PI. Following treatment with 7.0mg/kg BW of berenil[®] on day 16 PI, the values increased slightly ($p > 0.05$) without attaining its pre-infection value on day 24 PI (day 9 PT).




Fig.7. Mean absolute neutrophil counts ($\times 10^3/\text{mm}^3$) of Yankasa breed of sheep experimentally infected with *T. evansi* and treated with two different doses of berenil[®] and their controls.

3.8 Absolute Lymphocyte

The mean absolute lymphocyte changes of Yankasa breed of sheep and their controls are presented in Figure 8. In group A, the pre-infection value of 3.26 ± 0.23 continued to decline significantly ($p < 0.05$) without abating to a value as low as 1.0 ± 0.3 on day 28 PI. In group B, D and F, their pre-infection values of 3.45 ± 0.23 , 3.04 ± 0.22 and 3.60 ± 0.24 remained fairly

constant ($p>0.05$) throughout the experiment. In group C, the pre-infection value of 3.27 ± 0.23 declined significantly ($p<0.05$) to 1.9 ± 0.17 on day 16 PI. Following treatment with 3.5mg/kg BW of berenil[®] on day 16 PI, the values increased significantly ($P<0.05$), thereby attaining its pre-infection value on day 28 PI (day 13 PT). In group E the pre-infection value of 3.67 ± 0.24 declined significantly ($p<0.05$) to 1.6 ± 0.16 on day 16 PI. Following treatment with 7.0mg/kg BW of berenil[®] on day 16 PI, the pre-infection value was attained on day 24 PI (day 9 PT).




Fig.8. Mean absolute lymphocyte counts ($\times 10^3/\text{mm}^3$) of Yankasa breed of sheep experimentally infected with *T. evansi* and treated with two different doses of berenil[®] and their controls.

3.9 Absolute monocyte counts

The mean absolute monocyte changes of the sheep and their controls are presented in Figure 9. In group A, the pre-infection value of 4.5 ± 0.27 experienced a significant increase ($p<0.05$), which continued without abating to 47.8 ± 0.86 on day 28 PI. In group B, D and F, their pre-infection values of 4.6 ± 0.27 , 4.8 ± 0.27 and 4.0 ± 0.25 remained fairly constant ($p>0.05$) throughout the study. In group C, the pre-infection value of 4.7 ± 0.27 , rose significantly ($p<0.05$) to 20.7 ± 0.57 on day 16 PI. Following treatment with 3.5mg of berenil[®] on day 16 PI, the values

began to decline significantly ($P < 0.05$) to its pre-infection value on day 28 PI (day 13 PT). In group E, the pre-infection value of 4.5 ± 0.27 rose significantly ($p < 0.05$) to 23.2 ± 0.60 on day 12 PI. Following treatment with 7.0mg/kg BW of berenil[®] on day 16 PI, the value declined significantly ($p < 0.05$) to its pre-infection value on day 24 PI (day 9 PT).




Fig.9. Mean absolute monocyte counts ($\times 10^3/\text{mm}^3$) of Yankasa breed of sheep experimentally infected with *T. evansi* and treated with two different doses of berenil[®] and their controls.

4. Discussion

This study observed a short pre-patent (PP) of 8 days in the infected animals, which agrees and comparable to 7-11 days observed in Yankasa sheep infected with *T. evansi* [24, 25]. However, the result of PP in present study deviates from Wada et al. [26], who observed a much longer pre-patent period of 20 days in Yankasa breed of sheep infected with *T. evansi*. The variations in the pre-patent periods may be attributed to dose of infection, immunity, virulence of the strain of the parasite [27]. Uniform patency and parasitaemia was observed following inoculation of the animals, irrespective of the host susceptibility. Similar observation has been reported in *T. evansi* infected buffaloes, cow, and goats [28, 29], in *T. bruceigambiense* infected baboons (*Papioanubis*) [30]. Fluctuating parasitaemia were also observed generally, among the *T.*

evansi infected sheep. Fluctuations in parasitaemia are known features of trypanosomosis commonly caused by antigenic variation [31]. The ability of the host to limit the peak and number of each wave of parasitaemia is however, dependent on whether the infection is acute, sub-acute or chronic [32], and this may explain the reason why parasitaemia in sheep appreciated and declined following treatment on day 16 PI.

Evaluation of haematological parameters reveal that the infected sheep (group A, C and E) showed a significant decline in red blood cells (RBC), Packed cell volume (PCV), and haemoglobin concentrations (Hb) as the infection progressed. This was indicative of anaemia, which started at the onset of parasitaemia, from day 8 PI. This is in agreement with several reports that the anaemia in trypanosomosis often starts during the 1st wave of parasitaemia and is haemolytic in nature [33]. However, the haemolytic nature of the anaemia in most cases would depend on the species of trypanosomes involved as suggested by Mbaya et al.[34]. Trypanosome infection may cause haemolytic anaemia due to massive erythrophagocytosis by an expanded and active mononuclear phagocytic system (MPS) of the host[35, 36]. The presence of the MPS might have been associated with increased demand on the system to remove dead red blood cells, tissue cells, trypanosomes, antigen-antibody complexes and to participate in immune responses [37]. The fact that the red cell parameters (PVC, RBC, Hb) decreased sharply during wave of parasitaemia, but maintained a gradual increase during the period of low, or no parasitaemia, showed an inverse relationship between parasitaemia and anaemia [38, 39].

Also, the infected sheep exhibited leucopenia which was indicative of immunosuppression commonly encountered in African trypanosomosis [40]. Similar findings were observed in *T. evansi* infection in camels [41]. Lymphopenia was encountered among African giant rats infected with *T. brucei*[42]. This is probably associated with an increased demand on the system for lymphocytes, which is a common requirement in both immune and inflammatory responses in trypanosomosis [43]. Meanwhile, the significant neutropenia encountered among the infected sheep might be associated with splenic sequestration of leucocytes, which often suppressed the raising of neutrophil numbers [44].

The thrombocytopaenia encountered among the sheep is associated with coagulation defects due to low platelet numbers, which have been reported to be associated with disseminated intravascular coagulation in trypanosomosis [45]. This has been reported commonly in acute and chronic *T.brucei* infection of sheep [46] but it does not agree with the reports of Wada et al. [47], which perhaps is the only report of thrombocytosis in cattle trypanosomosis.

The infected sheep showed significant monocytosis. This has been reported as a common feature in trypanosomosis [48]. Monocytosis in trypanosomosis has been associated with a proliferation of tissue macrophages likely due to an increased demand on the system to remove dead red cells, antigen/antibody complexes and to participate in immune responses. Since macrophages are formed from blood monocytes, this increased need for macrophages may have been responsible for the consistent monocytosis encountered in this study. The result suggests that administered drugs improved blood components possibly by depletion of proliferating parasites, and all haematological parameters modulated to their pre-infection status with zero mortality.

CONCLUSION

In conclusion, our finding has provided evidence that the administered drug (berenil[®]) has the potentials for modulating the state of anaemia, immunosuppressive conditions induced by trypanosome infected sheep in a dose dependent manner.

ETHICAL APPROVAL

Experimental animals were used according to all local laws, guidelines, and policies at University of Maiduguri, Nigeria.

REFERENCES

1. Desquesnes M, Holzmuller P, Lai DH, Dargantes A, Lun ZR, Jittaplaong S. *Trypanosoma evansi* and Surra: a review and perspectives on origin, history, distribution, taxonomy, morphology, hosts, and pathogenic effects. *BioMedicine Research International*.2013; 22: p.194176.

2. Chandu AGS, Sengupta PP, Jacob SS, Suresh KP, Borthakur SK, Patra G, Roy P. Seroprevalence of *Trypanosoma evansi* in cattle and analysis of associated climatic risk factors in Mizoram. *Indian Journal of Parasitic diseases*. 2021; 45(1): 244–251.
3. Fesseha H, Eshetu E, Mathewos M, Tilante T. Study on Bovine Trypanosomiasis and Associated Risk Factors in Benatsemay District, Southern Ethiopia. *Environmental Health Insights*. 2022; 16: 1–10.
4. Kyari F, Mbaya AW, Biu AA, Adamu L, Dennis OO. Seroprevalence of *Trypanosoma evansi* in camels using CATT/T. evansi technique in Borno and Yobe states, Nigeria. *Parasite Epidemiology and Control*. 2021; 13. <https://doi.org/10.1016/j.parepi.2021.e00209>
5. Ngulde SI, Tijjani MB, Ihopo JM, Yauba AM. Antitrypanosomal potency of methanol extract of *Cassia arerehdelile* root bark in albino rats. *International Journal of Drug Research and Technology*. 2013; 3: 1-7.
6. Desquesnes M, Holzmuller P, Lai DH, Dargantes A, Lun ZR, Jittapalpong S. *Trypanosoma evansi* and Surra: a review and perspectives on origin, history, distribution, taxonomy, morphology, hosts, and pathogenic effects. *BioMedicine Research International*, (2013); 22: 194176.
7. Office International des Epizootic (OIE) (2004). Manual of diagnostic tests and vaccines for terrestrial animals. 2004; 5th Edition, chapter 2.3.11.
8. Office International des Epizootic (OIE). World Animal Health Organization: OIE-Listed diseases, infections and infestations in force in 2016. Paris, France. <http://www.oie.int/animal-health-in-the-world/oie-listed-diseases-2016>.
9. World Health Organisation (WHO) (1998). Control and surveillance of African trypanosomiasis. 1998; Geneva, Technical Report Series No.881.
10. Osman AS, Jennings FW, Holmes PH. The rapid development of drug resistance by *Trypanosoma evansi* in immunosuppressed mice. *Acta Tropical*, 1992; 50(3):249-257.
11. Desquesnes M, Gonzatti M, Sazmand A, Thévenon S, Bossard G, Boulangé A, Gimonneau G, Truc P, Herder S, Ravel S, Sereno D, Jamonneau V, Jittapalpong S, Jacquiet P, Solano P, Berthier D. A review on the diagnosis of animal trypanosomiasis. *Parasites and Vectors*. 2022; 15(64): 1-24.
12. Homeida AM, Elamin EA, Adam SEI, Mahmoud MM. Toxicity of diaminazene aceturate (Berenil) to camels. *Journal Comparative Pathology*. 1981; 91:355-360.
13. Tuntasuvan D, Trongwanichnam K, Sukruen A, Borisutsuvan S, Mohkaew K, Chompoochan T. Efficacy of diaminazene aceturate on the treatment of trypanosomiasis in pigs. *Journal of the Thai Veterinary Medicine Association*. 2003; 54(1-2), 49–55.
14. Dia ML, Desquesnes ML. Trypanosomoses: utilisation rationnelle des trypanocides. Fiche technique, Santé animale, CIRDES, BP454 Bobo-Dioulasso, Burkina Faso, 2004; 3:1–8.
15. Herbart WJ, Lumsden WHR. *Trypanosoma brucei*: A rapid matching method for estimating the host's parasitaemia. *Journal of Experimental Parasitology*. 1976; 40: 427-432.
16. Adeyeye AA, Ate IU, Lawal AI, Adamu S. Effects of experimental *Trypanosoma evansi* infection on pregnancy in Yankasa ewes. *Science direct: Theriogenology*. 2016; 85: 862–869.
17. Desquesnes M, Gonzatti M, Sazmand A, Thévenon S, Bossard G, Boulangé A, Gimonneau G, Truc P, Herder S, Ravel S, Sereno D, Jamonneau V, Jittapalpong S,

Comment [o5]: J

Comment [o6]: T

- Jacquet P, Solano P, Berthier D. A review on the diagnosis of animal trypanosomosis. *Parasites and Vectors*. 2022; 15(64): 1-24.
18. Herbart WJ, Lumsden WHR. *Trypanosoma brucei*: A rapid matching method for estimating the host's parasitaemia. *Journal of Experimental Parasitology*. 1976;40: 427-432.
 19. Coles EH. (1986). *Veterinary clinical pathology*. 2nd ed. W.B. Saunders Co., Pp 110-111.
 20. Brown BA. (1976). *Haematology principles and procedures*. 1986; 2nd. Ed. Lea and Febiger. Philadelphia. Pp 56- 81.
 21. Coles EH. (1986). *Veterinary clinical pathology*. 2nd ed. W.B. Saunders Co., Pp 110-111.
 22. Cheesbrough M. (2000). *District Laboratory Practice in Tropical Countries*. Law price edition Cambridge University Press. 2000;8:314-335.
 23. Graph Pad Instat. (2009). *Graph Pad Instat*® version 3: 10, bit for windows, Graph Pad.
 24. Audu PA, Esievo KAN, Mohammed G, Ajanusi JO. Studies on the infectivity of an isolate of *trypanosoma evansi* in yankasa sheep. *Veterinary Parasitology*. 1999;86:185-190.
 25. Shehu SA, Ibrahim NDG, Esievo KAN, Mohammed G. Role of erythrocyte surface sialic acid in inducing anaemia in Savannah Brown bucks experimentally infected with *Trypanosoma evansi*. *Veterinarski Arhiv*. 2010;76(6):521-530.
 26. Wada YA, Rekwot PI, Okubanjo OO, Mohammed B, Oniye SJ. Clinicopathological and Microscopic Features of *Trypanosoma brucei* and *Trypanosoma evansi* Induced Infections in Sheep II. *Nigerian Veterinary Journal*. 2020; 41(2): 144-160.
 27. Ndung'u K, Murilla GA, Thuita JK, Ngae GN, Auma JE, Gitonga PK. Differential virulence of *Trypanosoma brucei rhodesiense* isolates does not influence the outcome of treatment with antitrypanosomal drugs in the mouse model. *PLOS ONE*. 2020; 15(11): e022906.
 28. Mulengg GM, Gummow B. The detection of African trypanosomes in goats reared in tsetse infected villages of Eastern Zambia. *Tropical Animal Health and Production*. 2022; 54(6): 370.
 29. Mdachi RE, Ogolla KO, Auma JE, Wamwiri FN, Kurgat RK, Wanjala, KB, Mugunieri LG, Alusi PM, Chemuliti JK, Mukiria PW, Okoth SO. Variation of sensitivity of *Trypanosoma evansi* isolates from Isiolo and Marsabit counties of Kenya to locally available trypanocidal drugs. *PLOS ONE* | 2023; <https://doi.org/10.1371/journal.pone.0281180>.
 30. Odeniran PO, Macleod ET, Ademola IO, Welburn SC. Molecular identification of bloodmeal sources and trypanosomes in *Glossina spp.*, *Tabanus spp.* and *Stomoxys spp.* trapped on cattle farm settlements in southwest Nigeria. *Medical Veterinary Entomology*. 2019; 33(2): 269–81.
 31. Odeniran PO, Macleod ET, Ademola IO, Welburn SC. Molecular identification of bloodmeal sources and trypanosomes in *Glossina spp.*, *Tabanus spp.* and *Stomoxys spp.* trapped on cattle farm settlements in southwest Nigeria. *Medical Veterinary Entomology*. 2019; 33(2): 269–81.
 32. Wanjala KB, Mugunieri GL, Alusi PM, Kurgat RK, Mdachi RE, Chemuliti JK. Management of Camel Trypanosomiasis (Surra) among Pastoralists of Isiolo and Marsabit Counties, Kenya. *World Journal of Agricultural Research*. 2021; 9(1): 15–23.
 33. Elihu A. Effect of *Trypanosoma brucei* and *Trypanosoma evansi* on Ascorbic Acid and Semen Quality and its Relation with FREE Testosterone in Yankasa Rams. *Archives of Veterinary and Animal Sciences*. 2022; 4(1): 1-8.

34. Mbaya AW, Aliyu MM, Nwosu CO, Taiwo VO, Ibrahim UI. Effect of melarsamine hydrochloride (Cymelarsan[®]) and diaminazene aceturate (Berenil[®]) on the pathology of experimental *Trypanosoma brucei* infection in red fronted gazelles (*Gazellarufifrons*). *Veterinary Parasitology*.2009; 163(1-2): 140-143.
35. Mbaya AW, Aliyu MM., Nwosu CO, Taiwo VO, Ibrahim, UI. Effect of melarsamine hydrochloride (Cymelarsan[®]) and diaminazene aceturate (Berenil[®]) on the pathology of experimental *Trypanosoma brucei* infection in red fronted gazelles (*Gazellarufifrons*). *Veterinary Parasitology*.2009; 163(1-2): 140-143.
36. Mbaya AW, Aliyu MM, Nwosu CO, Taiwo VO, Ibrahim UI. Effect of melarsamine hydrochloride (cymelarsan) and diaminazene aceturate (Berenil) on the pathology of experimental *Trypanosomabrucei* in red fronted gazelle (*Gazellarufifrons*): short communication. *Veterinary parasitology*.2009; 163: 140-143.
37. Adeyeye AA, Ate IU, Lawal AI, Adamu S. Leukocyte Changes in Pregnant Yankasa Ewes Experimentally Infected with *Trypanosoma evansi*. *Nigerian Veterinary Journal*.2017; 38(2): 117-123.
38. Adeyeye AA, Ate IU, Lawal AI, Adamu S. Leukocyte Changes in Pregnant Yankasa Ewes Experimentally Infected with *Trypanosoma evansi*. *Nigerian Veterinary Journal*. 2017; 38(2): 117-123.
39. Mbaya AW, Aliyu MM. Nwosu CO, Taiwo VO, Ibrahim UI. Effect of melarsamine hydrochloride (cymelarsan) and diaminazene aceturate (Berenil) on the pathology of experimental *Trypanosomabrucei* in red fronted gazelle (*Gazellarufifrons*): short communication. *Veterinary parasitology*.2009; 163: 140-143.
40. Habeeb IF, Chechet GD, Kwaga JKP. Molecular identification and prevalence of trypanosomes in cattle distributed within the Jebba axis of the River Niger, Kwara state, Nigeria. *Parasites Vectors*.2021; 14: 560 <https://doi.org/10.1186/s13071-021-05054-0>.
41. Njiru ZK, Constatine CC, Ndung JM, Robertson I, Okaye S, Thonmpson RCA, Reid S. A. Detection of *Trypanosoma evansi* in camels using PCR and CATT/*T. evansi* tests in Kenya. *Veterinary Parasitology*. 2004;124: 187-199.
42. Mbaya AW, Kumshe HA, Geidam YA, Wiam AY. Effect of diaminazene aceturate (Berenil[®]) on the pathology of experimental *T. brucei* infected in African giant rats (*Criceetomysgambianus*). *Nigerian Veterinary Journal*.2011; 32(3): 192-198.
43. Igbokwe IO. Mechanisms of cellular injury in African trypanosomosis. *Veterinary bulletin*.1994; 64: 611-620.
44. Adeyeye AA, Ate IU, Lawal AI, Adamu S. Leukocyte Changes in Pregnant Yankasa Ewes Experimentally Infected with *Trypanosoma evansi*. *Nigerian Veterinary Journal*.2017; 38(2): 117-123.
45. Mbaya AW, Aliyu MM, Ibrahim UI. Clinico-pathology and mechanism of trypanosomosis in captive and free-living wild animals: A review. *Veterinary Research Communications*.2009; 33: 793-809.
46. Adeyeye AA, Ate IU, Lawal AI, Adamu S. Leukocyte Changes in Pregnant Yankasa Ewes Experimentally Infected with *Trypanosoma evansi*. *Nigerian Veterinary Journal*.2017; 38(2): 117-123.
47. Wada YA, Rekwot PI, Okubanjo OO, Mohammed B, Oniye SJ. Clinicopathological and Microscopic Features of *Trypanosoma brucei* and *Trypanosoma evansi* Induced Infections in Sheep II. *Nigerian Veterinary Journal*. 2020;41(2): 144-160.

Comment [o7]: P

Comment [o8]: P capital letter

Comment [o9]: B

48. Kasozi KI, MacLeod ET, Waiswa C, Mahero M, Ntulume I, Welburn SC. Systematic Review and Meta Analysis on Knowledge Attitude and Practices on African Animal Trypanocide Resistance. *Tropical Medicine and Infectious Disease*.2022; 7(9): 205.

UNDER PEER REVIEW