

## Use of selected chemical and biological insecticides to control lepidopteran pests of maize fields in central Côte d'Ivoire

### ABSTRACT

Control of lepidopteran larvae, the main pests in maize fields, has become necessary since the appearance and spread of *Spodoptera frugiperda*. The study was carried out at the Institut National Polytechnique Houphouët-Boigny in Yamoussoukro (Central Côte d'Ivoire). The study aimed to determine the effectiveness of three chemical - Viper 46 EC (Acetamiprid 16 g/l and Indoxacarb 30 g/l), K-Optimal 35 EC (Lambda-cyhalothrin 15 g/l and Acetamiprid 20 g/l) and Ampligo 150 ZC (Chlorantranilipol 100 g/l and Lambda-cyhalothrin 50 g/l)- and one biological insecticide Bio-Elit (Azadirachtin, Salanin, Nimbin and Melandriol) on lepidopteran larvae in maize fields using a randomized complete block design with five treatments and three repetitions. Data on insect identification, plant infestation, damage, and yield were collected. Insects' identification was based on morphology using identification keys. Plant damage was assessed by visually estimating the plant health status (unattacked and attacked plants). Grain dry weight was used to estimate field yield. The encountered maize field insects belonged to 10 orders: Heteroptera, Hymenoptera, Homoptera, Coleoptera, Diptera, Dictyoptera, Odonata, Orthoptera, Dermaptera, and Lepidoptera. Five lepidopteran pest larvae have been recorded. Three of them were classified as minor pests (*Eldanasaccharina*, *Ostrinia nubilalis*, and *Helicoverpa zea*), one as important (*Sesamia calamistis*), and one as a major pest (*Spodoptera frugiperda*). On untreated plots, more than 76% of plants were moderately to heavily attacked. However, on treated plots, plants showed isolated to moderate attacks. Insecticide spraying controlled pest populations, reduced damage, and increased yield. The yields obtained on untreated plots ( $2.26 \pm 0.21$  t/ha) were lower than those on treated plots ( $3.29 \pm 0.11$  to  $3.60 \pm 0.09$  t/ha). The yield increase rate ranged from 45.74 to 59.63%. The best control was recorded with Ampligo (59.63%) and Bio-Elit (50.83%) compared to Viper (49.41%) and K-Optimal (45.74%). Therefore, the alternating use of synthetic or biological insecticides, which are not very toxic for humans and the environment but are effective on insect pests, increases the effectiveness of the control and provides a positive response to the problem of pest resistance while protecting the environment.

**Keywords:** Lepidoptera, larvae, damage, insecticides, Maize, Côte d'Ivoire

UNDER PEER REVIEW

## 1. INTRODUCTION


Maize (*Zea mays* L.) is one of the world's most important crops, with over 150 million hectares cultivated and an annual production of around 800 million tons of grain (1). The crop has become a staple food in many parts of the world, with total production surpassing wheat or rice performances. In addition to direct human consumption, maize is also used for animal feed and producing ethanol, syrup, and corn starch (2). Maize is vital for most smallholders and is usually grown alongside horticultural crops and other cereals, which collectively provide food and income (3). There are over 3,500 products where maize is used. The increasing demand from the poultry feed sector, the largest consumer of maize, and increasing demand for specialty corn (sweet corn, baby corn, popcorn, and quality protein maize), enhance the scope of its production and farmers' income (4). However, maize crops suffer abiotic and biotic stresses. Abiotic stresses due to drought, salinity, and high and low temperatures (5; 6), together with biotic stresses caused by fungi, bacteria, and viruses (7; 8) and pests (9; 10) negatively affect maize growth, development, and eventually production (3). In addition, abiotic and biotic stresses are often present simultaneously and severely influence maize production (6).

Moreover, maize production is often subject to high losses caused by various insect pests (9; 3; 4; 11). More recently, an invasive insect, *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae), also known as the fall armyworm, became a major pest causing substantial yield losses on maize in the region (12; 13). This insect was detected in central and western Africa in early 2016 and quickly spread across all over within two years to more than forty African countries due to unscientific and uncontrolled trade. It is a highly polyphagous migratory lepidopteran pest species referring to the invasive behavior of larvae (12). Consequently, controlling pests is becoming a major concern for farmers. Multidisciplinary applied research and incentives for farmers to adopt innovative IPM strategies are essential to ensure the sustainability of pest management (14). Although insecticides have adverse consequences for the environment (15; 16), biodiversity and beneficial insects (17), human health (18; 19) leave food residues (20), and induce resistance phenomena in insects (21), chemical control has remained the most widely used due to the immediate effects observed after spraying. To overcome pest resistance, a wide range of insecticides is essential. The main objective of this study is to provide the best knowledge on lepidopteran pests of maize and to evaluate several insecticides to improve maize production.

## 2. MATERIAL AND METHODS

### 2.1. Experimental design

The trial was set up using a randomized complete block design containing 15 elementary plots with five treatments repeated thrice. The elementary plots covered 85.5 m<sup>2</sup> (19 m × 4.5 m) and were separated 2 m apart. Plots were spaced 0.75 m between rows and 0.5 m on the row, giving a density of 228 pots per elementary plot. The variety maize EV8728 SR was sowed in July 2021 at two seeds per pot. Subsequently, weeding was manually done when needed.



**Figure 1. Experimental design**

T0 : Untreated plots ; T1 : plots treated with Viper 46 EC ; T2 : plots treated with K-Optimal 35 EC ; T3 : plots treated with Ampligo 150 ZC ; T4 : plots treated with Bio-Elit

**Table 1. Insecticides used in this study**

Treatments	Trade name	Active substances concentration	Chemical Families
T0	Control	Untreated plots	
T1	Viper 46 EC	Acetamiprid 16 g/l Indoxacarb 30 g/l	Neonicotinoids Oxadiazins
T2	K-Optimal 35 EC	Lambdacyhalothrin 15 g/l Acetamiprid 20 g/l	Pyrethrinoids Neonicotinoids
T3	Ampligo 150 ZC	Chlorantanlipol 100 g/l Lambdacyhalothrin 50 g/l	Diamid Anthranilics Pyrethrinoids
T4	Bio-Elit	Azadirachtin, Salanin, Nimbin, Melandriol	Bio-insecticide

### 2.2. Data collection

Thirty (30) plants were observed on the central rows per elementary plot to avoid border effects. Two observations were made three days before and after insecticide spraying at the following plant development stages: emergence, growth (vegetative), flowering, and fruiting. At the ripening stage, an observation was made without insecticide spraying. Plant infestation, damage, and yield were collected. Insect identification was based on morphological characteristics, using a binocular microscope ( $G \times 50$ ). Observed characteristics were compared with the laboratory's collection and identification keys (22; 23; 24; 25; 26). In addition, the damage severity was assessed by visually evaluating the plant's health status (non-attacked and attacked plants). Six indices ranging from 0 to 5 (Table 2) were adapted from the scoring scale proposed by Sally-Sy(27).

The yield was assessed by weighing the dry grains on 30 plants per plot, using a 15 kg capacity electronic scale. Similarly, the yield increase rate was calculated using the formula adopted by Adja (28).

$$\text{Yield Increase Rate (\%)} = \frac{(\text{Yield of Treated Plot} - \text{Yield of Untreated plot}) \times 100}{\text{Yield of Untreated Plot}}$$

**Table 2. Damage severity index adapted from Sally-Sy (2013)**

<b>Indices</b>	<b>Severity damage</b>	<b>Observation</b>
<b>0</b>	No visible damage	No visible attack
<b>1</b>	From 1 to 9% of damage	Isolated attacks
<b>2</b>	From 10 to 24% of damage	Moderate attacks
<b>3</b>	From 25 to 49% of damage	Medium attacks
<b>4</b>	From 50 to 74% of damage	Heavy attacks
<b>5</b>	More than 75% of damage	Plants destroyed

### **2.3. Data Analysis**

Data were subjected to analysis of variance using STATISTICA 10.1. Where significant differences were found at a significance level of 0.05, Duncan's test was performed for multiple pairwise mean comparisons. The percentage of plants attacked was determined based on damage indices for each treatment.

### 3. RESULTS AND DISCUSSION

#### 3.1. Results

##### 3.1.1. Insect inventory

A total of 3,416 individuals representing 43 species, 26 families, and ten orders were collected in the field (Table 3). The most diversified orders were Heteroptera (8 species and four families), Hymenoptera (6 species and two families), Diptera (5 species and five families), Orthoptera (5 species and four families), and Lepidoptera (5 species and three families). These orders are followed by Coleoptera (4 species and three families), Homoptera (3 species and two families), Odonatoptera (3 species and one family), Dictyoptera (3 species and one family), Dermaptera (one species and one family) and Lepidoptera (2 species and one family). Dermaptera (one species and one family) (Table3) is the least diversified order in our test field.

Three trophic groups were identified: pests, predators, and pollinators. Pests were the most numerous and were found in Lepidoptera, Diptera, Coleoptera, Hymenoptera, Orthoptera, Homoptera, and Heteroptera. Predators belonged to Coleoptera, Hymenoptera, Heteroptera, Dermaptera, Dictyoptera, and Odonaptera. Pollinators were observed in Diptera, Hymenoptera, and Lepidoptera (Table 3).

**Table 3. Status of insects collected in the tested maize field**

Orders	Families	Species	Status	Relative Abundance (%)
<b>Lepidoptera</b>	Noctuidae	<i>Spodoptera frugiperda</i>	Pests	13.14
		<i>Sesamia calamistis</i>	Pests	1.64
		<i>Helicoverpa zea</i>	Pests	1.00
	Crambidae	<i>Ostrinia nubilalis</i>	Pests	0.94
	Pyralidae	<i>Eldana saccharina</i>	Pests	0.97
<b>Diptera</b>	Tachinidae	<i>Dexia rustica</i>	Pests	0.20
	Antomyiidae	<i>Delia platura</i>	Pests	0.18
	Diopsidae	<i>Diopsis thoracica</i>	Pests	0.23
	Muscidae	<i>Musca domestica</i>	Pollinators	3.22
	Tephritidae	<i>Dacus ciliatus</i>	Pests	0.35
<b>Coleoptera</b>	Coccinellidae	<i>Cheilomenes sulphurata</i>	Predators	2.08
		<i>Henosepilachna argus</i>	Pests	1.38
	Curculionidae	<i>Sphenophorus callosus</i>	Pests	0.20
	Tenebrionidae	<i>Lagriavillosa</i>	Pests	1.64
<b>Hymenoptera</b>	Formicidae	<i>Camponotus sp.</i>	Predators	30.94

		<i>Polyrhachis</i> sp.	Pests/Predators	1.93
		<i>Pachycondylasylvestri</i>	Pests/Predators	1.70
		<i>Crematogasterquadriformis</i>	Pests/Predators	3.89
		<i>Oecophyllalonginoda</i>	Pests/Predators	2.14
	Apidae	<i>Apis mellifera</i>	Pollinators	2.20
<b>Orthoptera</b>	Pyrgomorphidae	<i>Zonocerus variegatus</i>	Pests	0.41
		<i>Atractomorphaacutipennis</i>	Pests	0.85
	Gryllidae	<i>Gyllusbimaculatus</i>	Pests	0.32
	Tetrigidae	<i>Tetrixundulata</i>	Pests	0.85
	Acrididae	<i>Chorthippusalbomarginatus</i>	Pests	0.44
<b>Homoptera</b>	Aphididae	<i>Rhopalosiphummaidis</i>	Pests	1.17
		<i>Aphisgossipii</i>	Pests	2.05
	Cercopidae	<i>Deoisflavopicta</i>	Pests	1.17
<b>Heteroptera</b>	Pentatomidae	<i>Aspaviaarmigera</i>	Pests	8.34
		<i>Acrosternumheegeri</i>	Pests	0.50
		<i>Hediorcorisfasciatus</i>	Pests	1.08
		<i>Eysarcorisinconspicuus</i>	Pests	0.29
	Reduviidae	<i>Rhynocorishutsebauti</i>	Pests	0.50
	Pyrrhocoridae	<i>Dysdercuswoelkerii</i>	Pests	3.81
		<i>Dysdecussupertitiosus</i>	Pests	1.17
	Miridae	<i>Helopeltisantonii</i>	Pests	0.67
<b>Dermaptera</b>	Forficulidae	<i>Forficulasp.</i>	Predators	2.75
<b>Dictyoptera</b>	Mantidae	<i>Sphodromantissp.</i>	Predators	0.35
		<i>Tarachodesafzelii</i>	Predators	0.15
		<i>Mantisreligiosa</i>	Predators	0.76
<b>Odonoptera</b>	Libellulidae	<i>Sympetrumflaveolum</i>	Predators	0.97
		<i>Plathemislydia</i>	Predators	0.94
		<i>Libellulasp.</i>	Predators	0.50
<b>Total: 10</b>	<b>26</b>	<b>43</b>	<b>03</b>	<b>100</b>

### 3.1.2. Impact of insecticides on populations of Lepidopteran pests

#### Impact of insecticides on populations of *Spodopterafrugiperda*

*Spodopterafrugiperda* larvae were present on all plots at the time of emergence ( $4.33 \pm 0.57$  to  $5.33 \pm 0.57$  individuals on 30 plants per elementary plot). This population fluctuated in the untreated plots during all stages of plant development (Figure 2a). At emergence, after insecticide spraying, the larval population was significantly higher ( $F_{(4;14)} = 53.93$ ;  $p = 0.00001$ ) on control plots ( $8 \pm 1$  individuals) than on treated plots Viper ( $2 \pm 0$  individuals) and K-Optimal ( $3 \pm 1$  individuals), which were higher than those on other treated plots Ampligo ( $0.33 \pm 0.57$  individuals) and Bio-Elit ( $0.67 \pm 0.57$  individuals). From the growing to the ripening stage, *S. frugiperda* larvae on Control plots ( $8.33 \pm 0.57$  to  $14 \pm 1$  individuals) were

significantly higher than those on treated plots Viper, K-Optimal T2, T3Ampligo, and T4 Bio-Elit (less than  $3 \pm 1$  individuals). Analysis of variance showed significant differences ( $F_{(4;14)} = 43.25$  to  $257$ ;  $p = 0.00001$ ) between treatments at each plant development stage (Appendix 1).

#### **Impact of insecticides on populations of *Sesamiacalamistis***

During emergence, *Sesamiacalamistis* was present in the plots ( $0.33 \pm 0.57$  to  $1 \pm 1$  individual per plot). In control plots, *S. calamistis* populations increased from the growing (vegetative) to the flowering stage (Figure 2b). At emergence, after insecticide spraying, the *S. calamistis* population on control plots T0 ( $2 \pm 0$  individuals) was significantly higher ( $F_{(4;14)} = 34$ ;  $p = 0.000009$ ) than those on treated plots Viper, K-Optimal, Ampligo, and Bio-Elit (less than  $0.33 \pm 0.57$  individuals). From the growing to the flowering stage, *S. calamistis* populations on control plots T0 ( $2.33 \pm 0.57$  to  $3.33 \pm 0.57$  individuals) were significantly higher ( $F_{(4;14)} = 14.5$  to  $121$ ;  $p < 0.00036$ ) than those on treated plots T1, T2, T3, and T4 (less than  $0.33 \pm 0.57$  individuals). From fruiting to ripening, *S. calamistis* was absent.

#### **Impact of insecticides on the population of *Helicoverpazea***

*Helicoverpazea* was absent on all plots from the emergence to the growing stages. From flowering to ripening, the population of *H. zea* increased (Figure 2c). After insecticide spray, the population of *H. zea* on the control plot ( $0.66 \pm 0.57$  to  $3.33 \pm 1.15$  individuals per elementary plot) was significantly higher ( $F_{(4;14)} = 4$  to  $49$ ;  $p < 0.0343$ ) than those on the treated plots Viper, K-Optimal, Ampligo, and Bio-Elit (less than  $0.33 \pm 0.57$  individual).




#### **Impact of insecticides on the population of *Ostrinianubilalis***

*Ostrinianubilalis* was absent from all plots from the emergence to the growing or vegetative stage. From flowering to ripening, the *O. nubilalis* population increased (Figure 2d). After insecticide spraying, the *O. nubilalis* population on the control plot ( $0.66 \pm 0.57$  to  $2.66 \pm 0.57$  individuals per elementary plot) was significantly higher ( $F_{(4;14)} = 4$  to  $64$ ;  $p < 0.0343$ ) than on the treated plots Viper, K-Optimal, Ampligo, and Bio-Elit (less than  $0.33 \pm 0.57$  individuals).

#### **Impact of insecticides on the population of *Eldanasaccharina***

During the emergence, *Eldanasaccharina* was present on the plots ( $0.33 \pm 0.57$  to  $0.66 \pm 1.15$  individual per elementary plot). On control plots, the populations of *E. saccharina* increased from emergence to the growing stages and decreased at the flowering stage (Figure 2e). At emergence, after the insecticide spraying, the population of *E. saccharina* on control plots T0 ( $1.66 \pm 0.57$  to  $2.66 \pm 1.15$  individuals) was significantly higher ( $F_{(4;14)} = 12.2$  to  $49$ ;  $p <$

0.00073) than those on the treated plots Viper, K-Optimal, Ampligo, and Bio-Elit (less than  $0.33 \pm 0.57$  individual). *E. saccharina* was absent on all the plots, from the flowering to the ripening.



### Legend

- a) *Spodoptera frugiperda*,
- b) *Sesamia calamistis*,
- c) *Helicoverpa zea*,
- d) *Ostrinia nubilalis*,
- e) *Eldana saccharina*

Plant stage -3: three days before spraying;

Plant stage +3: three days after spraying

Ex: Emergence -3: three days before spraying during emergence, Emergence +3: three days after spraying during emergence.

## **Figure 2. Dynamic of lepidopteran larvae following insecticide treatments**

### **3.1.3. Impact of insecticides on insect plant damage**

At emergence, before insecticide spraying, 70 to 76.66% of all plots plants were healthy (index 0), while 13.33 to 16.66% of plants showed isolated attacks, 10 to 13.33% were moderately attacked, and 1.11% displayed medium attacks. After insecticide spraying, 66.66% of the plants were healthy on untreated plots, compared to 70 - 73.33% on treated plots. Moreover, the plants attacked on untreated plots showed isolated attacks (13.33%), moderate attacks (10%), and medium attacks (10%). On treated plots, 16.66 to 20% of plants showed isolated attacks and 10 to 13.33% moderate attacks.

During the growing phase, 60 to 63.33% of the plants on untreated plots were healthy, compared to those of treated plots (70 to 80%). On average, 13.33 to 20% of attacked plants on untreated plots presented isolated attacks, 13.33% moderate attacks, 3.33 to 6.66% showed medium attacks, and 3.33% were heavily attacked. On treated plots, attacked plants showed isolated attacks (13.33 to 16.66%) and moderated attacks (6.66 to 13.33%).

During flowering on untreated plots, 50 to 56.66% of plants were healthy, compared to those on treated plots (73.33 to 83.33%). Attacked plants on untreated plots presented isolated attacks (20%), moderate attacks (13.33 to 16.66%), medium attacks (6 to 10%), and heavy attacks (3.33%). On treated plots, 10 to 16.66% of attacked plants had isolated attacks, 6.66 to 13.33% were moderately attacked, and 3.33 to 16.66% displayed medium attacks.


During fruiting on untreated plots, 50% of plants were healthy compared to those on treated plots (76.66 to 86.66%). Attacked plants on untreated plots showed isolated attacks (16.66%), moderate attacks (16.66%), medium attacks (13.33%), and heavy attacks (3.33%). On treated plots, attacked plants had isolated attacks (6.66 to 13.33%) and moderate attacks (3.33 to 10%).

During ripening, 46.66% of plants were healthy on untreated plots compared to those on treated plots (80 to 90%). Attacked plants on untreated plots presented isolated attacks (20%), moderated attacks (16.66%), medium attacks (13.33%), and heavy attacks (3.33%). On treated plots, attacked plants showed isolated attacks (6.66 to 10%) and moderate attacks (3.33 to 10%).

### **3.1.4. Impact of the treatments on the maize yield**

There were significant differences ( $p < 0.05$ ) between the treatments. The yield on untreated plot T0 ( $2.26 \pm 0.21$  t/ha) was significantly lower than on treated plot T2 ( $3.29 \pm 0.11$  t/ha), which remained lower than treated plot T3 ( $3.61 \pm 0.22$  t/ha). However, the yield on treated

plots T1 ( $3.37 \pm 0.06$ ) and T4 ( $3.41 \pm 0.09$ ) was close and did not differ from those on T2 and T3 (Figure 3). The difference in yield between treated plots and untreated plots ranged from 1,034 to 1,348 t/ha. Then, the yield increase rate between the treated and untreated plots ranged from 45.74 to 59.63%. This rate is more important on T3 (Ampligo) than those on the other treated plots T1 (49.41%), T2 (45,74%), and T4 (50.83%) (Figure 3).



**Figure 3. Effect of insecticide treatments on the maize field yield**

## Discussion

The insects encountered on the studied maize field belong to 10 orders (Heteroptera, Hymenoptera, Homoptera, Coleoptera, Diptera, Dictyoptera, Odonata, Orthoptera, Dermaptera, and Lepidoptera). Five species of Lepidopteran larvae were attacking the maize field. Those Lepidopteran larvae were reported by several authors in Côte d'Ivoire (29; 30) and in Africa (7; 10). Three of the encountered Lepidoptera larvae were classified as minor pests (*Eldanasaccharina*, *Ostrinianubilalis*, and *Helicoverpazea*), while the two others, *Sesamiacalamistis*, and *Spodopterafrugiperda* were respectively important and major pests (11). *S. frugiperda* was the most abundant among the Lepidopteran larvae, which caused critical damage. This insect is a recently encountered pest in Africa (31; 12). The populations of Lepidopteran larvae fluctuated during the plant development stages on untreated plots. There is a synchronization between the plant's development stage and its pest, meaning that the pest outbreak occurs when the plant's stage is favorable to the pest's development (22).

This study revealed reductions in Lepidoptera larvae populations due to the tested insecticides. Lepidopteran larvae populations were significantly lower on treated plots than on untreated ones. The efficiency of the insecticides appears to be linked to their mode of action and doses. The most efficient insecticide is Ampligo. Its efficiency is due to its two active ingredients, chlorantraniliprole (anthranilicdiamides) and lambda-cyhalothrin (pyrethrinoids). Chlorantraniliprole is a new insecticide belonging to the anthranilicdiamide chemical class for use against Lepidoptera, Coleoptera, and certain dipteran pests in cash-crop agriculture on perennial and annual crops (32). Ingestion is the most effective route of entry and generally requires a lower dose to elicit a response. Chlorantraniliprole offers excellent plant protection, as affected insects cease feeding almost immediately after contact with the insecticide. Insects exposed to chlorantraniliprole show general lethargy and muscular paralysis, followed ultimately by death (33). Based on feeding cessation and reduction in feeding damage, chlorantraniliprole is among the fastest-acting insecticides available for lepidopteran pest control, comparable in speed of action with methomyl, lambda-cyhalothrin and esfenvalerate, and faster than emamectin benzoate, indoxacarb, methoxyfenozide and metaflumizone (32). Lambda-cyhalothrin (pyrethrinoids) is a systemic insecticide that also acts by contact with various larval pests (34). Chlorantraniliprole and lambda-cyhalothrin double-loaded nano-microcapsules for synergistic pest control have been reported by Feng et al. (35).

K-Optimal 35 EC and Viper 46 EC had a similar effect on lepidopteran larvae and were moderately effective on lepidopteran larvae. K-Optimal contains lambda-cyhalothrin (pyrethrinoids) and acetamiprid (neonicotinoids). Acetamiprid is a systemic chemical that acts on caterpillars via ingestion and contact (36). Neonicotinoids had an additive effect when mixed with the tested pyrethrinoids (37). Viper contains acetamiprid (neonicotinoids) and indoxacarb (oxadiazines). Indoxacarb is highly active on Lepidoptera larvae. Affected insects stop feeding, become paralyzed, and die (36). This combined mixture consisting of indoxacarb with outstanding translocation properties, and acetamiprid with excellent systemic properties, has a superb, long-lasting efficacy (36).

Bio-Elit is a bio-insecticide containing azadirachtin, salanine, nimbin, and melandriol. This insecticide has shown high control of larval Lepidoptera, close to that of Ampligo. Azadirachtin is used to control insects (Lepidoptera, Diptera, Coleoptera, Hymenoptera, Heteroptera, Homoptera, and Hemiptera), mites, and other arthropods (38; 39). In target species, azadirachtin has anti-feeding, sterilizing, and development-regulating effects (40; 41), induces cytotoxicity, tissue apoptosis, antimitotic effects, and growth abnormalities (inhibited,

abnormal, or delayed molting) (38). Azadirachtin also alters various reproductive processes, such as fecundity, fertility, and oviposition. *Azadirachtaindica* (Neem) is a source of several bioactive triterpenoids. However, only azadirachtins have been commercially exploited (42). Salanin and nimbin are the other major active potential bioactive compounds that can be used for insecticide development (43). Salanin and nimbin have been reported from neem seeds, kernels, and neem oil (44). Meliantiol and salanin are bioinsecticidal compounds that have been extracted from *Azadirachtaindica*. The efficacy of neem on insect pests of crops has been reported by several authors (45; 46; 38). Due to their relative selectivity, neem products can be recommended for many integrated pest management programs (41).

Bio-Elit, with natural neem substances (Azadirachtin, Salanin, Nimbin, Melandriol), was very effective against *S. frugiperda* larvae and the other lepidopteran larvae but showed a phytotoxic effect on young plants (at emergence). Before insecticide sprayings, infestations were medium on all the plots. Lepidopteran larvae populations were higher on untreated plots than on treated plots. The damage was caused by the larvae, which fed on leaves, stems, flowers, and fruits and reduced fruit quality and, subsequently, the yields (11). The damage was null to low on the plots treated with Ampligo and Bio-Elit, whereas the damage was low on the plots treated with Viper and K-Optimal. However, untreated plots showed very high damage. Feeding and stem tunneling by borer larvae on plants results in crop losses as a consequence of the destruction of the apical meristem, early leaf senescence, interference with translocation of metabolites and nutrients that result in malformation of the grain, stem breakage, plant stunting, lodging, and direct damage to ears (10). Infestations by stem borers increase the incidence and severity of stalk rots (47). Insecticide sprayings reduced insect pest populations and, consequently, the severity of their damage. In this study, grain yields obtained on untreated plot ( $2.26 \pm 0.21$  t/ha) was lower than those on treated plots. The yield on the Ampligo-treated plot ( $3.6 \pm 0.22$  t/ha) was higher than on the K-Optimal-treated plot ( $3.29 \pm 0.11$  t/ha). However, the yield on the Viper-treated plot ( $3.37 \pm 0.06$  t/ha) and Bio-Elit plot ( $3.41 \pm 0.09$  t/ha) did not significantly differ from those of Ampligo and K-Optimal. In Côte d'Ivoire, the yield of dry grains on untreated field varied between  $2.4 \pm 0.42$  t/ha to  $2.61 \pm 0.71$  t/ha (11) or between 1.64 to 2.34 t/ha (29). Maize yields were 2.46 t/ha in Nepal (8) and 1.84 to 1.88 t/ha in India (48). In addition, the computed yield increase rate was high on the plot treated with Ampligo (59.63%) and average on the plots treated with Bio-Elit (50.85%), Viper (49.41%), and K-Optimal (45.74%). In Kenya, stalk borer-induced crop yield losses vary from 10% to 88%, depending on the infestation intensity (49). On an economic

scale, 14% of Kenya's annual maize yields are estimated to be lost due to stalk borer damage, representing 0.4 million tonnes or \$25-60 million, enough to feed 3.5 million people yearly, with a 125 kg per capita maize consumption. (49). As with pests, predators and pollinators have also been observed in the field. The effectiveness of native predators on several species of borers has been the subject of several studies over the past decade (49; 50). In addition, agricultural practices such as organic and mineral fertilization, improved fallow cover crops, the use of repellent or insecticidal plants, intercropping, training farmers in pest recognition and damage estimation, and alert threshold treatments could improve pest control (14).

#### 4. CONCLUSION

This study was conducted to determine the effectiveness of three chemical and one biological insecticide on lepidopteran larvae in maize field. The insects recorded on maize field in this study belong to 10 orders: Heteroptera, Hymenoptera, Homoptera, Coleoptera, Diptera, Dictyoptera, Odonata, Orthoptera, Dermaptera, and Lepidoptera. Five lepidopteran larvae were pests. Three of them were minor pests (*Eldanasaccharina*, *Ostrinianubilalis*, and *Helicoverpazea*), one was classified as an important pest (*Sesamiacalamistis*), and the other one as a major pest (*Spodopterafrugiperda*). These insects caused considerable damage to the plants and crops. On untreated plots, more than 76% of plants were attacked with moderate to heavy attacks. However, on treated plots, attacked plants presented isolated to moderate attacks. Consequently, the spraying of insecticides controlled pest populations, reduced damage, and increased yield. Treated plots with insecticide presented higher yields than untreated plots. The yields obtained on untreated plots ( $2.26 \pm 0.21$  t/ha) were lower than those on treated plots ( $3.29 \pm 0.11$  to  $3.60 \pm 0.09$  t/ha). The yield increase rate ranged from 45.74 to 59.63%. All of these insecticides effectively controlled pests. Nevertheless, the best control of insect pests of maize in Yamoussoukro was obtained on plots treated with Ampligo and Bio-Elit compared with Viper and K-Optimal. So, the alternating use of synthetic or biological insecticides, which are not very toxic for humans and the environment but are effective on insect pests, increases the effectiveness of the control and provides a positive response to the problem of pest resistance while protecting the environment.

#### REFERENCES CITED

- 1- FAO (2022). Country summary reports. Global Information and Early Warning System (GIEWS).
- 2- Foley J. (2019). It's Time to Rethink America's Corn System". *Scientific American*. Retrieved February, 18 p.
- 3- Pratt F. C., Constantine K. L. & Murphy S. T. (2017). Economic impacts of invasive alien species on African smallholder livelihoods. *Global Food Security*, 14: 31-37.
- 4- Kumar P., Singh R., Suby S. B. · Kaur J., Sekhar J. C. &Soujanya P. L. (2018).An overview of crop loss assessment in maize. *Maize Journal*. 7(2): 56-63. DOI: 10.13140/RG.2.2.11554.17606. <https://www.researchgate.net/publication/335320605>.
- 5- Ahuja I., De Vos R. C. &Bones A. M. (2010). Plant molecular stress responses face climate change. *Trends Plant Sci*. 15, 664-674.
- 6- Gong F., Yang L., Tai F., Hu X. & Wang W. (2014). “Omics” of Maize Stress Response for Sustainable Food Production: Opportunities and Challenges. *OMICS. A Journal of Integrative Biology*, 18(12): 714–732. doi: [10.1089/omi.2014.0125](https://doi.org/10.1089/omi.2014.0125)
- 7- CIMMYT (2004). Maize Diseases: A Guide for Field Identification. 4th edition The CIMMYT Maize Program. Mexico. 124 p.
- 8- Subedi S. (2015). A review on important maize diseases and their management in Nepal. *Journal of Maize Research and Development* (2015) 1(1): 28-52. ISSN: 2467-9283 (Print)/ 2467-9291 (Online). DOI: 10.5281/zenodo.34292.
- 9- Groote H. D. (2002). Maize yield losses from stem borers in Kenya. *International Journal of Tropical Insect Science* , 22(2) : 89-96. DOI: <https://doi.org/10.1017/S1742758400015162>.
- 10- Kfir R., Overholt W. A., Khan Z. R. &Polaszek A. (2002). Biology and management of economically important Lepidopteran cereal stemborers in africa.*Annu. Rev. Entomol*. 47: 701-31.
- 11- Adja A. N., Bley-Atse A. B., Gnago J. A., Tienebo E.-O. &Danho M. (2023). Damage Caused by Lepidopteran Pests on Maize Fields in Yamoussoukro, Central Côte d'Ivoire. *International Journal of Plant & Soil Science*, 35(18), 815–823. <https://doi.org/10.9734/ijpss/2023/v35i183348>.
- 12- Goergen G., Kumar P.L., Sankung S.B., Togola A. &Tamò M. (2016). First report of outbreaks of the fall armyworm *Spodopterafrugiperda* (Smith JE) (Lepidoptera, Noctuidae), a

new alien invasive pest in West and Central Africa. *Plos One*. 2016; 11: e0165632. DOI: 10.1371/journal.pone.0165632.

13- Kumela T., Simiyu J., Sisay B., Likhayo P., Mendesil E., Gohole L. & Tefera T. (2018). Farmers' knowledge, perceptions, and management practices of the new invasive pest, fall armyworm (*Spodopterafrugiperda*) in Ethiopia and Kenya. *Int, J. Pest Manage*; 2018. Available: <https://doi.org/10.1080/09670874.2017.1423129>.

14- Vasileiadisa V.P., Sattin M., Van Der W. R. Y. (2011). Crop protection in European maize-based cropping systems: Current practices and recommendations for innovative Integrated Pest Management. *Agricultural Systems*, 104 (7): 533 – 540. ISSN 0308-521X.

15- Adam S., Edoth P., Totin H., Koumolou L., Amoussou E., Aklikokou K. & Boko M., (2011). Pesticides et métaux lourds dans l'eau de boisson, les sols et les sédiments de la ceinture cotonnière de Gogounou, Kandi et Banikoara (Bénin). *International Journal of Biological and Chemical Sciences* 4. <https://doi.org/10.4314/ijbcs.v4i4.63054>.

16- Konan Y. M., Benjamin Y. K., Albert T. & Bernard S. M. (2014). Assessment of Organic Pollution in Tropical Lagoon Bay Like Lake (Tiagba Bay Lagoon, Ébrié Lagoon, Côte D'Ivoire). *European Journal of Scientific Research*, 122 (3): 299-311. ISSN 1450-216X / 1450-202X. <http://www.europeanjournalofscientificresearch.com>

17- Chauzat M.-P., Carpentier P., Martel A.-C., Bougeard S., Cougoule N., Porta P., Lachaize J., Madec F., Aubert M., Faucon J.-P., 2009. Influence of pesticide residues on honey bee (Hymenoptera: Apidae) colony health in France. *Environmental Entomology*, 38: 514-523.

18- London L., Beseler C., Bouchard M. F., Bellinger D. C., Colosio C., Grandjean P., Harari R., Kootbodien T., Kromhout H., Little F., (2012). Neurobehavioral and neurodevelopmental effects of pesticide exposures. *Neurotoxicology*. 33: 887-896.

19- Sabarwal A., Kumar K. & Singh R.P., 2018. Hazardous effects of chemical pesticides on human health—Cancer and other associated disorders. *Environmental Toxicology and Pharmacology*. 63: 103–114. <https://doi.org/10.1016/j.etap.2018.08.018>

20- Kim H. S. & Hur S. J. (2018). Degradation of various insecticides in cooked eggs during in vitro human digestion. *Environmental Pollution*, 243: 437-443. <https://doi.org/10.1016/j.envpol.2018.09.021>.

- 21- Bass C. & Jones C. M. (2018). Editorial overview: Pests and resistance: Resistance to pesticides in arthropod crop pests and disease vectors: mechanisms, models and tools. *Current Opinion in Insect Science*. 27. <https://doi.org/10.1016/j.cois.2018.04.009>.
- 22- Bosque-Perez NA. (1995). Major insects pests of maize in Africa: biology and control. ITA Research Guide 30. Second edftion. Training Program, International Institute of Tropical Agriculture (IITA). Ibadan, Nigeria. 1995: 30.
- 23- Bordat D. &Arvanitakis L. (2004). Arthropods of vegetablecrops in West and Central Africa, Mayotte and Réunion. Cirad: Montpellier, France, 291 p.
- 24- Jean C. &Boisclair J. (2009). Pests and beneficial insects of maize: know them better. IRDA Ed. Quebec. Journal of Maize Research and development. 1(1): 28-52. ISSN2467-9305. (Online)/2467-9291 Print. DOI: 10.5281/zenodo.34292.
- 25- James B., Atcha-Ahowê C., Godonou I., Baimey H., Goergen G., Sikirou R. &Toko M. (2010). Integrated pest management in vegetable production: A guide for extension agents in West Africa. CTA IITA, Ibadan (Nigeria), 120 p.
- 26- Ayana W. (2019). Review on major pests of maize, their biology, ecology, damage and control methods. MSc in Agronomy. *Jimma, Ethiopia*. 2019: 32.
- 27- Sall-Sy D. (2013). Update on the monitoring of cabbage pests in the Niayes. ISRA/ CDH/ UCAD/ AUMN, Senegal. 2013: 25 p.
- 28- Adja N.A., Danho M., Alabi T.A.F., Gnago A.J., Zimmer J.Y., Francis F., Kouassi K.P., Baudoin J-P. &Zoro Bi I.A. (2014). Entomofauna associated with the cultivation of African oilseed cucurbits (*Lagenariasiceraria* Molina (Standl. 1930) and *Citrulluslanatus* thumb (Matsum&Nakai 1916)) and the impact of pests on production. *Annals of the Entomological Society of France (N.S.): International Journal of Entomology*. 2014 ; 50(3-4) : 301-310.
- 29- Nandjui J., Adja N.A., Kouadio K.T., N'gandi M-P.A. &Idrissou L. (2018). Impact of Soil Fertility Management Practices on Insect Pests and Diseases of Maize in Southwest Cote d'Ivoire. *Journal of Applied Biosciences*. 2018; 127: 12809-12820. ISSN 1997-5902.
- 30- Moyal P. & Tran M. (2023). Morphological study of the preimaginal stages of maize borer lepidoptera in the Savannah zone of Côte d'Ivoire. *Annals of the Entomological Society of France (N.S). International Journal of Entomology*. 2023; 25(4): 461-472.
- 31- IITA (2016). First report of outbreaks of the "Fall Armyworm" on the African continent. IITA Bulletin, No. 2330 ; Available : <http://bulletin.iita.org/index.php/2016/06/18/first-report-of-outbreaks-of-the-fallarmyworm-on-the-african-continent/>

- 32- Hannig G.T., Ziegler M. & Macron P.G. (2009). Feeding cessation effects of chlorantraniliprole, a new anthranilicdiamide insecticide, in comparison with several insecticides in distinct chemical classes and mode-of-action groups. *Pest Manag Sci.* 65(9): 969-974. Doi: 10.1002/ps.1781.
- 33- Bentley K. S., Fletcher J. L. & Woodward M. D. (2010). Chapter 102 - Chlorantraniliprole: An Insecticide of the AnthranilicDiamide Class. Hayes' Handbook of Pesticide Toxicology (Third Edition), Academic Press.: 2231-2242. <https://doi.org/10.1016/B978-0-12-374367-1.00102-6>.
- 34- Lewis K. A., Tzilivakis J., Warner D. & Green A. (2016). An international database for pesticide risk assessments and management. *Human and Ecological Risk Assessment: An International Journal*, 22 (4), 1050-1064. DOI: [10.1080/10807039.2015.1133242](https://doi.org/10.1080/10807039.2015.1133242). <https://doi.org/10.1080/10807039.2015.1133242>
- 35- Feng B, Zhi H, Chen H, Cui B, Zhao X, Sun C, Wang Y, Cui H, Zeng Z. (2021). Development of Chlorantraniliprole and Lambda Cyhalothrin Double-Loaded Nano-Microcapsules for Synergistical Pest Control. *Nanomaterials (Basel)*. 15; 11(10): 2730. doi: 10.3390/nano11102730. PMID: 34685168.
- 36- EFSA (2013). Scientific Opinion on the developmental neurotoxicity potential of acetamiprid and imidacloprid. EFSA Panel on Plant Protection Products and their Residues (PPR)2, 3European Food Safety Authority (EFSA), Parma, Italy. *EFSA Journal*. 2013: 11(12): 3471.
- 37- Kandil M.A., Fouad E.A. & Mokbel E-S M. S. (2022). Efficiency of certain Neonicotinoid Mixtures against the Cowpea Aphid, *Aphis craccivora* (Koch). *Egypt. Acad. J. Biolog. Sci.* 14(2):91-99. ISSN: 2092-0791. <http://ejbsf.journals.ekb.eg/>
- 38- Mordue A. J. L. & Nisbet A. J. (2000). Azadirachtin from the neem tree *Azadirachta indica*: its action against insects. *An Soc Entomol Bras*, 29. 615–632 (2000).
- 39- Benelli G., Canale A., Toniolo C., Toniolo C., Higuchi A., Murugan K., Pavela R. & Nicoletti M. (2017). Neem (*Azadirachta indica*): toward the ideal insecticide? *Nat Prod Res.* 017; 31(4): 369-86. DOI: [10.1080/14786419.2016.1214834](https://doi.org/10.1080/14786419.2016.1214834)
- 40- Chaudhary S., Kanwar R. K., Sehgal A., Cahill D. M., Barrow C., Sehgal R. & Kanwar J.R. (2017). Progress on *Azadirachta indica* based biopesticides in replacing synthetic toxic pesticides. *Front Plant Sci.* 2017 ;8 : 610. DOI: [10.3389/fpls.2017.00610](https://doi.org/10.3389/fpls.2017.00610)

- 41- Morgan E. D. (2009). Azadirachtin, a Scientific Gold Mine. *Bioorg. Med. Chem.* 17: 4096-4105. doi: 10.1016/j.bmc.2008.11.081. <https://doi.org/10.1016/j.bmc.2008.11.081>.
- 42- Mordue A. (2004). Present concepts of the mode of action of azadirachtin from neem, in *Neem: Today and in the New Millennium*, ed Opende K. W. (Dordrecht: Kluwer Academic Publishers, 229–242).
- 43- Sidhu O.P., Kumar V. & Behl H. M. (2004). Variability in triterpenoids (nimbin and salanin) composition of neem among different provenances of India. *Industrial Crops and Products*, 19 (1) : 69-75. <https://doi.org/10.1016/j.indcrop.2003.07.002>.
- 44- Govindachari T. R., Sandhya G. & Ganesh Raj S.P. (2010). Structure of Azadirachtin K, a New Tetranortriterpenoid from *Azadirachta indica*. *Chem Inform* 23 (37) : 275-275. DOI:[10.1002/chin.199237275](https://doi.org/10.1002/chin.199237275).
- 45- Gnago A. J., Danho M., Agneroh A. T., Fofana I. K. & Kohou G. A., 2010. Efficacy of neem (*Azadirachta indica*) and papaya (*Carica papaya*) extracts in controlling insect pests of okra (*Abelmoschus esculentus*) and cabbage (*Brassica oleracea*) in Côte d'Ivoire. *International Journal of Biological and Chemical Sciences*, 4(4): 953-966.
- 46- Riba M., Marti J. & Sans A. (2003). Influence of Azadirachtin on development and reproduction of *Nezaraviridula* L. (Het., Pentatomidae). *J Appl Entomol* 127, 37–41 (2003).
- 47- Bosque-Perez N.A. & Mareck J.H. (1991). Effect of the stem borer *Eldanasaccharina* (Lepidoptera: Pyralidae) on the yield of maize. *Bull. Entomol. Res.* 81 : 243-247.
- 48- Balla A., Bhaskar M., Bagade P. & Rawal N. (2019). Yield losses in maize (*Zea mays*) due to fall armyworm infestation and potential IoT-based interventions for its control. *Journal of Entomology and Zoology Studies*. JEZS. 2019; 7(5): 920-927. E-ISSN: 2320-7078 P-ISSN: 2349-6800.
- 49- Odendo M., Ouma J., Wachira S. & Wanyama J. (2003). Economic assessment of maize yield loss due to stem borer in major maize agro-ecological zones of Kenya. African crop science conference proceedings. *African Crop Science Society*, 6: 683-687. ISSN 1023-070X \$ 4.00.
- 50- Scovgard H & Päts P. 1996. Effects of intercropping on maize stem borers and their natural enemies. *Bull. Entomol. Res.* 86: 599-607.

51- Bonhof M.J. (2000). The impact of predators on maize stem borers in coastal Kenya. PhD thesis. Wageningen Univ., Neth. 181 p.

## Appendix1

### List 1: Data statistics

	Emergen ce-3	Emerge nce +3	Growt h -3	Growt h +3	Flowri ng-3	Flowerin g+3	Fructificati on-3	Fructificatio n+3	Maturat ion
<i>Spodopterafrugiperda</i>									
F	0.5	53.93	160.72	79.81	43.25	194.56	190	112.05	257
p	0.736	0.00000	0.0000	0.0000	0.00003	0.000001	0.000001	0.000001	0.000001
		1	01	1					
<i>Sesamiaclamistis</i>									
F	0.2692	34	14.5	100	48	121			
p	0.8911	0.00000	0.0003	0.0000	0.00000	0.000001			
		9	6	1	2				
<i>Helicoverpaarmigera</i>									
F					4	25	49	19.37	25
P					0.0343	0.00003	0.000002	0.0001	0.00003
<i>Ostrinianubilalis</i>									
F				4	16	25	11.75	49	64
P				0.0343	0.00024	0.00003	0.00085	0.000001	0.000001
<i>Eldanasaccharina</i>									
F	0.1875	25	12.2	49	16				
P	0.939	0.00003	0.0007	0.0000	0.00024				
		4	3	01					